

University of Groningen

Magnitude of neonatal near miss in public hospitals in Eastern Ethiopia

Tola, Melese Adugna; Semahegn, Agumasie; Tiruye, Getahun; Tura, Abera Kenay

Published in:
SAGE Open Medicine

DOI:
[10.1177/20503121221108926](https://doi.org/10.1177/20503121221108926)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Tola, M. A., Semahegn, A., Tiruye, G., & Tura, A. K. (2022). Magnitude of neonatal near miss in public hospitals in Eastern Ethiopia: A cross-sectional study. *SAGE Open Medicine*, 10. <https://doi.org/10.1177/20503121221108926>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Magnitude of neonatal near miss in public hospitals in Eastern Ethiopia: A cross-sectional study

SAGE Open Medicine
Volume 10: 1–8
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/20503121221108926
journals.sagepub.com/home/smo



Melese Adugna Tola^{1,2} , Agumasie Semahegn^{2,3},
Getahun Tiruye²  and Abera Kenay Tura^{2,4}

Abstract

Objectives: Although neonatal near miss is an emerging concept and a tool for improving neonatal care, its magnitude and associated factors are less researched in Ethiopia. Thus, this study was aimed to uncover the magnitude of neonatal near miss and its associated factors in public hospitals in Eastern Ethiopia.

Methods: A facility-based cross-sectional study was employed on a randomly selected 405 mother–neonate pairs. An interview using a structured questionnaire accompanied by review of medical records was used to collect data from the mothers and records of the neonates. Neonatal near miss was defined as having any of the pragmatic (gestational age < 33 weeks, birth weight < 1750 g, and fifth minutes Apgar score < 7) or management criteria. Crude and adjusted logistic regression analysis was done to identify associated factors and presented with adjusted odds ratio with 95% confidence interval.

Results: Of 401 mother–neonate pairs included in the study, 126 (31.4%, 95% confidence interval = [26.9, 36.2]) neonates had at least one neonatal near miss event at discharge. Neonatal near miss was more likely among neonates from referred women (adjusted odds ratio = 2.24, 95% confidence interval = [1.25, 4.03]), no antenatal care (adjusted odds ratio = 2.08, 95% confidence interval = [1.10, 3.93]), antepartum hemorrhage (adjusted odds ratio = 4.29, 95% confidence interval = [2.16, 8.53]), premature rupture of membrane (adjusted odds ratio = 4.07, 95% confidence interval = [2.05, 8.07]), obstructed labor (adjusted odds ratio = 2.61, 95% confidence interval = [1.23, 5.52]), non-vertex presentation (adjusted odds ratio = 3.03, 95% confidence interval = [1.54, 5.95]), and primiparous (adjusted odd ratio = 2.67, 95% confidence interval = [1.49, 4.77]).

Conclusions: In this study, we found that neonatal near miss is higher than previous findings in Ethiopia. Improving neonatal near miss requires promoting antenatal care, maternal referral system, and early identification and management of obstetric complications.

Keywords

Eastern Ethiopia, public hospitals, neonatal near miss

Date received: 28 January 2022; accepted: 6 June 2022

Introduction

Despite all the efforts, the decline in neonatal mortality rate (NMR) is slow worldwide. In 2018, 2.5 of 5.3 million under-five deaths were among the neonates, with sub-Saharan Africa accounting for the highest magnitude (28 per 1000 live births).¹ As such, neonatal mortality remained a major challenge in several low- and middle-income countries.^{2,3} Although neonatal mortality is an indicator of neonatal health, it shows only the iceberg tip of neonatal ill-health, with majority of neonates surviving the complications going unnoticed.² Therefore, understanding the true magnitude of neonatal ill-health requires studying neonates who survived

¹Department of Midwifery, College of Health Sciences, Mettu University, Mettu, Ethiopia

²School of Nursing and Midwifery, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia

³Department of Population, Family and Reproductive Health, School of Public Health, University of Ghana, Accra, Ghana

⁴Department of Obstetrics and Gynecology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

Corresponding author:

Melese Adugna Tola, Department of Midwifery, College of Health Sciences, Mettu University, Mettu, Ethiopia.
Email: melluadugna@gmail.com



from severe complications (neonatal near miss (NNM)) in addition to neonatal deaths.⁴

NNM refers to a neonate that almost died but survived from severe complications that occurred within the first 28 days of life.^{5,6} Analogous to maternal near miss, NNM is an emerging approach and is becoming a key indicator for assessing quality of neonatal care to reduce preventable neonatal morbidity and mortality.^{5,7} Study of NNM has an invaluable contribution to reduce neonatal mortality.^{2,5,6} Although there is no commonly agreed identification criteria for NNM, the Latin American Center for Perinatology (CLAP) proposed a standard definition and identification criteria for NNM based on the World Health Organization multi-country study.^{6,7} As such, NNM identification criteria consist of two set of criteria: pragmatic (gestational age < 33 weeks, birth weight < 1750 g, and fifth minutes Apgar score < 7) and management based (use of continuous positive air pressure (CPAP), use of vasoactive drugs, bag and mask ventilation, use of phototherapy, use of anticonvulsant drugs, respiratory distress, any intubation, etc. (listed in the subsection “measurement and variables”).^{2,6,7}

Despite the wider acceptance of NNM as a concept, there is a paucity of information and the need for more studies in different settings was previously indicated.^{2,5,6} Studying NNM in Ethiopia, where reduction in neonatal mortality is still slow,³ would generate data on important factors for neonatal survival.⁸ However, the magnitude of NNM remained uncovered.^{9–13} In addition, methodological limitations, such as non-random sampling^{14,15} or use of secondary data,¹⁶ may underestimate or overestimate NNM cases and made it difficult to generalize findings. In this study, we assessed the magnitude of NNM and its associated factors among neonates born in major public hospitals in Harari Region, Eastern Ethiopia.

Methods

Study settings

This study was conducted in public hospitals in Harar, Eastern Ethiopia. Harar is the capital city of Harari Region, located 526 km away from the capital city, Addis Ababa. According to population projection from the 2007 census, the region has an estimated population of 263,656 in 2020/2021.¹⁷ There are two public hospitals, one private hospital, one police hospital, eight health centers, 54 private clinics, and 24 health posts in the region—majority of which are located in Harar. The study was conducted in Hiwot Fana Comprehensive Specialized University Hospital (HFSUH) and Jugel hospital (JH). HFSUH is the tertiary hospital affiliated with Haramaya University while JH is a regional hospital located in the same town. Both hospitals provide maternal neonatal and child health services. HFSUH provides comprehensive specialized care including the only well-established neonatal intensive care unit (NICU) in the region. It is serving as a major referral center serving more than 5 million

people in Eastern Ethiopia. The study was conducted from 20 June to 20 August 2021.

Study design and population

A facility-based cross-sectional study design was conducted. All neonates born in public hospitals in Harari Region were the source population, while all live births in both hospitals during the study period were the study population. All randomly selected live births with their mothers during the study period were the study units.

Inclusion criteria

All live births with their mothers were included in the study.

Exclusion criteria

Self-discharges or discharges against medical advice and neonates referred out were excluded from the study.

Sample size and sampling techniques

Sample size was calculated using single population proportion formula and factors associated with NNM using Epi-Info Version 7.2.4.0 Stat Calc. computer software using the assumptions like power 80%, confidence interval 95%, percent of cases among exposed and unexposed, and finally, the largest sample size was considered ($n=405$). After sample was allocated to each hospital proportionally based on the expected births over the study period, a systematic random sampling technique was employed to select eligible neonate–mother pairs and interviewed at discharge. Every third live birth from the delivery registration book were identified and interviewed at discharge.

Data collection

Data were collected through interview of the mother using a structured questionnaire supplemented with review of maternal and neonatal medical records.⁶ The questionnaire consists of data on maternal and fetal sociodemographic, obstetric conditions, and identification of the presence of any of the pragmatic or management-based NNM criteria.^{6,7} Data were collected by four well-trained BSc midwives and supervised by two MSc midwives in maternity and neonatal nursing, who are fluent in the local languages (Afan Oromo and Amharic). A pre-tested structured interviewer-administered questionnaire was used to collect sociodemographic data complemented with review of maternal and neonatal records at discharge. Onsite daily supervision was carried out throughout the data collection period.

Measurement and variables

The outcome of the study, NNM, was categorized as “Yes” (code as 1), if they fulfill any of the pragmatic or/and

management criteria at discharge^{2,6,7} or No (coded as 0) otherwise. The independent variables included socio-demographic characteristics (age, education, occupation, residence, referral status, smoking status), medical, and obstetrics conditions (diabetes mellitus, syphilis, anemia in pregnancy, parity, history of abortion, history of neonatal loss, antenatal care (ANC), birth interval, onset of labor, pregnancy induced hypertension, premature rupture of

membrane, antepartum hemorrhage, obstetric complications, and neonatal conditions).

Operational definition

NNM: any neonate identified with at least one of the following pragmatic or/and management criteria but survived either by chance or treatment (Box 1).⁶

Box 1. Neonatal near miss identification criteria.^{2,6,7}

| Pragmatic criteria | Management criteria |
|--|---|
| <ul style="list-style-type: none"> • Birth weight < 1750 g • Gestational age < 33 weeks • 5th minutes Apgar score < 7; validated by world health organization multi-country studies (WHOMCS)⁶ | <ul style="list-style-type: none"> • Parenteral therapeutic antibiotics • Positive pressure ventilation • Parenteral nutrition • Nasal continuous positive airway pressure • Any intubations • Phototherapy use • Use of anticonvulsant drugs • Use of vasoactive drugs • Cardiopulmonary resuscitation (CPR) • Use of steroid for the treatment of refractory hypoglycemia • Use of any blood products and any surgery done |

Statistical analysis

All collected data were cross-checked for completeness and consistency, coded, and double-entered to Epi-Data 3.1 and exported to SPSS 26 for analysis. Descriptive summary measures such as frequency, percentages, means, and standard deviation were used to describe the characteristics of the participants. Bivariable logistic analysis was used to identify variables associated with NNM, and variables with *p* value of <.25 in the bivariable analysis were candidates for multivariable analysis after checking for multicollinearity. Multicollinearity was checked using variance inflation factor (VIF) or tolerance test and standard error. No variables with VIF > 10 or standard error of >2 or tolerance test <0.1 were identified with the test. Model fitness was also checked using Hosmer-Lemeshow tests, and the model was fitted. Association was described using adjusted odds ratio (AOR) along with 95% confidence interval (CI). Finally, *p* < .05 in the multivariable analysis was considered as a cut-off point for statistically significant association.

Results

Characteristics of study participants

Of 405 women–neonate pairs approached, 401 (99%) were included in the study. The mean age of the mothers was 26.7 years (\pm 5.5 years), ranging from 16 to 40 years. The majority of the mothers were 20–34 years old (81%), married (93.3%), and housewives (78.8%; Table 1). A fifth of the mothers had pregnancy induced hypertension (20.2%), history of abortion (22%), and prolonged labor (19.2%). One in

four (38.7%) births was by cesarean section (Table 2). Majority of the neonates had vertex presentation (83.8%) and delivered at term (78.6%). However, a third of the neonates had non-reassurance fetal heart rate (34.9%) and were admitted to NICU (29.7%; Table 3).

Magnitude of NNM. At discharge, 126 (31.4%; 95% CI = [26.9, 36.2]) neonates had at least one NNM event. Accordingly, 3 out of 10 neonates delivered in public hospitals developed NNM at discharge. The most frequent pragmatic criteria of NNM were low fifth minutes Apgar score (54.8%) while use of parenteral antibiotics (61.9%) and cardiopulmonary resuscitation (61.9%) were the major NNM events as per the management-based criteria (Table 4).

Factors associated with NNM

After controlling for confounding factors, antenatal care, parity, antepartum hemorrhage, premature rupture of membrane, obstructed labor, non-vertex presentation, and maternal referral status were significantly associated with NNM. The odds of NNM among neonates born to mothers who were referred from other facilities were 2 times (AOR=2.24, 95% CI = [1.25, 4.03]) compared with those from mothers who were not referred. Neonates of primiparous women were almost 3 times (AOR=2.65, 95% CI = [1.49, 4.77]) more likely to have NNM compared with their counterparts. Moreover, NNM was 2 times (AOR=2.08, 95% CI = [1.10, 3.93]) and 4 times (AOR=4.29, 95% CI = [2.16, 8.53]) higher among mothers with no antenatal care and who had antepartum hemorrhage (APH), respectively, compared

Table 1. Sociodemographic characteristics of mothers of neonates born in public hospitals in Harari Region, Eastern Ethiopia, 2021 (n = 401).

| Characteristics | Frequency (n) | Percent (%) |
|---|---------------|-------------|
| Age of mothers in years | | |
| <20 | 40 | 10.0 |
| 20–34 | 325 | 81.0 |
| 35–49 | 36 | 9.0 |
| Marital status | | |
| Married | 374 | 93.3 |
| Others | 27 | 6.7 |
| Maternal education | | |
| No formal education | 269 | 67.1 |
| 1–8 | 58 | 14.4 |
| 9–12 | 36 | 9.0 |
| Diploma and above | 38 | 9.5 |
| Paternal education level (n = 396) | | |
| No formal education | 210 | 53.0 |
| 1–8 | 60 | 15.2 |
| 9–12 | 57 | 14.4 |
| Diploma and above | 69 | 17.4 |
| Maternal occupation | | |
| Housewife | 316 | 78.8 |
| Merchant | 50 | 12.5 |
| Non-government employee | 29 | 7.2 |
| Daily laborer | 6 | 1.5 |
| Paternal occupation (n = 396) | | |
| Farmer | 184 | 46.5 |
| Merchant | 91 | 23.0 |
| Non-government employee | 76 | 19.2 |
| Daily laborer | 45 | 11.3 |
| Residence | | |
| Urban | 174 | 43.4 |
| Rural | 227 | 56.6 |
| Mother's smoking status in pregnancy | | |
| Yes | 14 | 3.5 |
| No | 387 | 96.5 |
| Maternal referred from other facilities | | |
| Yes | 222 | 55.4 |
| No | 179 | 46.6 |

with their counterparts. Neonates born from mothers with premature rupture of membrane and obstructed labor were 4 (AOR=4.07, 95% CI = [2.05, 8.07]) and 2 (AOR=2.61, 95% CI = [1.23, 5.52]) times more likely to develop NNM, respectively, compared with their counterparts. In addition, neonates with non-vertex presentation were 3 times (AOR=3.03, 95% CI = [1.54, 5.95]) more likely to become near miss than their counterparts (Table 5).

Discussion

In this study, we assessed the magnitude of NNM and its associated factors among neonates born in major public

Table 2. Maternal medical and obstetrics conditions among neonates born in public hospitals in Harari Region, Eastern Ethiopia, 2021 (n = 401).

| Variables | Frequency (n) | Percent (%) |
|--|---------------|-------------|
| Diabetes mellitus | | |
| Yes | 14 | 3.5 |
| No | 387 | 96.5 |
| Anemia in pregnancy | | |
| Yes | 27 | 6.7 |
| No | 374 | 93.3 |
| Syphilis in pregnancy | | |
| Yes | 12 | 3.0 |
| No | 389 | 97.0 |
| Parity | | |
| 1 | 149 | 37.2 |
| 2–4 | 183 | 45.6 |
| ≥5 | 69 | 17.2 |
| History of abortion | | |
| Yes | 88 | 22.0 |
| No | 313 | 78.0 |
| History of stillbirth | | |
| Yes | 42 | 10.5 |
| No | 359 | 89.5 |
| History of neonatal loss | | |
| Yes | 24 | 6.0 |
| No | 377 | 94.0 |
| Had ANC visit | | |
| Yes | 321 | 80.0 |
| No | 80 | 20.0 |
| Number of ANC visits (n = 321) | | |
| 1–3 times | 189 | 58.9 |
| ≥4 times | 132 | 41.1 |
| Gestational age at first ANC visit (321) | | |
| ≤12 weeks | 71 | 22.1 |
| >12 weeks | 250 | 78.9 |
| Onset of labor | | |
| Spontaneous | 294 | 73.3 |
| Induced | 76 | 19.0 |
| CS before labor | 31 | 7.7 |
| Obstetrics complications | | |
| Pregnancy-induced hypertension | 81 | 20.2 |
| Antepartum hemorrhage | 65 | 16.2 |
| Premature rupture of membrane | 58 | 14.5 |
| Prolonged labor | 77 | 19.2 |
| Obstructed labor | 49 | 12.2 |
| Oligohydramnios | 17 | 4.2 |
| Mode of delivery | | |
| SVD | 200 | 49.9 |
| Instrumental assisted vaginal delivery | 46 | 11.5 |
| CS | 155 | 38.6 |

ANC: antenatal care; CS: cesarean section; SVD: spontaneous vertex delivery.

Table 3. Neonatal characteristics among neonates born in public hospitals in Harari Region ($n=401$).

| Characteristics | Frequency (n) | Percent (%) |
|---|---------------|-------------|
| Fetal presentation at birth | | |
| Non-vertex | 65 | 16.2 |
| Vertex | 336 | 83.8 |
| Non-reassurance fetal heart rate pattern | | |
| Yes | 140 | 34.9 |
| No | 261 | 65.1 |
| Sex of neonate | | |
| Male | 205 | 51.0 |
| Female | 196 | 49.0 |
| Gestational age at birth | | |
| <37 weeks | 75 | 18.7 |
| 37–42 weeks | 315 | 78.6 |
| >42 weeks | 11 | 2.7 |
| 5th minutes Apgar score | | |
| <7 | 69 | 17.20 |
| ≥7 | 332 | 82.80 |
| Birth weight (grams) | | |
| <2500 | 41 | 10.2 |
| 2500–3999 | 341 | 85.0 |
| ≥4000 | 19 | 4.8 |
| Neonatal complication | | |
| Yes | 147 | 36.7 |
| No | 254 | 63.3 |
| Type of complication identified ($n=147$) | | |
| Birth asphyxia | 86 | 58.5 |
| Respiratory distress syndrome | 89 | 60.5 |
| Meconium aspiration syndrome | 105 | 71.4 |
| Birth trauma | 18 | 12.2 |
| Hypothermia | 43 | 29.3 |
| Hypoglycemia | 12 | 8.2 |
| Congenital malformation | 7 | 4.8 |
| Admitted to NICU | | |
| Yes | 119 | 29.7 |
| No | 282 | 70.3 |

NICU: neonatal intensive care unit.

hospitals in Harari Region, Eastern Ethiopia. We found that the overall magnitude of NNM was 31.4% (95% CI = [26.9, 36.2]), which is relatively higher than other countries report and much higher than world health organization multi-country studies (WHOMCS) (7.25%).⁶ NNM was found to be more likely among women who were referred from other facilities, primiparous, had no prenatal care, had antepartum hemorrhage, obstructed labor, premature rupture of membrane, and neonates with non-vertex fetal presentations at birth.

Our finding is in line with findings in Ethiopia [Hawassa (33.4%)¹⁵ and Debre Tabor (32.9%)¹⁶] and Brazilian university hospitals (30.3%).¹⁸ However, it is higher than some other studies in Ethiopia [Jimma (26.7%)¹⁴ and Injibara (23.3%)¹⁹], Nepal (7.9%),²⁰ Brazil [(8.7%),²¹ (22.2%),²²

Table 4. Description of pragmatic and management events among NNM cases ($n=126$).

| Neonatal near miss events | Frequency (n) | Percent (%) |
|--|---------------|------------------|
| Pragmatic criteria | | |
| Birth weight < 1750 g | 9 | 7.1 |
| 5th minutes Apgar score < 7 | 69 | 54.8 |
| Gestational age < 33 weeks at birth | 15 | 11.9 |
| Management criteria | | |
| Use of therapeutic parenteral antibiotics | 78 | 61.9 |
| Use of nasal continuous positive air pressure | 49 | 38.9 |
| Positive pressure ventilation | 68 | 54.0 |
| Use of any intubation | 29 | 23.0 |
| Use of parenteral nutrition | 25 | 19.8 |
| Use of phototherapy | 19 | 15.1 |
| Cardiopulmonary resuscitation | 78 | 61.9 |
| Use of any vasoactive drugs | 33 | 26.2 |
| Use anticonvulsants | 26 | 20.6 |
| Use of steroids for the treatment of refractory hypoglycemia | 6 | 4.8 |
| Any surgery done | 6 | 4.8 |
| Use of any blood products | 5 | 4.0 |
| Any pragmatic or management markers of severity | 126 | 100 ^a |

NNM: neonatal near miss.

^aPercentage > 100% since some neonates have more than one neonatal near miss event.

(3.3%)²³], and the WHOMCS (7.3%).⁶ This might be related with differences in sociodemographic conditions and mothers' ability of early recognition of complications and health-seeking behavior, study settings, and NNM identification criteria. However, our study is lower than a finding from Uganda (36.7%)²⁴ and Ghana (70%).²⁵ This might be related with differences in inclusion criteria and settings.

Consistent with previous studies, NNM was more likely among neonates born to women who were referred from other facilities, primiparous, did not have prenatal care, had obstetric complications, and non-vertex fetal presentations. The high burden of NNM among neonates born to mothers who were referred from other facilities was also reported in Ethiopia^{11,19} and India.²⁶ This might be related to the delays in reaching facilities on time and the obstetric complications leading to referral.²⁷ Given problems with our referral system—delays, poor communication, and safety of roads and distances—it is more likely that the women will reach the appropriate facilities after it is too late for preventing maternal or neonatal complications.²⁸ In addition, effect of obstetric complications on adverse neonatal outcomes was previously reported.²⁹

In line with several studies,^{16,19,22,30} we found that NNM was higher among primiparous mothers. This might be

Table 5. Factors associated with NNM in public hospitals in Harari Region, Eastern Ethiopia, 2021.

| Variables category | Neonatal near miss (NNM) | | COR (95% CI) | AOR (95% CI) |
|--------------------------|--------------------------|------------|--------------------|---------------------------|
| | Yes (%) | No (%) | | |
| Maternal age in years | | | | |
| <20 | 18 (45.0) | 22 (55.0) | 2.20 [1.13, 4.30] | 1.13 [0.48, 2.69] |
| ≥35 | 20 (55.6) | 16 (44.4) | 3.37 [1.70, 6.80] | 2.70 [0.99, 6.70] |
| 20–34 | 88 (27.1) | 237 (72.9) | 1 | 1 |
| Maternal referral status | | | | |
| No | 27 (15.1) | 152 (84.9) | 1 | 1 |
| Yes | 99 (44.6) | 123 (55.4) | 4.53 [2.80, 7.40] | 2.24 [1.25, 4.03]* |
| Anemia in pregnancy | | | | |
| No | 112 (29.9) | 262 (70.1) | 1 | 1 |
| Yes | 14 (51.9) | 13 (48.1) | 2.52 [1.15, 5.53] | 1.68 [0.61, 4.59] |
| Parity | | | | |
| 1 | 59 (39.6) | 90 (60.4) | 1.81 [1.18, 2.79] | 2.67 [1.49, 4.77]* |
| ≥2 | 67 (26.6) | 185 (74.4) | 1 | 1 |
| Antenatal care visit | | | | |
| No | 43 (53.8) | 37 (46.2) | 3.33 [2.01, 5.53] | 2.08 [1.10, 3.93]* |
| Yes | 83 (25.9) | 238 (74.1) | 1 | 1 |
| History of abortion | | | | |
| No | 86 (27.5) | 227 (72.5) | 1 | 1 |
| Yes | 40 (45.5) | 48 (54.5) | 2.20 [1.35, 3.58] | 1.48 [0.79, 2.76] |
| Antepartum hemorrhage | | | | |
| No | 84 (25.0) | 252 (75.0) | 1 | 1 |
| Yes | 42 (64.6) | 23 (35.4) | 5.48 [3.11, 9.64] | 4.29 [2.16, 8.53]* |
| Obstructed labor | | | | |
| No | 100 (28.4) | 252 (71.6) | 1 | 1 |
| Yes | 26 (53.1) | 23 (46.9) | 2.85 [1.55, 5.23] | 2.61 [1.23, 5.52]* |
| PROM | | | | |
| No | 91 (26.5) | 252 (73.5) | 1 | 1 |
| Yes | 35 (60.3) | 23 (39.7) | 4.21 [2.36, 7.51] | 4.07 [2.05, 8.07]* |
| Fetal presentation | | | | |
| Vertex | 83 (24.7) | 253 (75.3) | 1 | 1 |
| Non-vertex | 43 (66.2) | 22 (33.8) | 5.96 [3.37, 10.54] | 3.03 [1.54, 5.95]* |

NNM: neonatal near miss; COR: crude odds ratio; AOR: adjusted odds ratio; CI: confidence interval; PROM: premature rupture of membrane.

*Indicates that significant at $p < .05$, 1, reference.

related with limited experience of pregnancy, danger signs, labor, and delivery, delay in health-seeking, and high risk for prolonged labor, induction of labor, and birth asphyxia among primiparous.^{31,32} Moreover, our finding shows those neonates born to mothers with no antenatal care were more likely to develop NNM in agreement with other studies.^{9,10,16,28} Having prenatal care will enable women to get information about their pregnancy, knowledge regarding danger signs, and where to seek care when needed,^{33,34} thus reducing risk of NNM. We also found that complications in pregnancy or labor and delivery are significant contributors of NNM. Consistent with other studies, NNM is more likely among neonates born to a woman who had APH.^{16,30} Given APH increases the risk of compromised fetal blood perfusion, which leads to uteroplacental insufficiency, this might result in preterm birth and birth asphyxia and thereby NNM.^{31,35–37} Similarly, neonates born to a mother with premature rupture of membrane (PROM) were more likely to

have NNM, a finding supported by previous studies.^{10,16,19} Since association of PROM with preterm birth, respiratory distress syndrome, intravascular hemorrhage, cord prolapse, reduced amniotic fluid volume, chorioamnionitis, neonatal sepsis, and birth asphyxia was reported,^{38–40} this finding was expected. Similarly, obstructed labor^{14,16,19} and non-vertex presentation were associated with NNM.^{10,14}

Strengths and limitations of the study

The strength of this study is the use of combined data collection technique (face-to-face interview and review of medical records) which enabled us to identify basic sociodemographic and obstetric factors. However, our study has also some limitations. *First*, our follow-up was limited to hospital discharges or 28 days, whichever comes first. As a result, NNM after discharge or among neonates referred out might be missed. *Second*, since we limited our observation to only

hospital births, NNM might be overestimated since majority of births—especially low risk births are occurring at home or health centers—since mainly women with high risk pregnancies are giving birth in facilities.

Conclusion

We found that NNM among neonates delivered in public hospitals in Eastern Ethiopia is higher than findings from similar settings in Ethiopia. NNM was more likely among neonates born to a woman who is referred, primiparous, and had no prenatal care, had non-vertex presentation, and obstetrics complications (antepartum hemorrhage, premature rupture of membrane, and obstructed labor). Improving NNM and attaining sustainable development goals require addressing major obstetric complications resulting in NNM. Further study on the quality of care provided to neonates and delays in the referral system is warranted to identify missed opportunities in the care.

Acknowledgements

The authors would like to thank managers of both hospitals for facilitation of the study in their respective hospitals. The authors also thank Haramaya University for funding the study as part of MSC study to MAT.

Author contributions

M.A.T., A.K.T., and A.S. conceived study. M.A.T. analyzed and interpreted the data under the close mentorship and supervision of A.K.T., A.S., and G.T. M.A.T. wrote the original draft of the manuscript with continuous input from A.K.T., A.S., and G.T. All authors contributed to the writing and reviewed the article and approved the final manuscript to be published.

Declaration of conflicting interests


The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was funded by Haramaya University as part of MSC study by MAT.

ORCID iDs

Melese Adugna Tola  <https://orcid.org/0000-0001-7728-2980>

Getahun Tiruye  <https://orcid.org/0000-0002-0441-0440>

References

1. UN, WHO, UNICEF, et al. *Levels and trends in child mortality: report 2018: United Nation Population Division; United Nations on behalf of the UN Inter-agency Group for Child Mortality Estimation*. New York: UN, 2019.
2. Surve S, Sanjay C and Ragini K. Neonatal near miss review: tracking its conceptual evolution and way forward. *Curr Pediatr Res* 2017; 21, <https://www.alliedacademies.org/articles/neonatal-near-miss-review-tracking-its-conceptual-evolution-and-wayforward.html>
3. EMDHS. *Ethiopia mini demographic and health survey 2019: key indicators*. Rockville, MD: EPHI and ICF, 2019.
4. Da Silva AAM, Leite ÁJM, Lamy ZC, et al. Morbidade neonatal near miss na pesquisa Nascer no Brasil. *Cadern Saúde Públ* 2014; 30: S182–S91.
5. Santos JP, Pileggi-Castro C, Camelo JS Jr, et al. Neonatal near miss: a systematic review. *BMC Pregn Childb* 2015; 15: 320.
6. Pileggi-Castro C, Camelo JS Jr, Perdoná GC, et al. Development of criteria for identifying neonatal near-miss cases: analysis of two WHO multicountry cross-sectional studies. *BJOG* 2014; 121(Suppl. 1): 110–118.
7. Santos JP, Cecatti JG, Serruya SJ, et al. Neonatal Near Miss: the need for a standard definition and appropriate criteria and the rationale for a prospective surveillance system. *Clinics* 2015; 70(12): 820–826.
8. Belachew A, Tewabe T and Dessie G. Neonatal mortality and its association with antenatal care visits among live births in Ethiopia: a systematic review and meta-analysis. *J Matern Fetal Neonatal Med* 2020; 35: 348–355.
9. Yohannes E, Assefa N and Dessie Y. Determinants of neonatal near miss among neonates admitted to Ambo University Referral Hospital and Ambo General Hospital, Ethiopia, 2019. *J Pregn Child Health* 2020; 7: 11–18.
10. Mersha A, Bante A and Shibiru S. Factors associated with neonatal near-miss in selected hospitals of Gamo and Gofa zones, southern Ethiopia: a nested case-control study. *BMC Pregn Childb* 2019; 19: 516.
11. Abebe H, Wasie A, Yeshaneh A, et al. Determinant factors of neonatal near miss among neonates in Gurage Zone Hospitals, Ethiopia: a case-control study. *Pediatric Health Med Ther* 2021; 12: 129–139.
12. Nugussie F, Alemayehu M and Mariam KG. A case-control study examining determinants of neonatal near-miss in public hospitals in Tigray Region, Northern Ethiopia. *Res Rev: J Med Sci Technol* 2019; 7: 1–11.
13. Tekelab Tesfalidet Chojenta et al. Incidence and determinants of neonatal near miss in south Ethiopia: a prospective cohort study. *BMC Pregn Childb* 2020; 20: 1–13.
14. Wondimu M, Balcha F, Bacha G, et al. The magnitude of neonatal near miss and associated factors among live births in public hospitals of Jimma Zone, Southwest Ethiopia, 2020: a facility-based cross-sectional study. *PLoS ONE* 2021; 16(5): e0251609.
15. Tekola Anteneh F, Baye G, Amaje E, et al. Neonatal near misses and associated factors among mothers who give a live neonate at Hawassa City governmental hospitals, 2019: a facility-based cross-sectional study design. *BMC Pregn Childb* 2021; 21: 1–9.
16. Tassew HA, Kassie FY and Mihret MS. Neonatal near miss and its predictors among neonates delivered at Debretabor General Hospital, Northern Ethiopia; a retrospective analysis. *Int J Pediatr* 2020; 2020: 1092479.
17. Ethiopia Central Statistical Agency. *Health survey 2007 Addis Ababa*. Addis Ababa, Ethiopia: Ethiopia Central Statistical Agency, 2012.
18. Morais LR, Patz BC, Campanharo FF, et al. Neonatal near miss among newborns of women with Type 1 diabetes mellitus. *Obstet Gynecol Int* 2019; 2019: 8594158.

19. Gebrehana B, Habtamu L, Simachew K, et al. Neonatal near miss and its associated factors at Injibara General Hospital, Awi Zone, Northwest Ethiopia, 2019. *Exp Res Hyp Med* 2020; 5: 62–69.
20. Sushma R, Norhayati N and Nik Hazlina NH. The prevalence of neonatal near miss and associated factors in Nepal: a cross-sectional study. *BMC Pregn Childb* 2020; 21: 422.
21. De Araujo Brasilia DRP, Vilela MBR, de França KEX, et al. Morbidade neonatal near Miss em Hospitais Terciários de Uma Capital do nordeste do Brasilia. *Rev Paul Pediatr* 2019; 37: 275–282.
22. De Lima THB, Katz L, Kassab SB, et al. Neonatal near miss determinants at a maternity hospital for high-risk pregnancy in Northeastern Brazil: a prospective study. *BMC Pregn Childb* 2018; 18: 401.
23. Silva GA, Rosa KA, Saguier ESF, et al. A populational based study on the prevalence of neonatal near miss in a city located in the South of Brazil: prevalence and associated factors. *Rev Brasil Saúde Mater Infant* 2017; 17: 159–167.
24. Nakimuli A, Mbalinda SN, Nabirye RC, et al. Still births, neonatal deaths and neonatal near miss cases attributable to severe obstetric complications: a prospective cohort study in two referral hospitals in Uganda. *BMC Pediatr* 2015; 15: 44.
25. Bakari A, Bell AJ, Oppong SA, et al. Neonatal near-misses in Ghana: a prospective, observational, multi-center study. *BMC Pediatr* 2019; 19: 509.
26. Shroff BD and Ninama NH. A call for eminence obstetrics care by way of “neonatal near miss” events (NNM): a hospital-based case–control study. *J Obstet Gynaecol India* 2019; 69(1): 50–55.
27. Yeshialem E, Abera M and Tesfay A. Determinants of adverse pregnancy outcomes among mothers who gave birth from Jan 1-Dec 31/2015 in Jimma University specialized hospital, case-control study, 2016. *Ethiop J Reprod Health* 2019; 11: 10.
28. Carvalho OMC Jr, Viana AB Jr, Augusto MCC, et al. Delays in obstetric care increase the risk of neonatal near-miss morbidity events and death: a case-control study. *BMC Pregn Childb* 2020; 20: 437.
29. Tura AK, Scherjon S, van Roosmalen J, et al. Surviving mothers and lost babies—burden of stillbirths and neonatal deaths among women with maternal near miss in eastern Ethiopia: a prospective cohort study. *J Glob Health* 2020; 10(1): 01041310.
30. Kale PL, de Mello-Jorge MHP, da Silva KS, et al. Neonatal near miss and mortality: factors associated with life-threatening conditions in newborns at six public maternity hospitals in Southeast Brazil. *Cad Saúde Públ* 2017; 33: e00179115.
31. Tasew H, Zemicheal M, Teklay G, et al. Risk factors of birth asphyxia among newborns in public hospitals of Central Zone, Tigray, Ethiopia. *BMC Res Notes* 2018; 2018: 496.
32. Kassahun EA, Aweke AM, Getu AA, et al. Proportion and associated factors of nonreassuring fetal heart rate patterns in Finote Selam Primary Hospital, North West Ethiopia. *Biomed Res Int* 2020; 2020: 6948972.
33. Shiferaw K, Mengiste B, Gobena T, et al. The effect of antenatal care on perinatal outcomes in Ethiopia: a systematic review and meta-analysis. *PLoS ONE* 2021; 16(1): e0245003.
34. Tolossa T, Fekadu G, Mengist B, et al. Impact of antenatal care on neonatal mortality among neonates in Ethiopia: a systematic review and meta-analysis. *Arch Publ Health* 2020; 78: 1–11.
35. Gebremedhin MM, Gebremichael MW, Gebremichael BG, et al. Determinants of low APGAR score in newborns delivered at Lemlem Karl General Hospital, Northern Ethiopia, 2018: a case-control study, 2020, <https://assets.researchsquare.com/files/rs-32019/v2/471e4d83-4c3e-4038-9e0a-c1d71684c7c9.pdf?c=1631856465>
36. Muchie KF, Lakew AM, Teshome DF, et al. Epidemiology of preterm birth in Ethiopia: systematic review and meta-analysis. *BMC Pregn Childb* 2020; 20: 1–12.
37. Zanardi DM, Parpinelli MA, Haddad SM, et al. Adverse perinatal outcomes are associated with severe maternal morbidity and mortality: evidence from a national multicentre cross-sectional study. *Arch Gynecol Obstet* 2019; 299(3): 645–654.
38. Alemu AY, Belay GM, Berhanu M, et al. Determinants of neonatal mortality at neonatal intensive care unit in Northeast Ethiopia: unmatched case-control study. *Trop Med Health* 2020; 48: 40–10.
39. Debelew GT, Afework MF and Yalew AW. Determinants and causes of neonatal mortality in Jimma zone, southwest Ethiopia: a multilevel analysis of prospective follow-up study. *PLoS ONE* 2014; 9(9): e107184.
40. Masanja PP, Kibusi SM and Mkhoyi ML. Predictors of early-onset neonatal sepsis among neonates in Dodoma, Tanzania: a case-control study. *J Trop Pediatr* 2020; 66: 257–266.