

University of Groningen

Shedding light on active species in Fe, Ni and Cu catalysis

Draksharapu, Apparao

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Draksharapu, A. (2013). *Shedding light on active species in Fe, Ni and Cu catalysis: photochemical, spectroscopic and electrochemical studies*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Summary

Many of the transformations that chemists struggle with in the laboratory are often carried out by nature with better activity, selectivity and with bio-degradable waste. In sharp contrast to biological systems, manmade structural and functional models of biological systems usually suffer severely from a lack of reactivity and selectivity in their catalytic reactions. Matching the capabilities of biological systems with mimics in the laboratory is a seemingly impossible task. In biological systems complex structures are involved in chemical reactions, for example enzymes, where chemical reactions are carried out at a so called active site. Apart from the active site, the remaining part of the protein provides a second coordination sphere, which helps to achieve better selectivities towards chemical reactions by allowing the substrate to enter into the active site preferentially in only one orientation. These advantages are not readily available to biomimetic systems. However, in a few biomimetic systems a controlled second coordination sphere environment has been created successfully.¹ One such system is the topic of chapter 8, however, albeit that the catalytic reactions are not discussed there. In this thesis a number of biomimetic systems including the Fe^{II}-N4Py group of complexes (Figure 1), which are functional mimics for Fe^{II}-BLM (BLM = bleomycin) (chapter 2), and Fe^{III}-OCl/Ni^{II}-OCl species being mimics for haloperoxidases (chapter 6 and 7), are discussed.




Figure 1 Structures of the complexes employed in this thesis.

When chemists refer to complexes as biomimetic, they generally mean that their systems are able to generate intermediates and, hence, catalyse chemical reactions similar to those carried out by the biological systems referred to. Generation of active intermediates, their characterisation by various spectroscopic techniques and

understanding their role in catalysis have direct relevance to understanding their role in biological systems. The synthesis and characterisation of several intermediates such as $\text{Fe}^{\text{IV}}=\text{O}$ (chapter 4), $\text{Fe}^{\text{III}}-\text{OR}$ (chapter 5), $\text{Fe}^{\text{III}}-\text{OCl}$ (chapter 6), $\text{Ni}^{\text{II}}-\text{OCl}$ and $\text{Ni}^{\text{II}}-\text{O}^{\cdot}$ (chapter 7) are discussed in this thesis. The multi-technique approach taken in this work allows for the detailed characterisation of several of these intermediates. In particular, Raman spectroscopy plays a pivotal role in the studies described here.

The aqueous chemistry of complexes **1** and **2** was explored extensively in chapter 2. A multi-technique approach was taken to establish the speciation of $\text{Fe}^{\text{II}}(\text{N4Py})$ complexes of relevance to DNA cleavage studies reported earlier.² Exchange of the ligand CH_3CN with aqua/hydroxido ligands occurs immediately upon dissolution of $[(\text{N4Py})\text{Fe}^{\text{II}}(\text{CH}_3\text{CN})]^{2+}$ in water. Moreover, depending on the pH, distinct species exist in aqueous solution and interchange reversibly with changes in pH. Furthermore, the pentadentate N4Py ligand is tetradentate at high pH. A further complication to speciation analysis of this system is that the complexes present in solution at equilibrium are a mixture of low and high spin states and show rapid spin interconversion. The equilibrium between singlet (low spin) and quintet (high spin) spin states in aqueous solution was proposed to be at the origin of the activity of $\text{Fe}^{\text{II}}(\text{N4Py})$ complexes in cleaving DNA with oxygen ($^3\text{O}_2$) as terminal oxidant, through the transient population of a triplet state.

The photo-induced oxidation of complexes **1** and **2** in various solvents (*i.e.* H_2O and MeOH) with oxygen as terminal oxidant is discussed in chapter 3. Interestingly, both complexes are photochemically inactive in acetonitrile, reflecting their high positive oxidation potentials (1.1 V *vs.* SCE). Irradiation with UV and visible light induces the oxidation of iron(II) complexes to the corresponding iron(III) complexes with oxygen as terminal oxidant. Moreover, the primary photoproduct of **2** in methanol is also photochemically active. Photochemical enhancement of the rate of oxidation of iron(II) complexes by oxygen ($^3\text{O}_2$) through outer sphere electron transfer was proposed. These results are important in understanding the photo enhanced cleavage of DNA with **1**. As mentioned above, the interconversion between the singlet and higher spin states would be expected to involve an intermediate triplet state, which would facilitate electron transfer to $^3\text{O}_2$, the first step in the cleavage of DNA by **1** with oxygen ($^3\text{O}_2$).

The photochemistry of iron complexes was extended further to the $\text{Fe}^{\text{IV}}=\text{O}$

complexes, which is the subject of chapter 5. In sharp contrast to **1** and **2**, complexes **1a** and **2a** show photochemical activity in acetonitrile. These complexes undergo photoreduction to the corresponding $\text{Fe}^{\text{II}}\text{-CH}_3\text{CN}$ complexes under irradiation with UV and visible light (400 nm). Partial photoreduction to Fe^{III} and to Fe^{II} complexes was observed even in the solid state (dispersed in KCl) under irradiation with visible light. These photochemical properties were explored with respect for enhancing catalysis, *e.g.*, alkane hydroxylation and alkene epoxidation. The photochemical reduction of the $\text{Fe}^{\text{IV}}=\text{O}$ complexes is faster in the presence of substrates such as ethyl benzene, benzyl alcohol, fluorene and indane. There are two steps involved in the photochemical process, the first step is the conversion of $\text{Fe}^{\text{IV}}=\text{O}$ to $\text{Fe}^{\text{III}}\text{-OH/OR}$ followed by a second photochemical step in which the $\text{Fe}^{\text{III}}\text{-OR}$ converts to $\text{Fe}^{\text{III}}\text{-CH}_3\text{CN}$. This latter species is a powerful enough oxidant to oxidise $\text{Fe}^{\text{III}}\text{-OH/OR}$ to the $\text{Fe}^{\text{IV}}=\text{O}$ state and form the $\text{Fe}^{\text{II}}\text{-CH}_3\text{CN}$ complex also. In addition, the $\text{Fe}^{\text{III}}\text{-OMe}$ (**3**) complex is photochemically active in acetonitrile. In sharp contrast, the $\text{Fe}^{\text{III}}\text{-OH}$ complex in acetonitrile is photochemically much less active. So the primary photoproduct in the reduction of $\text{Fe}^{\text{IV}}=\text{O}$ is not an $\text{Fe}^{\text{III}}\text{-OH}$ species, but instead an $\text{Fe}^{\text{III}}\text{-OR}$ ($\text{R} = \text{alkyl}$) species. These data suggests that the C-H hydroxylation is via a rebound mechanism. The enhanced reactivity of $\text{Fe}^{\text{IV}}=\text{O}$ species, under irradiation, towards substrate oxidation is attributed to the transient population of an $\text{S} = 2$ state.

Knowledge of the redox properties of $\text{Fe}^{\text{IV}}=\text{O}$ species benefits the understanding of their reactivity towards substrate oxidation. Unfortunately, determination of the $\text{Fe}^{\text{IV}}/\text{Fe}^{\text{III}}$ reduction potential by cyclic voltammetry is hampered by the sluggish rate of heterogeneous electron transfer encountered in the reduction. These problems are dealt with in depth in chapter 4. The chemical and electrochemical synthesis and characterisation of a non-heme $\text{Fe}^{\text{IV}}=\text{O}$ species in water is reported. A clean conversion of complex **2** to **2a** in water was achieved electrochemically. In acetonitrile, conversion of complex **2** to **2a** was observed only in the presence of water. Moreover, the required potential (1.25 V *vs.* SCE) to generate the $\text{Fe}^{\text{IV}}=\text{O}$ species is higher in acetonitrile compared to the value of (0.85 V *vs.* SCE) in water. This is attributed to the necessity of a high potential to oxidise $\text{Fe}^{\text{II}}\text{-CH}_3\text{CN}$ to $\text{Fe}^{\text{III}}\text{-CH}_3\text{CN}$. Once $\text{Fe}^{\text{III}}\text{-CH}_3\text{CN}$ is generated, immediate ligand exchange takes place to form $\text{Fe}^{\text{III}}\text{-OH}_2$, which oxidises to $\text{Fe}^{\text{IV}}=\text{O}$ spontaneously at such high applied potentials. The available data suggested that the oxidation potential of the $\text{Fe}^{\text{IV}}/\text{Fe}^{\text{III}}$ redox couple is around ~ 0.85 V, and not 0.41 V *vs.* SCE in water as concluded by Que and co-workers³ for $[(\text{N4Py})\text{Fe}^{\text{IV}}(\text{O})]^{2+}$.

It has been proposed that in haloperoxidases a M-O-X (M = metal, X = Cl, Br and I) adduct is generated and is responsible for the halogenation of the substrates. Direct spectroscopic evidence for such intermediates had not yet been reported for any non-heme system. In chapter 6, the synthesis and spectroscopic characterisation of a non-heme Fe^{III}-OCl species is reported. Addition of aqueous NaOCl to acidic aqueous solutions of **1** or **2** at room temperature generates the corresponding Fe^{III}-OCl species. Over time the Fe^{III}-OCl species converts to the corresponding high valent Fe^{IV}=O species. Assignment of these intermediates was carried out by UV/Vis absorption, (resonance)Raman and EPR spectroscopy and by Cryo ESI-MS. Raman spectroscopic characterisation of the Fe^{III}-O and O-Cl modes was aided by bromine (Fe^{III}-O-Br) and with ¹⁸O labelling. These intermediates react with styrene to yield the corresponding halohydrin and diol.

In chapter 7, generation of non-porphyrinic Ni^{II}-O[•] and Ni^{II}-OCl species and their spectroscopic characterisation is described. Addition of aqueous NaOCl/Ca(OCl)₂ to an acetonitrile solution of **4** at room temperature generates a putative Ni^{II}-O[•] species. Addition of Brønsted acids promotes the formation of such intermediates and increases the rate of their formation also. Characterisations of these intermediates were carried out with UV/Vis absorption, (resonance)Raman, NMR and EPR spectroscopy and by Cryo ESI-MS. The formation and loss of the Ni^{II}-O[•] species was followed by (resonance)Raman spectroscopy. Formation of the Ni^{II}-O[•] species was confirmed by Raman spectroscopy, again with the use of bromine and with ¹⁸O labelling and evidence for the formation of Ni^{II}-OCl species was obtained by Cryo ESI-MS. The former species is proposed to activate C-H bonds and the latter species is proposed to insert the halogen into the substrate. To the best of my knowledge, this is the first direct spectroscopic evidence for the involvement of Ni^{II}-O[•] and Ni^{II}-OCl species in catalysing the halogenation of aliphatic moieties.

Complexes **5** to **8** catalyse enantioselective C-C bond forming reactions in water with DNA as the source of chirality. Complex **5** provides the highest enantioselectivities in DNA-based asymmetric catalysis, whereas complex **6** yields the opposite enantiomer.⁴ In chapter 8, the interaction with DNA of a series of Cu(II) polypyridyl complexes was studied by various spectroscopic techniques in the absence of substrate. For the first time UV resonance Raman spectroscopy was employed to probe the interaction of Cu(II) complexes with DNA. In the presence of DNA, complexes **7** and **8**, which are intercalators, show clear shifts in their

Raman spectrum whereas the groove binder **5** does not. These spectroscopic observations point out those complexes that induce the highest *ee* in DNA-based asymmetric catalysis are primarily groove binders. These findings provide important insights into the complex structure of these DNA-based catalysts and set the stage for establishing the effect of substrates on the interaction between DNA and the complexes.


Finally in chapter 9 the work described in the thesis is placed in perspective.

References

- (1) (a) J. Steinreiber, T. R. Ward, *Coord. Chem. Rev.*, **2008**, *252*, 751-766; (b) V. Köhler, Y. M. Wilson, C. Lo, A. Sardo, T. R. Ward, *Curr. Opin. Biotech.*, **2010**, *21*, 744-752; (c) A. J. Boersma, R. P. Megens, B. L. Feringa, G. Roelfes, *Chem. Soc. Rev.*, **2010**, *39*, 2083-2092.
- (2) (a) Q. Li, W. R. Browne, G. Roelfes, *Inorg. Chem.*, **2010**, *49*, 11009-11017; (b) Q. Li, W. R. Browne, G. Roelfes, *Inorg. Chem.*, **2010**, *50*, 8318-8325.
- (3) D. Wang, M. Zhang, P. Bühlmann, L. Que, Jr., *J. Am. Chem. Soc.*, **2010**, *132*, 7638-7644.
- (4) (a) G. Roelfes, A. J. Boersma, B. L. Feringa, *Chem. Commun.*, **2006**, 635-637; (b) A. J. Boersma, J. E. Klijn, B. L. Feringa, G. Roelfes, *J. Am. Chem. Soc.*, **2008**, *130*, 11783-11790.

Samenvatting

Vele chemische omzettingen waar chemici in het laboratorium mee worstelen worden door natuurlijke systemen vaak uitgevoerd met hogere activiteit, betere selectiviteit en onder vorming van biologisch afbreekbare bijproducten. In scherpe tegenstelling tot biologische systemen, lijden de door mensen gemaakte structurele en functionele modellen van biologische systemen aan een ernstig gebrek aan reactiviteit en selectiviteit in de door hun gekatalyseerde reacties. Het evenaren van de capaciteiten van biologische systemen met modellen in het laboratorium is een op het eerste gezicht onmogelijke taak. In biologische systemen zijn complexe structuren betrokken bij het uitvoeren van chemische reacties; in enzymen vinden de chemische reacties bijvoorbeeld plaats in het zogenaamde actieve centrum. Naast het actieve centrum vormt een enzym een secundaire coordinatie-omgeving. Dit helpt om een betere selectiviteit van de gekatalyseerde chemische reactie te verkrijgen aangezien hierdoor het substraat bij voorkeur maar in één orientatie in het actieve centrum past. Deze voordelen zijn niet gemakkelijk te bewerkstelligen in een biomimetisch systeem. Het is echter in een aantal biomimetische systemen gelukt om een gedefinieerde secundaire coordinatie omgeving te creëren.¹ Een van deze systemen vormt het onderwerp van hoofdstuk 8, hoewel de katalytische reacties zelf daar niet beschreven worden. In dit proefschrift worden een aantal biomimetische systemen beschreven, waaronder een serie Fe^{II}-N4Py complexen (Figuur 1) die functionele modellen zijn voor Fe^{II}-BLM (BLM = bleomycine) (hoofdstuk 2) en Fe^{III}-OCl/Ni^{II}-OCl deeltjes die model staan voor haloperoxidasen (hoofdstuk 6 en 7).



Figuur 1 De structuren van de complexen die in dit proefschrift beschreven worden.

Wanneer chemici complexen biomimetisch noemen, dan bedoelen ze in het algemeen dat ze in staat zijn om met hun systemen intermediären te genereren en

dat deze, dientengevolge, chemische reacties kunnen katalyseren die normaliter uitgevoerd worden door de nagebootste biologische systemen. Het genereren van actieve intermediaren, de karakterisatie ervan met behulp van diverse spectroscopische technieken en het begrijpen van hun rol in de katalyse zijn direct van belang met betrekking tot het begrijpen van hun rol in biologische systemen. De synthese en karakterisatie van diverse intermediaren zoals $\text{Fe}^{\text{IV}}=\text{O}$ (hoofdstuk 4), $\text{Fe}^{\text{III}}-\text{OR}$ (hoofdstuk 5), $\text{Fe}^{\text{III}}-\text{OCl}$ (hoofdstuk 6), $\text{Ni}^{\text{II}}-\text{OCl}$ en $\text{Ni}^{\text{II}}-\text{O}^{\cdot}$ (hoofdstuk 7) worden in dit proefschrift beschreven. De in dit proefschrift genomen benadering, dat wil zeggen gebruikmakend van een groot aantal technieken, heeft geresulteerd in de gedetailleerde karakterisatie van een aantal van deze intermediaren. Met name Ramanspectroscopie speelt een cruciale rol in het onderzoek dat beschreven wordt in dit proefschrift.

Het onderzoek van de chemie in water van de complexen **1** en **2** wordt uitgebreid beschreven in hoofdstuk 2. Door gebruik te maken van een groot aantal technieken was het mogelijk om vast te stellen welke $\text{Fe}^{\text{II}}(\text{N4Py})$ complexen relevant zijn in verband met eerdere studies hiermee met betrekking tot het knippen van DNA.² Uitwisseling van het CH_3CN ligand met water/hydroxido liganden vindt meteen plaats als $[(\text{N4Py})\text{Fe}^{\text{II}}(\text{CH}_3\text{CN})]^{2+}$ opgelost wordt in water. Afhankelijk van de pH zijn er bovendien duidelijk verschillende deeltjes in water te onderscheiden en deze deeltjes kunnen reversibel in elkaar overgaan als gevolg van veranderingen in de pH. Tevens treedt het, in principe, pentadentaat ligand N4Py bij hoge pH op als tetradentaat ligand. Een verdere complicatie bij het analyseren van welke deeltjes in evenwicht aanwezig zijn in dit systeem is dat de in oplossing aanwezige complexen in zowel lage als hoge spintoestand voorkomen en dat deze snel in elkaar overgaan. Er wordt gepostuleerd dat het evenwicht tussen de singlet (lage spin) en quintet (hoge spin) spintoestanden in oplossingen van $\text{Fe}^{\text{II}}(\text{N4Py})$ in water ten grondslag ligt aan de activiteit van $\text{Fe}^{\text{II}}(\text{N4Py})$ complexen met betrekking tot het knippen van DNA met zuurstof ($^3\text{O}_2$) als oxidator, via de tijdelijke populatie van een triplet toestand.

In hoofdstuk 3 wordt de foto-geïnduceerde oxidatie van de complexen **1** en **2** in diverse oplosmiddelen (*i.e.* H_2O en MeOH) met zuurstof als uiteindelijk oxidant beschreven. Het is interessant dat beide complexen niet fotochemisch actief zijn in acetonitril, hetgeen weerspiegeld wordt door hun hoge positieve oxidatie potentiaal (1.1 V *vs.* SCE). Bestraling met UV en zichtbaar licht brengt de oxidatie van de ijzer(II) complexen naar de corresponderende ijzer(III) complexen teweeg waarbij

zuurstof als uiteindelijke oxidator optreedt. Bovendien is het primaire fotoproduct van complex **2** in methanol ook fotochemisch actief. De fotochemische versnelling van de oxidatiesnelheid van ijzer(II) complexen door zuurstof ($^3\text{O}_2$) door middel van ‘outer sphere’ elektronen-overdracht wordt gepostuleerd. Deze resultaten zijn van belang om de versnelling van het knippen van DNA door **1** met behulp van licht te kunnen begrijpen. Zoals hierboven vermeld wordt, valt het te verwachten dat er bij de overgang tussen de singlet en de hogere spintoestanden een triplet toestand als intermediair betrokken is, hetgeen elektronoverdracht naar $^3\text{O}_2$, de eerste stap in het knippen van DNA door **1** met zuurstof ($^3\text{O}_2$), mogelijk zou maken,

De fotochemie van ijzer complexen is verder uitgebreid naar de $\text{Fe}^{\text{IV}}=\text{O}$ complexen, hetgeen het onderwerp is van hoofdstuk 5. In scherpe tegenstelling tot **1** en **2**, laten de complexen **1a** en **1b** fotochemische activiteit zien in acetonitril. Deze complexen ondergaan foto-reductie naar de corresponderende $\text{Fe}^{\text{II}}-\text{CH}_3\text{CN}$ complexen als ze bestraald worden met UV en zichtbaar licht (400 nm). Gedeeltelijke foto-reductie naar Fe(III) en naar Fe(II) complexen door middel van bestraling met zichtbaar licht werd zelfs in vaste toestand (als dispersie in KCl) gezien. De fotochemische eigenschappen met betrekking tot het versnellen van de katalyse, bijvoorbeeld de hydroxylering van alkanen en de epoxidatie van alkenen, werd onderzocht. De fotochemische reductie van de $\text{Fe}^{\text{IV}}=\text{O}$ complexen verloopt sneller in de aanwezigheid van substraten zoals ethylbenzeen, benzylalcohol, fluoreen en indaan. Er zijn twee stappen betrokken in het fotochemische proces: de eerste stap is de overgang van $\text{Fe}^{\text{IV}}=\text{O}$ naar $\text{Fe}^{\text{III}}-\text{OH/OR}$ en dit wordt gevolgd door een tweede fotochemische stap waarin $\text{Fe}^{\text{III}}-\text{OR}$ omgezet wordt in $\text{Fe}^{\text{III}}-\text{CH}_3\text{CN}$. Dit laatste deeltje is een voldoende sterke oxidator om $\text{Fe}^{\text{III}}-\text{OH/OR}$ te oxideren naar $\text{Fe}^{\text{IV}}=\text{O}$ en hierbij wordt tevens het $\text{Fe}^{\text{II}}-\text{CH}_3\text{CN}$ complex gevormd. Ook het $\text{Fe}^{\text{III}}-\text{OMe}$ (**3**) complex is fotochemisch actief in acetonitril. Het $\text{Fe}^{\text{III}}-\text{OH}$ complex is daarentegen fotochemisch veel minder actief in acetonitril. Het primaire fotoproduct van de reductie van $\text{Fe}^{\text{IV}}=\text{O}$ is dus niet een $\text{Fe}^{\text{III}}-\text{OH}$ deeltje maar een $\text{Fe}^{\text{III}}-\text{OR}$ (R = alkyl) deeltje. Deze gegevens suggereren dat de C-H oxidatie plaatsvindt via een zogenaamd ‘rebound mechanisme’. De door bestraling verhoogde reactiviteit van $\text{Fe}^{\text{IV}}=\text{O}$ met betrekking tot het oxideren van substraten wordt toegeschreven aan de tijdelijke populatie van een S = 2 toestand.

Kennis van de redox eigenschappen van $\text{Fe}^{\text{IV}}=\text{O}$ deeltjes is van belang om de reactiviteit van deze deeltjes met betrekking tot de oxidatie van substraten te begrijpen. Helaas wordt het vaststellen van de $\text{Fe}^{\text{IV}}/\text{Fe}^{\text{III}}$ reductie potentiaal door

middel van cyclovoltammetrie belemmerd door de traagheid van de heterogene elektronenoverdracht die optreedt bij de reductie. Deze problemen worden uitgebreid behandeld in hoofdstuk 4. De chemische en electrochemische synthese en karakterisatie van een niet-heem $\text{Fe}^{\text{IV}}=\text{O}$ deeltje in water wordt beschreven. Complex **2** kan in water electrochemisch omgezet worden naar complex **2a**. In acetonitril werd de omzetting van complex **2** in **2a** alleen gezien in de aanwezigheid van water. Bovendien was de benodigde potentiaal om het $\text{Fe}^{\text{IV}}=\text{O}$ deeltje te genereren hoger in acetonitril (1.25 V vs. SCE) dan in water (0.85 V vs. SCE). Dit wordt toegeschreven aan de vereiste dat een hoge potentiaal nodig is om $\text{Fe}^{\text{II}}-\text{CH}_3\text{CN}$ te oxideren naar $\text{Fe}^{\text{III}}-\text{CH}_3\text{CN}$. Zodra $\text{Fe}^{\text{III}}-\text{CH}_3\text{CN}$ is gegenereerd vindt onmiddellijk ligand uitwisseling plaats resulterend in de vorming van $\text{Fe}^{\text{III}}-\text{OH}_2$ en dit deeltje wordt spontaan geoxideerd naar $\text{Fe}^{\text{IV}}=\text{O}$ onder dergelijk hoge toegepaste potentialen. De beschikbare data suggereert dat de oxidatie potentiaal van het het $\text{Fe}^{\text{IV}}/\text{Fe}^{\text{III}}$ redox-koppel in water rond ~ 0.85 V ligt en niet rond 0.41 V vs. SCE, zoals geconcludeerd wordt door Que *et al.*³ voor $[(\text{N4Py})\text{Fe}^{\text{IV}}(\text{O})]^{2+}$.

Het is voorgesteld dat een M-O-X (M = metaal, X = Cl, Br en I) adduct gegenereerd wordt in haloperoxidasen en dat dit deeltje verantwoordelijk is voor de halogenering van de substraten. Direct spectroscopisch bewijs voor het bestaan van dergelijke intermediaren is tot dusver nog niet gerapporteerd voor niet-heem systemen. In hoofdstuk 6 wordt de synthese en spectroscopische karakterisatie van een niet-heem $\text{Fe}^{\text{III}}-\text{OCl}$ deeltje behandeld. Door toevoeging van een oplossing van NaOCl in water aan zure oplossingen van **1** en **2** in water wordt bij kamertemperatuur het corresponderende $\text{Fe}^{\text{III}}-\text{OCl}$ deeltje gevormd. Na verloop van tijd verandert het $\text{Fe}^{\text{III}}-\text{OCl}$ deeltje in het corresponderende hoog-valente $\text{Fe}^{\text{IV}}=\text{O}$ deeltje. De karakterisatie van deze intermediaren is uitgevoerd met behulp van UV/Vis absorptie-, (resonantie) Raman- en EPR-spectroscopie en met cryo ESI-MS. Bij de toekenning van de $\text{Fe}^{\text{III}}-\text{O}$ en O-Cl pieken in de Ramanspectra werd gebruik gemaakt van labelling met broom ($\text{Fe}^{\text{III}}-\text{O}-\text{Br}$) en ^{18}O . Deze intermediaren reageren met styreen onder vorming van het corresponderende halohydrine en het diol.

In hoofdstuk 7 wordt de vorming en de spectroscopische karakterisatie van niet-porphyrine $\text{Ni}^{\text{II}}-\text{O}^{\cdot}$ en $\text{Ni}^{\text{II}}-\text{OCl}$ deeltjes beschreven. Bij kamertemperatuur zorgt toevoeging van een oplossing van NaOCl/ $\text{Ca}(\text{OCl})_2$ in water aan een oplossing van **4** in acetonitril voor de vorming van, vermoedelijk, een $\text{Ni}^{\text{II}}-\text{O}^{\cdot}$ deeltje. De toevoeging van Brønsted-zuren bevordert de vorming van dergelijke intermediaren

en versneld tevens de vorming ervan. De karakterisatie van deze intermediären werd uitgevoerd met behulp van UV/Vis absorptie-, (resonantie) Raman-, NMR- en EPR-spectroscopie en met cryo ESI-MS. Zowel de vorming als het verdwijnen van het Ni^{II}-O· deeltje werd gevolgd met behulp van (resonantie) Ramanspectroscopie. De formatie van het Ni^{II}-O· deeltje werd bevestigd met Ramanspectroscopie, wederom met behulp van labelling met broom en ¹⁸O, en bewijs voor de vorming van het Ni^{II}-OCl deeltje werd verkregen door middel van cryo ESI-MS. Er wordt gepostuleerd dat het eerstgenoemde deeltje C-H bindingen activeert en dat het laatstgenoemde deeltje verantwoordelijk is voor de insertie van een halogeen in het substraat. Voor zover mij bekend is is dit het eerste directe spectroscopische bewijs voor de betrokkenheid van Ni^{II}-O· en Ni^{II}-OCl deeltjes bij de gekatalyseerde halogenering van alifatische verbindingen.

De complexen **5** en **8** katalyseren de enantioselectieve vorming van C-C bindingen in water waarbij DNA de bron van de chiraliteit is. Complex **5** zorgt voor de hoogste enantioselectiviteit in DNA-gebaseerde asymmetrische katalyse, terwijl bij gebruik van complex **6** het tegenovergestelde enantiomeer verkregen wordt.⁴ In hoofdstuk 8 wordt de interactie van een serie Cu(II) polypyridyl complexen met DNA in de afwezigheid van substraten bestudeerd met behulp van diverse spectroscopische technieken. Voor het eerst is er gebruik gemaakt van UV resonantie Ramanspectroscopie om de interactie van Cu(II) complexen met DNA te onderzoeken. In de aanwezigheid van DNA laten de intercalerende complexen **7** en **8** duidelijke verschuivingen zien in hun Ramanspectra, terwijl dit niet het geval is bij complex **5**, dat in de groeve van het DNA bindt. Het spectroscopische bewijs duidt erop dat de complexen die de hoogste *ee* geven in de DNA-gebaseerde asymmetrische katalyse voornamelijk binden in de groeve van het DNA. Deze bevindingen geven belangrijke inzichten in de complexe structuur van deze DNA-gebaseerde katalysatoren en zijn een belangrijke basis voor het vaststellen van het effect van de substraten op de interactie tussen DNA en de complexen.

Tenslotte wordt in hoofdstuk 9 het werk dat in dit proefschrift beschreven wordt in perspectief geplaatst.

Referenties

-
- (1) J. Steinreiber, T. R. Ward, *Coord. Chem. Rev.*, **2008**, 252, 751-766; (b) V. Köhler, Y. M. Wilson, C. Lo, A. Sardo, T. R. Ward, *Current Opinion in Biotech.*, **2010**, 21, 744-752; (c) A. J. Boersma, R. P. Megens, B. L. Feringa, G. Roelfes, *Chem. Soc. Rev.*, **2010**, 39, 2083-2092.

-
- (2) (a) Q. Li, W. R. Browne, G. Roelfes, *Inorg. Chem.*, **2010**, *49*, 11009-11017; (b) Q. Li, W. R. Browne, G. Roelfes, *Inorg. Chem.*, **2010**, *50*, 8318-8325.
- (3) D. Wang, M. Zhang, P. Bühlmann, L. Que, Jr., *J. Am. Chem. Soc.*, **2010**, *132*, 7638-7644.
- (4) (a) G. Roelfes, A. J. Boersma, B. L. Feringa, *Chem. Commun.*, **2006**, 635-637; (b) A. J. Boersma, J. E. Klijn, B. L. Feringa, G. Roelfes, *J. Am. Chem. Soc.*, **2008**, *130*, 11783-11790.

సారాంశము

మన నిత్యజీవితంలో నిరంతరం ఎన్నో రసాయన చర్యలు మన ప్రమేయం లేకుండానే జరుగుతాయి. ఈ రసాయన చర్యలను ప్రకృతిలోని జీవనిర్మాణాలు అత్యంత సూక్ష్మనిర్మాణాలైన ఎంజైమ్ లు మరియు ప్రోటీనుల సహాయంతో ఎటువంటి హానికరమైన వ్యర్థ పదార్థాలను విడుదల చేయకుండా నిర్వహిస్తాయి. ఈ చర్యలలో వివిధ రకములైన లోహాలు పాల్గొంటాయి. ఉదాహరణకు కిరణజన్యసంయోగక్రియలో 'మాంగనీస్' అనే లోహము నీటిని, ఆక్సిజన్ మరియు హైడ్రోజన్ గా మార్చే శక్తిని విడుదల చేస్తుంది. చాలా మంది రసాయన శాస్త్రవేత్తలు ప్రకృతిలో ఇటువంటి జీవ నిర్మాణాల నుండి ప్రేరణపొంది, అటువంటి నిర్మాణాలను ప్రయోగశాలల్లో అనుకరించి, వాటి సహాయంతో రసాయన చర్యలను ఉత్పేరణ చెయ్యడానికి కలలు కంటారు. కానీ ఇప్పటివరకు ఏ ఒక్క జీవ నిర్మాణాన్ని ఖచ్చితంగా ప్రయోగశాలల్లో ఏ రసాయన శాస్త్రవేత్త అనుకరించలేదు. దీనికి గల కారణం ప్రకృతి ఉపయోగించే సంక్లిష్టమైన నిర్మాణాలను ప్రయోగశాలల్లో తయారు చెయ్యడం కష్టసాధ్యము. కానీ శాస్త్రవేత్తలు అనుకరించిన కొన్ని జీవాణుకరణులు మంచి ఉత్పేరకాలుగా పనిచేసి ఆశాజనకమైన ఫలితాలను ఇస్తున్నాయి. ఇటువంటి కొన్ని జీవాణుకరణులను (ఉత్పేరకాలను) ఈ పరిశోధనలో భాగంగా జరిగింది. ఉదాహరణకు ఐరన్ (II) - N4py అనే సంక్లిష్ట సమ్మేళనం ఐరన్(II) - బ్లయోమైసిన్ అనే జీవ నిర్మాణానికి ఒక అనుకరణ. ఈ జీవాణుకరణులను అధ్యయనం చేసిన ఫలితాల ఆధారంగా ఆ జీవ నిర్మాణాల పనితీరును అర్థం చేసుకోవడానికి కృషి చేస్తున్నాము.

ఈ పరిశోధనలో భాగంగా ఐరన్(II) రసాయన లక్షణాలను మరియు వాటి యొక్క స్పిన్ సమతాస్థితులను నీరు మరియు ఎసిటోనైట్రేట్ ద్రావణాలలో అధ్యయనం చేయడం జరిగింది. ఈ ఫలితాల ఆధారంగా ఈ సమ్మేళనాలు ఆక్సిజన్ తో చర్య నొందుటకు స్పిన్ సమతాస్థితి ఎంతో దోహదం చేస్తుందనీ నిర్ధారించితిమి. ఈ పుస్తకం లో వివరించిన మిగిలిన వివిధ పరిశోధనలకు ఎంతో ఉపయోగపడ్డాయి. ఈ ఐరన్ (II) రసాయన సమ్మేళనాలు ఆక్సిజన్ సమక్షంలో జన్యువులను నాశనం చెయ్యడం ద్వారా క్యాన్సర్ ను నిర్మూలిస్తాయి. ఈ చర్యను కాంతి సమక్షంలో మరింత సమర్థవంతంగా జరుగుతాయనీ ఈ మధ్యకాలంలో కనుగొనడం జరిగింది. దీనికి కారణం తెలుసుకునేందుకు ఈ ఐరన్(II) సమ్మేళనాల కాంతి రసాయన చర్యలను వేర్వేరు ద్రావణాలలో అధ్యయనం చేయడం జరిగింది. ఈ ఫలితాలను అధ్యయనం - 3 లో విశదీకరించాను. కాంతి సమక్షంలో ఈ ఐరన్(II) రసాయన సమ్మేళనాలు ఆక్సికరణం చెంది ఐరన్(II) సమ్మేళనాలను ఏర్పరుస్తాయి. ఈ కాంతి రసాయన చర్యలకు ముఖ్యకారణం నీటిలో గల స్పిన్ సమతాస్థితి కారణంగానే ఈ ఐరన్(II) సమ్మేళనాలు ఆక్సిజన్ తో సమర్థవంతంగా చర్య నొంది జన్యువులను నాశనం చెయ్యడం ద్వారా క్యాన్సర్ ను నిర్మూలిస్తాయనీ నిర్ధారించితిమి.

అదే విధంగా వివిధ చర్యామాధ్యమాలు ఉదాహరణకు ఐరన్(IV) సమ్మేళనాల యొక్క కాంతి రసాయన చర్యలను ఉపయోగించి అనేక రసాయన చర్యలను వేగవంతం చేయడం జరిగింది. అంతే కాకుండా మరికొన్ని చర్యామాధ్యమాలను బ్లీచింగ్ పౌడర్ నీటిని ఐరన్(II) సమ్మేళనాలకు కలపడం ద్వారా తయారు చేయడం జరిగింది. ఈ చర్యా మాధ్యమాలను ఉపయోగించి అనేక రసాయన చర్యలను సమర్థవంతంగా ఉత్పేరణ చేయడం జరిగింది.

చివరిగా ఈ పరిశోధనలను కాపర్(II) సమ్మేళనాలు మరియు జన్యువుల మధ్య చర్యను అవగాహన చేసుకోవడంపై దృష్టి పెట్టడం జరిగింది. ఈ పరిశోధనా ఫలితాలు, కష్టతరమైన ఉత్పేరణ చర్యలను అర్థం చేసుకోవడానికి ఉపయోగపడతాయని భావిస్తున్నాను. చివరిగా ఈ పరిశోధనలో నిర్మించిన ఎన్నో జీవాణుకరణులు మరియు వాటి నుండి తయారు చేసిన రసాయన మాధ్యమాల జీవనిర్మాణాలను మరియు వాటి యొక్క పనితీరును విజయవంతంగా పూర్తి చేయడం జరిగింది.

- అప్పారావు ద్రాక్షారపు.

Acknowledgements

Of course, plenty of people helped me, in a direct and indirect manner, to finish this thesis.

It's my great pleasure to thank Amma and Ayya on this memorable occasion. Your support and love, has truly helped me through the journey of my PhD, successfully. You are both strong and encouraged me to work in the Netherlands (a very foreign country). Amma and Ayya, I am not able express my gratitude in words, but I am sure you can understand.

Even before Wesley, I would like to thank Sir C. V. Raman for his great discovery of the Raman Effect. It is a great technique and makes my life a lot easier in characterising many of the intermediates discussed in this thesis.

I express my sincere gratitude to my promoter Prof. Wesley R. Browne. Being an Indian student with an Irish boss is great fun. In the beginning of my PhD half of the time I didn't understand what you were saying. Over time, things changed, you are so kind and spent a lot of time with me to teach me all the things... and killed my fear. It's my great pleasure to have worked with an enthusiastic boss (always, even at weekends and Christmas holidays) and I remember all of the discussions we have had over the last four years. You gave me a lot of freedom to explore my own field in chemistry. I thank you very much for everything that you have done for me. I am really excited to be the first to be awarded a doctorate by your words.

I thank Prof. Ben L. Feringa for all your comments, interest and enthusiasm towards chemistry. I really feel proud when I think that I am working with a truly great scientist. I thank Prof. Miquel Costas for our active collaboration in many of projects discussed in this thesis. I am amazed by your hospitality and time for friendly discussions on the projects, even though you have a very busy schedule. My next big thanks goes to Prof. Gerard Roelfes for the synthesis of such an amazing complex and our collaboration on many projects described in this thesis. Thank you very much for your critical comments on my research from which I have learned a lot.

I express my sincerest gratitude to Prof. Peter Comba, Prof. Bas De Bruin and Prof. Gerard Roelfes for the time you have spent on reading my thesis and for the many critical comments that have polished this thesis. If there are any mistakes remaining in this thesis, full responsibility is on me of course.

I am very grateful to Prof. J. J. McGarvey, for his valuable suggestions and fruitful discussions over the years. John, I learnt a lot from you personally and scientifically.

Dr. Hans de Boer, especially in our scientific discussions, I really appreciated your critical

comments. Thank you very much for the Samenvatting. Of course, we have fought so many times over the superiority of iron and manganese. I have to admit that manganese is a wonderful metal for catalysis as iron for spectroscopy. It's a choice. I also thank Dr. Ronald Hage for your enthusiasm and fruitful discussion over the last four years.

My thanks to Dr. Gabor Molnár, Prof. Gábor Speier, Dr. József Kaizer and Dr. Alvin Holder for their collaboration on different projects. For X-ray crystallographic analysis, I thank Dr. Edwin Otten. We almost got the crystal structure of an $\text{Fe}^{\text{IV}}=\text{O}$ complex, but it was unfortunate that the crystal quality was insufficient to generate good analysis. Drs. A. Meetsma is kindly acknowledged for the crystal structures of iron and Copper complexes in this thesis. I thank Peter Spanning and Bors István for collaborating on various research topics good luck in your own final push in completing your theses.

I had a great time in Girona, in our collaboration on nickel catalysis with the Costas's group. I am very grateful to all colleagues in Girona for making my scientific life easy, especially to Laura, Anna, Zoel, Marc and Isaac. I have to admit that Girona's paella is awesome. My non-scientific friends Albert and Josep are kindly acknowledged for the wonderful trip to Begur beach in Catalonia and dinners.

My special thanks goes to Jeffrey Bos for designing the cover page of this thesis. Davide and Eswar are kindly acknowledged as being my paranimfen. I thank Sandeep for correcting my thesis for little details (and for carrying on some of the projects enthusiastically).

Davide, I really appreciate our non-scientific discussions. Thank you very much for hosting me on a nice trip to Italy (in our colleagues view, a romantic trip). My sincere thanks also goes to your family members and especially to Laura. I hope you enjoyed our great trip to India with Oleksii, at ICORS conference as much as I did. We collaborated on many projects discussed in this thesis. I thank you for your great contribution on the DFT side. My dear physicist Oleksii, I appreciate your analogy of the end of the world in 2012. I enjoyed our conversations and parties we had together. I thank you very much for your suggestions (I have to say that I stand my promise).

My sincere thanks go to Ashoka, Rama Kumar and Vijay for their guidance and help to enter into research in the Netherlands. Especially, to Ashoka for being there to solve my problems at the beginning of my PhD, of course later as well. Adi annayya, I am amazed how you manage to know so many things, I strongly agree with Ashoka's saying 'Adipedia'.

Eswar, we started our PhD journey on the same day by taking the same flight. I really enjoyed your company and discussions about science, movies and psychology. I thank you

Acknowledgements

very much for your support and help in the last four years. I have so many memories to remember for the rest of my life. I hope Sandeep will survive in that house under your guidance. I express my gratitude to Vamsi (Saroo), Arun and Goutham for the happy moments we shared while cooking and eating spicy Indian food. Sandeep tammudu, I know you want to try everything, but please be a master in at least one thing. Pranov tammudu good luck in finding a PhD position in your desired field.

My housemates Rose, Gabor and Pascal are kindly acknowledged for their help at the start of my PhD.

I have to admit that I have a great team (Brownies) with Great Spirit towards science. I thank Hella in many regards, helping in translating Dutch letters and also in the lab. Pat, thank you very much for my first and last column in the Netherlands and I appreciate your help and nice smile always. Heloise (my philosopher), thank you very much for the philosophical analysis about life, which made me strong. Nikki (my English teacher 1), maybe only a thank you is not enough for the help you gave me. I thank Shaghayegh (cookie monster) and Jia Jia for many things that we shared (dinners and gossip). Peter thank you very much for hosting many international dinners. My sincere thanks goes to the Italians (Davide, Francesco and Federico), Peter, Emma (my English teacher 2), Duenpen, Sandeep and students (Pim, Maggie, Chris, Luuk, (mini)Hans, Tjalling, Kasper) for the pleasant working environment. Duenpen and Sandeep good luck for your research career.

‘Cricket’ there goes my weekends in the summer. I really wonder how we managed to turn the car parking area into a so called cricket ground. We even managed to play on a frozen lake. I would like to thank Ashoka, Adi, Shiva, Kiran, Eswar (never ending cricket lover for watching on TV as well), Arun, Sandeep (story teller), Goutham (Tinku baba, weather prediction guru), Pranov, Praneeth, Raju, Malli, Suresh (the chef), Milon, Afzal, Khayum (daactare) and many friends whom I have failed to mention. I also thank to GCC (Groningen Cricket Club) players (friends) for having Great Spirit for the game. I also thank my tennis companions Ganesh and Gopi, Arun and Eswar. Suresh tammudu, my special thanks to you for our squash war, and of course for the delicious Indian food also.

In adjacent labs I have many friends; it is very difficult to name all of you. It’s my great pleasure to work with colleagues from Ben’s group, Adri’s group, Gerard’s group, Sijbren’s group, Syzy’s group, Anna’s group, Katalin’s group and Edwin’s group.

I thank my friends in India, especially Santu (sabbu), Ajay (president) and Arun (kaaki) for their constant support. I thank Naveen (blaky) and Keerthi (gummadikaay) for our discussions on the current research. I thank Gani for all your technical and mental support. My sincere thanks goes to Vijay Kumar and Kukka Ashok for Telugu summary.

My dear Pavani, we met almost 5 years back over a phone call. After that, I never missed your call; simply I like to chat with you all the time. It's you who gave me a lot of courage, mental support and sharing all of my feelings during this period. I thank you very much for everything that you have done for me.

I am fully aware that my short term memory is so bad that I am not able to recall all the names of you (there are too many of you), who made my life pleasant and joyful during my PhD. If I forgot to mention your name please accept my apologies and, thank you very much for everything you did for me.