

University of Groningen

**Genome Sequence of *Lactobacillus saerimneri* 30a (Formerly *Lactobacillus* sp. Strain 30a), a Reference Lactic Acid Bacterium Strain Producing Biogenic Amines**

Romano, Andrea; Trip, Hein; Campbell-Sills, Hugo; Bouchez, Olivier; Sherman, David; Lolkema, Juke S.; Lucas, Patrick M.

*Published in:*  
Genome Announcements

*DOI:*  
[10.1128/genomea.00097-12](https://doi.org/10.1128/genomea.00097-12)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2013

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Romano, A., Trip, H., Campbell-Sills, H., Bouchez, O., Sherman, D., Lolkema, J. S., & Lucas, P. M. (2013). Genome Sequence of *Lactobacillus saerimneri* 30a (Formerly *Lactobacillus* sp. Strain 30a), a Reference Lactic Acid Bacterium Strain Producing Biogenic Amines. *Genome Announcements*, 1(1), 1-2. [0097]. <https://doi.org/10.1128/genomea.00097-12>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# Genome Sequence of *Lactobacillus saerimneri* 30a (Formerly *Lactobacillus* sp. Strain 30a), a Reference Lactic Acid Bacterium Strain Producing Biogenic Amines

Andrea Romano,<sup>a</sup> Hein Trip,<sup>b</sup> Hugo Campbell-Sills,<sup>c</sup> Olivier Bouchez,<sup>d,e</sup> David Sherman,<sup>f,g</sup> Juke S. Lolkema,<sup>b</sup> Patrick M. Lucas<sup>c</sup>

Research and Innovation Centre, Fondazione Edmund Mach, S. Michele all'Adige, Italy<sup>a</sup>; Molecular Microbiology, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Nijenborgh, Groningen, the Netherlands<sup>b</sup>; University of Bordeaux, ISVV, Unit of Research in Oenology (EA 4577), Villenave d'Ornon, France<sup>c</sup>; Institut National de la Recherche Agronomique (INRA), UMR444 Laboratoire de Génétique Cellulaire, INRA Auzeville, Castanet-Tolosan, France<sup>d</sup>; GeT-PlaGe, Genotoul, INRA Auzeville, Castanet-Tolosan, France<sup>e</sup>; Inria, Joint Project Team MAGNOME (INRIA, CNRS, University of Bordeaux), Talence, France<sup>f</sup>; UMR 5800 LaBRI (CNRS, University of Bordeaux), Talence, France<sup>g</sup>

***Lactobacillus* sp. strain 30a (*Lactobacillus saerimneri*) produces the biogenic amines histamine, putrescine, and cadaverine by decarboxylating their amino acid precursors. We report its draft genome sequence (1,634,278 bases, 42.6% G+C content) and the principal findings from its annotation, which might shed light onto the enzymatic machineries that are involved in its production of biogenic amines.**

Received 4 November 2012 Accepted 12 December 2012 Published 7 February 2013

**Citation** Romano A, Trip H, Campbell-Sills H, Bouchez O, Sherman D, Lolkema JS, Lucas PM. 2013. Genome sequence of *Lactobacillus saerimneri* 30a (formerly *Lactobacillus* sp. strain 30a), a reference lactic acid bacterium strain producing biogenic amines. *Genome Announc.* 1(1):e00097-12. doi:10.1128/genomeA.00097-12.

**Copyright** © 2013 Romano et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Patrick M. Lucas, [patrick.lucas@univ-bordeauxsegalen.fr](mailto:patrick.lucas@univ-bordeauxsegalen.fr).

*Lactobacillus* sp. strain 30a (ATCC 33222) was isolated from horse stomach in the early 1950s as the first strain of the genus *Lactobacillus* that produced biogenic amines (1). This is the only strain described thus far that forms all three biogenic amines—histamine, putrescine, and cadaverine—from histidine, ornithine, and lysine, respectively (1, 2). *Lactobacillus* sp. 30a has been used as a reference strain in many laboratories and in many studies relating to the production of biogenic amines by lactic acid bacteria (LAB). *Lactobacillus* sp. 30a carries a pyruvoyl-dependent histidine decarboxylase and a pyridoxal-phosphate-dependent ornithine decarboxylase that have been characterized extensively (3–10). Their genes have been identified (4), but their overall genomic environment remains unknown. *Lactobacillus* sp. 30a also possesses a pyridoxal-phosphate-dependent lysine decarboxylase (10), although this enzyme has not been identified in this strain or in any other LAB.

Here, we report the genome sequence of *Lactobacillus* sp. strain 30a, which was grown in deMan, Rogosa, and Sharpe (MRS) broth at 37°C. Genomic DNA was extracted using the Wizard genomic DNA purification kit (Promega). Whole-genome sequencing was performed at Genotoul (Toulouse, France) using single-read analysis of a fragment library with the 454 GS-FLX Titanium pyrosequencing system (Roche Diagnostics). A total of 213,826 reads were obtained and assembled using Newbler (454 Life Sciences), with an average coverage of 47-fold. Annotation of genes and rRNA was performed using the Prokaryotic Genome Annotation Pipeline (PGAAP) (11). tRNAs were identified with tRNAscan-SE (12).

The draft genome has 1,634,278 bases in 24 contigs (N<sub>50</sub>, 150,234) and a G+C content of 42.6%. It contains 1,519 predicted coding sequences, two 16S-23S-5S operons, and 55 tRNAs. No plasmids were detected in the sequenced DNA. *Lactobacillus* sp.

30a was attributed to the species *Lactobacillus saerimneri* on the basis of 16S rRNA gene analysis (>99% sequence identity with that of *L. saerimneri*).

The gene encoding the histidine decarboxylase is surrounded by the three genes typically encountered in the histamine-producing pathway in LAB (13). The ornithine decarboxylase gene stands alone, in contrast to in other LAB strains, where it is associated with an ornithine/putrescine exchanger gene (14, 15). *Lactobacillus* sp. 30a also contains a biosynthetic ornithine decarboxylase, which may account for its intracellular production of putrescine (15). A third gene that codes for a putative ornithine decarboxylase is also present and is associated with a predicted amino acid transporter; this likely represents the lysine decarboxylase pathway genes (unpublished results).

**Nucleotide sequence accession numbers.** This Whole Genome Shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [ANAG00000000](https://www.ncbi.nlm.nih.gov/nuccore/ANAG00000000). The version described in this article is the first version, [ANAG01000000](https://www.ncbi.nlm.nih.gov/nuccore/ANAG01000000).

## ACKNOWLEDGMENTS

This work was funded by the EU commission in the framework of the BIAMFOOD project (Controlling Biogenic Amines in Traditional Food Fermentations in Regional Europe) (project no. 211441).

## REFERENCES

1. Rodwell AW. 1953. The occurrence and distribution of amino-acid decarboxylases within the genus *Lactobacillus*. *J. Gen. Microbiol.* 8:224–232.
2. Coton M, Romano A, Spano G, Ziegler K, Vetrana C, Desmarais C, Lonvaud-Funel A, Lucas P, Coton E. 2010. Occurrence of biogenic amine-forming lactic acid bacteria in wine and cider. *Food Microbiol.* 27:1078–1085.
3. Copeland WC, Vanderslice P, Robertus JD. 1987. Expression and characterization of *Lactobacillus* 30a histidine decarboxylase in *Escherichia coli*. *Protein Eng.* 1:419–423.

4. Hackert ML, Carroll DW, Davidson L, Kim SO, Momany C, Vaaler GL, Zhang L. 1994. Sequence of ornithine decarboxylase from *Lactobacillus* sp. strain 30a. *J. Bacteriol.* 176:7391–7394.
5. Huynh QK, Snell EE. 1986. Histidine decarboxylase of *Lactobacillus* 30a. Hydroxylamine cleavage of the -seryl-seryl- bond at the activation site of prohistidine decarboxylase. *J. Biol. Chem.* 261:1521–1524.
6. Momany C, Ernst S, Ghosh R, Chang NL, Hackert ML. 1995. Crystallographic structure of a PLP-dependent ornithine decarboxylase from *Lactobacillus* 30a to 3.0 Å resolution. *J. Mol. Biol.* 252:643–655.
7. Momany C, Hackert ML. 1989. Crystallization and molecular symmetry of ornithine decarboxylase from *Lactobacillus* 30a. *J. Biol. Chem.* 264:4722–4724.
8. Parks EH, Ernst SR, Hamlin R, Xuong NH, Hackert ML. 1985. Structure determination of histidine decarboxylase from *Lactobacillus* 30a at 3.0 Å resolution. *J. Mol. Biol.* 182:455–465.
9. Rodwell AW. 1953. Factors affecting the activation of the ornithine apodecarboxylase of a strain of *Lactobacillus*. *J. Gen. Microbiol.* 8:238–247.
10. Rodwell AW. 1953. The histidine decarboxylase of a species of *Lactobacillus*; apparent dispensability of pyridoxal phosphate as coenzyme. *J. Gen. Microbiol.* 8:233–237.
11. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75.
12. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 25:955–964.
13. Lucas PM, Wolken WA, Claisse O, Lolkema JS, Lonvaud-Funel A. 2005. Histamine-producing pathway encoded on an unstable plasmid in *Lactobacillus hilgardii* 0006. *Appl. Environ. Microbiol.* 71:1417–1424.
14. Coton E, Mulder N, Coton M, Pochet S, Trip H, Lolkema JS. 2010. Origin of the putrescine-producing ability of the coagulase-negative bacterium *Staphylococcus epidermidis* 2015B. *Appl. Environ. Microbiol.* 76:5570–5576.
15. Romano A, Trip H, Lonvaud-Funel A, Lolkema JS, Lucas PM. 2012. Evidence of two functionally distinct ornithine decarboxylation systems in lactic acid bacteria. *Appl. Environ. Microbiol.* 78:1953–1961.