

University of Groningen

Quasinormal modes of scalar fields on small Reissner-Nordström- AdS5 black holes

Amado, Julián Barragán; Da Cunha, Bruno Carneiro; Pallante, Elisabetta

Published in:
Physical Review D

DOI:
[10.1103/PhysRevD.105.044028](https://doi.org/10.1103/PhysRevD.105.044028)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Amado, J. B., Da Cunha, B. C., & Pallante, E. (2022). Quasinormal modes of scalar fields on small Reissner-Nordström- AdS5 black holes. *Physical Review D*, 105(4), Article 044028. <https://doi.org/10.1103/PhysRevD.105.044028>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Quasinormal modes of scalar fields on small Reissner-Nordström-AdS₅ black holes

Julián Barragán Amado^{1,2,3,*}, Bruno Carneiro da Cunha^{2,†} and Elisabetta Pallante^{3,4,‡}

¹*Department of Mathematics, University of Sherbrooke, 2500, boulevard de l'Université, J1K 2R1 Sherbrooke, Quebec, Canada*

²*Departamento de Física, Universidade Federal de Pernambuco, 50670-901 Recife, Pernambuco, Brazil*

³*Van Swinderen Institute for Particle Physics and Gravity, University of Groningen, 9747 AG Groningen, Netherlands*

⁴*Nikhef, Science Park 105, 1098 XG Amsterdam, Netherlands*



(Received 12 November 2021; accepted 26 January 2022; published 14 February 2022)

We study the quasinormal modes (QNMs) of a charged scalar field on a Reissner-Nordström-anti-de Sitter (RN-AdS₅) black hole in the small radius limit by using the isomonodromic method. We also derive the low-temperature expansion of the fundamental QNM frequency. Finally, we provide numerical evidence that instabilities appear in the small radius limit for large values of the charge of the scalar field.

DOI: [10.1103/PhysRevD.105.044028](https://doi.org/10.1103/PhysRevD.105.044028)

I. INTRODUCTION

The energy extraction from a rotating uncharged black hole can be realized by the scattering of a wave of frequency ω and axial quantum number m that satisfies the superradiance condition $\omega < m\Omega$, where Ω is the angular velocity of the black hole. It turns out that the reflected wave is amplified with respect to the original incoming wave [1]. By imposing generic boundary conditions at the horizon and infinity, multiple amplifications and reflections of the scattered wave can occur and lead to superradiant instabilities.

An analogous superradiance condition for charged fields on charged black holes can be given by $\omega < e\Phi$, where e is the charge of the field and Φ the electrostatic potential difference between the horizon and the spatial infinity [2–4]. Then, by imposing an asymptotic AdS spacetime or a mirrorlike boundary condition, Reissner-Nordström black holes can develop such instabilities, as it has been shown in four dimensions [5,6] and $d \geq 4$ dimensions in the small outer horizon radius, r_+ , limit [7].

Black-hole superradiance is of relevance in numerous areas, from high-energy physics to astrophysics, and to fundamental questions of general relativity, such as the stability of a black hole solution and the existence of hairy

black hole configurations. In [8] it was suggested that a Reissner-Nordström-AdS₄ black hole coupled to a Higgs field Lagrangian can exhibit a spontaneous Abelian gauge symmetry breaking due to a scalar condensate near the horizon, thus implying that superradiant instabilities can occur for a certain range of values of the charges and the masses of the black hole and the scalar field. This observation may also suggest a holographic description of spontaneous symmetry breaking and superfluidity through the gauge-gravity duality—for comprehensive reviews see [9,10]. In five dimensions, the phase transition between the RN-AdS₅ black holes and hairy black hole solutions at the onset of superradiant instabilities has been explored in [11,12].

We study the occurrence of instabilities in RN-AdS₅ by computing the QNMs frequencies of a charged scalar field scattering on a RN-AdS₅ black hole. Due to the boundary conditions at the black hole horizon and at spatial infinity, the QNMs are generically complex frequencies, where the real part is related to the oscillation frequency and the imaginary part gives the inverse of the damping time. A superradiant mode thus corresponds to an eigenfrequency with a positive imaginary part.

In this context, we apply the isomonodromic method to recast the associated boundary value problem into an initial value problem for the corresponding isomonodromic tau function. Since the radial second-order ordinary differential equation possesses four regular singular points, see (18), one should expect that the initial conditions are given in terms of the Painlevé VI (PVI) tau function [13,14]. Interestingly, in recent years, the Painlevé transcendents have been employed to solve a variety of problems such as correlation functions in Ising and Ashkin-Teller model (see [15] and references therein), conformal maps [16],

*Jose.Julian.Barragan.Amado@USherbrooke.ca

†bruno.ccunha@ufpe.br

‡e.pallante@rug.nl

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

black hole physics [17,18] and the Rabi model in quantum optics [19]. The Painlevé transcendents implement the Riemann-Hilbert map relating the accessory parameters of the differential equations to the monodromy properties of its solutions, and [15,20] showed that their general expansion is expressible in terms of $c = 1$ conformal blocks. We should mention at this point that an alternative proposal directly using four-dimensional supersymmetric gauge theories in the Nekrasov-Shatashvili phase, corresponding to semiclassical conformal blocks, was developed in [21,22]. Also worthy of note is the effort to relate quantities of physical interest to monodromy data in [23–25], where the authors considered greybody factors, QNMs and Love numbers for different gravity backgrounds.

As argued in [14], in the small r_+ limit the first correction to the (fundamental) QNMs frequencies receives contributions from all the intermediate channels of the $c = 1$ Virasoro conformal blocks expansion. Nevertheless, these contributions can be resummed by using the generating function for the Catalan numbers, allowing us to obtain an asymptotic expression for the frequencies [14].

This manuscript is organized as follows. In Sec. II we study charged scalar perturbations of the RN-AdS₅ black hole, and recast the eigenvalue problem into a set of transcendental equations involving the PVI tau function. Section III is devoted to the analysis of the QNMs in the small r_+ limit. We discuss two different regimes for the temperature of the black hole: the high-temperature and the low-temperature limit for small black holes. Firstly, we show that at high temperature the QNMs are given by the zeros of the PVI tau function, and compute an asymptotic expression in the small r_+ limit for the $\ell = 0$ modes in Sec. III A. Secondly, in Sec. III B we provide an asymptotic expression for the fundamental QNM frequency in the low-temperature limit by applying the confluence limit of the PVI tau function. Section III C compares the asymptotic analytic expressions and the numerical solutions. We close in Sec. IV with some remarks on the implications of the obtained results. Finally, in Appendix we review the Nekrasov expansion of the PVI tau function.

II. SCALAR FIELDS IN REISSNER-NORDSTRÖM-AdS₅

The line element of the Reissner-Nordström-AdS₅ black hole is

$$ds^2 = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2d\Omega_3^2, \quad (1)$$

where $d\Omega_3^2$ is the metric of the unit three-sphere and the function $f(r)$ is given by

$$f(r) = 1 - \frac{M}{r^2} + \frac{Q^2}{r^4} + r^2 = \frac{\Delta_r}{r^4} = \frac{(r^2 - r_+^2)(r^2 - r_-^2)(r^2 - r_0^2)}{r^4}, \quad (2)$$

with the AdS radius $L = 1$ and the parameters M, Q are related to the black hole mass and charge, respectively. There is a correspondence between the roots of Δ_r and the Killing horizons of the black hole, whose radial positions will be parametrized by the largest real root r_+ , corresponding to the outer horizon. The roots corresponding to r_-^2 and r_0^2 can be written as

$$\begin{aligned} r_-^2 &= \frac{1}{2} \left(-1 - r_+^2 + \sqrt{(1 + r_+^2)^2 + \frac{4Q^2}{r_+^2}} \right), \\ r_0^2 &= \frac{1}{2} \left(-1 - r_+^2 - \sqrt{(1 + r_+^2)^2 + \frac{4Q^2}{r_+^2}} \right). \end{aligned} \quad (3)$$

The temperature at each horizon is

$$T_k = \frac{1}{4\pi} \frac{d}{dr} \left(\frac{\Delta_r}{r^4} \right) \Big|_{r=r_k} = \frac{1}{2\pi} \frac{(r_k^2 - r_i^2)(r_k^2 - r_j^2)}{r_k^3}, \quad i, j \neq k, \quad (4)$$

and specifically, at the outer horizon

$$T_+ = \frac{1}{2\pi} \frac{(r_+^2 - r_-^2)(r_+^2 - r_0^2)}{r_+^3}, \quad (5)$$

which by means of (3) reads

$$T_+ = \frac{1}{2\pi} \left[\frac{1}{r_+} - \frac{Q^2}{r_+^5} + 2r_+ \right]. \quad (6)$$

We define the critical charge Q_c as the maximal charge at extremality ($T_+ = 0$)

$$Q_c = r_+^2 \sqrt{1 + 2r_+^2}, \quad (7)$$

and require $Q \leq Q_c$ for a regular outer horizon. We then parametrize the accessible charge values as $Q = qQ_c$, where $q \leq 1$ is an extremality parameter for fixed $r_+ > 0$, and in terms of q the roots r_-^2 and r_0^2 in (3) read

$$\begin{aligned} r_-^2 &= \frac{1}{2} \left(-1 - r_+^2 + \sqrt{1 + 2(1 + 2q^2)r_+^2 + (1 + 8q^2)r_+^4} \right), \\ r_0^2 &= \frac{1}{2} \left(-1 - r_+^2 - \sqrt{1 + 2(1 + 2q^2)r_+^2 + (1 + 8q^2)r_+^4} \right). \end{aligned} \quad (8)$$

As we will see in Sec. III, one can think of small black holes near the critical charge ($q \lesssim 1$) both at low temperature and at high temperature.

The electromagnetic potential of the charged black hole reads

$$A_\mu dx^\mu = \left(-\frac{\sqrt{3}Q}{2r^2} + C \right) dt, \quad (9)$$

where the choice of the constant C is determined by the gauge choice [26]. We assume $C = 0$ for a vanishing potential at the spatial infinity, while it has also been shown in [6], for the case of asymptotically AdS, that the role of C is merely a shift in the real part of the quasinormal frequency ω , resulting in $\text{Re}(\omega) \rightarrow \text{Re}(\omega) + eC$.

We consider massive charged scalar field perturbations of the RN-AdS₅ black hole. The field equation is the Klein-Gordon equation

$$\frac{1}{\sqrt{-g}} D_\mu (\sqrt{-g} g^{\mu\nu} D_\nu) \Phi - \mu^2 \Phi = 0, \quad (10)$$

where $D_\mu = \nabla_\mu - ieA_\mu$, and e and μ are the field charge and mass, respectively. The scalar field solution of (10) can be decomposed by the Ansatz

$$\Phi_{\omega,\ell}(t, \theta, \varphi, \psi, r) = e^{-i\omega t} Y_\ell^{m_1, m_2}(\theta, \varphi, \psi) R_{\omega,\ell}(r), \quad (11)$$

where $Y_\ell^{m_1, m_2}(\theta, \varphi, \psi)$ are the scalar (spinless) spherical harmonics on the three-sphere, with eigenvalues determined by the equation

$$\Delta Y_\ell^{m_1, m_2}(\theta, \varphi, \psi) = -\ell(\ell + 2) Y_\ell^{m_1, m_2}(\theta, \varphi, \psi), \quad (12)$$

with ℓ the angular momentum quantum number and m_1 and m_2 are integers. By means of (12), Eq. (10) reduces to a second-order ordinary differential equation for the radial function $R_{\omega,\ell}(r)$ of the form

$$\left[\frac{1}{r} \frac{d}{dr} \left(\frac{\Delta_r}{r} \frac{d}{dr} \right) + \frac{r^6}{\Delta_r} \left(\omega - \frac{\sqrt{3} e Q}{2 r^2} \right)^2 - \mu^2 r^2 - \ell(\ell + 2) \right] R_{\omega,\ell}(r) = 0. \quad (13)$$

Equation (13) possesses four regular singular points, located at the roots of Δ_r in (2) and infinity. The characteristic exponents β_k, β_∞ are determined by the asymptotic behavior of the solution in (13) near the singular points

$$\beta_k = \pm \frac{1}{2} \theta_k, \quad \beta_\infty = \frac{1}{2} (2 \pm \theta_\infty), \quad k = +, -, 0, \quad (14)$$

where

$$\theta_k = \frac{i}{2\pi T_k} \left(\omega - \frac{\sqrt{3} e Q}{2 r_k^2} \right), \quad \theta_\infty = 2 - \Delta, \quad k = +, -, 0, \quad (15)$$

and θ_+ is the variation of the entropy δS of the black hole as it absorbs a quantum of frequency ω and electric charge e at the outer horizon

$$\theta_+ = \frac{i}{2\pi} \delta S. \quad (16)$$

When $\delta S < 0$, i.e., $\text{Im}\theta_+ < 0$, unstable modes can occur and the wave function grows exponentially with time, which means that the black hole is unstable. Furthermore, θ_∞ is related to Δ —the conformal dimension of the CFT primary field associated to the AdS₅ scalar, i.e., $\Delta(\Delta - 4) = \mu^2$, with μ the mass of the scalar field.

One can reduce (13) to the canonical Heun form by performing the following change of variables:

$$z = \frac{r^2 - r_-^2}{r^2 - r_0^2}, \quad (17a)$$

$$R_{\omega,\ell}(z) = z^{-\theta_-/2} (z - z_0)^{-\theta_+/2} (z - 1)^{\Delta/2} F(z), \quad (17b)$$

that leads to an equation for $F(z)$

$$\frac{d^2 F}{dz^2} + \left[\frac{1 - \theta_-}{z} + \frac{1 - \theta_+}{z - z_0} + \frac{\Delta - 1}{z - 1} \right] \frac{dF}{dz} + \left(\frac{\kappa_1 \kappa_2}{z(z - 1)} - \frac{z_0(z_0 - 1) K_0}{z(z - z_0)(z - 1)} \right) F(z) = 0, \quad (18)$$

with

$$\begin{aligned} \kappa_1 &= \frac{1}{2} (\theta_- + \theta_+ - \Delta - \theta_0), \\ \kappa_2 &= \frac{1}{2} (\theta_- + \theta_+ - \Delta + \theta_0), \end{aligned} \quad (19)$$

and the accessory parameters $\{z_0, K_0\}$ are

$$z_0 = \frac{r_+^2 - r_-^2}{r_+^2 - r_0^2}, \quad (20a)$$

$$\begin{aligned} 4z_0(z_0 - 1)K_0 &= -\frac{\ell(\ell + 2) + \Delta(\Delta - 4)r_-^2 - \omega^2}{(r_+^2 - r_0^2)} \\ &\quad - (z_0 - 1)[(\theta_- + \theta_+ - 1)^2 - \theta_0^2 - 1] \\ &\quad - z_0[2(\theta_+ - 1)(1 - \Delta) + (2 - \Delta)^2 - 2]. \end{aligned} \quad (20b)$$

The QNMs are solutions of (18) obeying specific boundary conditions, in the sense that there is only an incoming wave at the horizon r_+ ($z = z_0$) and regularity at spatial infinity ($z = 1$). For the radial equation in (13) with $\mu^2 > 0$, the conditions read as follows:

$$R_{\omega,\ell}(r) \sim \begin{cases} (r - r_+)^{-\frac{1}{2}\theta_+}, & r \rightarrow r_+, \\ r^{2-\Delta}, & r \rightarrow \infty, \end{cases} \quad (21)$$

and $F(z)$ in (18) is a regular function at the boundaries. The conditions in (21) will enforce the quantization of the (not necessarily real) frequencies ω , the QNMs frequencies.

Following the seminal works of the Kyoto school [27–29], one of the authors has described in [18], by

means of the method of isomonodromic deformations, the Riemann-Hilbert map between invariants of the representation of the monodromy group of the four-punctured Riemann sphere and Fuchsian 2×2 linear systems with four regular singular points. The monodromy group action on the solutions of the Fuchsian system is represented by matrices M_i describing the analytic continuation as one circles once around a singular point z_i of the system. The map gives a procedure to compute the accessory parameters $\{z_0, K_0\}$ of the Heun equation (18) in terms of the monodromy data $\{\vec{\theta}, \vec{\sigma}\} = \{\theta_1, \theta_2, \theta_3, \theta_4; \sigma_{12}, \sigma_{23}, \sigma_{13}\}$, where the single monodromy parameters are defined as follows:

$$2 \cos \pi \theta_i = \text{Tr} M_i, \quad k = 1, 2, 3, 4, \quad (22)$$

and the composite monodromy parameters σ_{ij} are given by

$$2 \cos \pi \sigma_{ij} = \text{Tr} M_i M_j, \quad i, j = 1, 2, 3, \quad (23)$$

where $M_i M_j$ represents the analytic continuation around two singular points. Following [28], only two of the three composite monodromy parameters $\sigma_{ij} = \{\sigma_{12}, \sigma_{23}, \sigma_{13}\}$ are independent, and we choose the pair $(\sigma_{12}, \sigma_{23})$ in the solution of the differential equation (18). The parameter σ_{12} from now on will be denoted by σ .

As shown in [18] the Riemann-Hilbert map between the accessory parameters of the Heun differential equation (18) and the monodromy parameters can be expressed implicitly via the PVI tau function

$$\tau(\vec{\theta}; \sigma, s; z_0) = 0, \quad (24a)$$

$$\frac{\partial}{\partial t} \log \tau(\vec{\theta}^-; \sigma - 1, s; t) \Big|_{t=z_0} - \frac{(\theta_2 - 1)\theta_3}{2(z_0 - 1)} - \frac{(\theta_2 - 1)\theta_1}{2z_0} = K_0. \quad (24b)$$

where $\vec{\theta}^- = \{\theta_1, \theta_2 - 1, \theta_3, \theta_4 + 1\}$ and s is a function of the monodromy parameters defined by (A5).

Repeating the analysis of [14], in order to solve the first condition (24a), one can invert the series expansion of the PVI tau function (A7), for $t = z_0$ sufficiently close to zero, to write

$$\chi(\sigma; z_0) = \kappa z_0^\sigma, \quad (25)$$

where κ is given in terms of the monodromy data by (A8). From the structure of the tau function expansion, we can see that χ is analytic in z_0 , and its expansion can be computed recursively from (A7), yielding

$$\chi(\sigma; z_0) = \left[\frac{((\theta_2 + \sigma)^2 - \theta_1^2)((\theta_3 + \sigma)^2 - \theta_4^2)}{16\sigma^2(\sigma - 1)^2} z_0 \right] \left(1 + (1 - \sigma) \frac{(\theta_1^2 - \theta_2^2)(\theta_3^2 - \theta_4^2) + \sigma^2(\sigma - 2)^2}{2\sigma^2(\sigma - 2)^2} z_0 + \mathcal{O}(z_0^2) \right). \quad (26)$$

We can then determine the logarithmic derivative of the tau function in (24b) using the expansion (A7) and write the expansion of the accessory parameter K_0 in terms of the monodromy data and z_0

$$\begin{aligned} 4z_0 K_0 = & (\sigma - 1)^2 - (\theta_1 + \theta_2 - 1)^2 + \left[2(\theta_3 - 1)(\theta_2 - 1) + \frac{((\sigma - 1)^2 + \theta_2^2 - \theta_1^2 - 1)((\sigma - 1)^2 + \theta_3^2 - \theta_4^2 - 1)}{2\sigma(\sigma - 2)} \right] z_0 \\ & + \left[\frac{13}{32}\sigma(\sigma - 2) + 2(\theta_3 - 1)(\theta_2 - 1) - \frac{1}{32}(5 + 14(\theta_1^2 + \theta_4^2) - 18(\theta_2^2 + \theta_3^2)) + \frac{(\theta_1^2 - \theta_2^2)^2(\theta_3^2 - \theta_4^2)^2}{64} \left(\frac{1}{\sigma^3} - \frac{1}{(\sigma - 2)^3} \right) \right. \\ & \left. - \frac{((\theta_1^2 - \theta_2^2)(\theta_3^2 - \theta_4^2) + 8)^2 - 2(\theta_1^2 + \theta_2^2)(\theta_3^2 - \theta_4^2)^2 - 2(\theta_1^2 - \theta_2^2)^2(\theta_3^2 + \theta_4^2) - 64}{32\sigma(\sigma - 2)} \right. \\ & \left. + \frac{((\theta_1 - 1)^2 - \theta_2^2)((\theta_1 + 1)^2 - \theta_2^2)((\theta_3 - 1)^2 - \theta_4^2)((\theta_3 + 1)^2 - \theta_4^2)}{32(\sigma + 1)(\sigma - 3)} \right] z_0^2 + \mathcal{O}(z_0^3). \quad (27) \end{aligned}$$

Equations (26) and (27) were derived under the assumption that σ lies in the strip $0 < \text{Re} \sigma < 1$. Formulas for $\text{Re} \sigma < 0$ can be obtained by the replacement $\sigma \rightarrow -\sigma$. The expansion for K_0 is symmetric under the replacement $\sigma \rightarrow 2 - \sigma$ and has the generic structure

$$4z_0 K_0 = k_0 + k_1 z_0 + k_2 z_0^2 + \dots + k_n z_0^n + \dots, \quad (28)$$

where k_n is a rational function of the monodromy parameters, the numerator is a polynomial in the monodromy parameters θ_i and σ , and the denominator is a polynomial of σ alone. The coefficient k_n has single poles at integer $\sigma = 3, \dots, n + 1$, as well as poles of order up to $2n - 1$ at $\sigma = 2$. The leading behavior of k_n near $\sigma = 2$ is

$$k_n = (-1)^{n-1} C_{n-1} \frac{(\theta_2^2 - \theta_1^2)^n (\theta_3^2 - \theta_4^2)^n}{4^{2n-1} (\sigma - 2)^{2n-1}} + \dots, \quad n \geq 1, \quad (29)$$

where C_n is the n th Catalan number. A structure analogous to (28) exists for $\chi(\sigma; z_0)$ in (26), and the reader is referred to [14] for further details. Recently, similar analytic expansions involving irregular conformal blocks have been analyzed in [30,31].

For our particular problem of the radial system, the single monodromy parameters θ_i can be read directly from (18) as follows:

$$\theta_1 = \theta_-, \quad \theta_2 = \theta_+, \quad \theta_3 = 2 - \Delta, \quad \theta_4 = \theta_0, \quad (30)$$

and then the conditions (24a) and (24b) define for us the monodromy parameters σ and κ (or, rather, s) as functions of the modulus z_0 and the accessory parameter K_0 . As it

was shown in [17], physically relevant quantities such as scattering coefficients can be expressed in terms of σ and s .

For the QNM problem at hand, we have to enforce the boundary conditions in (21). As shown in [14], these can be encoded with the requirement that the composite monodromy matrix around the two singular points r_+ and ∞ is triangular. In turn, this implies that the composite monodromy parameter σ_{23} satisfies

$$\sigma_{23} = \theta_2 + \theta_3 + 2n, \quad n \in \mathbb{Z}, \quad (31)$$

which will be referred to as the quantization condition for the radial function. With this restriction, we can solve for the s parameter in (A5) to yield

$$s = \frac{\sin \frac{\pi}{2}(\theta_2 + \theta_1 + \sigma) \sin \frac{\pi}{2}(\theta_2 - \theta_1 + \sigma) \sin \frac{\pi}{2}(\theta_3 + \theta_4 + \sigma) \sin \frac{\pi}{2}(\theta_3 - \theta_4 + \sigma)}{\sin \frac{\pi}{2}(\theta_2 + \theta_1 - \sigma) \sin \frac{\pi}{2}(\theta_2 - \theta_1 - \sigma) \sin \frac{\pi}{2}(\theta_3 + \theta_4 - \sigma) \sin \frac{\pi}{2}(\theta_3 - \theta_4 - \sigma)}, \quad (32)$$

which, given the relation between κ and s in (A8), can then be substituted in (25) to give

$$\frac{\Gamma^2(1 - \sigma) \Gamma(1 + \frac{1}{2}(\theta_2 + \theta_1 + \sigma)) \Gamma(1 + \frac{1}{2}(\theta_2 - \theta_1 + \sigma)) \Gamma(1 + \frac{1}{2}(\theta_3 + \theta_4 + \sigma)) \Gamma(1 + \frac{1}{2}(\theta_3 - \theta_4 + \sigma))}{\Gamma^2(1 + \sigma) \Gamma(1 + \frac{1}{2}(\theta_2 + \theta_1 - \sigma)) \Gamma(1 + \frac{1}{2}(\theta_2 - \theta_1 - \sigma)) \Gamma(1 + \frac{1}{2}(\theta_3 + \theta_4 - \sigma)) \Gamma(1 + \frac{1}{2}(\theta_3 - \theta_4 - \sigma))} s z_0^\sigma = \chi(\sigma; z_0). \quad (33)$$

Equations (27) and (33) can be interpreted as an overdetermined system for σ in terms of the parameters of the differential equation, namely z_0 in (20a), K_0 in (20b) and $\vec{\theta}$ in (30), which will only admit solutions for specific (discrete) values of ω . As we will see in the following, these equations can be solved perturbatively in the small r_+ limit to provide us with the QNMs frequencies.

Similar methods have been employed to study scalar and vector perturbations in Kerr-AdS₅ backgrounds [14,32], and more recently in the near-extremal limit [33]. In four dimensions, rotating black holes in asymptotically flat and de-Sitter spacetimes were analyzed in [30,31,34].

III. SMALL RN-AdS₅ BLACK HOLES VIA ISOMONODROMY

We proceed to study the small r_+ limit of RN-AdS₅ black holes. By means of (8), the accessory parameter z_0 in (20a) can be rewritten in terms of r_+ and q only

$$z_0 = \frac{r_+^2 - r_-^2}{r_+^2 - r_0^2} = \frac{1 + 3r_+^2 - \sqrt{1 + 2(1 + 2q^2)r_+^2 + (1 + 8q^2)r_+^4}}{1 + 3r_+^2 + \sqrt{1 + 2(1 + 2q^2)r_+^2 + (1 + 8q^2)r_+^4}}. \quad (34)$$

For small r_+ and arbitrary $0 \leq q \leq 1$

$$z_0 = (1 - q^2)r_+^2 + \mathcal{O}(r_+^4), \quad (35)$$

so that $r_+ \rightarrow 0$ implies $z_0 \rightarrow 0$ in (35). Hence, the analytic expansions for $\chi(\sigma; z_0)$ in (26) and the accessory parameter K_0 in (27) for small z_0 apply in the small r_+ limit.

By means of (8) and $q = Q/Q_c$, the temperature (5) reduces to

$$\begin{aligned} 2\pi T_+ &= \frac{1 - q^2}{r_+} + 2(1 - q^2)r_+ \\ &= \frac{\epsilon(2 - \epsilon)}{r_+} + 2\epsilon(2 - \epsilon)r_+, \end{aligned} \quad (36)$$

where in the second equality we introduced the extremality parameter $\epsilon = 1 - q$. For small black holes $r_+ \ll 1$ and ϵ fixed, the terms of order $1/r_+$ in the last equality of (36) dominate as $r_+ \rightarrow 0$.

Equation (36) also suggests that we can identify two different regimes, depending on the behavior of ϵ :

- (1) The first regime corresponds to $0 < \epsilon < 1$ fixed and small black hole, $r_+ \ll 1$, hence high temperature $T_+ \sim 1/r_+$. Specifically, we should set $r_+ \ll \epsilon < 1$ in our expansions. This regime is investigated in Sec. III A.
- (2) The second regime describes the scaling limit for $\epsilon \ll r_+ \ll 1$, as both ϵ and r_+ vary. This is the low-temperature regime and it is explored in Sec. III B.

A. QNMs frequencies at high temperature

The single monodromy parameters θ_\pm and θ_0 can be expanded for small r_+ as follows:

$$\theta_- = -i \frac{q^2(2q\omega - \sqrt{3}e)}{2(1-q^2)} r_+ + \mathcal{O}(r_+^3), \quad \theta_- = -i\vartheta_- r_+, \quad \theta_+ = i\vartheta_+ r_+, \quad (38)$$

$$\theta_+ = i \frac{(2\omega - \sqrt{3}eq)}{2(1-q^2)} r_+ + \mathcal{O}(r_+^3),$$

$$\theta_0 = \omega + \frac{1}{2}(\sqrt{3}eq - 3(1+q^2)\omega)r_+^2 + \mathcal{O}(r_+^4), \quad (37)$$

with

and the coefficients ϑ_{\pm} are finite for $q < 1$ fixed, such that θ_{\pm} approach zero as $r_+ \rightarrow 0$. The expansion (27) for $z_0 K_0$ allows us to compute the composite monodromy parameter ($\sigma = \sigma_{\ell}$) by inverting the series in (27). For $\ell \geq 2$ the result is

$$\sigma_{\ell} = \sum_{n=0}^{\infty} c_n r_+^n = \ell + 2 - \frac{((1+q^2)(3\omega^2 + 3\ell(\ell+2) - \Delta(\Delta-4)) - 2\sqrt{3}eq\omega)}{4(\ell+1)} r_+^2 + \mathcal{O}(r_+^4), \quad \ell \geq 2. \quad (39)$$

For $\ell = 0$ and $\ell = 1$, the pole structure of k_n described in (28) renders the derivation of the coefficients c_n more complicated, and we will deal separately with the $\ell = 0$ case below. At any rate, σ_{ℓ} always admits the generic structure

$$\sigma_{\ell} = \ell + 2 - \nu_{\ell} r_+^2 + \mathcal{O}(r_+^4). \quad (40)$$

By substituting (39) in (26), one obtains an expansion for small r_+ for $\chi(\sigma_{\ell}; z_0)$

$$\chi_{\ell} \equiv \chi(\sigma_{\ell}; z_0) = -(1-q^2) \frac{\omega^2 - (\Delta - \ell - 4)^2}{16(\ell+1)^2} \left(1 + i \frac{2\vartheta_+}{(\ell+2)} r_+ \right) r_+^2 + \mathcal{O}(r_+^4), \quad \ell \geq 2, \quad (41)$$

where $\ell \geq 2$ avoids the poles in (26).

We now focus on the $\ell = 0$ case, and defer the analysis of higher ℓ modes to future work. The $\ell = 0$ case is peculiar, since the leading behavior of $\sigma_0 - 2$ in (40) is of order r_+^2 and the scaling limit of θ_{\pm} in (38) suggests that the accessory parameter K_0 receives contributions from all orders in z_0 in the small r_+ limit. The left-hand side of (27) is given by (20b) for $\ell = 0$, and for small r_+ we have

$$4z_0 K_0 = -2i(\vartheta_- - \vartheta_+)r_+ + ((\vartheta_- - \vartheta_+)^2 + \Delta^2 - 2(1+q^2)\Delta + \omega(\sqrt{3}eq - 3(1+q^2)\omega) + (1+2q^2)\omega^2)r_+^2 + \mathcal{O}(r_+^3), \quad (42)$$

whereas the right-hand side of (27) for $\ell = 0$ is

$$4z_0 K_0 = -2i(\vartheta_- - \vartheta_+)r_+ + \left[(\vartheta_- - \vartheta_+)^2 - 2\nu_0 - \frac{1}{2}(1-q^2)(\omega^2 - \Delta^2) \right. \\ \left. + \frac{(1-q^2)(\omega^2 - (\Delta-2)^2)(\vartheta_-^2 - \vartheta_+^2)}{4\nu_0} + \frac{(1-q^2)^2(\omega^2 - (\Delta-2)^2)^2(\vartheta_-^2 - \vartheta_+^2)^2}{64\nu_0^3} \right. \\ \left. + \frac{(1-q^2)^3(\omega^2 - (\Delta-2)^2)^3(\vartheta_-^2 - \vartheta_+^2)^3}{512\nu_0^5} + \frac{5(1-q^2)^4(\omega^2 - (\Delta-2)^2)^4(\vartheta_-^2 - \vartheta_+^2)^4}{16384\nu_0^7} + \dots \right] r_+^2 + \mathcal{O}(r_+^3), \quad (43)$$

where $\nu_0 = \nu_{\ell}(\ell = 0)$ in (40) and the dots are higher orders in z_0 that contribute to the r_+^2 term. It was shown in [14] that all the contributions from the conformal blocks expansion can be resummed using the generating function for the Catalan numbers to compute the first correction to σ_0 of order r_+^2 . Defining

$$X = \frac{(1-q^2)(\omega^2 - (\Delta-2)^2)(\vartheta_-^2 - \vartheta_+^2)}{16\nu_0^2}, \quad (44)$$

and resumming all the terms in the series will lead to an equation for the coefficient ν_0

$$\nu_0 \sqrt{1-4X} = \frac{1}{2} \omega (3(1+q^2)\omega - \sqrt{3}eq) - \frac{1}{4} (1+q^2)(3\omega^2 - \Delta(\Delta-4)) + \mathcal{O}(r_+^2). \quad (45)$$

Then, ν_0 in (45) admits the following small r_+ expansion:

$$\begin{aligned} \nu_0 = & \frac{1}{4} [(1+q^2)^2((3\omega^2 - \Delta(\Delta-4))^2 - 4\omega^2(\omega^2 - (\Delta-2)^2)) + 4q^2\omega^2(\omega^2 - (\Delta-2)^2) \\ & + 3e^2q^2(3\omega^2 + (\Delta-2)^2) - 8\sqrt{3}(1+q^2)eq\omega(\omega^2 + 2)]^{1/2} + \mathcal{O}(r_+^2). \end{aligned} \quad (46)$$

Hence, by means of (44) and resumming all the contributions of order r_+^2 , we get for χ_0 in (41)

$$\chi_0 = -(1-q^2) \frac{(\omega^2 - (\Delta-4)^2)}{64} r_+^2 (1 + i\vartheta_+ r_+) \left(1 + \sqrt{1 - \frac{(1-q^2)(\omega^2 - (\Delta-2)^2)(\vartheta_-^2 - \vartheta_+^2)}{4\nu_0^2}} \right)^2 + \mathcal{O}(r_+^4), \quad (47)$$

where

$$\vartheta_- = \frac{q^2(2q\omega - \sqrt{3}e)}{2(1-q^2)}, \quad \vartheta_+ = \frac{(2\omega - \sqrt{3}eq)}{2(1-q^2)}. \quad (48)$$

The knowledge of the parameter s in (33) is now sufficient to determine the QNMs frequencies. By expanding for small r_+ and employing (47) in (33), we obtain

$$s = \Sigma_0 \left(1 + \frac{i2\nu_0\vartheta_+ r_+}{(\vartheta_+^2 - \vartheta_-^2)} \right) + \mathcal{O}(r_+^2, r_+^2 \log r_+), \quad (49)$$

with

$$\Sigma_0 = \nu_0^2 \frac{\Gamma(\frac{1}{2}(2-\omega-\Delta))\Gamma(\frac{1}{2}(2+\omega-\Delta))}{\Gamma(\frac{1}{2}(6-\omega-\Delta))\Gamma(\frac{1}{2}(6+\omega-\Delta))} \frac{(\omega^2 - (\Delta-4)^2)}{4(1-q^2)(\vartheta_+^2 - \vartheta_-^2)} \left(1 + \sqrt{1 + \frac{(1-q^2)(\omega^2 - (\Delta-2)^2)(\vartheta_+^2 - \vartheta_-^2)}{4\nu_0^2}} \right)^2. \quad (50)$$

Equation (49) is a nonanalytic expansion in r_+ due to terms $\mathcal{O}(r_+^2 \log r_+)$, which are originated by the term $z_0^{\nu_0}$ in (33). However, the limit as $r_+ \rightarrow 0$ is finite.

We now parametrize the QNMs frequencies $\omega_{n,\ell}$ with $n \geq 0$ and $\ell = 0$ in terms of a perturbation to the normal modes in empty AdS₅

$$\omega_{n,0} = 2n + \Delta + \eta_{n,0} r_+^2, \quad (51)$$

under the assumption that $\eta_{n,0}$ has a finite limit as $r_+ \rightarrow 0$. Then, by replacing (51) in (37) for ω , one obtains

$$\begin{aligned} \theta_- = & -i \frac{q^2(2q(2n+\Delta) - \sqrt{3}e)}{2(1-q^2)} r_+ + \mathcal{O}(r_+^3), \\ \theta_+ = & i \frac{(2(2n+\Delta) - \sqrt{3}eq)}{2(1-q^2)} r_+ + \mathcal{O}(r_+^3), \\ \theta_0 = & 2n + \Delta + \beta_{n,0} r_+^2 + \mathcal{O}(r_+^4), \end{aligned} \quad (52)$$

where $\beta_{n,0}$ encodes the correction $\eta_{n,0}$ to the QNMs as follows:

$$\beta_{n,0} = \eta_{n,0} + \frac{1}{2} (\sqrt{3}eq - 3(1+q^2)(2n+\Delta)). \quad (53)$$

On the other hand, $\beta_{n,0}$ can also be computed by equating (32) to (49), where $\Sigma_{n,0} \equiv \Sigma_0(\omega = \omega_{n,0})$ and the radial monodromies are given by (52), and by expanding for small r_+ we have

$$\begin{aligned} \beta_{n,0} = & \nu_{n,0} \frac{1 + \Sigma_{n,0}}{1 - \Sigma_{n,0}} + 4i \frac{\vartheta_+ \nu_{n,0}^2}{(\vartheta_-^2 - \vartheta_+^2)} \frac{\Sigma_{n,0}}{(\Sigma_{n,0} - 1)^2} r_+ \\ & + \mathcal{O}(r_+^2, r_+^2 \log r_+), \end{aligned} \quad (54)$$

with $\nu_{n,0} \equiv \nu_0(\omega = \omega_{n,0})$ in (46). By means of (54), Eq. (53) yields for $\eta_{n,0}$

$$\begin{aligned} \eta_{n,0} = & \frac{1}{2} (3(1+q^2)(2n+\Delta) - \sqrt{3}eq - Z_{n,0}^{1/2} \\ & - i(n+1)(\Delta+n-1)(2(2n+\Delta) - \sqrt{3}eq)r_+ \\ & + \mathcal{O}(r_+^2, r_+^2 \log r_+), \end{aligned} \quad (55)$$

where

$$Z_{n,0} = \left(\frac{1}{2}(1+q^2)(3(2n+\Delta)^2 - \Delta(\Delta-4)) - \sqrt{3}eq(2n+\Delta) \right)^2. \quad (56)$$

Finally, by employing (55), Eq. (51) admits the small r_+ expansion

$$\begin{aligned} \omega_{n,0} = 2n + \Delta - \frac{1}{2}(Z_{n,0}^{1/2} - 3(1+q^2)(2n+\Delta) + \sqrt{3}eq)r_+^2 - i\frac{1}{2}(n+1)(\Delta+n-1)(2(2n+\Delta) - \sqrt{3}eq)r_+^3 \\ + \mathcal{O}(r_+^4, r_+^4 \log r_+). \end{aligned} \quad (57)$$

In fact, Eq. (57) is different from the analytical result in [7] due to the correction to the real part of $\omega_{n,0}$ of order r_+^2 . The asymptotic expansion (57) for $n=0$ reproduces the numerical results for the real part of $\omega_{0,0}$ of Table I in [7]. In addition, the imaginary part of $\omega_{n,0}$ for $n=0$ in (57) for specific values of e , Δ , q and r_+ falls in between the numerical and analytical values given in Tables II, IV, V for the five-dimensional case in [7]. Importantly, for $q > \frac{2(2n+\Delta)}{\sqrt{3}e}$, the imaginary part in (57) becomes positive, and the superradiance condition is satisfied.

Figure 1 shows our numerical solutions for the imaginary part of $\omega_{0,0}$ and illustrates explicitly that for $q = \{0.5775, 0.57775, 0.578, 0.57825\}$ the superradiance condition is satisfied in a finite interval of r_+ , and the stability is eventually restored as the black hole radius increases. As $q \rightarrow 1$, the superradiance condition is always satisfied in the small r_+ limit. The QNMs frequencies shown in Fig. 1 have been obtained through the numerical implementation of the initial conditions for the PVI tau function using an arbitrary-precision library in Julia programming language [35]. For the fast convergence of the numerical results, we have introduced the Fredholm determinant formulation of the PVI tau function, instead of its combinatorial




FIG. 1. Numerical results for the imaginary part of the fundamental QNM $\omega_{0,0}$ as a function of r_+ for increasing values of q in the interval $[0.1, 0.9]$. The charge and the scaling dimension of the scalar field are $e = 8$ and $\Delta = 4$, respectively.

description. We refer to previous works [13,14,16] for a more detailed discussion of the numerical methods.

Finally, in terms of $\epsilon = 1 - q$ and for $n=0$, (57) reads

$$\begin{aligned} \omega_{0,0} = \Delta - \frac{1}{2}(\Delta-1)(\Delta(\epsilon^2 - 2\epsilon + 2) - \sqrt{3}e(1-\epsilon))r_+^2 \\ - i\frac{1}{2}(\Delta-1)(2\Delta - \sqrt{3}e(1-\epsilon))r_+^3 \\ + \mathcal{O}(r_+^4, r_+^4 \log r_+), \end{aligned} \quad (58)$$

where $e < 2(\Delta+2)/\sqrt{3}$. Equation (58) should provide a good description of the high-temperature regime $r_+ \ll \epsilon < 1$. We perform a detailed comparison of (58) with the numerical solution in Sec. III C (Fig. 3).

B. The low-temperature limit of the fundamental QNM

The description of the low-temperature regime requires a more careful approach since the single monodromy parameters (37) may diverge *a priori*. This issue is solved by means of a double series expansion for small r_+ and ϵ , and by considering the scaling regime $\epsilon \ll r_+ \ll 1$, which indeed corresponds to the low-temperature regime of small black holes, i.e., near-extremal small black holes. In the low-temperature limit (36) is given by

$$2\pi T_+ = \frac{2\epsilon}{r_+} + \mathcal{O}\left(\frac{\epsilon^2}{r_+}, \epsilon r_+\right), \quad (59)$$

and the single monodromies in terms of ϵ and r_+ are

$$\begin{aligned} \theta_+ = i\frac{(2\omega - \sqrt{3}e)r_+}{4} \frac{1}{\epsilon} + i\frac{1}{16}(2\omega + \sqrt{3}e)(2+\epsilon)r_+ \\ + \mathcal{O}\left(\epsilon^2 r_+, \frac{r_+^3}{\epsilon}, r_+^3, \epsilon r_+^3\right), \\ \theta_- = -i\frac{(2\omega - \sqrt{3}e)r_+}{4} \frac{1}{\epsilon} + i\frac{1}{8}(10\omega - 3\sqrt{3}e)r_+ \\ - i\frac{1}{16}(14\omega - \sqrt{3}e)\epsilon r_+ + \mathcal{O}\left(\epsilon^2 r_+, \frac{r_+^3}{\epsilon}, r_+^3, \epsilon r_+^3\right), \\ \theta_0 = \omega - \frac{1}{2}(6\omega - \sqrt{3}e)(1-\epsilon)r_+^2 + \mathcal{O}(\epsilon^2 r_+^2, r_+^4, \epsilon r_+^4), \end{aligned} \quad (60)$$

where θ_{\pm} diverge as $\epsilon \rightarrow 0$, as it is also evident from the pole at $q^2 = 1$ in (37).

The initial conditions for the confluent limit of the PVI tau function were derived in [33]. The procedure is further analogous to the rotating case in [33]. We need to compute the correction of order r_+^2 , as well as the first nontrivial correction in ϵ for $\omega_{0,0} = \Delta + \eta_{0,0}r_+^2$ in (51) and

$$\sigma_{0,0} \equiv \sigma_0(\omega = \omega_{0,0}) = 2 - \nu_{0,0}r_+^2. \quad (61)$$

For this purpose, we expand the left-hand side of (27) in the small ϵ and r_+ limit to obtain

$$4z_0K_0 = \frac{1}{4}i(6\Delta - \sqrt{3}e)(2 - \epsilon)r_+ + \frac{1}{8}i(2\Delta + \sqrt{3}e)\epsilon^2r_+ + \frac{1}{16}(3e^2 + 4\sqrt{3}e\Delta + 4\Delta(\Delta - 16))(1 - \epsilon)r_+^2 + \mathcal{O}(\epsilon^3r_+, \epsilon^2r_+^2, r_+^3, \epsilon r_+^3) \quad (62)$$

while the right-hand side of (27) gives

$$4z_0K_0 = \frac{1}{4}i(6\Delta - \sqrt{3}e)(2 - \epsilon)r_+ + \frac{1}{8}i(2\Delta + \sqrt{3}e)\epsilon^2r_+ - \left[2\nu_{0,0} - \frac{1}{16}(\sqrt{3}e - 6\Delta)^2 - 4\nu_{0,0}x(1 + x + 2x^2 + 5x^3 + \dots) \right] r_+^2 - \left[\frac{1}{16}(\sqrt{3}e - 6\Delta)^2 + 8\nu_{0,0}x(1 + 2x + 6x^2 + 20x^3 + \dots) \right] \epsilon r_+^2 + \mathcal{O}(\epsilon^3r_+, \epsilon^2r_+^2, r_+^3, \epsilon r_+^3), \quad (63)$$

where the dots are the higher order poles in $\sigma_{0,0}$ that contribute to the r_+^2 and ϵr_+^2 terms, and we have defined

$$x = \frac{(2\Delta - \sqrt{3}e)(\sqrt{3}e - 6\Delta)(\Delta - 1)}{16\nu_{0,0}^2}. \quad (64)$$

Equating (62) and (63) leads to the following equation for $\nu_{0,0}$:

$$\frac{2\nu_{0,0}}{\sqrt{1 - 4x}} - \frac{8\nu_{0,0}x(1 - \epsilon)}{\sqrt{1 - 4x}} = \Delta(2(\Delta + 2) - \sqrt{3}e)(1 - \epsilon) + \mathcal{O}(\epsilon^2), \quad (65)$$

that can be solved to yield

$$\nu_{0,0} = \frac{1}{2}(1 - \epsilon)\sqrt{4\Delta^2(\Delta^2 + \Delta + 7) - 4\sqrt{3}e\Delta(\Delta^2 + 2) + 3e^2(\Delta^2 - \Delta + 1)} + \mathcal{O}(\epsilon^2). \quad (66)$$

Hence, $\sigma_{0,0}$ in (61) reads

$$\sigma_{0,0} = 2 - \nu_{0,0}r_+^2 = 2 - \tilde{\nu}_{0,0}(1 - \epsilon)r_+^2 + \mathcal{O}(\epsilon^2r_+^2), \quad (67)$$

where

$$\tilde{\nu}_{0,0} = \frac{1}{2}\sqrt{4\Delta^2(\Delta^2 + \Delta + 7) - 4\sqrt{3}e\Delta(\Delta^2 + 2) + 3e^2(\Delta^2 - \Delta + 1)}. \quad (68)$$

The asymptotic expansion for $\chi_{0,0} \equiv \chi_0(\omega = \omega_{0,0})$ now receives contributions from all orders in z_0 due to the pole structure at $\sigma = 2$ in (26). Nevertheless, one can again use the generating function for the Catalan numbers to obtain an analytic expansion for $\epsilon \ll r_+ \ll 1$ given by

$$\chi_{0,0} = \frac{1}{2} + \frac{(\Delta - 1)(2\Delta - \sqrt{3}e)(6\Delta - \sqrt{3}e)}{16\tilde{\nu}_{0,0}^2} + \frac{1}{2}\sqrt{1 + \frac{(\Delta - 1)(2\Delta - \sqrt{3}e)(6\Delta - \sqrt{3}e)}{4\tilde{\nu}_{0,0}^2}} + \mathcal{O}(\epsilon^2, r_+^2, \epsilon r_+^2). \quad (69)$$

By means of the expansion for θ_0 in (60), with $\omega = \omega_{0,0}$ in (51), we obtain

$$\theta_0 = \Delta - \beta_{0,0} r_+^2 = \Delta - \left(\frac{1}{2} (6\Delta - \sqrt{3}e)(1 - \epsilon) - \eta_{0,0} \right) r_+^2, \quad (70)$$

where

$$\eta_{0,0} = \frac{1}{2} (1 - \epsilon)(6\Delta - \sqrt{3}e) - \beta_{0,0}. \quad (71)$$

Inserting (69) in the right-hand side of (33) for $n = \ell = 0$ and expanding the left-hand side of (33) for $\epsilon \ll r_+ \ll 1$, we obtain

$$\begin{aligned} & \frac{(\Delta - 1)(6\Delta - \sqrt{3}e)(2\Delta - \sqrt{3}e)\beta_{0,0} + \tilde{\nu}_{0,0}}{16\tilde{\nu}_{0,0}^2} \frac{\beta_{0,0} + \tilde{\nu}_{0,0}}{\beta_{0,0} - \tilde{\nu}_{0,0}} \left[1 - \frac{2\tilde{\nu}_{0,0}\beta_{0,0}\epsilon}{\beta_{0,0}^2 - \tilde{\nu}_{0,0}^2} + \frac{i4\tilde{\nu}_{0,0}r_+}{(6\Delta - \sqrt{3}e)} - \frac{i2\tilde{\nu}_{0,0}(\beta_{0,0}^2 + 4\beta_{0,0}\tilde{\nu}_{0,0} - \tilde{\nu}_{0,0}^2)\epsilon r_+}{(6\Delta - \sqrt{3}e)(\beta_{0,0}^2 - \tilde{\nu}_{0,0}^2)} \right. \\ & \quad \left. + \mathcal{O}(\epsilon^2, \epsilon^2 r_+, r_+^2, r_+^2 \log r_+, \epsilon r_+^2, \epsilon r_+^2 \log r_+) \right] \\ & = \frac{1}{2} + \frac{(\Delta - 1)(2\Delta - \sqrt{3}e)(6\Delta - \sqrt{3}e)}{16\tilde{\nu}_{0,0}^2} + \frac{1}{2} \sqrt{1 + \frac{(\Delta - 1)(2\Delta - \sqrt{3}e)(6\Delta - \sqrt{3}e)}{4\tilde{\nu}_{0,0}^2}} + \mathcal{O}(\epsilon^2, r_+^2, \epsilon r_+^2) \end{aligned} \quad (72)$$

whose perturbative solution for $\beta_{0,0}$ reads

$$\beta_{0,0} = \frac{1}{2} \Delta (2(\Delta + 2) - \sqrt{3}e)(1 - \epsilon) + i \frac{1}{4} (\Delta - 1)(2\Delta - \sqrt{3}e)(2 - 3\epsilon)r_+ + \mathcal{O}(\epsilon^2, \epsilon^2 r_+, r_+^2, r_+^2 \log r_+, \epsilon r_+^2, \epsilon r_+^2 \log r_+). \quad (73)$$

Therefore, by means of (71) and (73), the fundamental QNM frequency is

$$\omega_{0,0} = \Delta - \frac{1}{2} (\Delta - 1)(2\Delta - \sqrt{3}e)(1 - \epsilon)r_+^2 - i \frac{1}{4} (\Delta - 1)(2\Delta - \sqrt{3}e)(2 - 3\epsilon)r_+^3 + \mathcal{O}(\epsilon^2 r_+^2, r_+^4, r_+^4 \log r_+, \epsilon r_+^4, \epsilon r_+^4 \log r_+), \quad (74)$$

which is valid for small black holes in the low-temperature limit, i.e., $\epsilon \ll r_+ \ll 1$. The imaginary part of $\omega_{0,0}$ in (74), at this order of perturbation theory, suggests that the black hole is stable ($\text{Im}\omega_{0,0} < 0$) under perturbations with charge $e < 2\Delta/\sqrt{3}$ and $\Delta > 1$. The imaginary part has a leading correction in ϵ of order ϵr_+^3 , indicating that contributions from the black hole temperature are not sufficiently enhanced to trigger superradiant instabilities for $e < 2\Delta/\sqrt{3}$. Such instabilities can instead be triggered by the charge of the scalar field provided that $\Delta > 1$. Incidentally, we notice that although Eqs. (57) and (74) were derived in two different regimes, they coincide in the extremal limit $\epsilon = 0$ ($q = 1$) at least at the given order in r_+ .

C. Comparison with numerical results

It is interesting to compare the analytical expression (74) for small black holes at low temperature with the numerical prediction for a given charge of the scalar field e as a function of the radius of the outer horizon r_+ , as well as the ϵ parameter. We recall that the numerical results have been obtained by solving the transcendental equations (24) for the Fredholm determinant of the PVI tau function [20]. The initial conditions have been implemented in Julia, using an

arbitrary-precision library. Then, the QNM frequencies have been found by applying a root-finding algorithm which employs the Newton-Raphson method.

Figure 2 shows the comparison between the numerical results (in red, solid line) and the low-temperature asymptotic formula (74) for the fundamental QNM frequency as a function of r_+ for fixed small $\epsilon = 5 \times 10^{-6}$ and the charge of the scalar field $e = 0.005$. We observe a small discrepancy between the two curves, both for the real and imaginary part of $\omega_{0,0}$, increasing to about 2% for $r_+ \simeq 0.01$. In Fig. 3 we display the transition from low to high temperatures—small and large ϵ , respectively—by varying ϵ while keeping $r_+ = 0.001$ fixed. The numerical results (again in red, solid line) are compared with the asymptotic formulas (in blue, dashed line) for high and low temperature, given by (58) and (74), respectively. For low temperatures, we have $\epsilon \ll r_+ \ll 1$ and thus r_+^3 corrections to (74) are more relevant for the approximation than the ϵr_+^2 terms. In other words, one should resum the series in r_+ at each given order in ϵ for a full comparison. Nevertheless, the ϵ corrections encode the temperature dependence, and can be checked to agree with the numerical prediction by comparing the slope of the two curves for small ϵ in the imaginary part of $\omega_{0,0}$ in Fig. 3.




FIG. 2. Analytical result (blue dashed line) in (74) for the real (left) and imaginary (right) part of the fundamental QNM frequency $\omega_{0,0}$ compared with the numerical result (red solid line) as function of r_+ for fixed $\epsilon = 5 \times 10^{-6}$ and charge of the scalar field $e = 0.005$.




FIG. 3. Analytical result (blue dashed line) in (74) for the real (left) and imaginary (right) part of the fundamental QNM frequency $\omega_{0,0}$ compared with the numerical result (red solid line) as function of ϵ for $e = 2$ and $r_+ = 0.001$. The ascending and descending dashed lines for the $\text{Im } \omega_{0,0}$ in the right panel reproduce (74) and (58), respectively. The ascending line describes the same slope of the numerical result; however, the shift with respect to the numerical result is due to higher orders in r_+ , for $\epsilon = 0$ in (74).

IV. DISCUSSION

In this paper, we have investigated the quasinormal modes of a charged scalar field scattering on small RN-AdS₅ black holes, where the outer horizon radius r_+ is much smaller than the AdS radius, via the isomonodromy method. We analytically explored two regimes in terms of r_+ and the parameter $\epsilon = 1 - q$, with $0 \leq q \leq 1$ the charge

of the black hole in units of the critical charge Q_c . In Sec. III A, we have considered the small black hole approximation at high temperatures ($r_+ \ll \epsilon < 1$) and provided an asymptotic expression for the QNMs frequencies for n generic and $\ell = 0$ that generalizes previous results found in the literature [7]. These asymptotic expressions display instabilities for the fundamental QNM mode for large enough q , and we verified

numerically (for $e = 8$ and $\Delta = 4$) that there is a super-radiance window whose width in r_+ increases with q . We leave open the question whether, in parallel to the Hawking-Page transition for uncharged black holes [36], stability might be restored as the radius of the black hole increases.

A different approach is required in the low-temperature limit $\epsilon \ll r_+ \ll 1$ and we have presented the analysis of this case in Sec. III B. By considering a double series expansion for small ϵ and r_+ , we have computed the first nontrivial correction in ϵ to the fundamental QNM frequency. We found that small black holes at low temperature are stable under scalar perturbations with Δ satisfying the unitarity bound $\Delta > 1$, and charge $e < 2\Delta/\sqrt{3}$. When the latter conditions are not met, the imaginary part of (74) becomes positive, meaning that the black hole is unstable.

In [7], the correction to the empty AdS frequency is derived from the matching condition between the analytical solutions constructed in terms of hypergeometric functions. In contrast, we have shown via the isomonodromy method that the asymptotic expansion for the fundamental QNM frequency receives contributions from all descendants of the CFT primaries in the vacuum module. These contributions can be resummed in terms of the generating functions of the Catalan numbers and the result suggests that in the matching approach one should consider all levels in the Frobenius expansion. It is worth mentioning that for numerical purposes the Fredholm determinant representation of the PVI tau function can reach a faster convergence time than direct instanton counting methods based on Nekrasov functions.

This work can be seen as an application to RN-AdS₅ of the ideas exposed in [33], where the authors have written the initial conditions for the Painlevé V tau function to compute an asymptotic expansion for the fundamental QNM frequency in the low-temperature regime.

Finally, in [5] it was argued that large RN-AdS₄ black holes, i.e., $r_+ \gg 1$ in units of the AdS curvature radius, become unstable below a critical temperature. Since one can control ϵ and r_+ in the accessory parameter expansion, it would be interesting to explore this regime and the nature of these instabilities in terms of the PVI tau function and its confluent limits. It is also interesting to study the hydrodynamical modes in RN-AdS₅ black holes, since scalar perturbations in this background have been introduced as dual CFT states of strongly correlated holographic models with finite density, for which the bound on the speed of sound can be violated [37]. The method presented here and in [33] works for higher spin perturbations on rotating black holes [32] and can be used to study those perturbations in RN-AdS₅.

ACKNOWLEDGMENTS

J. B. A. is grateful to Marco Bertola, Dmitry Korotkin, and Vasilisa Shramchenko for the support through their NSERC Discovery grants, as well as to the Mathematical Physics Laboratory at the Centre de Recherches Mathématiques in Montreal and to the Department of Mathematics at the University of Sherbrooke for financial support and hospitality.

APPENDIX: NEKRASOV EXPANSION FOR THE PVI TAU FUNCTION

The series representation of the PVI tau function, written in [15,38], around the critical point $t = 0$ is given by

$$\tau(t) = \sum_{n \in \mathbb{Z}} C(\vec{\theta}, \sigma + 2n) s^n t^{\frac{1}{2}((\sigma+2n)^2 - \theta_1^2 - \theta_2^2)} \mathcal{B}(\vec{\theta}, \sigma + 2n; t), \quad (\text{A1})$$

where $\vec{\theta} = \{\theta_1, \theta_2, \theta_3, \theta_4\}$ are the single monodromy parameters, and the parameters σ, s are two integration constants. The structure constants $C(\vec{\theta}, \sigma)$ are given in terms of Barnes's functions

$$C(\vec{\theta}, \sigma) = \frac{\prod_{\alpha, \beta = \pm} G(1 + \frac{1}{2}(\theta_3 + \alpha\theta_4 + \beta\sigma)) G(1 + \frac{1}{2}(\theta_2 + \alpha\theta_1 + \beta\sigma))}{G(1 + \sigma) G(1 - \sigma)}, \quad (\text{A2})$$

and $\mathcal{B}(\vec{\theta}, \sigma; t)$ in (A1) coincides with the $c = 1$ Virasoro conformal blocks, which are explicitly given by a combinatorial series in terms of Nekrasov functions

$$\mathcal{B}(\vec{\theta}, \sigma + 2n; t) = (1-t)^{\frac{1}{2}\theta_2\theta_3} \sum_{\lambda, \mu \in \mathbb{Y}} \mathcal{B}_{\lambda, \mu}(\vec{\theta}, \sigma + 2n) t^{|\lambda| + |\mu|}, \quad (\text{A3})$$

where \mathbb{Y} denotes the space of Young diagrams, λ and μ are two of its elements, with number of boxes $|\lambda|$ and $|\mu|$, and the coefficients in (A3) are

$$\begin{aligned} \mathcal{B}_{\lambda,\mu}(\vec{\theta}, \sigma) &= \prod_{(i,j) \in \lambda} \frac{((\theta_i + \sigma + 2(i-j))^2 - \theta_0^2)((\theta_1 + \sigma + 2(i-j))^2 - \theta_\infty^2)}{16h_\lambda^2(i,j)(\lambda'_j - i + \mu_i - j + 1 + \sigma)^2} \\ &\times \prod_{(i,j) \in \mu} \frac{((\theta_i - \sigma + 2(i-j))^2 - \theta_0^2)((\theta_1 - \sigma + 2(i-j))^2 - \theta_\infty^2)}{16h_\lambda^2(i,j)(\mu'_j - i + \lambda_i - j + 1 - \sigma)^2}. \end{aligned} \quad (\text{A4})$$

For each box situated at (i, j) in λ , λ_i is the number of boxes at row i of λ and λ'_j is the number of boxes at column j of λ ; $h(i, j) = \lambda_i + \lambda'_j - i - j + 1$ is the hook length of the box at (i, j) . The parameter s can be determined in terms of the monodromy matrices $\{\sigma, \sigma_{23}\}$ from the formula (3.48a) in [39]:

$$\begin{aligned} \sin^2 \pi \sigma \cos \pi \sigma_{23} &= \cos \pi \theta_1 \cos \pi \theta_4 + \cos \pi \theta_2 \cos \pi \theta_3 - \cos \pi \sigma (\cos \pi \theta_1 \cos \pi \theta_3 + \cos \pi \theta_2 \cos \pi \theta_4) \\ &- \frac{1}{2} (\cos \pi \theta_4 - \cos \pi (\theta_3 - \sigma)) (\cos \pi \theta_1 - \cos \pi (\theta_2 - \sigma)) s \\ &- \frac{1}{2} (\cos \pi \theta_4 - \cos \pi (\theta_3 + \sigma)) (\cos \pi \theta_1 - \cos \pi (\theta_2 + \sigma)) s^{-1}. \end{aligned} \quad (\text{A5})$$

For t sufficiently close to zero,¹ and generic monodromy parameters in the sense that

$$\sigma \notin \mathbb{Z}, \quad \sigma \pm \theta_1 \pm \theta_2 \notin \mathbb{Z}, \quad \sigma \pm \theta_3 \pm \theta_4 \notin \mathbb{Z}, \quad (\text{A6})$$

we have

$$\begin{aligned} \tau(t) &= C_0 t^{\frac{1}{4}(\sigma^2 - \theta_1^2 - \theta_2^2)} (1-t)^{\frac{1}{2}\theta_3\theta_2} \left\{ 1 + \left[\frac{\theta_3\theta_2}{2} + \frac{(\theta_1^2 - \theta_2^2 - \sigma^2)(\theta_4^2 - \theta_3^2 - \sigma^2)}{8\sigma^2} \right. \right. \\ &\left. \left. - \frac{(\theta_1^2 - (\theta_2 - \sigma)^2)(\theta_4^2 - (\theta_3 - \sigma)^2)}{16\sigma^2(1+\sigma)^2} \kappa t^\sigma - \frac{(\theta_1^2 - (\theta_2 + \sigma)^2)(\theta_4^2 - (\theta_3 + \sigma)^2)}{16\sigma^2(1-\sigma)^2} \frac{1}{\kappa t^\sigma} \right] t + \dots \right\}, \end{aligned} \quad (\text{A7})$$

where $0 < \text{Re} \sigma < 1$, C_0 is a constant independent of t , and κ is a known function of the monodromy parameters:

$$\kappa = s \frac{\Gamma^2(1-\sigma) \Gamma(1+\frac{1}{2}(\theta_2+\theta_1+\sigma)) \Gamma(1+\frac{1}{2}(\theta_2-\theta_1+\sigma)) \Gamma(1+\frac{1}{2}(\theta_3+\theta_4+\sigma)) \Gamma(1+\frac{1}{2}(\theta_3-\theta_4+\sigma))}{\Gamma^2(1+\sigma) \Gamma(1+\frac{1}{2}(\theta_2+\theta_1-\sigma)) \Gamma(1+\frac{1}{2}(\theta_2-\theta_1-\sigma)) \Gamma(1+\frac{1}{2}(\theta_3+\theta_4-\sigma)) \Gamma(1+\frac{1}{2}(\theta_3-\theta_4-\sigma))}. \quad (\text{A8})$$

¹One can choose different series expansions around the other critical points of the PVI tau function at $t = 1$ and $t = \infty$ [29].

-
- [1] R. Brito, V. Cardoso, and P. Pani, Superradiance: New frontiers in black hole physics, *Lect. Notes Phys.* **906**, 1 (2015).
- [2] J. D. Bekenstein, Extraction of energy and charge from a black hole, *Phys. Rev. D* **7**, 949 (1973).
- [3] G. Denardo and R. Ruffini, On the energetics of Reissner Nordström geometries, *Phys. Lett.* **45B**, 259 (1973).
- [4] G. W. Gibbons, Vacuum polarization and the spontaneous loss of charge by black holes, *Commun. Math. Phys.* **44**, 245 (1975).
- [5] K. Maeda, S. Fujii, and J.-i. Koga, Final fate of instability of Reissner-Nordström-anti-de Sitter black holes by charged complex scalar fields, *Phys. Rev. D* **81**, 124020 (2010).
- [6] N. Uchikata and S. Yoshida, Quasinormal modes of a massless charged scalar field on a small Reissner-Nordström-anti-de Sitter black hole, *Phys. Rev. D* **83**, 064020 (2011).
- [7] M. Wang and C. Herdeiro, Superradiant instabilities in a D-dimensional small Reissner-Nordström-anti-de Sitter black hole, *Phys. Rev. D* **89**, 084062 (2014).
- [8] S. S. Gubser, Breaking an Abelian gauge symmetry near a black hole horizon, *Phys. Rev. D* **78**, 065034 (2008).
- [9] S. A. Hartnoll, C. P. Herzog, and G. T. Horowitz, Holographic superconductors, *J. High Energy Phys.* **12** (2008) 015.
- [10] S. A. Hartnoll, Lectures on holographic methods for condensed matter physics, *Classical Quantum Gravity* **26**, 224002 (2009).

- [11] P. Basu, J. Bhattacharya, S. Bhattacharyya, R. Loganayagam, S. Minwalla, and V. Umesh, Small hairy black holes in global AdS spacetime, *J. High Energy Phys.* **10** (2010) 045.
- [12] O. J. C. Dias, P. Figueras, S. Minwalla, P. Mitra, R. Monteiro, and J. E. Santos, Hairy black holes and solitons in global AdS₅, *J. High Energy Phys.* **08** (2012) 117.
- [13] J. B. Amado, B. Carneiro da Cunha, and E. Pallante, On the Kerr-AdS/CFT correspondence, *J. High Energy Phys.* **08** (2017) 094.
- [14] J. Barragán Amado, B. Carneiro da Cunha, and E. Pallante, Scalar quasinormal modes of Kerr-AdS₅, *Phys. Rev. D* **99**, 105006 (2019).
- [15] O. Gamayun, N. Iorgov, and O. Lisovyy, How instanton combinatorics solves Painlevé VI, V and IIIs, *J. Phys. A* **46**, 335203 (2013).
- [16] T. Anselmo, R. Nelson, B. Carneiro da Cunha, and D. G. Crowdy, Accessory parameters in conformal mapping: Exploiting the isomonodromic tau function for Painlevé VI, *Proc. R. Soc. A* **474**, 20180080 (2018).
- [17] F. Novaes and B. Carneiro da Cunha, Isomonodromy, Painlevé transcendents and scattering off of black holes, *J. High Energy Phys.* **07** (2014) 132.
- [18] B. Carneiro da Cunha and F. Novaes, Kerr-de Sitter greybody factors via isomonodromy, *Phys. Rev. D* **93**, 024045 (2016).
- [19] B. Carneiro da Cunha, M. C. de Almeida, and A. R. de Queiroz, On the existence of monodromies for the Rabi model, *J. Phys. A* **49**, 194002 (2016).
- [20] P. Gavrylenko and O. Lisovyy, Fredholm determinant and Nekrasov sum representations of isomonodromic tau functions, *Commun. Math. Phys.* **363**, 1 (2018).
- [21] G. Aminov, A. Grassi, and Y. Hatsuda, Black hole quasinormal modes and Seiberg-Witten theory, [arXiv:2006.06111](https://arxiv.org/abs/2006.06111).
- [22] M. Bershtein, P. Gavrylenko, and A. Grassi, Quantum spectral problems and isomonodromic deformations, [arXiv:2105.00985](https://arxiv.org/abs/2105.00985).
- [23] G. Bonelli, C. Iossa, D. P. Lichtig, and A. Tanzini, Exact solution of Kerr black hole perturbations via CFT₂ and instanton counting, [arXiv:2105.04483](https://arxiv.org/abs/2105.04483).
- [24] M. Bianchi, D. Consoli, A. Grillo, and J. F. Morales, QNMs of branes, BHs and fuzzballs from quantum SW geometries, *Phys. Lett. B* **824**, 136837 (2022).
- [25] M. Bianchi, D. Consoli, A. Grillo, and J. F. Morales, More on the SW-QNM correspondence, *J. High Energy Phys.* **01** (2022) 024.
- [26] G. T. Horowitz, Introduction to holographic superconductors, *Lect. Notes Phys.* **828**, 313 (2011).
- [27] M. Jimbo, T. Miwa, and K. Ueno, Monodromy preserving deformation of linear ordinary differential equations with rational coefficients: I. General theory and τ -function, *Physica (Amsterdam)* **2D**, 306 (1981).
- [28] M. Jimbo and T. Miwa, Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. II, *Physica (Amsterdam)* **2D**, 407 (1981).
- [29] M. Jimbo, Monodromy problem and the boundary condition for some Painlevé equations, *Publ. RIMS* **18**, 1137 (1982).
- [30] B. Carneiro da Cunha and J. P. Cavalcante, Confluent conformal blocks and the Teukolsky master equation, *Phys. Rev. D* **102**, 105013 (2020).
- [31] B. Carneiro da Cunha and J. P. Cavalcante, Teukolsky master equation and Painlevé transcendents: Numerics and extremal limit, *Phys. Rev. D* **104**, 084051 (2021).
- [32] J. B. Amado, B. Carneiro da Cunha, and E. Pallante, Vector perturbations of Kerr-AdS₅ and the Painlevé VI transcendent, *J. High Energy Phys.* **04** (2020) 155.
- [33] J. Barragán Amado, B. Carneiro da Cunha, and E. Pallante, Remarks on holographic models of the Kerr-AdS₅ geometry, *J. High Energy Phys.* **05** (2021) 251.
- [34] F. Novaes, C. Marinho, M. Lencsés, and M. Casals, Kerr-de Sitter quasinormal modes via accessory parameter expansion, *J. High Energy Phys.* **05** (2019) 033.
- [35] <https://github.com/strings-ufpe/painleve>.
- [36] S. W. Hawking and D. N. Page, Thermodynamics of black holes in anti-de Sitter space, *Commun. Math. Phys.* **87**, 577 (1983).
- [37] C. Hoyos, N. Jokela, D. Rodríguez Fernández, and A. Vuorinen, Breaking the sound barrier in AdS/CFT, *Phys. Rev. D* **94**, 106008 (2016).
- [38] O. Gamayun, N. Iorgov, and O. Lisovyy, Conformal field theory of Painlevé VI, *J. High Energy Phys.* **10** (2012) 038; *J. High Energy Phys. Erratum*, **10** (2012) 183.
- [39] A. Its, O. Lisovyy, and A. Prokhorov, Monodromy dependence and connection formulae for isomonodromic tau functions, *Duke Math. J.* **167**, 1347 (2018).