

University of Groningen

Peroxisomes in yeast ageing

Kumar, Sanjeev

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:
2015

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kumar, S. (2015). *Peroxisomes in yeast ageing*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Peroxisomes in yeast ageing

© 2015 Sanjeev Kumar

ISBN: 978-94-6182-631-2

Cover: Sanjeev Kumar

Layout and printing: Off Page, www.offpage.nl



**rijksuniversiteit
 groningen**



The studies presented in this thesis were performed in the research group Molecular Cell Biology of the Groningen Biomolecular Sciences and Biotechnology Institute (GBB) of the University of Groningen, The Netherlands.

This project was carried out within the research program of the Systems Biology Centre for Energy Metabolism and Aging (SBC-EMA), which is one of three national systems biology centres funded by the Dutch Science Foundation (NWO).



university of
 groningen

Peroxisomes in yeast ageing

PhD thesis

to obtain the degree of PhD at the
 University of Groningen
 on the authority of the
 Rector Magnificus Prof. E. Sterken
 and in accordance with
 the decision by the College of Deans.

This thesis will be defended in public on

Friday 4 December 2015 at 12.45 hours

by

Sanjeev Kumar

born on 18 December 1983
 in Bihar, India

Supervisor

Prof. I.J. van der Klei

Assessment Committee

Prof. O.P. Kuipers

Prof. B.M. Bakker

Prof. H.A.B. Wosten

Acknowledgements

I would like to take this opportunity to acknowledge people who in various capacities contributed to the preparation of this thesis.

Firstly, I would like to express my sincere gratitude to Prof. Ida J. van der Klei for giving me the opportunity to work in the Molecular Cell Biology group. I am very grateful for your supervision, guidance and encouragement throughout my stay in the laboratory. You were open to new ideas and also allowed me to work independently on diverse research projects that helped immensely in my training as a researcher.

I would like to acknowledge Prof. Marten Veenhuis for his valuable suggestions on different research projects.

I extend my special thanks to the members of thesis reading committee Prof. O.P. Kuipers, Prof. B.M. Bakker and Prof. H.A.B. Wosten for evaluating my thesis and giving valuable suggestions.

I take this opportunity to thank my paranimfs Ritika and Chris for their help in preparation of my defense. I also thank Anita for translating my thesis summary in Dutch.

Evelyne, Barbara, Frank, Ludmilla and Bhagawati, I feel privileged that I had the opportunity to work with you during your internship and learn a lot in the process. I would like to thank you all for your contributions to my research.

My special thanks to Sophie for her support during early days of my PhD and also for introducing me to yeast ageing research.

I am thankful to Adam for being a good friend and colleague. I thoroughly enjoyed numerous scientific discussions and our time together in the lab.

I earnestly express my gratitude to fellow labmates, past and present, in the Molecular Cell Biology group for providing a pleasant ambience at the workplace. I am thankful and indebted to each one of you for innumerable insightful discussions and also for extending your expertise to my research.

My sincere thanks to Jannet for her help in handling official procedures during my stay in the Netherlands.

Outside the lab, I would like to thank Jan Maat, Lea and Nate for all the good times that we spent together that made my stay in the Netherlands even more memorable.

I am extremely thankful to my friend and soon-to-be my wife Runa for her love and encouragement. Your constant support made this journey much easier. I am so lucky to have you in my life.

Finally, I am grateful to my beloved parents, brothers and entire family for their encouragement, support and faith in me. I am especially thankful to my father for making all the sacrifices and never letting failures cloud my thoughts. You always have been an inspiration to me.

Thank you!

Sanjeev

Contents

Aim and outline	9
Chapter 1 Introduction	13
Chapter 2 Peroxisome heterogeneity in yeast	39
Chapter 3 Stress exposure results in increased peroxisomal levels of yeast Pnc1 and Gpd1, which are imported via a piggy-backing mechanism	61
Chapter 4 Extension of yeast chronological lifespan by methylamine	85
Chapter 5 Inhibition of peroxisome fission, but not mitochondrial fission, increases yeast chronological lifespan	107
Summary	125
Samenvatting	133
Biography of the author	141

Aim and outline

Eukaryotic cells are subdivided into specialized membrane bound organelles that are characterized by their unique structure and functions. Among these, peroxisomes are single membrane bound, ubiquitous organelles which display a wide range of physiological functions depending on organism, environmental conditions and developmental stage. An important feature of peroxisomes is the presence of oxidases, which generate hydrogen peroxide that has been shown to be important in cellular ageing. Ageing is defined as a multifactorial process that is characterized by a progressive decline in cellular and organismal functions. Various model organisms, among which yeast, have been instrumental in the advancement of our understanding of the molecular principles of the ageing process.

Using yeast as model organism, the aim of the research described in this thesis was to understand the significance of peroxisomes in cellular ageing processes.

In **Chapter 1** an overview of our current understanding of peroxisome biology has been presented. Special emphasis has been given to our knowledge of peroxisomal quality control mechanisms and the roles of peroxisomes in ageing.

Chapter 2 describes the heterogeneity of peroxisomes in wild type cells of the yeast *Saccharomyces cerevisiae* that exists with respect to their matrix protein composition. By using specific fluorescent proteins that display different maturation times, we show that peroxisomes containing relatively old or new matrix protein content are present in one and the same cell. We showed that peroxisomes harboring a relatively old matrix protein content have a reduced capacity to import newly synthesized matrix proteins. This was related to reduced levels of Pex14, a peroxisomal membrane protein involved in matrix protein import, on mature peroxisomes. Further analysis revealed that mature peroxisomes are retained by the mother cells whereas peroxisomes with a younger peroxisomal matrix protein content are preferentially transported to the daughter cells.

Chapter 3 describes studies on the sorting of two stress related proteins, Pnc1 (nicotinamidase) and Gpd1 (glycerol-3-phosphate dehydrogenase) to peroxisomes in yeast. We show that Pnc1 is transported to peroxisomes by piggy-backing on the peroxisomal targeting signal-2 (PTS2) containing enzyme Gpd1. Our data indicate that Gpd1 and Pnc1 physically interact and most likely form a transient complex during import into peroxisomes. Previous reports indicated that Gpd1 relocates to the cytosol upon exposure of cells to osmotic stress. Quantitative analysis of the distribution of Gpd1 and Pnc1 over the cytosol and peroxisomes revealed that both proteins are predominantly localized to peroxisomes in unstressed cells. Upon exposure of cells to stress the levels of both peroxisomal and cytosolic Gpd1 and Pnc1 increased. A similar behaviour was observed for the non-stress related

peroxisomal control protein thiolase produced under control of the *GPD1* promoter. This observation suggests that presence of peroxisomal matrix proteins in the cytosol of cells exposed to stress is a result of reduced matrix protein import efficiency.

Chapter 4 describes the effects of different carbon and nitrogen sources on the chronological lifespan of the methylotrophic yeast *Hansenula polymorpha*. We found that this lifespan is enhanced when cells are grown on methanol or ethanol as sole carbon sources, compounds that require peroxisomal enzymes for their metabolism, relative to cultivation on glucose, which is metabolized independent of peroxisomal enzymes. We also observed that the use of methylamine as sole nitrogen source has a positive effect on yeast chronological lifespan relative to ammonium sulphate. Methylamine is initially metabolized inside peroxisomes by the enzyme amine oxidase. The lifespan extension by the utilization of methylamine was shown to be due to the surplus energy generated by the oxidation of methylamine during the stationary phase.

Mitochondrial fragmentation mediated by the Dnm1/Fis1 fission machinery has been suggested to accelerate cellular ageing. However, this Dnm1/Fis1 fission machinery is also involved in fission of peroxisomes. In **Chapter 5** we describe studies on the importance of peroxisomal fission in yeast ageing. We show that lifespan extension in *FIS1* deletion cells is mainly caused by the defects in peroxisome fission and not due to blockage of mitochondrial fragmentation. The deletion of *FIS1* in peroxisome deficient *pex3* cells did not result into lifespan extension, which further underlines the importance of peroxisome fission in yeast ageing.

