

University of Groningen

Sorting and trafficking of proteins in oligodendrocytes during myelin membrane biogenesis

Klunder, Lammert

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2007

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Klunder, L. (2007). *Sorting and trafficking of proteins in oligodendrocytes during myelin membrane biogenesis*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

References

- Ainger, K., Avossa, D., Morgan, F., Hill, S.J., Barry, C., Barbarese, E. and Carson, J.H. (1993): Transport and localization of exogenous myelin basic protein mRNA microinjected into oligodendrocytes. *J. Cell Biol.* 123: 431-441
- Ainger, K., Avossa, D., Diana, A.S., Barry, C., Barbarese, E. and Carson, J.H. (1997): Transport and localization elements in myelin basic protein mRNA. *J. Cell Biol.* 138: 1077-1087
- Aït Slimane, T. and Hoekstra, D. (2002): Sphingolipid trafficking and protein sorting in epithelial cells. *FEBS Lett.* 529: 54-59
- Aït Slimane, T., Trugnan, G., van IJzendoorn, S. C. D. and Hoekstra, D. (2003): Raft-mediated trafficking of apical resident proteins occurs in both direct and transcytotic pathways in polarized hepatic cells: role of distinct lipid microdomains. *Mol. Biol. Cell* 14: 611-624
- Allan, B.B., Moyer, B.D. and Balch, W.E. (2000): Rab1 recruitment of p115 into a cis-SNARE complex: programming budding COPII vesicles for fusion. *Science* 289: 444-448
- Allinquant, B., Staugaitis, S.M., D'Urso, D. and Colman, D.R. (1991): The ectopic expression of myelin basic protein isoforms in shiverer oligodendrocytes: implications for myelinogenesis. *J. Cell Biol.* 113: 393-403
- Alonso, M.A., Fan, L. and Alarcon, B. (1997): Multiple sorting signals determine apical localization of a nonglycosylated integral membrane protein. *J. Biol. Chem.* 272: 30748-30752
- Amiguet, P., Gardinier, M.V., Zanetta, J.P. and Matthieu, J.M. (1992): Purification and partial structural and functional characterization of mouse myelin oligodendrocyte glycoprotein (MOG). *J. Neurochem.* 58: 1676-1682
- Anitei, M., Ifrim, M., Ewart, M-A., Cowan, A.E., Carson, J.H., Bansal, R. and Pfeiffer, S.E. (2006): A role for Sec8 in oligodendrocyte morphological differentiation. *J. Cell Sci.* 119: 807-818
- Apodaca, G., Cardone, M.H., Whiteheart, S.W., DasGupta, B.R. and Mostov, K.E. (1996): Reconstitution of transcytosis in SLO-permeabilized MDCK cells: existence of an NSF-dependent fusion mechanism with the apical surface of MDCK cells. *EMBO J.* 15: 1471-1481
- Aroeti, B., Kosen, P.A., Kuntz, I.D., Cohen, F.E. and Mostov, K.E. (1993): Mutational and secondary structural analysis of the basolateral sorting signal of the polymeric immunoglobulin receptor. *J. Cell Biol.* 123: 1149-1160

- Aronov, E. and Gerst, J.E. (2004): Involvement of the late secretory pathway in actin regulation and mRNA transport in yeast. *J. Biol. Chem.* 35: 36962-36971
- Arroyo, E.J. and Scherer, S.S. (2000): On the molecular architecture of myelinated fibers. *Histochem. Cell Biol.* 113: 1-18
- Asotra, K. and Macklin, W.B. (1993): Protein kinase C activity modulates myelin gene expression in enriched oligodendrocytes. *J. Neurosci. Res.* 34: 571-588
- Bansal, R., Gard, A.L., and Pfeiffer, S.E. (1988): Stimulation of oligodendrocyte differentiation in culture by growth in the presence of a monoclonal antibody to sulfated glycolipid. *J. Neurosci. Res.* 21: 260-267
- Bansal, R. and Pfeiffer, S.E. (1994): Inhibition of protein and lipid sulfation in oligodendrocytes blocks biological responses to FGF-2 and retards cytoarchitectural maturation, but not developmental lineage progression. *Dev. Biol.* 162: 511-524
- Bansal, R., Winkler, S. and Bheddah, S. (1999): Negative regulation of oligodendrocyte differentiation by galactosphingolipids. *J. Neurosci.* 19: 7913-7924
- Barbarese, E., Koppel, D.E., Deutcher, M.P., Ainger, K., Morgan, F. and Carson, J.H. (1995): Protein translation components are colocalized in granules in oligodendrocytes. *J. Cell Sci.* 108: 2781-2790
- Barlett, P.F. and Mackay, I.R. (1983): The oligodendroglial cell: biology and immunology and relationship to multiple sclerosis. *J. Clin. Lab. Immunol.* 11: 1-7
- Barnett, M.H. and Prineas, J.W. (2004): Relapsing and remitting multiple sclerosis: pathology of the newly forming lesion. *Ann. Neurol.* 55: 458-468
- Baron, W., De jonge, J. C., de Vries, H. and Hoekstra, D. (1998): Regulation of oligodendrocyte differentiation: Protein K activation prevents differentiation of O2A progenitor cells towards oligodendrocytes. *Glia* 22: 121-129
- Baron, W., de Vries, E.J., de Vries, H. and Hoekstra, D. (1999): Protein kinase C prevents oligodendrocyte differentiation: modulation of actin cytoskeleton and cognate polarized membrane traffic. *J. Neurobiol.* 41: 385-398.
- Baron, W., Decker, L., Colognato, H. and French-Constant, C. (2003): Regulation of integrin growth factor interactions in oligodendrocytes by lipid raft microdomains. *Curr. Biol.* 13: 151-155
- Baron, W., Colognato, H. and French-Constant, C. (2005): Integrin-growth factor interactions as regulators of oligodendrocyte development and function. *Glia* 49: 467-479

References

- Baumann, N. and Pam-Dinh, D. (2001): Biology of oligodendrocyte and myelin in the mammalian central nervous system. *Physiol. Rev.* 81: 871-927
- Bello, V., Goding, J.W., Greengrass, V., Sali, A., Dubljevic, V., Lenoir, C., Trugnan, G. and Maurice, M. (2001): Characterization of a di-Leucine-based signal in the cytoplasmic tail of the nucleotide-pyrophosphatase NPP1 that mediates basolateral targeting but not endocytosis. *Mol. Biol. Cell* 10: 3004-3015
- Benting, J.H., Rietveld, A.G. and Simons, K. (1999): N-glycans mediate the apical sorting of a GPI-anchored, raft-associated protein in Madin-Darby canine kidney cells. *J. Cell Biol.* 146: 313-320
- Bernards, A. (2003): GAPs galore! A survey of putative Ras superfamily GTPase activating proteins in man and *Drosophila*. *Biochim. Biophys. Acta* 1603: 47-82
- Birling, M.C., Roussel, G. Nussbaum, F. and Nussbaum, J.L. (1993): Biochemical and immunohistochemical studies with specific polyclonal antibodies directed against bovine myelin/oligodendrocyte glycoprotein. *Neurochem. Res.* 18: 937-945
- Blaschuk, K.L., Frost, E.E. and French-Constant, C. (2000): The regulation of proliferation and differentiation in oligodendrocyte progenitor cells by αV integrins. *Development* 127: 1961-1969.
- Bo, L., Quarles, R.H., Fujita, N., Bartoszewicz, Z., Sato, S. and Trapp, B.D. (1995): Endocytic depletion of L-MAG from CNS myelin in quaking mice. *J. Cell Biol.* 131: 1811-1820
- Boccaccio, G.L. and Colman, D.R. (1995): Myelin basic protein mRNA localization and polypeptide targeting. *J. Neurosci. Res.* 42: 277-286.
- Bonifacino, J.S. and Dell-Angelica, E.C. (1999): Molecular bases for the recognition of tyrosine-based sorting signals. *J. Cell Biol.* 145: 923-926.
- Bouveratt, B.P., Krueger, W.H., Coetzee, T., Bansal, R. and Pfeiffer, S.E. (2000): Expression of Rab GTP-binding proteins during oligodendrocyte differentiation in culture. *J. Neurosci. Res.* 59: 446-453.
- Braakman, I., Hoover-Litty, H., Wagner, K.R. and Helenius A. (1991): Folding of influenza haemagglutinin in the endoplasmic reticulum. *J. Cell Biol.* 114: 401-411.
- Braun, P.E., Sandillon, F., Edwards, A., Matthieu, J.M. and Privat, A. (1988): Immunocytochemical localization by electron microscopy of 2'3'-cyclic nucleotide 3'-phosphodiesterase in developing oligodendrocytes of normal and mutant brain. *J. Neurosci.* 8: 3057-3066.

- Bravo-Zehder, M., Orio, P., Norambuena, A., Wallner, M., Meera, P., Toro, L., Latorre, R. and Gonzalez, A. (2000): Apical sorting of a voltage- and Ca²⁺-activated K⁺ channel a subunit in Madin-Darby canine kidney is independent of N-glycosylation. *Proc. Natl. Acad. Sci. USA* 97: 13114-13119
- Breuzza, L., Fransen, J. and Le Bivic, A.(2000): Transport and function of syntaxin 3 in human epithelial intestinal cells. *Am. J. Physiol. Cell. Physiol.* 279: C1239–C1248
- Brewer, C.B. and Roth, M.G. (1991): A single amino acid change in the cytoplasmic domain alters the polarized delivery of influenza viral haemagglutinin. *J. Cell Biol.* 114: 413–421.
- Bronstein, J.M., Tiwari-Woodruff, S., Buznikov, A.G. and Stevens, D.B. (2000): Involvement of OSP/claudin-11 in oligodendrocyte membrane interactions: role in biology and disease. *J. Neurosci. Res.* 59: 706–711
- Brown, D.A. and London, E. (1998): Functions of lipid rafts in biological membranes. *Annu. Rev. Cell Dev. Biol.* 14: 111–136
- Brown, M.C., Besio Moreno, M., Bongaroze, E.R., Cohen, P.D. Soto, E.F. and Pasquini, J.M. (1993): Vesicular transport of myelin proteolipid and cerebroside sulfates to the myelin membrane. *J. Neurosci. Res.* 35: 402-408
- Burcelin, R., Rodriguez-Gabin, A.G., Charron, M.J., Almazan, G. and Larocca, J.N. (1997): Molecular analysis of the monomeric GTB-binding proteins of oligodendrocytes. *Molec. Brain Res.* 50: 9-15
- Buttery, P.C. and ffrench-Constant, C. (1999): Laminin-2/integrin interactions enhance myelin formation by oligodendrocytes. *Mol. Cell Neurosci.* 14: 199-212
- Campagnoni, A.T. (1988): Molecular biology of myelin proteins from the central nervous system. *J. Neurochem.* 51: 1-14
- Campagnoni, A.T. (1993): Structure and development regulation of Golli-mbp, a 105-kilobase gene that encompasses the myelin basic protein gene and is expressed in cells in the oligodendrocyte lineage in the brain. *J. Biol. Chem.* 268: 4930-4938
- Cardone, M.H., Smith, B.L., Song, W., Mochly-Rosen, D. and Mostov, K.E. (1994): Phorbol myristate acetate-mediated stimulation of transcytosis and apical recycling MDCK cells. *J. Cell Biol.* 124: 717-727.
- Carson, J.H., Nielson, M.L. and Barbarese, E. (1983): Developmental regulation of myelin basic protein expression in mouse brain. *Dev. Biol.* 96: 485-492

References

- Carson, J.H., Worboys, K., Ainger, K. and Barbarese, E. (1997): Translocation of myelin basic protein mRNA in oligodendrocytes requires microtubules and kinesin. *Cell Motil. Cytoskel.* 38: 318-328
- Carson, J.H., Kwon, S. and Barbarese, E. (1998): RNA trafficking in myelinating cells. *Curr. Opin. Neurobiol.* 8: 607-612
- Chen, Y. and Scheller, R.H. (2001): SNARE-mediated membrane fusion. *Nat. Rev. Mol. Cell Biol.* 2: 98-106
- Cheong, K.H., Zacchetti, D., Schneeberger, E.E. and Simons, K. (1999): VIP17/MAL, a lipid raft-associated protein, is involved in apical transport in MDCK cells. *Proc. Natl. Acad. Sci. USA* 11: 6241-6248
- Coetzee, T., Fujita, N., Dupree, J., Shi, R., Blight, A., Suzuki, K. and Popko, B. (1996): Myelination in the absence of galactocerebroside and sulfatide: normal structure with abnormal function and regional stability. *Cell* 86: 209-219
- Colman, D.R., Kreibich, G., Frey, A.B. and Sabatini, D.D. (1982): Synthesis and incorporation of myelin polypeptides into CNS myelin. *J. Cell Biol.* 95: 598-608
- Colman, D.R., Kreibich, G. and Sabatini, D.D. (1983): Biosynthesis of myelin-specific proteins. *Methods Enzymol.* 96: 378-385
- Colognato, H., Baron, W., Avellana-Adalid, V., Relvas, J.B., Baron-Van Evercooren, A., Georges-Labouesse, E. and ffrench-Constant, C. (2002): CNS integrins switch growth factor signaling to promote target-dependent survival. *Nat. Cell Biol.* 4: 833-841
- Colognato, H., Galvin, J., Whang, Z., Relucio, J., Nguyen, T., Harrison, D., Yurchenco, P.D. and ffrench-Constant, C. (2007): Identification of dystroglycan as a second laminin receptor in oligodendrocytes, with a role in myelination. *Development* 134: 1723-1736
- Corley, S.M., Ladiwala, U., Besson, A. and Yong, V.W. (2001): Astrocytes attenuate oligodendrocyte death in vitro through an alpha (6) integrin-laminin-dependent mechanism. *Glia* 36: 281-294
- De Angelis, D.A. and Braun, P.E. (1996) 2',3'-Cyclic nucleotide 3'-phosphodiesterase binds to actin-based cytoskeletal elements in an isoprenylation-dependent manner. *J. Neurochem.* 67: 943-951
- Dell'Angelica, E.C., Ohno, H., Ooi, C.E., Rabinovich, E., Roche, K.W., and Bonifacino, J.S. (1997): AP-3: an adaptor-like protein complex with ubiquitous expression. *EMBO J.* 16: 917-928

- Delacour, D. and Jacob, R. (2006): Apical protein transport. *Cell. Mol. Life. Sci.* 63: 2491-2505.
- De Maat, M.P., Laurens, N. and Koolwijk, P. (2006): Fibrin structure and wound healing. *J. Thromb. Haemost.* 4: 932-939
- De Vries, H., Schrage, C., Hoekstra, K., KoK, J.W., Van der Haar, M.E., Kalicharan, D., Liem, R.S.B., Copray, J.C.V.M. and Hoekstra, D. (1993): Outstations of the Golgi complex are present in the processes of cultured rat oligodendrocytes. *J. Neurosci. Res.* 36: 336-343
- De Vries, H. and Hoekstra, D. (2000): On the biogenesis of the myelin sheath: cognate polarized trafficking pathways in oligodendrocytes. *Glycoconj. J.* 17: 181-190
- De Vries, H., Schrage, C. and Hoekstra, D. (1998): An apical-type of trafficking pathway is present in cultured oligodendrocytes but the sphingolipid-enriched myelin membrane is the target of a basolateral pathway. *Mol. Biol. Cell* 9: 599-609
- De Vries, H. de Jonge, J.C., Schrage, C., van der Haar, M.E. and Hoekstra, D. (1997): Differential and cell development-dependent localization of myelin mRNAs in oligodendrocytes. *J. Neurosci. Res.* 47: 479-488
- De Winter, F., Hollmaat, A.J. and Verhaagen, J. (2002): Neurophilin and class 3 semaphorins in nervous system regeneration. *Adv. Exp. Med. Biol.* 515: 115-139
- Dimou, L., Klugmann, M., Werner, H., Jung, M., Griffiths, I.R. and Nave, K.-A. (1999): Dysmyelination in mice and the proteolipid protein gene family. *Adv. Exp. Med. Biol.* 468: 261-271
- Dotzauer, A., Brenner, M., Gebhardt, U. and Vallbracht, A. (2005): IgA-coated particles of *Hepatitis A virus* are translocated antivectorially from the apical to the basolateral site of polarized epithelial cells via the polymeric immunoglobulin receptor. *J. General Virology* 86: 2747-2751
- Drubin, D.G. and Nelson, W.J. (1996): Origins of cell polarity. *Cell* 84: 335-344.
- Erb, M., Flueck, B., Kern, F., Erne, B., Steck, A.J. and Schaeren-Wiemers, N. (2006): Unraveling the differential expression of the two isoforms of myelin-associated glycoprotein in a mouse expressing GFP-tagged S-MAG specifically regulated and targeted into the different myelin compartments. *Mol. Cell Neurosci.* 31: 613-627.
- Fasshauer, D., Sutton, R.B., Brunger, A.T. and Jahn, R. (1998): Conserved structural features of the synaptic fusion complex. *Proc. Natl. Acad. Sci. USA* 95: 15781-15786
- Fiedler, K. and Simons, K. (1995): Characterization of VIP36, an animal lectin homologous to leguminous lectins. *J. Cell Sci.* 109: 271-276

References

- Folch, J. and Lees, M. (1951): Proteolipids, a new type of tissue lipoproteins. Their isolation from brain. *J. Biol. Chem.* 191: 807-817
- Foster, L.J. and Klip, A. (2000): Mechanism and regulation of GLUT-4 vesicle fusion in muscle and fat cells. *Am. J. Physiol. Cell Physiol.* 279: C877–C890
- Frail, D.E. and Braun, P.E. (1984): Two developmentally regulated messenger RNAs differing in their coding region may exist for the myelin-associated glycoprotein. *J. Biol. Chem.* 259: 14857-14862
- Frail, D.E. and Braun, P.E. (1985): Abnormal expression of the myelin-associated glycoprotein in the central nervous system of dysmyelinating mutant mice. *J. Neurochem.* 45: 1071-175.
- Frank, M., van der Haar, M.E., Schaeren-Wiemers, N. and Schwab M.E. (1998): rMAL is a glycosphingolipid-associated protein of myelin and apical membranes of epithelial cells in kidney and stomach. *J. Neurosci.* 18: 4901-4913
- Frank, M., Schaeren-Wiemers, N., Schneider, R. and Schwab, M.E. (1999): Developmental expression pattern of the myelin proteolipid MAL indicates different functions of MAL for immature Schwann cells and in a late step of CNS myelinogenesis. *J. Neurochem.* 73: 587-597
- Frank, M. (2000): A MAL proteolipid in glycosphingolipid enriched domains: Functional implications in myelin and beyond. *Prog. Neurobiol.* 60: 531-544
- Frank, M., Atanasoski, S., Sancho, S., Magyar, J.P., Rüllicke, T., Schwab, M.E. and Sluter, U. (2000): Progressive segregation of unmyelinated axons in peripheral nerves, myelin alterations in the CNS, and cyst formation in the kidneys of myelin and lymphocyte protein-overexpressing mice. *J. Neurochem.* 75: 1927-1939
- Frost, E.E., Buttery, P.C., Milner, R. and French-Constant, C. (1999): Integrins mediate a neuronal survival signal for oligodendrocytes. *Curr. Biol.* 9: 1251-1254
- Fujita, N., Kemper, A., Dupree, J., Nakayasu, H., Bartsch, U., Schachner, M., Maeda, N., Suzuki, K. and Popko, B. (1998): The cytoplasmic domain of the large myelin-associated glycoprotein isoform is needed for proper CNS but not peripheral nervous system myelination. *J. Neurosci.* 18: 1970-1978
- Fujita, H., Tuma, P.L., Finnegan, C.M., Locca, L. and Hubbard, A.L. (1998): Endogenous syntaxin 2, 3 and 4 exhibit distinct but overlapping patterns of expression at the hepatocyte plasma membrane. *Biochem. J.* 329: 527-538

- Fuja, T.J., Ostrem, E.M., Probst-Fuja, M.N. and Titze, I.R. (2006): Different cell adhesion to vocal fold extracellular matrix constituents. *Matrix Biol.* 25: 240-251
- Garbern, J., Cambi, F., Shy, M., and Kamholz, J. (1999): The molecular pathogenesis of Pelizaeus-Merzbacher disease. *Arch. Neurol.* 56: 1210-1214
- Garcion, E., Halilagic, A., Faissner, A and ffrench-Constant, C. (2004): Generation of an environmental niche for neural stem cell development by the extracellular matrix molecule tenascin C. *Development* 131: 3423-3432
- Gerst, J.E. (2003): SNARE regulators: matchmakers and matchbreakers. *Biochim. Biophys. Acta* 1641: 1251-1264
- Gielen, E., Baron, W., van de Ven, M., Steels ,P., Hoekstra, D. and Ameloot, M. (2006): Rafts in oligodendrocytes: evidence and structure-function relationship. *Glia* 54: 499-512.
- Gillespie, C.S. Bernier, L., Brophy, P.J. and Colman, D.R. (1990): Biosynthesis of 2',3'-cyclic nucleotide 3'-phosphodiesterases. *J. Neurochem.* 54: 656-661
- Gow, A., Southwood, C.M. and Lazzarini, R.A. (1998): Disrupted proteolipid protein trafficking results in oligodendrocyte apoptosis in an animal model of Pelizaeus/Merzbacher disease. *J. Cell Biol.* 140: 825-834
- Gow, A., Southwood, C.M., Li, J.S., Pariali, M., Riordan, G.P., Brodie, S.E., Danias, J., Bronstein, J.M., Kachar, B. and Lazzarini, R.A. (1999): CNS myelin and sertoli cell tight junction are absent in Osp-claudin/11 null mice. *Cell* 99: 649-659
- Gravotta, D., Deora, A., Perret, E., Oyanadel, C., Soza, A., Schreiner, R., Gonzalez, A. and Rodriguez-Boulan, E. (2007): AP1B sorts basolateral proteins in recycling and biosynthetic routes of MDCK cells. *Proc. Natl. Acad. Sci. USA* 104: 1564-1569
- Greenfield, E.A., Reddy, J., Lees, A., Dyer, C.A., Koul, O., Nguyen, K., Bell, S., Kassam, N., Hinojoza, J., Eaton, M.J., Lees, M.B., Kuchroo, V.K. and Sobel, R.A. (2006): Monoclonal antibodies to distinct regions of human myelin proteolipid protein simultaneously recognize central nervous system myelin and neurons of many vertebrate species. *J. Neurosci. Res.* 83: 415-431
- Griffiths, I.R., Schneider, A., Anderson, T. and Nave, K.-A. (1995): Transgenic and natural mouse models of proteolipid protein (PLP)- related dysmyelination and demyelination. *Brain Pathol.* 5: 275-281
- Griffiths, I.R., Klugmann, M., Anderson, T.J., Thomson, C., Vouyiouklis, D. and Nave, K.-A. (1998): Current concepts of PLP and its role in the nervous system. *Microsc. Res. Tech.* 41: 344-538

References

- Grima, B., Zelenika, D., and Pessac, B. (1992): A novel transcript overlapping the myelin basic protein gene. *J. Neurochem.* 59: 2318-2323
- Gruenberg, J. (2003): Lipids in endocytic membrane transport and sorting. *Curr. Opin. Cell Biol.* 15: 382-388
- Gudz, T.I., Komuro, H. and Macklin, W.B. (2006): Glutamate stimulates oligodendrocyte progenitor migration mediated via an alpha v integrin/myelin proteolipid protein complex. *J. Neurosci.* 26: 2458-2466
- Gut, A., Kappeler, F., Hyka, N., Balda, M.S., Hauri, H.P. and Matter, K. (1998): Carbohydrate-mediated Golgi to cell surface transport and apical targeting of membrane proteins. *EMBO J.* 17: 1919-1929
- Hao, M. Mukherjee, S. and Maxfield, F.R. (2001): Cholesterol depletion induces large scales domain segregation in living cell membranes. *Proc. Natl. Acad. Sci. USA* 98: 13072-13077
- Hammer, J.A., and Wu, X.S. (2002): Rabs grab motors: defining the connections between GTPases and motor proteins. *Curr. Opin. Cell Biol.* 14: 69-75
- Hay, J.C. and Scheller, R.H. (1997): SNAREs and NSF in targeted membrane fusion. *Curr. Biol.* 9: 505-512
- Hepp, R., Perraul, M., Chasserot-Golaz, S., Galli, T., Aunis, D., Langley, K. and Grant, N.J. (1999): Cultured glial cells express the SNAP-25 analogue SNAP-23. *Glia* 27: 181-187
- Hirahara, Y., Bansal, R., Honke, K., Ikenaka, K. and Wada, Y. (2004): Sulfatide is a negative regulator of oligodendrocyte differentiation: development in sulfatide-null mice. *Glia* 45: 269-277
- Hirst, J., Bright, N.A., Rous, D. and Robinson, M.S. (1999): Characterization of a fourth adaptor-related protein complex. *Mol. Biol. Cell* 10: 2787-2802
- Hoek, K.S., Kidd, G.J., Carson, J.H. and Smith, R. (1998): hnRNP2 A2 selectively binds the cytoplasmic transport sequence of myelin basic protein mRNA. *Biochemistry* 37: 7021-7029
- Hoekstra, D., Tyteca, D. and van IJzendoorn S. C. D. (2004): The subapical compartment: a traffic center in membrane polarity development. *J. Cell Sci.* 117: 2183-2192
- Hoppe, C.A., Connolly, T.P. and Hubbard, A.L. (1985): Transcellular transport of polymeric IgA in the rat hepatocyte: biochemical and morphological characterization of the transport pathway. *J. Cell Biol.* 101: 2113-2123

- Hsu, S.C., Ting, A.E., Hazuka, C.D., Davanger, S., Kenny, J.W., Kee, Y. and Scheller, R.H. (1996): The mammalian brain rsec6/8 complex. *Neuron* 17: 1209-1219
- Hunziker, W., Harter, C., Matter, K. and Mellman, I. (1991): Basolateral sorting in MDCK cells requires a distinct cytoplasmic domain determinant. *Cell* 66: 907-920.
- Ikonen, E., Tagaya, M., Ullrich, O., Montecucco, C. and Simons, K. (1995): Different requirements for NSF, SNAP, and Rab proteins in apical and basolateral transport in MDCK cells. *Cell* 81: 571-580
- Ikonen, E. and Simons, K. (1998): Protein and lipid sorting from the trans-Golgi network to the plasma membrane in polarized cells. *Semin. Cell Dev. Biol.* 9: 503-509
- Iwabuchi, K., Handa, K. and Hakomori, S. (1998): Separation of "Glycosphingolipid signaling domain" from caveolin-containing membrane fraction in mouse melanoma B16 cells and its role in cell adhesion coupled with signaling. *J. Biol. Chem.* 273: 33766-33773
- Jacob, R., Heine, M., Alfalah, M. and Naim, H.Y. (2003): Distinct cytoskeletal tracks direct individual vesicle populations to the apical membrane of epithelial cells. *Curr. Biol.* 13: 607-612.
- Jahn, R. and Sudhoff, T.C., (1999): Membrane fusion and exocytosis. *Annu. Rev Biochem.* 68: 863-911
- Kaiser, C.A. and Schekman, R. (1990): Distinct sets of SEC genes govern transport vesicle formation and fusion early in the secretory pathway. *Cell* 61: 723-733
- Karcher, R.L., Deacon, S.W. and Gelfand, V. (2002): Motor/cargo interactions: the key of to transport specificity. *Trends Cell Biol.* 12: 21/27
- Karram, K., Chatterjee, N. and Trotter, J. (2005): NG2-expressing cells in the nervous system: role of proteoglycan in migration and glial-neuron interaction. *J. Anat.* 207: 735-744
- Kagawa, T., Ikenaka, K., Inoue, Y., Kuriyama, S., Tsuji, T., Nakao, J., Nakajima, K., Aruga, J., Okana, H. and Mikoshiba, K. (1994): Glial cell degeneration and hypomyelination caused by overexpression of myelin proteolipid protein gene. *Neuron* 13: 427-442
- Kee, Y., Yoo, J-S., Hazuka, C.D., Peterson, K.E., Hsu, S-C. and Scheller, R.H. (1997): Subunit structure of the mammalian exocyst complex *Proc. Natl. Acad. Sci. USA* 94: 14438-14443
- Keller, P. and Simons, K. (1998): Cholesterol is required for surface transport of influenza virus hemagglutinin. *J. Cell Biol.* 140: 1357-1367

References

- Kidd, G.J., Yadav, V.K., Huang, P., Brand, S.L., Low, S.H., Weimbs, T. and Trapp, B.D. (2006): A dual tyrosine-leucine motif mediates myelin protein P₀ targeting in MDCK cells. *Glia* 54: 135-145
- Kim, T., Fiedler, K., Madison, D.L., Krueger, W.H., and Pfeiffer, S.E (1995): Cloning and characterization of MVP17: a developmentally regulated myelin protein in oligodendrocytes. *J. Neurosci. Res.* 42: 413-22
- Kim, T. and Pfeiffer, S.E. (1999): Myelin glycosphingolipid/cholesterol-enriched microdomains selectively sequester the non-compact myelin proteins CNP and MOG. *J. Neurocytol.* 28: 281-293
- Kim, T. and Pfeiffer, S.E. (2002): Subcellular localization and detergent solubility of MVP17/rMAL, a lipid raft-associated protein in oligodendrocytes and myelin. *J. Neurosci. Res.* 69: 217-226
- Klugmann, M., Schwab, M.H., Puhlhofer, A., Schneider, A., Zimmermann, F., Griffiths, I.R. and Nave, K.-A. (1997): Assembly of CNS myelin in the absence of proteolipid protein. *Neuron* 18: 59-70
- Korngrurth, S. and Anderson, L. (1965): Localization of a basic protein of various species with the aid of fluorescence and electron microscopy. *J. Cell Biol.* 26: 157-166
- Krämer, E.M., Koch, T., Niehaus, A. and Trotter, J. (1997): Oligodendrocytes direct glycosylphosphatidylinositol-anchored proteins to the myelin sheath in glycosphingolipid-rich complexes. *J. Biol. Chem.* 272: 8937-8945
- Krämer, E.M., Schardt, A., and Nave, K.-A. (2001): Membrane traffic in myelinating oligodendrocytes. *Microsc. Res. Tech.* 52: 656-671
- Kroepfl, J.L. and Gardinier, M.V. (2001): Mutually exclusive apicobasolateral sorting of two oligodendroglial membrane proteins, proteolipid protein and myelin/oligodendrocyte glycoprotein, in Madin-Darby canine kidney cells. *J. Neurosci. Res.* 66: 1140-1148
- Kundu, A., Avalos, R.T., Sanderson, C.M. and Nayak, D.P. (1996): Transmembrane domain of influenza virus neuraminidase type II protein possesses an apical sorting signal in polarized MDCK cells. *J. Virol.* 70: 6508-6515
- Lafont, F., Verkade, P., Galli, T., Wimmer, C., Louvard, D. and Simons, K. (1999): Raft association of SNAP receptors acting in apical trafficking in Madin-Darby canine kidney cells. *Proc.Natl. Acad. Sci. USA* 96: 3734-3738.

- Lalive, P.H., Memge, T., Delarasse, C., Della Gaspera, B., Pham-Dinh, D., Villoslada, P. Von Budingen, H.C. and Genain, C.P (2006): Antibodies to native myelin oligodendrocyte glycoprotein are serologic markers of early inflammation in multiple sclerosis. *Proc. Natl. Acad. Sci. USA* 103: 2280-2285
- Larocca, J.N. and Rodriguez-Gabin, A.G. (2002): Myelin Biogenesis Vesicle Transport in Oligodendrocytes. *Neurochem. Res.* 27: 1313-1329
- Lazarovits, J. and Roth, M. (1988): A single amino acid change in the cytoplasmic domain allows the influenza virus hemagglutinin to be endocytosed through coated pits. *Cell* 53: 743-752.
- Ledesma, M.D., Brugger, B., Bunning, C., Wieland, F.T. and Dotti, C.G. (1999): Maturation of the axonal plasma membrane requires upregulation of sphingomyelin synthesis and formation of protein-lipid complexes. *EMBO J.* 18: 1761-1771
- Lee, J., Gravel, M., Zhang, R., Thibault, P and Braun, P.E. (2005): Process outgrowth in oligodendrocytes is mediated by CNP, a novel microtubule assembly myelin protein. *J. Cell Biol.* 170: 661-673
- Lees, M.B. and Brostoff, S.W. (1984): Proteins and Myelin. Pages 197-214, in Morell, P. (ed.), Myelin, Plenum Press, New York.
- Li, J., Hertzberg, E.L. and Nagy, J.I. (1997): Connexin32 in oligodendrocytes and association with myelinated fibers in mouse and rat brain. *J. Comp. Neurol.* 379: 571-91
- Lin, S., Naim, H.Y. and Roth, M.G. (1997): Tyrosine-dependent basolateral sorting signals are distinct from tyrosine-dependent internalization signals. *J. Biol. Chem.* 272: 26300-26305
- Linnington, L.B., Webb, M. and Woodhams, P.L. (1984): A novel myelin-associated glycoprotein defined by a mouse monoclonal antibody. *J. Neuroimmunol.* 6: 387-396
- Lipschutz, J.H., Guo, W., O'Brien, L.E., Nguyen, Y.H., Novick, P. and Mostov, K.E. (2000): Exocyst is involved in cystogenesis and tubulogenesis and acts by modulating synthesis and delivery of basolateral plasma membrane and secretory proteins. *Mol. Biol. Cell* 11: 4259-4275
- Lipschutz, J.H., Lingappa, V.R. and Mostov, K.E. (2003): The exocyst affects protein synthesis by acting on the translocation machinery of the endoplasmic reticulum. *J. Biol. Chem.* 278: 20954-20960
- Lisanti, M.P., Caras, I.W., Davitz, M.A. and Rodriguez-Boulan, E. (1989): A glycopospholipid membrane anchor acts as an apical targeting signal in polarized epithelial cells. *J. Cell Biol.* 109: 2145-2156

References

- Low, S.H., Chapin, S.J., Wimmer, C., Whiteheart, S.W., Komuves, L.G., Mostov, K.E. and Weimbs, T. (1998): The SNARE machinery is involved in apical plasma membrane trafficking in MDCK Cells. *J. Cell Biol.* 141: 1503-1513
- Low, S.H., Chapin, S.J., Weimbs, T., Komuves, L.G., Bennett, M.K. and Mostov, K.E. (1996): Differential localization of syntaxin isoforms in polarized Madin-Darby canine kidney cells. *Mol. Biol. Cell* 7: 2007-2018
- Ludger, J. and Galli, T. (1998): Exocytosis: SNAREs drum up! *Eur. J. Neurosci.* 10: 415-422
- Luzio, J.P., Brake, B., Banting, G., Howell, K.E., Braghetta, P. and Stanley, K.K. (1990): Identification, sequencing and expression of an integral membrane protein of the trans-Golgi network (TGN-38). *Biochem. J.* 270: 97-102
- Madison, D.L., Krueger, W.H., Cheng, D., Trapp, B. and Pfeiffer S.E. (1999): SNARE complex proteins, including the cognate pair VAMP-2 and syntaxin-4 are expressed in cultured oligodendrocytes. *J. Neurochem.* 72: 988-998
- Madore, N., Smith, K.L., Graham, C.H., Jen, A., Brady, K., Hall, S. and Morris, R. (1999): Functionally different GPI proteins are organized in different domains on the neuronal surface. *EMBO J.* 18: 6917-6926
- Maggipinto, M., Rabiner, C., Kidd, G.J., Hawkins, A.J., Smith, R. and Barbarese, E. (2004): Increased expression of the MBP mRNA binding protein HnRNP A2 during oligodendrocyte differentiation. *J. Neurosci. Res.* 75: 614-623
- Maier, O., van der Heide, T., van Dam, A.M., Baron, W., de Vries, H. and Hoekstra, D. (2005): Alteration of the extracellular matrix interferes with raft association of neurofascin in oligodendrocytes. Potential significance for multiple sclerosis? *Mol. Cell Neurosci.* 28: 390-401
- Maier, O., van der Heide, T., Johnson, R., de Vries, H., Baron, W. and Hoekstra, D. (2006): The function of neurofascin155 in oligodendrocytes is regulated by metalloprotease-mediated cleavage and ectodomain shedding. *Exp. Cell Res.* 312: 500-511.
- Marks, M.S., Ohno, H., Kirchhausen, T. and Bonifacino, J. (1997): Protein sorting by tyrosine-based signals: adapting to the Ys and wherefore. *Trends Cell Biol.* 7: 124-128
- Martin-Belmonte, F., Kremer, L., Albar, J.P., Marazuela, M. and Alonzo, M.A. (1998): Expression of the MAL gene in thyroid: the Mal proteolipid, a component of glycolipid-enriched membranes, is apically distributed in thyroid follicles. *Endocrinology* 139: 2077-2084.

- Martin-Belmonte, F., Puertollano, R., Millan, J. and Alonso, M.A. (2000): The MAL proteolipid is necessary for the overall apical delivery of membrane proteins in the polarized epithelial Madin-Darby Canine Kidney and Fischer rat Thyroid cell lines. *Mol. Biol. Cell* 11: 2033-2045
- Martin-Belmonte, F., Arvan, P. and Alonso, M.A. (2001): MAL mediates apical transport of secretory proteins in polarized epithelial Madin-Darby Canine Kidney cells. *J. Biol. Chem.* 276: 49337-49342.
- Matter, K. and Mellman, I. (1994): Mechanisms of cell polarity: sorting and transport in epithelial cells. *Curr. Opin. Cell Biol.* 6: 545-554.
- Matter, K. (2000): Sorting out the sorters. *Curr. Biol.* 10: R39-R42.
- Matute, C. and Pérez-Cerdá, F. (2005): Multiple sclerosis: novel perspectives on newly forming lesions. *Trends Neurosci.* 28: 173-175
- McCarthy, K.D. and de Vellis, J. (1980): Preparation of separate astroglial and oligodendroglial cell cultures from rat cerebral tissue. *J. Cell Biol.* 85: 890-902.
- McLoon, S.C., McLoon, L.K., Palm, S.L. and Furcht L.T. (1988): Transient expression of laminin in the optic nerve of the developing rat. *J. Neurosci.* 8: 1981-1990
- Meerson, N.R., Valerie, B., Delaunay, J-L., Slimane, T.A., Delautier, D., Lenoir, C., Trugnan, G. and Maurice, M. (2000): Intracellular traffic of the ecto-nucleotide pyrophosphodiesterase NPP3 to the apical plasma membrane of MDCK and Caco-2 cells: apical targeting occurs in the absence of N-glycosylation. *J. Cell Sci.* 113: 4193-4202.
- Mellman, I. and Warren, G. (2000): The road taken: past and future foundations of membrane traffic. *Cell* 100: 99-112.
- Millan, J., Puertollano, R., Fan, L., Rancano, C. and Alonso, M.A. (1997): The MAL proteolipid is a component of the detergent-insoluble membrane subdomains of human T-lymphocytes. *J. Biochem.* 321: 247-52
- Milner R.J., Lai C., Nave, K.-A., Lenoir, D., Ogata, J., and Sutcliffe, J.G. (1985): Nucleotide sequences of two mRNAs for rat brain myelin proteolipid protein. *Cell* 41: 931-939
- Minuk, J. and Braun, P.E. (1996): Differential intracellular sorting of the myelin-associated glycoprotein isoforms. *J. Neurosci. Res.* 44: 411-420
- Morell, P. and Radin, N.S. (1969): Synthesis of cerebroside by brain from uridine diphosphate galactose and ceramide containing hydroxy fatty acid. *Biochemistry* 8: 506-512

References

- Morita, K., Sasaki, H., Fujimoto, K., Furuse, M. and Tsukita, S. (1999): Claudin-11/OSP-based tight junctions of myelin sheaths in brain and Sertoli cells in testis. *J. Cell Biol.* 145: 579-588
- Moyer, B.D., Allan, B.D. and Balch, W.E. (2001): Rab1 interaction with a GM130 effector complex regulates COPII vesicle cis-Golgi tethering. *Traffic* 2: 268-276
- Munro, T.P., Magee, R.J., Kidd, G.J., Carson, J.H., Barbarese, E., Smith, L.M. and Smith, R. (1999): Mutational analysis of a heterogeneous nuclear ribonucleoprotein A2 response element for RNA trafficking. *J. Biol. Chem.* 274: 34389-34395
- Muth, T.R. and Caplan, M.J. (2003): Transport protein trafficking in polarized cells. *Annu. Rev. Cell Dev. Biol.* 19: 333-366
- Nave, K.-A., Lai, C., Bloom, F.E. and Milner, R.J. (1987): Splice site selection in the proteolipid protein (PLP) gene transcript and primary structure of the dm-20 protein of central nervous system myelin. *Proc. Natl. Acad. Sci. USA* 84: 5665-5669
- Nelson, W.J. and Yeaman, C. (2001): Protein trafficking in the exocytic pathway of polarized epithelial cells. *Trends Cell Biol.* 11: 483-486
- Norton, W.T. and Cammer, W. (1984): Isolation and characterization of myelin. 147-195, in Morell P. (ed.), Myelin Plenum Press, New York.
- Novick, P., Field, C. and Schekman, R. (1980): Identification of 23 complementation groups required for posttranslational events in the yeast secretory pathway. *Cell* 21: 205-215
- Novick, P., Ferro, S. and Schekman, R. (1981): Order of events in the yeast secretory pathway. *Cell* 25: 461-469
- Nussbaum, J.L. and Roussel, G. (1983): Immunocytochemical demonstration of the transport of myelin proteolipids through the Golgi apparatus. *Cell Tissue Res.* 234: 547-559
- Odorizzi, G. and Trowbridge, I.S. (1997): Structural requirements for Major histocompatibility complex class II invariant chain trafficking in polarized Madin-Darby Canine kidney cells. *J. Biol. Chem.* 272: 1157-11762
- O'Shea, K.S., Rheinheimer J.S.T. and Dixit, W.M. (1990): Deposition and role of thrombospondin in the histogenesis of the cerebellar cortex. *J. Cell Biol.* 110: 1275-1283
- Pasquini, J.M., Guarna, M.M., Besio-Moreno, M.A., Iturregui, M.T., Oteiza, P.I. and Soto, E.F. (1989): Inhibition of the synthesis of glycosphingolipids affects the translocation of proteolipid protein to the myelin membrane. *J. Neurosci. Res.* 22: 289-296

- Pedraza, L., Fidler, L., Staugaitis, S.M. and Coleman, D.R. (1997): The active transport of myelin basic protein into the nucleus suggests a regulatory role in myelination. *Neuron* 18: 579-589
- Pedraza, L. (1997): Nuclear transport of myelin basic protein. *J. Neurosci. Res.* 50: 258-264
- Pereyra, P.M., Horvath, E. and Braun, P.E. (1988): Triton X-100 extractions of central nervous system myelin indicate a possible role for the minor myelin proteins in stability of lamellae. *Neurochem. Res.* 13: 583-595
- Pfeffer, S.R. (1996): Transport vesicle docking: SNAREs and associates. *Annu. Rev. Cell Dev. Biol.* 12: 441-461
- Pfeffer, S.R. (2001): Rab GTPases: specifying and deciphering organelle identity and function. *Trends Cell Biol.* 11: 487-491
- Pfeiffer, S.E, Warrington, A.E. and Bansal, R. (1993): The oligodendrocyte and its many cellular processes. *Trends Cell Biol.* 3: 191-197
- Pham-Dinh, D., Popot, J.L., Boespflug-Tanguy, O., Landrieu, P., Deleuze, J.F., Boué, J., Jollès, P. and Dautigny, A. (1991): Pelizaeus-Merzbacher disease: a valine to phenylalanine point mutation in a putative extracellular loop of myelin proteolipid. *Proc. Natl. Acad. Sci. USA* 88: 7562-7566.
- Pike, L.J. (2006): Rafts defined: a report on the Keystone symposium on lipid rafts and cell function. *J. Lip. Res.* 47: 1597-1598
- Pimplikar, S.W. and Simons, K. (1994): Activators of protein kinase A stimulate apical but not basolateral transport in epithelial Madin-Darby canine kidney cells. *J. Biol. Chem.* 269: 19054-19059.
- Pires Neto, M.A., Braga-de-Souza, S. and Lent, R. (1999): Extracellular matrix molecules play diverse roles in the growth and guidance of central nervous system axons. *Braz. J. Med. Biol. Res.* 243: 633-638
- Polverini, E., Fasano A., Zito, F., Riccio, P. and Cavatorta, P. (1999): Confirmation of bovine myelin basic protein purified with bound lipids. *Eur. Biophys.* 28: 351-355
- Pombo, I., Rivera, J. and Blank, U. (2003): Munc18-2/ syntaxin-3 complexes are spatially separated from syntaxin-3-containing SNARE complexes. *FEBS Letters* 550: 144-148
- Privat, A., Jacque, C., Bourre, J.M., Dupouey, P. and Baumann, N. (1979): Absence of the major dense line in myelin of the mutant mouse 'shivere'. *Neurosci. Lett.* 12: 107-112

References

- Probstmeier, R., Stichel, C.C., Muller, H.W., Asou, H. and Pesheva, P. (2000): Chondroitin sulfates expressed on oligodendrocyte-derived tenascin-R are involved in neural cell recognition. Functional implications during CNS development and regeneration. *J. Neurosci. Res.* 60: 21-36
- Puertollano, R., Martin-Belmonte, F., Millan, J., de Marco, M.C., Albar, J.P., Kremer, L. and Alonso, M.A. (1999): The Mal proteolipid is necessary for normal apical transport and accurate sorting of the influenza virus Hemagglutinin in Madin-Darby Canine Kidney cells. *J. Cell Biol.* 145: 141-151
- Puertollano, R. and Alonso, M.A. (1999): MAL, an Integral element of the apical sorting machinery, is an itinerant protein that cycles between the Trans-Golgi network and the plasma membrane. *Mol. Biol. Cell* 10: 3435-3447
- Puertollano, R., Martínez-Menárguez, J.A., Batista, A., Ballesta, J. and Alonso, M.A. (2001): An intact dilysine-like motif in the carboxyl terminus of MAL is required for normal apical transport of the influenza virus hemagglutinin cargo protein in epithelial Madin-Darby canine kidney cells. *Mol. Biol. Cell* 12: 1869-1883
- Quarles, R.H., Everly, J.L. and Brady, R.O. (1972): Demonstration of a glycoprotein which is associated with a purified myelin fraction from rat brain. *Biochem. Biophys. Res. Comm.* 47: 491-497
- Quarles, R.H., Everly, J.L. and Brady, R.O. (1973): Evidence for the close association of a glycoprotein with myelin in rat. *J. Neurochem.* 21: 1177-1191
- Rangaswami, H., Bulbule, A. and Kundu, G.C. (2006): Osteopontin: role in cell signalling and cancer progression. *Trends Cell Biol.* 16: 97-87
- Ranscht, B., Clapshaw, P.A., Price, J., Noble, M. and Seifert, W. (1982): Development of oligodendrocytes and Schwann cells studied with a monoclonal antibody against galactocerebroside. *Proc Natl Acad Sci USA* 79: 2709-2713.
- Rasband, M.N., Tayler, J., Kaga, Y., Yang, Y., Lappe-Siefke, C., Nave, K.-A. and Bansal, R. (2005): CNP is required for maintenance of axon/glia interactions at nodes of Ranvier in the CNS. *Glia* 50: 86-90
- Readhead, C., Schneider, A., Griffiths, I. and Nave, K.-A. (1994): Premature arrest of myelin formation in transgenic mice with increased proteolipid protein gene dosage. *Neuron* 12: 583-595

- Relvas, J.B., Setzu, A., Baron, W., Buttery, P.C., LaFlamme, S.E., Franklin, R J.M. and ffrench-Constant, C. (2001): Expression of dominant-negative and chimeric subunits reveals an essential role for β 1 integrin during myelination. *Curr. Biol.* 11: 1039-1043
- Richter-Landsberg, C. (2000) The oligodendroglia cytoskeleton in health and disease *J. Neurosci. Res.* 59:11-18
- Richter-Landsberg, C. and Heinrich, M. (1996): OLN-93: A new permanent oligodendroglia cell line derived from primary rat brain glial cultures. *J. Neurosci. Res.* 45: 161-173
- Roberts, D.D., and Ginsburg, V.(1988): Sulfated glycolipids and cell adhesion. *Arch. Biochem. Biophys.* 267: 405-415
- Robinson, M.S. and Bonifacino, J.S. (2001): Adaptor-related proteins. *Curr. Opin. Cell Biol.* 4: 444-453
- Rodriguez-Boulan E. and Gonzalez A. (1999): Glycans in post-Golgi apical targeting: sorting signals or structural props? *Trends Cell Biol.* 9: 291–294
- Rodriguez-Boulan, E., Musch, A. and Le Bivic, A. (2004): Epithelial trafficking: new routes to familiar places. *Curr. Opin. Cell Biol.* 16: 436-442
- Rodriguez-Gabin, A.G., Cammer, M., Almazan, G., Charron, M. and Larocca, J.N. (2001): Role of rRAB22b, an oligodendrocyte protein, in regulation of transport from trans-Golgi to endocytic compartments. *J. Neurosci. Res.* 66: 1149-1160
- Rodriguez-Gabin, A.G., Almazan, G. and Larocca, J.N. (2004): Vesicle transport on oligodendrocytes: probable role of rab40c protein. *J. Neurosci. Res.* 76: 758-770
- Roper, K., Corbeil, D. and Huttner, W. B. (2000): Retention of prominin in microvilli reveals distinct cholesterol-based lipid microdomains within the apical plasma membrane of epithelial cells. *Nature Cell Biol.* 2: 582-592
- Rosenbluth, J., Nave, K.-A., Mierzwa, A. and Schiff, R. (2006): Subtle myelin defects in PLP-null mice. *Glia* 54: 172-182
- Roussel, G., Neskovic, N.M., Trifilieff, E., Artault, J.C. and Nussbaum, J.C. (1987): Arrest of proteolipid transport through the golgi apparatus in jimpy brain. *J. Neurocytol.* 16: 195-204
- Sambrook, J., Frisch, E.F. and Maniatis, T. (1989): Molecular cloning; A laboratory manual, 2nd ed, Cold Spring Harbor Laboratory Press, Cold Spring harbor, N.Y.

References

- Saravanan, K., Schaeren-Wiemers, N., Klein, D., Sandhoff, R., Schwarz, A., Yaghoofam, A., Gieselmann, V. and Franken, S. (2004): Specific downregulation and mistargeting of the lipid raft-associated protein MAL in a glycolipid disorder. *Neurobiol. Dis.* 16: 396-406
- Schaeren-Wiemers, N., Valenzuela, D.M., Frank, M. and Schwab, M.E. (1995): Characterization of a rat gene, rMAL, encoding a protein with four hydrophobic domains in central and peripheral myelin. *J. Neurosci.* 15: 5753-5764
- Schaeren-Wiemers, N., Bonnet, A., Erb, M., Erne, B., Bartsch, U., Kern, F., Mantei, N., Sherman, D. and Suter, U. (2004): The raft-associated protein MAL is required for maintenance of proper axon-glia interactions in the central nervous system. *J. Cell Biol.* 5: 731-740
- Scheiffele, P., Roth, M. G. and Simons, K. (1997): Interaction of influenza virus hemagglutinin with sphingolipid-cholesterol membrane domains via its transmembrane domain. *EMBO J.* 16: 5501-5508.
- Schneider, A., Länder, H., Schulz, G., Wolburg, H., Nave, K-A., Schulz, J.B. and Simons, M. (2005): Palmitoylation is a sorting determinant for transport to the myelin membrane. *J. Cell Sci.* 118: 2415-2423.
- Schuck, S. and Simons, K. (2004): Polarized sorting in epithelial cells: raft clustering and the biogenesis of the apical membrane. *J. Cell. Sci.* 117: 5955–5964.
- Schwob, V.S., Clark, H.B., Agrawal, D. and Agrawal, H.C. (1985): Electron microscopic immunocytochemical localization of myelin proteolipid and myelin basic protein to oligodendrocytes in rat brain during myelination. *J. Neurochem.* 45: 559-571
- Šišková, Z., Baron, W., de Vries, H. and Hoekstra, D. (2006): Fibronectin impedes “myelin” sheet-directed flow in oligodendrocytes: a role for a beta 1 integrin-mediated PKC signaling pathway in vesicular trafficking. *Mol. Cell Neurosci.* 33: 150-159.
- Simons, K. and Ikonen, E. (1997): Functional rafts in cell membranes. *Nature* 387: 569–572
- Simons, M., Krämer, E-M., Thiele, C., Stoffel, W., and Trotter, J. (2000): Assembly of myelin by association of proteolipid protein with cholesterol- and galactosylceramide-rich membrane domains. *J. Cell Biol.* 151: 143-153
- Simons, M., Krämer, E-M., Macchi, P., Rathke-Hartlieb, S., Trotter, J., Nave, K-A., and Shulz, J.B. (2002): Overexpression of the myelin proteolipid protein leads to accumulation of cholesterol and proteolipid proteins in endosomes/ lysosomes: implications for Pelizaeus-Merzbacher disease. *J. Cell Biol.* 157: 327-336

- Sinoway, M.P., Kitagawa, K., Timsit, S. Hashim, G.A. and Coleman, D.R. (1994): Proteolipid interactions in transfectants: implications for myelin assembly. *J. Neurosci. Res.* 37: 551-562
- Smith, R. (1992): The basic protein of CNS myelin: Its structure and ligand binding. *J. Neurochem.* 59: 1589-1608
- Sobel, R.A. and Mitchell, M.E. (1989): Fibronectin in multiple sclerosis lesions. *Am. J. Path.* 135: 161-168
- Spang, A. (2004): Vesicle transport: a close collaboration of Rabs and effectors. *Curr. Biol.* 14: R33-34
- Stahira, R.L. and Kim, S.U. (2001): Protein kinase C and mitogen-activated protein kinase signalling in oligodendrocytes. *Micros. Res. Tech.* 52: 680-688.
- Staugaitis, S.M., Smith, P.R. and Coleman, D.R. (1990): Expression of myelin basic protein isoforms in nonglial cells *J. Cell Biol.* 110: 1719-1727
- Sternberger, N.H., Quarles, R.H., Itoyama, Y. and Webster, H.D. (1979): Myelin-associated glycoproteins demonstrated immunocytochemically in myelin and myelin-forming cells of developing rat. *Proc. Natl. Acad. Sci. USA* 76: 1510-1514
- Stoffel, W. and Bosio, A. (1997): Myelin glycolipids and their functions. *Curr. Op. Neurobiol.* 7: 654-661
- Stuart, B.H. (1996): A Fourier transform infrared spectroscopy study of the secondary structure of myelin basic protein in reconstituted myelin. *Biochem. Mol. Biol. Int.* 38: 839-845
- Tait, S., Gunn-Moore, F., Collinson, J.M., Huang, J., Lubetzki, C., Predaza, L., Sherman, D.L., Colman, D.R. and Brophy, P.J. (2000): An oligodendrocyte cell adhesion molecule at the site of assembly of the paranodal axo-glial junction. *J. Cell Biol.* 150: 657-666
- Taylor, C.M., Coetzee, Y. and Pfeiffer, S.E. (2002): Detergent-insoluble glycosphingolipid/cholesterol microdomains of the myelin membrane. *J. Neurochem.* 81: 993-1004
- Ter Beest, M.B.A. and Hoekstra, D. (1993): Interaction of myelin basic protein with artificial membranes. Parameters governing binding, aggregation and dissociation. *Eur. J. Biochem.* 211: 689-696
- Ter Beest, M.B.A, Chapin, S.J., Avrahami, D. and Mostov, K.E. (2005): The role of syntaxins in the specificity of vesicle targeting in polarized epithelial cells. *Mol. Biol. Cell* 16: 5784-5792

References

- Terbush, D.R. and Novick, P. (1995): Sec6, Sec8, and Sec15 are components of a multisubunit complex which localizes to small bud tips in *Saccharomyces cerevisiae*. *J. Cell Biol.* 130: 299-312
- Thomas, D.C. and Roth, M.G. (1993): Vesicular stomatitis virus glycoprotein contains a dominant cytoplasmic basolateral sorting signal critically dependent upon a tyrosine. *J. Biol. Chem.* 268: 3313-3320.
- Thomas, D.C. and Roth, M.G. (1994): The basolateral targeting signal in the cytoplasmic domain of glycoprotein G from vesicular stomatitis virus resembles a variety of intracellular targeting motifs related by primary sequence but having diverse targeting activities. *J. Biol. Chem.* 269: 15732-15739.
- Ting, A. E., Hazuka, C. D., Hsu, S.-C., Kirk, M. D., Bean, A. J. and Scheller, R. H. (1995): rSec6 and rSec8, mammalian homologs of yeast proteins essential for secretion. *Proc. Natl. Acad. Sci. USA* 92: 9613-9617
- Trajkovic, K., Dhaunchak, A.S., Goncalves, J.T., Wenzel, D., Schneider, A., Bunt, G., Nave, K.-A. and Simons, M. (2006): Neuron to glia signalling triggers myelin membrane exocytosis from endosomal storage sites. *J. Cell Biol.* 172: 937-948
- Trapp, B.D., Bernier, L., Andrews, S.B. and Colman, D.R. (1988): Cellular and subcellular distribution of 2',3'-cyclic nucleotide 3'-phosphodiesterase and its mRNA in the rat central nervous system. *J. Neurochem.* 51: 859-868.
- Trapp, B.D., Andrews, S.B. Cootauco, C. and Quarles, R. (1989): The myelin-associated glycoprotein is enriched in multivesicular bodies and periaxonal membranes of actively myelinating oligodendrocytes. *J. Cell Biol.* 109: 2417-2426
- Trapp, B.D. (1990): Myelin-associated glycoprotein. Location and potential functions. *Ann. N.Y. Acad. Sci.* 605: 29-43
- Traub, L.M. and Apodaca, G. (2003): AP-1B: polarized sorting at the endosome. *Nat. Cell Biol.* 5: 1045-1047
- Tropak, M.B., Johnson, P.W., Dunn, R.J. and Roder, J.C. (1988): Differential splicing of MAG transcripts during CNS and PNS development. *Mol. Brain Res.* 4: 143-155
- Tyteca, D., van Ijendoorn, S.C.D. and Hoekstra, D. (2005): Calmodulin modulates hepatic membrane polarity by protein kinase C-sensitive steps in the basolateral endocytic pathway. *Exp. Cell Res.* 310: 293-302.

- van der Haar, M.E., Visser, H.W., de Vries, H. and Hoekstra, D. (1998): Transport of proteolipid protein to the plasma membrane does not depend on glycosphingolipid cotransport in oligodendrocyte cultures. *J. Neurosci. Res.* 51: 371-381
- Vos, J.P., Lopez-Cardozo, M. and Gadella, B.M. (1994): Metabolic and functional aspects of sulfogalactolipids. *Biochim. Biophys. Acta.* 1211: 125-149
- Weimbs, T. and Stoffel, W. (1992): Proteolipid protein (PLP) of CNS myelin: position of free, disulfide-bonded, and fatty acid thioester-linked cysteine residues and implications for the membrane topology of PLP. *Biochemistry* 31: 12289-1229
- Weimbs, T., Low, S.H., Chapin, S. and Mostov, K.E.(1997): Protein traffic in polarized epithelial cells. *Trends Cell Biol.* 7: 393-399
- Weimbs, T., Low, S.H. Chapin, Mostov, K.E., Bucher, P. and Hofmann, K. (1997): A conserved domain is present in different families of vesicular fusion proteins. *Proc. Natl. Acad. Sci. USA* 94: 3046-3051
- Weisner, S., Legate, K.R. and Fassler, R. (2005): Integrin-actin interactions. *Cell Mol. Life Sci.* 62: 1081-1099
- Yang, X. and Skoff, R.P. (1997): Proteolipid protein regulates the survival and differentiation of oligodendrocytes. *J. Neurosci.* 17: 2056-2070
- Yool, D.A., Edgar, J.M., Montague, P. and Malcolm, S. (2000): The proteolipid protein gene and myelin disorders in man and animal models. *Hum. Mol. Genet.* 9: 987-992
- Zacchetti, D., Peranen, J., Murata, M., Fiedler, K. and Simons, K. (1995): VIP17/MAL, a proteolipid in apical transport vesicles. *FEBS Lett.* 377: 465-469
- Zegers, M.M.P. and Hoekstra, D. (1997): Sphingolipid transport to the apical plasma membrane in human hepatoma cells is controlled by PKC and PKA activity: A correlation with cell polarity in hepG2 cells. *J. Cell Biol.* 138: 307-321
- Zegers, M.M.P. and Hoekstra, D. (1998): Actin filaments and microtubules are involved in different membrane traffic pathways that transport sphingolipids to the apical surface of polarized HepG2 cells. *Mol. Biol. Cell* 9: 1939-1949

References
