

University of Groningen

Improving long-term outcome of major depression in primary care

Smit, Jannetje

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2007

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Smit, J. (2007). *Improving long-term outcome of major depression in primary care: the role of recurrence prevention*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

- Ackermann RT, Williams JW jr (2002): Rational treatment choices for non-major depressions in primary care: an evidence-based review. *Journal of General and Internal Medicine* 17: 293-301.
- Agency for Health Care Policy and Research, Depression Guideline Panel (1993): Clinical Practice Guideline number 5: Depression in primary care. Volume 2: Treatment of Major Depression. Rockville, Md: US Dept of Health and Human Services, Agency for Health Care Policy and Research (AHCPR).
- American Psychiatric Association (1994): Diagnostic and Statistical Manual of mental disorders, 4th edition . Washington, DC: American Psychiatric Association.
- Andrews GA, Peters L (1998): The psychometric properties of the Composite International Diagnostic Interview. *Social Psychiatry and Psychiatric Epidemiology* 33: 80-88.
- Andrews G & Slade T (1998): Depression, dythymia and substance use disorders: sources of dissonance between ICD-10 and DSM-IV. *International Journal of Methods in Psychiatric Research* 7: 116-120.
- Andrews G, Sanderson K, Slade T & Issakidis C (2000): Why does the burden of illness persist? Relating the burden of anxiety and depression to effectiveness of treatment. *Bulletin of the World Health Organization* 78 : 446-454.
- Andrews G (2001): Should depression be managed as a chronic disease? *British Medical Journal* 322: 19-21.
- Andrews G, Henderson S & Hall W (2001): Prevalence, comorbidity , disability and service utilization. Overview of the Australian Mental Health Survey. *British Journal of Psychiatry* 178: 145-153.
- Angst J (1992): How recurrent and predictable is depressive illness? In: Long-term treatment of depression. Perspectives in psychiatry volume 3. Montgomery S, Rouillon F (editors). Chichester: J Wiley & Sons; 1-13.
- Arrindell WA & Ettema JHM (1986): SCL-90. Handleiding bij een multidimensionele psychopathologie-indicator (Nederlandse vertaling van de Manual of the Symptom Checklist, SCL-90). Lisse: Swets & Zeitlinger.
- Arroll B, MacGillivray S, Ogston S, Reid I, Sullivan F, Williams B, Crombie I (2005): Efficacy and tolerability of tricyclic antidepressants and SSRIs compared with placebo for treatment of depression in primary care: a meta-analysis. *Annals of Family Medicine* 3: 449-456.
- Badamgarav E, Weingarten SR, Henning JM, Knight K, Hasselblad V, Gano A. & Ofman J J (2003). Effectiveness of disease management programs in depression: a systematic review. *American Journal of Psychiatry* 160: 2080-2090.
- Baker R, Reddish S, Robertson N, Hearnshaw H & Jones B (2001): Randomized controlled trial of tailored strategies to implement guidelines for the management of patients with depression in general practice. *British Journal of General Practice* 51: 737-741.
- Barrett B, Byford S & Knapp M (2005): Evidence of cost-effective treatments for depression: a systematic review. *Journal of Affective Disorders* 84: 1-13.
- Bebbington PE, Brugha TS, Meltzer H, Farrell M, Ceresa C, Jenkins R, Lewis G. (2000): Psychiatric disorder and dysfunction in the UK National Survey of Psychiatric Morbidity. *Social Psychiatry and Psychiatric Epidemiology* 35:191-197.
- Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J (1961): An inventory for measuring depression. *Archives of General Psychiatry* 4: 53 - 63.

- Beck, AT, Rush AJ, Shaw BF & Emery G (1979): Cognitive therapy of depression. New York: Wiley.
- Beck AT, Steer RA & Garbin A (1988): Psychometric properties of the BDI: 25 years of evaluation. *Clinical Psychology Review* 8: 77 - 100.
- Belsher G & Costello CG (1988) : Relapse after recovery from unipolar depression: a critical review. *Psychological Bulletin* 104: 84-96.
- Berto P, D'Ilario D, Ruffo P, Di Virgilio R & Rizzo F (2000): Depression: Cost-of-illness Studies in the International Literature, a Review. *Journal of Mental Health Policy and Economics* 3: 3-10.
- Bijl RV, Ravelli A & Zessen G van (1998): Prevalence of psychiatric disorder in the general population: results of The Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Social Psychiatry and Psychiatric Epidemiology* 33: 587 - 595
- Black WC (1990): The CE plane: a graphic representation of cost-effectiveness. *Medical Decision Making* 10: 212-214.
- Blackburn IM, Eunson KM & Bishop S (1986): A two-year naturalistic follow-up of depressed patients treated with cognitive therapy, pharmacotherapy, and a combination of both. *Journal of Affective Disorders* 10: 67-75.
- Bockting CLH, Schene AH, Spinhoven P, Koeter MWJ, Wouters LW, Huyser J & Kamphuis JH (2005): Preventing relapse/recurrence in recurrent depression with cognitive therapy: a randomized controlled trial. *Journal of Consulting and Clinical Psychology* 73: 647-657.
- Boelens W (1997): Protocolaire behandeling van depressieve patiënten: Cognitieve gedragstherapie. In: Protocolaire behandelingen in de ambulante GGZ. Keijsers GPJ, van Minnen A & Hoogduin CAL(eds), p 15-42. Houten: Bohn Stafleu van Loghum.
- Bosmans JE (2006): Cost-effectiveness of treatment of depression in primary care. Thesis. VU University Medical Center: Amsterdam.
- Bower P, Byford S, Sibbald B, Ward E, King M, Lloyd M et al (2000): RCT of non-directive counselling, cbt and usual general practioner care for patients with depression. II: Cost-effectiveness. *British Medical Journal* 321: 1389-1392.
- Bower P & Sibbald B (2000) : Systematic review of the effect of on-site mental health professionals on the clinical behavior of general practioners. *British Medical Journal* 320: 614-617.
- Bower P, Rowland N, Mellor Clark J, Heywood P, Godfrey C , Hardy R (2002): Effectiveness and cost effectiveness of counselling in primary care . *Cochrane Database Syst Rev* ; CD001025.
- Bower P, Sibbald B (2002) : On site mental health workers in primary care: effects on professional practice (Cochrane review). In : *The Cochrane Library*, issue 3.
- Briggs A, Clark T, Wolstenholme J & Clarke P (2003): Missing.... presumed at random: cost-analysis of incomplete data. *Health Economics* 12: 377-392.
- Brilman EI, Brink Wvd, Giel R,Ormel J (1992): Voorkomen en lange-termijnbeloop van psychische stoornissen in de huisartspraktijk. *Tijdschrift voor Psychiatrie* 34 (6):385-399
- Brink RHS van den, Ormel J, Meer K van der, Tiemens BG, Os TWDP van, Smit A, Jenner JA (2001): Accuracy of general practioner's prognosis of the 1-year course of depression and generalised anxiety. *British Journal of Psychiatry* 178: 18 - 22.
- Brink RHS van den , Ormel J, Tiemens B, Smit A, Jenner J, Meer K van der, Os TWDP van (2002): Predictability of the 1 year course of depression and generalized anxiety in primary care. *General Hospital Psychiatry* 24:156-163.

- Broadhead WE, Blazer DG, George LK, Tse CK (1990): Depression, disability days, and days lost from work in a prospective epidemiologic survey. *JAMA* 264: 2524 – 2528.
- Brook OH, Hout H van , Stalman W, Nieuwenhuyse H, Bakker B, Heerdink E, Haan M de (2005): A pharmacy-based coaching program to improve adherence to antidepressant treatment among primary care patients. *Psychiatric Services* 56: 487-489.
- Brown C, Schulberg HC (1995): The efficacy of psychosocial treatments in primary care. A review of randomized clinical trials. *General Hospital Psychiatry* 17:414 – 424.
- Brown C, Schulberg HC, Prigerson HG (2000): Factors associated with symptomatic improvement and recovery from major depression in primary care patients. *General Hospital Psychiatry* 22: 242 - 250.
- Bryk AS & Raudenbush SW (1987): Application of hierarchical linear models to assessing change. *Psychological Bulletin* 101: 147-158.
- Buist-Bouwman MA, Graaf R de, Vollebergh WAM, Alonso J, Bruffaerts R & Ormel J (2006): Functional disability of mental disorders and comparison with physical disorders: a study among the general population of six European countries. *Acta Psychiatrica Scandinavica* 113: 492-500.
- Bush T, Ludman E, Von Korff M, Lin E (2001): Self-efficacy for managing depression: a reliable and valid self-report measure. Poster presentation American Psychological Society, Toronto, June 2001.
- Chilvers C, Dewey M, Fielding K, Gretton V , Miller P , et al (2001): Antidepressant drugs and generic counselling for treatment of depression in primary care: randomized trial with patient preference arms. *British Medical Journal* 322:772-775.
- Churchill R, Dewey M , Gretton V, Duggan C, Chilvers C, Lee A (1999): Should general practitioners refer patients with major depression to counsellors? A review of current published evidence. *British Journal of General Practice* 49: 737 – 743.
- Conradi HJ, Jonge P de, Kluiters H, Smit A, Meer K van der , Jenner JA, Van Os TWDP, Emmelkamp PMG & Ormel J: Enhanced treatment of depression in primary care: long-term outcomes of a psycho-educational prevention program alone and enriched with psychiatric consultation or cognitive behavioral therapy. In press.
- Corney RH (1981): Social work effectiveness in the management of depressed women: A clinical trial (preliminary communication). *Psychological Medicine* 11: 417 – 423.
- Corney RH (1984): The effectiveness of attached social workers in the management of depressed female patients in general practice. *Psychological Medicine ; Monograph Supplement 6; Cambridge University Press*.
- Corney RH (1987): Marital problems and treatment outcome in depressed women. A clinical trial of social work intervention. *British Journal of Psychiatry* 151: 652 – 659.
- CVZ , College voor Zorgverzekeringen (2003): Gebruik van antidepressiva 1996-2001. GIP-signaal 4. Genees- en hulpmiddelen project, Diemen.
- Davidson JR & Meltzer-Brody (1999): The underrecognition and undertreatment of depression: what is the breadth and depth of the problem? *Journal of clinical psychiatry* 60: 4-9.
- De Maeseneer J, van Driel M, Green L, van Weel C (2003): The need for research in primary care. *Lancet* 362;1314-1319.
- De Maeseneer J & De Sutter (2004) Why research in family medicine? A superfluous question. *Annals of Family Medicine* 2 (suppl): 17-22.

DeRubeis RJ, Hollon SD, Amsterdam JD, Shelton RC, Young PR, Salomon RM, O'Reardon JP, Lovett ML, Gladis MM, Brown LL & Gallop R (2005): Cognitive therapy vs. medications in the treatment of moderate to severe depression. *Archives of General Psychiatry* 62: 409-416.

Dietrich AJ, Oxman TE, Williams JW, Schulberg HC, Bruce ML, Lee PW et al (2004): Re-engineering systems for the treatment of depression in primary care: cluster randomized controlled trial. *British Medical Journal* 329: 602-605.

Dolan P (1997): Modeling valuations for EuroQol health states. *Medical Care* 35: 1095-1108.

Donald-Sherbourne C, Unutzer J, Schoenbaum M, Duan N, Lenert LA, Sturm R & Wells KB (2001): Can utility-weighted health-related quality-of-life estimates capture health effects of quality improvement for depression? *Medical Care* 39: 1246-1259.

Drummond MF, O'Brien BJ, Stoddart GL & Torrance GW (1997): Methods for the economic evaluation of Healthcare Programmes. Oxford University Press: Oxford.

Efron B & Tibshirani RJ (1993): An introduction to the bootstrap. Chapman & Hall: New York.

Ellis PM & Smith DA (2002) : Treating depression: the beyond blue guidelines for treating depression in primary care. *Medical Journal of Australia* 176: S77-S83.

Emmelkamp PMG (2004): Behavior therapy with adults. In M. Lambert (Ed.), Bergin and Garfield's Handbook of Psychotherapy and Behavior Change (5 ed., pp. 393-446). New York: Wiley.

Emanuels-Zuurveen L & Emmelkamp PMG (1996): Individual behavioural-cognitive therapy v. marital therapy for depression in maritally distressed couples. *British Journal of Psychiatry* 169: 181-188.

ESEMeD/MHEDEA 2000 consortium (2004a): Prevalence of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. *Acta Psychiatrica Scandinavica Supplement 109: 21-27.*

ESEMeD/MHEDEA 2000 consortium (2004b): Use of mental health services in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. *Acta Psychiatrica Scandinavica Supplement 109: 47-54.*

EuroQol Group (1990): EuroQol: a new facility for the measurement of health-related quality of life. *Health Policy* 16: 199-208.

Evans MD, Hollon SD, DeRubeis RJ, Piasecki JM, Grove WM, Garvey MJ, Tuason VB (1992): Differential relapse following cognitive therapy and pharmacotherapy for depression. *Archives of General Psychiatry* 49: 802-808.

Fairhurst K, Dowrick C (1996) : Problems with recruitment in a randomized controlled trial of counselling in general practice: causes and implications. *Journal of Health Services Res Policy* 1: 77-80.

Fava GA, Grandi S, Zielezny M, Rafanelli C, Canestrari R (1996): Four-year outcome for cognitive behavioral treatment of residual symptoms in major depression. *American Journal of Psychiatry* 153: 945-947.

Fava GA, Rafanelli C, Grandi S et al (1998 a): Six-year outcome for CBT of residual symptoms in major depression. *American Journal of Psychiatry* 155:1443 – 1445.

Fava GA, Rafanelli C, Grandi S, Conti S and Belluardo P (1998 b): Prevention of recurrent depression with cognitive behavioral therapy. Preliminary findings. *Archives of General Psychiatry* 55:816 – 820.

Fava GA, Ruine C, Rafanelli C, Finos L, Conti S, Grandi S (2004): Six-year outcome of cognitive behavior therapy for prevention of recurrent depression. *American Journal of Psychiatry* 161:1872-1876.

- Fava GA (2006) : The intellectual crisis of psychiatric research. *Psychotherapy and Psychosomatics* 75: 202-208.
- Fenwick E, O'Brien BJ & Briggs A (2004): Cost-effectiveness acceptability curves - facts, fallacies and frequently asked questions. *Health Economics* 13: 405-415.
- Frank E, Kupfer DJ, Perel JM, Cornes C, Jarrett DB et al (1990) : Three-year outcomes for maintenance therapies in recurrent depression. *Archives of General Psychiatry* 47:1093-1099.
- Frank E, Kupfer DJ, Wagner EF, McEachran AB, Cornes C (1991): Efficacy of interpersonal psychotherapy as a maintenance treatment of recurrent depression. Contributing factors. *Archives of General Psychiatry* 48: 1053-1059.
- Frank E, Prien RF, Jarrett RB, Keller MB, Kupfer DJ, Lavori PW, Rush AJ, Weismann MM (1991): Conceptualization and rationale for consensus definitions of terms in major depressive disorder. Remission, recovery, relapse, and recurrence. *Archives of General Psychiatry* 48: 851-55.
- Frasure-Smith N, Lesperance F, Prince RH, Verrier P, Garber RA, Juneau M, et al (1997): Randomised trial of home-based psychosocial nursing intervention for patients recovering from myocardial infarction. *Lancet* 350:473-479.
- Geddes JR, Carney SM, Davies C, Furukawa TA, Kupfer DJ, Frank E, Goodwin GM (2003) : Relapse prevention with antidepressant drug treatment in depressive disorders: a systematic review. *Lancet* 361:653-661.
- Gensichen J , Beyer M, Muth C, Gerlach FM, Von Korff M & Ormel J (2006) : Case management to improve major depression in primary health care: a systematic review. *Psychological Medicine* 36:7-14.
- Gilbody S, Whitty P, Grimshaw J & Thomas R (2003) : Educational and organizational interventions to improve the management of depression in primary care. A systematic review. *JAMA* 289: 3145 -3151.
- Greenberg PE, Kessler RCK, Birnbaum HG, Leong SA, Lowe SW, Berglund PA. & Corey-Lisle PK (2003): The economic burden of depression in the United States: how did it change between 1990 and 2000? *Journal of Clinical Psychiatry* 64: 1465-1475.
- Goldberg D, Privett M, Ustun B, Simon G, Linden M. (1998): The effects of detection and treatment on the outcome of major depression in primary care: a naturalistic study in 15 cities. *British Journal of General Practice* 48: 1840 – 1844
- Hammen, C (1997): Depression. East Sussex: Psychology Press.
- Hasin DS, Goodwin RD, Stinson FS, Grant BF (2005) : Epidemiology of major depressive disorder. Results from the national epidemiologic survey on alcoholism and related conditions. *Archives of General Psychiatry* 62: 1097-1106.
- Hays RD, Wells KB, Sherbourne C, Rogers W, Spritzer K (1995): Functioning and well-being outcomes of patients with depression compared with chronic general medical illnesses. *Archives of General Psychiatry* 52 : 11-19
- Hensley PL, Nadiga D & Uhlenhuth EH (2004): Long-term effectiveness of cognitive therapy in major depressive disorder. *Depression and Anxiety* 20: 1-7.
- Hermens MLM , Hout HPJ van, Terluin B, Ader HJ, Penninx BWJH, Marwijk HWJ, Bosmans JE, Dyck R van & Haan M de (2005) : Clinical effectiveness of counselling with or without antidepressant medication for primary care patients with minor or mild-major depression: a randomised non-inferiority trial. (submitted)

Hirschfeld RM, Keller MB, Panico A, Arons BS, Barlow D, Davidoff F, Endicott J, et al (1997): The national and manic-depressive association statement on the undertreatment of depression. *JAMA* 277:333-340.

Hodges B, Inch C & Silver I (2001) : Improving the psychiatric knowledge, skills and attitudes of primary care physicians, 1950-2000: A review. *American Journal of Psychiatry* 158:1579-1586.

Hollon SD & Beck AT (2004): Cognitive and cognitive behavioral therapies. In M.Lambert (Ed.), Bergin and Garfield's Handbook of Psychotherapy and Behavior Change (5 ed., pp. 447-492). New York: Wiley.

Hollon SD, DeRubeis RJ, Shelton RC, Amsterdam JD, Salomon RM, O'Reardon JP, Lovett ML, Young PR, Haman KL, Freeman BB & Gallop R (2005): Prevention of relapse following cognitive therapy vs. medications in moderate to severe depression. *Archives of General Psychiatry* 62: 417-422.

Holman H & Lorig K (2000): Patients as partners in managing chronic disease. *British Medical Journal* 320: 526-527.

Hout BA van, Al MJ, Gordon GS & Rutten FF (1994): Costs, effects and C/E ratios alongside a clinical trial. *Health Economics* 3: 309-319.

Hunkeler EM, Meresman JF, Hargreaves WA, Fireman B, Berman WH et al (2000): Efficacy of nurse telehealth care and peer support in augmenting treatment of depression in primary care. *Archives of Family Medicine* 9: 700 – 708.

Hunt CJ , Shepherd LM & Andrews G (2001): Do doctors know best? Comments on a failed trial. *Medical Journal of Australia* 174:144-146.

Hutschemaekers G, Tiemens B & Smit A (2006) : Weg van Professionalisering. Paradoxe bewegingen in de geestelijke gezondheidszorg. GRIP, de Gelderse Roos.

Jakob-Tacke KHM, Koopmanschap MA, Meerding WJ & Severens JL (2005): Correcting for compensating mechanisms related to productivity costs in economic evaluations of healthcare programmes. *Journal of Health Economics* 14: 435-443.

Jarrett RB, Kraft D, Doyle J, Foster BM, Eaves GG, Silver PC (2001): Preventing recurrent depression using cognitive therapy with and without a continuation phase - A randomized clinical trial. *Archives of General Psychiatry* 58: 381- 388.

Jenner JA, Meer K van der, Os TWDP van, Brink RHS van den, Tiemens BG, Ormel J (1995): Psychosociale klachten in de huisartspraktijk: een trainingsprogramma voor huisartsen. Project Interventie Studie Eerste Lijn (INSTEL). Assen: van Gorcum.

Judd LL (1997): The clinical course of unipolar major depressive disorders. Commentary. *Archives of General Psychiatry* 54:989-991.

Judd LL, Akiskal HS, Maser JD, Zeller PJ, Endicott J , Coryell et al (1998): Major depressive disorder: a prospective study of residual symptoms as predictor of rapid relapse. *Journal of Affective Disorders* 50: 97-108.

Judd LL, Paulus MJ, Schettler PJ, Akiskal HS, Endicott J, Leon AC et al (2000) : Does incomplete recovery from first lifetime episode herald a chronic course of illness? *American Journal of Psychiatry* 157: 1501-1504.

Kaasenbrood A, Werf van der B, Hanneman P (2004): Indicatie: geen behandeling. Voorspelbare schade als gevolg van psychiatrisch behandelen. *Maandblad Geestelijke Volksgezondheid* 59: 291-302.

Katon W, Gonzales J (1994): A review of randomized trials of psychiatric consultation-liaison studies in primary care. *Psychosomatics* 35: 268-278.

Katon W, Von Korff M, Lin E, Walker E, Simon GE et al (1995): Collaborative management to achieve treatment guidelines. Impact on depression in primary care. *JAMA* 273: 1026–1031.

Katon W, Robinson P, Von Korff M, Lin E, Bush T, Ludman E, Simon G, Walker E (1996): A multi-faceted intervention to improve treatment of depression in primary care. *Archives of General Psychiatry* 53: 924 – 932.

Katon W, Ludman E, Simon G, Lin E, Sullivan M, Walker E, Bush T, Von Korff M & Carter L (1996): Depression: self-care companion for better living. Group Health Cooperative of Puget Sound, Center for Health Studies, University of Washington, Seattle. Time Life Medical.

Katon W, Von Korff M, Lin E, Simon G, Walker E, Unutzer J, Bush T, Russo J, Ludman E (1999) : Stepped collaborative care for primary care patients with persistent symptoms of depression. A randomized trial. *Archives of General Psychiatry* 56: 1109 – 1115.

Katon W, Rutter C, Ludman EJ, Von Korff M, Lin E, Simon G, Bush T, Walker E, Unutzer J (2001): A randomized trial of relapse prevention in primary care. *Archives of General Psychiatry* 58:241-247.

Katon W, Russo J, Von Korff M, Lin E, Simon G, Bush T, Ludman E , Walker E (2002) : Long-term effects of a collaborative care intervention in persistently depressed primary care patients. *Journal of General Internal Medicine* 17: 741-748.

Keller MB, Lavori PW , Mueller TI , Endicott J, Coryell WH, Hirschfeld RMA , Shea MT (1992) : Time to recovery, chronicity, and levels of psychopathology in major depression. A 5-year prospective follow-up of 431 subjects. *Archives of General Psychiatry* 49: 809 – 816.

Keller MB (1994) : Depression: A long-term illness. *British Journal of Psychiatry Suppl* 165: 9 -15.

Keller MB, McCullough JP, Klein DN, Arnow B, Dunner DL , Gelenberg AJ, Markowitz JC, Nemeroff CB, Russell JM, Thase ME, Trivedi MH & Zajecka J (2000): A comparison of nefazodone, the cognitive behavioral–analysis system of psychotherapy, and their combination for the treatment of chronic depression. *The New England Journal of Medicine* 342: 1462 – 1470.

Keller MB (2003): Past, present and future directions for defining optimal treatment outcome in depression. Remission and beyond. *JAMA* 289:3152-3160.

Kennedy N & Paykel ES (2003): Residual symptoms at remission from depression : impact on long-term outcome. *Journal of Affective Disorders* 80: 135-144.

Kessing LV, Hansen MG, Andersen PK, Angst J (2004) : The predictive effect of episodes on the risk of recurrence in depressive and bipolar disorders – a life long perspective . *Acta Psychiatrica Scandinavica* 109: 339-344.

Kessler RC, Berglund P, Demler O et al (2003): The epidemiology of major depressive disorder: results from the national comorbidity survey replication (NCS-R). *JAMA* 289:3095-3105.

King M, Sibbald B, Ward E, Bower P, Lloyd M, Gabbay M , Byford S (2000): RCT of non-directive counselling, cognitive behavior therapy and usual general practioner care in the management of depression as well as mixed anxiety depression in primary care. *Health Technology Assessment* 4 (19).

Klein DN, Santiago NJ, Vivian D, Arnow BA, Blalock JA et al (2004) : Cognitive-Behavioral analysis system of psychotherapy as a maintenance treatment for chronic depression. *Journal of Consulting and Clinical Psychology* 72 (4): 681-688.

Klerman GL, Weismann MM, Rounsaville BJ, Chevron E (1984) : Interpersonal psychotherapy for depression. New York: Academic Press & Basic Books Inc, 1984.

Klinkmann MS (1997): Competing demands in psychosocial care. A model for the identification and treatment of depressive disorders in primary care. *General Hospital Psychiatry* 19: 98-111.

Kluiter H, Wiersma D (1996) : Randomized controlled trials of programmes. In: Mental Health Service Evaluation. Knudsen HC, Thornicroft G. (editors). p. 259 - 280. Cambridge: Cambridge University Press.

Koopmanschap MA, Rutten FFH, Van Ineveld BM & Van Roijen L (1995): The friction cost method for measuring indirect costs of disease. *Journal of Health Economics* 14: 171-189.

Kroenke K, Taylor-Vaisey A, Dietrich AJ , Oxman TE (2000): Interventions to improve provider diagnosis and treatment of mental disorders in primary care. *Psychosomatics* 41 : 39 – 52.

Laurant M, van Lin N, Braspenning J, Grol R (2004): Geestelijke gezondheid. In: Linden et al, Tweede Nationale studie naar ziekten en verrichtingen in de huisartspraktijk. Utrecht/ Bilthoven: NIVEL, RIVM.

Lave JR, Frank RG, Schulberg HC & Kamlet MS (1998): Cost-effectiveness of treatments for major depression in primary care practice. *Archives of General Psychiatry* 55: 645-651.

Lawrenson RA, Tyrer F, Newson RB & Farmer RDT (2000): The treatment of depression in UK general practice: selective serotonin reuptake inhibitors and tricyclic antidepressants compared. *Journal of Affective Disorders* 59: 149 – 157.

Lewinsohn PM (1975): The behavioral study and treatment of depression. In M. Hersen, R. M. Eisler, & P. M. Miller (Eds.). Progress in behavior modification, volume I, (pp. 19-64). New York: Academic Press.

Lin EHB, Katon WJ, Simon GE, Von Korff M, Bush TM et al (1997): Achieving Guidelines for the treatment of depression in primary care. Is physician education enough? *Medical Care* 35:831-842.

Lin E, Katon WJ, Von Korff M, Russo JE, Simon GE et al (1998): Relapse of depression in primary care. Rate and clinical predictors. *Archives of Family Medicine* 7: 443 – 449.

Lin EHB, Simon GE, Katon WJ, Russo JE, Von Korff M et al (1999): Can enhanced acute-phase treatment of depression improve long-term outcomes? A report of randomized trials in primary care. *American Journal of Psychiatry* 156: 643 – 645.

Linden MW van der, Westert GP, Bakker DH de, Schellevis FG (2004): Tweede Nationale studie naar ziekten en verrichtingen in de huisartspraktijk. Symptomen en ziekte in de algemene bevolking. Utrecht, Bilthoven: Nivel, RIVM.

Little RJA & Rubin DB (1987): Statistical analysis with missing data. John Wiley and Sons: New York.

Lorig K (1993) : Self-management of chronic illness: a model for the future. *Generations* 19: 11-14.

Lorig KR, Mazonson PD & Holman HR (1993) : Evidence suggesting that health education for self-management for patients with chronic arthritis has sustained health benefits while reducing health care costs. *Arthritis Rheumatism* 36: 439-446.

Lorig K (2003) : Self-management education: more than a nice extra. *Medical Care* 41: 699-701.

Ludman E, Von Korff M, Katon W, Lin E, Simon G, Walker E, Unützer J, Bush T, Wahab S (2000): The design, implementation and acceptance of a primary care-based intervention to prevent depression relapse. *International Journal Psychiatry in Medicine* 30 (3); 229 - 245.

Ludman E, Katon W, Bush T, Rutter C, Lin E, Simon G, Von Korff M & Walker E (2003): Behavioral factors associated with symptom outcomes in a primary care based depression prevention intervention trial. *Psychological Medicine* 33: 1061-1070.

Luteijn F & Bouman TK (1988): De validiteit van Beck's Depression Inventory. *Nederlands Tijdschrift voor de Psychologie* 4: 340-343.

Ma SH & Teasdale J (2004) : Mindfulness-based cognitive therapy for depression: replication and exploration of differential relapse prevention effects. *Journal of Consulting and Clinical Psychology* 72 : 31-40.

MacGillivray S, Arroll B, Hatcher S, Ogston S, Reid I, Sullivan F, Williams B & Crombie I (2003) : Efficacy and tolerability of selective serotonin reuptake inhibitors compared with tricyclic antidepressants in depression treated in primary care: systematic review and meta-analysis. *British Medical Journal* 326: 1014-1017.

Marlatt GA & Gordon JR (1985): Relapse prevention: maintenance strategies in the treatment of addictive behaviors. Guilford Press: New York.

Marwijk HWJ van, Grundmeijer HGLM, Brueren MM, Sigling HO, Stolk J & van Gelderen MG (1994): NHG standaard Depressie. *Huisarts en Wetenschap* 37:482-490.

Marwijk HW van, Bijl D , Ader HJ, Haan M de (2001) : Antidepressant prescription for depression in general practice in the Netherlands. *Pharm World Sci* 23; 2: 46 – 49.

Marwijk HWJ van, Grundmeijer HGLM, Bijl D, van Gelderen MG, de Haan M, van Weel-Baumgarten EM, Burgers JS, Boukes FS, Romeijnders ACM (2003) : NHG standaard Depressieve Stoornis, eerste herziening. *Huisarts en Wetenschap* 46:416-633.

Mauskopf J, Rutten F & Schonfeld W (2003): Cost-effectiveness league tables: valuable guidance for decision makers? *Pharmacoeconomics* 21: 991-1000.

McCrone P, Knapp M, Proudfoot J, Ryden C, Cavanagh K, Shapiro DA, Ilson S, Gray JA, Goldberg D, Mann A, Marks I, Everitt B & Tylee A (2004): Cost-effectiveness of computerised cognitive-behavioural therapy for anxiety and depression in primary care: randomised controlled trial. *British Journal of Psychiatry* 185: 55-62.

McCullough JP (2000): Treatment for chronic depression : Cognitive behavioral analysis system of psychotherapy (CBASP). New York: Guilford Press, 2000.

McIntyre RS & O' Donovan C (2004): The human cost of not achieving full remission in depression. *Canadian Journal of Psychiatry* 49 (supp) : 10-16.

Meijer SA , Zantinge EM, Verhaak PFM, Scholte M, Pols J, Valenkamp MW, Kornalijslijper M, Smeets KAPW (2003) : Evaluatie versterking eerstelijns GGZ ; tweede interim rapportage. Utrecht: NIVEL .

Melfi CA, Chawla AJ, Croghan TW, Hanna MP, Kennedy S, Sredl K (1998): The effects of adherence to antidepressant treatment guidelines on relapse and recurrence of depression. *Archives of General Psychiatry* 55:1128-1132.

Miller WR, Rollnick S (Eds); (1991): Motivational interviewing: Preparing people to change addictive behavior. New York: The Guildford Press.

Miller WR, Rollnick S (Eds); (2002): Motivational interviewing: Preparing people for change. New York: The Guildford Press.

Monroe S & Harkness KL (2005): Life stress, the 'Kindling' hypothesis, and the recurrence of depression: Considerations from a life stress perspective. *Psychological Review* 112: 417-445.

Mueller TI, Leon AC, Keller MB, Solomon DA, Endicott J, Coryell W, Warshaw M & Maser JD (1999): Recurrence after recovery from major depressive disorder during 15 years of observational follow-up. *American Journal of Psychiatry* 156:1000-1006.

Mulrow CD, Williams JW , Chiquette E, Aguilar C, Hitchcock-Noel P, et al (2000) : Efficacy of newer medications for treating depression in primary care patients. *American Journal of Medicine* 108:54-64.

Mynors-Wallis LM, Gath DH, Lloyd-Thomas AR & Tomlinson D (1995): Randomised controlled trial comparing problem solving treatment with amitriptyline and placebo for major depression in primary care. *British Medical Journal* 310: 441-445.

Mynors-Wallis LM, Davies I, Gray A, Barbour F, Gath D (1997): A randomized controlled trial and cost analysis of problem solving treatment for emotional disorders given by community nurses in primary care. *British Journal of Psychiatry* 170: 113 – 119.

Mynors-Wallis LM , Gath DH, Day A, Baker F (2000): Randomized controlled trial of problem solving treatment, antidepressant medication , and combined treatment for major depression in primary care. *British Medical Journal* 320: 26 –30.

NICE (2004): Depression –management of depression in primary and secondary care. Clinical guideline 23. London: National Institute for Clinical Excellence.

Nolen-Hoeksema S (1987): Sex differences in unipolar depression: evidence and theory. *Psychological Bulletin* 102: 252-282.

Olfson M, Fireman B, Weismann MM, Leon AC, Sheehan DV, Kathol RG, Hoven C, Farber L (1997) : Mental disorders and disability among patients in a primary care group practice. *American Journal of Psychiatry* 154: 1734-1740.

Oostenbrink JB, Koopmanschap MA & Rutten FFH (2000): Manual for cost studies, methods and standard prices for economic evaluation in healthcare (In Dutch). Healthcare Insurance Council: Amstelveen.

Oostenbrink JB & Al MJ (2005): The analysis of incomplete cost data due to dropout. *Health Economics* 14: 763-76.

Ormel J, Brink W van den, Koeter MWJ, Giel R, Meer K van der, Willige G van de, Wilink FW (1990): Recognition, management and outcome of psychological disorders in primary care: a naturalistic follow-up study. *Psychological Medicine* 20: 909-923.

Ormel J, Koeter MWJ, van den Brink W, van de Willige G (1991): Recognition, management, and course of anxiety and depression in general practice. *Archives of General Psychiatry* 48:700 – 706.

Ormel J, Oldehinkel AJ, Brillman E, van den Brink W (1993) : Outcome of depression and anxiety in primary care: a three wave, 3.5 year study of psychopathology and disability. *Archives of General Psychiatry* 50: 759-766.

Ormel J, Von Korff M, Üstün T, Pini S, Korten A & Oldehinkel T (1994) : Common mental disorders and disabilities across cultures: results from the WHO Collaborative primary care study. *JAMA* 272: 1741-1748.

Ormel J , Tiemens BG (1995) : Recognition and treatment of mental illness in primary care: towards a better understanding of a multifaceted problem. *General Hospital Psychiatry* 17 : 160-164.

Ormel J, Tiemens BG, van Os TWDP, van der Meer K, Jenner J, van den Brink RHS (1998): Betere herkenning en behandeling van depressie en soms sneller herstel na nascholing van huisartsen. *Nederlands Tijdschrift voor Geneeskunde* 142: 47, 2572- 2577.

Ormel J, Oldehinkel AJ, Nolen WA & Vollebergh (2004a) : Psychosocial disability before, during and after a major depressive episode: a 3-wave population based study of state, scar and trait effects. *Archives of General Psychiatry* 61:387-392.

Ormel J, Oldehinkel AJ, Vollebergh W. (2004). Vulnerability before, during and after a major depressive episode. A three-wave population based study. *Archives of General Psychiatry* 61: 990-996.

Os TWDP van, Ormel J, Brink RHS van den, Jenner JA, Meer K van der, Tiemens BG, van der Doorn W, Smit A, Brink W van den (1999) : Training primary care physicians improves the management of depression. *General Hospital Psychiatry* 21:168-176.

Os TWDP van, Brink RHS van den, Jenner JA , van der Meer K , Tiemens BG & Ormel J (2002) : Effects on depression pharmacotherapy of a Dutch general practioner training program. *Journal of Affective Disorders* 71:105-111.

Os TWDP van (2003): Pandora's Box. How to improve treatment and outcomes of depression in primary care? Thesis: University of Groningen, the Netherlands.

Os TWDP van, Brink RHS van den, Meer K van der, Ormel J (2006) : The care provided by general practioners for persistent depression. *European Psychiatry* 21: 87-92.

Pampallona S, Bollini P, Tibaldi G, Kupelnick B & Munizza C (2002): Patient adherence in the treatment of depression. *British Journal of Psychiatry* 180: 104-109.

Parker G (2000): Classifying depression: should paradigms lost be regained? *American Journal of Psychiatry* 157:1195-1203.

Parker G (2005): Beyond major depression. *Psychological Medicine* 35(4):467-474.

Paykel ES, Priest RG (1992): Recognition and management of depression in general practice: consensus statement . *British Medical Journal* 305: 1198–1202.

Paykel ES (1994) : Historical overview of outcome of depression.*British Journal of Psychiatry* 165: 6-8.

Paykel ES, Ramana R, Cooper Z, Hayhurst H, Kerr J & Barocka A (1995): Residual symptoms after partial remission: an important outcome in depression. *Psychological Medicine* 25: 1171-1180.

Paykel ES, Scott J, Teasdale JD, Johnson AL , Garland A, Moore R, Jenaway A, Cornwall PL, Hayhurst H, Abbott R & Pope M (1999): Prevention of relapse in residual depression by cognitive therapy – a controlled trial. *Archives of General Psychiatry* 56: 829 – 835.

Paykel ES, Scott J, Cornwall PL, Abbott R, Crane C, Pope M & Johnson AL (2005) : Duration of relapse prevention after cognitive therapy in residual depression : follow-up of controlled trial. *Psychological Medicine* 35: 59-68.

Polder JJ, Takkern J, Meerding WJ, Kommer GJ & Stokx LJ (2002): Cost of illness in the Netherlands. RIVM: Bilthoven.

Post RM & Weiss SRB (1999): Neurobiological models of recurrence in mood disorders. In: Neurobiology of mental illness (ed. D. S. Charney, EJ Nestler and BS Bunney), pp. 365-384. Oxford University Press: New York.

Regier DA, Narrow WE, Rae DS, Manderscheid RW, Locke BZ, Goodwin FK (1993): The defacto US mental and addictive disorders service system. Epidemiologic catchment area prospective 1-year prevalence rate of disorders and services. *Archives of General Psychiatry* 50: 85-94.

Revicki DA, Siddique J, Frank L, Chung JY, Green BL, Krupnick J, Prasad M & Miranda J (2005): Cost-effectiveness of evidence-based pharmacotherapy or cognitive behavior therapy compared with community referral for major depression in predominantly low-income minority women. *Archives of General Psychiatry* 62: 868-875.

Richter P, Werner J, Heerlein A, Kraus A, Sauer H (1998) : On the validity of the BDI: A review. *Psychopathology* 31: 160 – 168 .

Roland M, Torgerson DJ (1998) : What are pragmatic trials? *British Medical Journal* 316: 285.

- Rost K, Zhang M, Fortney J, Smith J, Coyne J, Smith GR, Jr. (1998) : Persistently poor outcomes of undetected major depression in primary care. *General Hospital Psychiatry* 20:12-20.
- Rost KM, Nutting PA , Smith J and Werner JJ (2000): Designing and implementing a primary care intervention trial to improve the quality and outcome of care for major depression. *General Hospital Psychiatry* 22: 66 - 77.
- Rost KM, Duan N, Rubinstein LV, Ford DE, Sherbourne C, Meredith LS, Wells KB (2001): The Quality Improvement for Depression Collaboration : general analytic strategies for a coordinated study of quality improvement in depression care. *General Hospital Psychiatry* 23: 239 - 253.
- Rost K, Nutting P, Smith J, Werner J, Duan N(2001): Improving depression outcomes in community primary care practice - a randomized trial of the QuEST intervention. *Journal General Internal Medicine* 16:143 - 149.
- Rost K, Nutting P, Smith JL, Elliott CE, Dickinson M (2002) : Managing depression as a chronic disease: a randomized trial of ongoing treatment in primary care. *British Medical Journal* 325: 934 - 939
- Roth A & Fonagy P (eds) : What works for whom? 2nd edition (2004). New York: Guildford Publications Inc.
- Sapin C, Fantino B, Nowicki ML & Kind P (2004): Usefulness of EQ-5D in assessing health status in primary care patients with major depressive disorder. *Health and Quality of Life Outcomes* 5: 2-20.
- Schaik DJF van, Klijn AFJ, Hout, HPJ van , Marwijk HWJ van , Beekman ATF, Haan M de & Dijck R van (2004): Patient preferences in the treatment of depressive disorder in primary care. *General Hospital Psychiatry* 26: 184-189.
- Schulberg HC, Block MR, Madonia MJ, Scott CP , Rodriguez E et al (1996): Treating Major Depression in Primary Care Practice. Eight-month clinical outcomes. *Archives of General Psychiatry* 53: 913 – 919.
- Schulberg HC, Block MR, Madonia MJ, Scott CP, Lave JR, Rodrigues E, Coulehan JL (1997): The 'usual care' of depression in primary care practice. *Archives of Family Medicine* 6:334-339.
- Schulberg HC, Pilkonis PA, Houck P (1998): The severity of major depression and choice of treatment in primary care practice. *Journal of Consulting and Clinical Psychology* 66 (6):932 – 938.
- Schulberg HC, Katon W , Simon GE, Rush J (1998) : Treating major depression in Primary Care Practice. An update of the Agency for Health Care Policy and Research Practice Guidelines. *Archives of General Psychiatry* 55: 1121 – 1127.
- Schulberg HC, Raue PJ, Rollman BL.(2002) : The effectiveness of psychotherapy in treating depressive disorders in primary care practice: clinical and cost perspectives. *General Hospital Psychiatry* 24 :203 – 212.
- Schwappach DL (2002): Resource allocation, social values and the QALY: a review of the debate and empirical evidence. *Health Expectations* 5: 210-222.
- Scott AIF, Freeman CPL (1992) : Edinburgh primary care depression study: treatment outcome, patient satisfaction and cost after 16 weeks. *British Medical Journal* 304: 883 – 887.
- Scott C, Tacchi MJ, Jones R, Scott J (1997): Acute and one-year outcome of a randomized controlled trial of brief cognitive therapy for major depressive disorder in primary care. *British Journal of Psychiatry* 171: 131 – 134.
- Scott J, Teasdale JD, Paykel ES , Johnson AL , Abbott R, Hayhurst H, Moore R & Garland A (2000): Effects of cognitive therapy on psychological symptoms and social functioning in residual depression. *British Journal of Psychiatry* 177: 440 – 446.

Scott J, Palmer S, Paykel E, Teasdale J & Hayhurst H (2003): Use of cognitive therapy for relapse prevention in chronic depression. Cost-effectiveness study. *British Journal of Psychiatry* 182:221-227.

Segal ZV, Williams JMG, Teasdale JD (2002): Mindfulness-based cognitive therapy for depression: a new approach to preventing relapse. New York: Guildford Press.

Sherbourne CD, Wells KB, Duan N, Miranda J, Unützer J, Jaycox I, Schoenbaum M, Meredith LS, Rubinstein LV (2001): Long-term effectiveness of disseminating quality improvement for depression in primary care. *Archives of General Psychiatry* 58:697 - 703.

Simon GE (2000): Long-term prognosis of depression in primary care. *Bulletin of the World Health Organization* 78 (4) : 439-445.

Simon GE, Von Korff M, Rutter C, Wagner E (2000) : Randomized trial of monitoring, feedback and management of care by telephone to improve treatment of depression in primary care. *British Medical Journal* 320: 550– 554.

Simon GE (2002) : Evidence review: efficacy and effectiveness of antidepressant treatment in primary care. *General Hospital Psychiatry* 24: 213 – 224.

Simon GE, Von Korff M, Ludman EJ, Katon WJ, Rutter C, Unutzer J, Lin EH, Bush T & Walker E (2002): Cost-effectiveness of a program to prevent depression relapse in primary care. *Medical Care* 40: 941-950.

Simon GE, Ludman EJ, Tutty S, Operskalski B & Von Korff M (2004): Telephone psychotherapy and telephone care management for primary care patients starting antidepressant treatment, a randomized controlled trial. *JAMA* 292:935–942.

Simpson S, Corney R, Fitzgerald P, Beecham J (2000) : A randomized controlled trial to evaluate the effectiveness and cost-effectiveness of counselling patients with chronic depression. *Health Technology Assessment* 4 (36).

Simpson S, Corney R, Fitzgerald P, Beecham J (2003) : A randomized controlled trial to evaluate the effectiveness and cost-effectiveness of psychodynamic counselling for general practice patients with chronic depression. *Psychological Medicine* 33:229-239.

Smit A, Tiemens BG, Ormel J, Kluiters H, Jenner JA, Meer K v an der, Os TWDP Van & Conradi HJ (2005). Enhanced treatment for depression in primary care: first year results on compliance, self-efficacy, the use of antidepressants and contacts with the primary care physician. *Primary Care and Community Psychiatry* 10: 39-49.

Smit A, Kluiters H, Conradi HJ, Meer K van der, Tiemens BG, Jenner JA, Os TWDP van & Ormel, J (2006). Short-term effects of enhanced treatment for depression in primary care: Results from a randomised controlled trial. *Psychological Medicine* 36: 15-26.

Smitten MH ter, Smeets RM & Brink W van den (1998) : Composite International Diagnostic Interview (CIDI). Basisversie 2.1. Lifetime manual (in Dutch). Amsterdam / Geneva: World Health Organization.

Solomon DA, Keller MB, Leon AC, Mueller TI, Lavori PW, Shea T, Coryell W, Warshaw M, Turvey C, Maser JD & Endicott J (2000) : Multiple recurrences of depressive disorder. *American Journal of Psychiatry* 157:229 - 233.

Spijker J, Bijl RV, Graaf R de, Nolen WA (2001) : Care utilization and outcome of DSM-III-R major depression in the general population. Results from the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Acta Psychiatrica Scandinavica* 104: 19 – 24.

Spijker J, Bijl RV, Graaf R de, Nolen WA (2001): Determinants of poor 1-year outcome of DSM-III-R major depression in the general population: results of the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Acta Psychiatrica Scandinavica* 103: 122 - 130 .

Spijker J, Graaf R de , Bijl RV, Beekman ATF, Ormel J & Nolen WA (2002) : Duration of major depressive episodes in the general population: results from the Netherlands mental Health survey and Incidence Study (NEMESIS) . *British Journal of Psychiatry* 181:208-213.

Stant AD, TenVergert EM, Kluiters H, Conradi HJ, Smit A & Ormel J : Cost-effectiveness of a depression recurrence prevention program for depression in primary care. *Submitted*.

Teasdale JD, Fennell MJV, Hibbert GA, Amies PL (1984): Cognitive therapy for depression in primary care . *British Journal of Psychiatry* 144:400- 406.

Teasdale JD, Segal ZV , Williams JMG, Ridgeway VA, Soulsby JM & Lau MA (2000): Prevention of Relapse / Recurrence in major depression by mindfulness-based cognitive therapy. *Journal of Consulting and Clinical Psychology* 68 : 615 – 623.

Thompson C, AL Kinmonth, L Stevens, RC Peveler et al (2000): Effects of a clinical practice guideline and practice-based education on detection and outcome of depression in primary care: Hampshire Depression Project RCT. *Lancet* 355:185-191.

Tiemens BG, Brink W van den , Meer K van der, Ormel J (1995) : Results from the Groningen centre. In: Üstun TB, Sartorius N (eds). Mental illness in general healthcare. An international study. Chichester: Wiley publishers.

Tiemens BG, Ormel J , Simon GE (1996): Occurrence, recognition and outcome of psychological disorders in primary care. *American Journal of Psychiatry* 153: 636 - 644.

Tiemens BG, Jenner JA, Kluiters H, van der Meer K (1998): Trek je eigen plan - een wegwijzer voor zelfzorg bij depressie. Afdeling Psychiatrie Rijksuniversiteit Groningen. Vertaalde en bewerkte versie van Katon et al (1996): Depression, self-care companion for better living. Group Health Cooperative of Puget Sound, Center for Health Studies, University of Washington. Seattle: Time Life Medical.

Tiemens BG, Ormel J, Jenner JA, Meer K van der, Os TWDP van, Brink RHS van den, Smit A & Brink W van den (1999): Training primary-care physicians to recognize, diagnose and manage depression: does it improve patient outcomes? *Psychological Medicine* 29 : 833-845.

Tiemens BG (1999): Management of mental health problems in primary care. The doctor, the patient and the medical model. Groningen: State University.

Üstün TB & Sartorius N (eds); (1995): Mental illness in general health care. An international study. Chichester: Wiley publishers.

Üstün TB, Ayuso-Mateos JL, Chatterji S, Mathers C & Murray CJL (2004): Global burden of depressive disorders in the year 2000. *British Journal of Psychiatry* 184: 386-392.

Verhaak PFM , Lisdonk EH, Bor JHJ , Hutschemaekers GJM (2000): GPs' referral to mental health care during the past 25 years. *British Journal of Family Practice* 50: 307-308.

Von Korff M, Gruman J , Schaefer J, Curry SJ & Wagner EH (1997): Collaborative management of chronic illness. *Annals of Internal Medicine* 127: 1097-1102.

Von Korff M & Goldberg D (2001): Improving outcomes in depression (editorial). *British Medical Journal* 323: 948-949.

Von Korff M, Glasgow RE, Sharpe M (2002) : Organising care for chronic illness. *British Medical Journal* 325: 92-94.

Vos T, Corry J, Haby MM, Carter R & Andrews G (2005) : Cost-effectiveness of cognitive-behavioural therapy and drug interventions for major depression. *The Australian and New Zealand Journal of Psychiatry* 39: 683-692.

- Wagner EH, Austin BT, VonKorff M (1996): Improving outcomes in chronic illness. *Managed Care Quarterly* 4: 12-25.
- Walker EA, Katon WJ, Russo J, Von Korff M, Lin E, Simon G, Bush T, Ludman E, Üñützer J (2000): Predictors of outcome in a primary care depression trial. *Journal of General Internal Medicine* 15: 859-867.
- Ward E, King M, Lloyd M, Bower P, Sibbald B et al (2000): RCT of non-directive counselling, cognitive behavior therapy and usual general practitioner care for patients with depression. I: Clinical effectiveness. *British Medical Journal* 321: 1383 – 1388
- Ware JE, Sherbourne CD (1992): the Rand-36 Short Form Health Status Survey. Conceptual framework and item selection. *Medical Care* 30: 473-481.
- Weel C van (2001): Context en medisch handelen. Een visie vanuit de huisartspraktijk. *Huisarts en Wetenschap* 11: 494-497.
- Weel E van-Baumgarten, Bosch W van den, Hoogen H van den, Zitman FG (1998): Ten year follow-up of depression after diagnosis in general practice. *British Journal of General Practice* 48: 1643-1646.
- Weissman MM & Markowitz JC (1994): Interpersonal psychotherapy. Current status. *Archives of General Psychiatry* 51:599-606.
- Wells KB, Stewart A, Hays RD, Burnam A, Rogers W, Daniels M, Berry S, Greenfield S & Ware J (1989): The functioning and well-being of depressed patients. Results from the Medical Outcome Study. *Journal of the American Medical Association* 262: 914-919.
- Wells KB, Sherbourne CD (1999) : Functioning and utility for current health of patients with depression or chronic medical conditions in managed, primary care practices. *Archives of General Psychiatry* 56: 897-904.
- Wells KB, Sherbourne C, Schoenbaum M, Duan N, L Meredith et al (2000) : Impact of disseminating quality improvement programs for depression in managed primary care. A randomized controlled trial. *JAMA* 283: 212 – 220.
- Wells KB, Sherbourne C, Schoenbaum M, Ettner S, Duan N, Miranda J, Unützer J, Rubinstein L (2004) : Five-year impact of Quality improvement for depression: results of a group-level randomized controlled trial. *Archives of General Psychiatry* 61:378-386.
- World Health Organization (1992): The International Classification of Diseases and related health problems, 10th revision. Geneva: WHO.
- World Health Organization (1997): The Composite International Diagnostic Interview (CIDI). Geneva: World Health Organisation.
- Wittchen HU (1994): Reliability and validity studies of the WHO-Composite International Diagnostic interview: a critical review. *Journal of Psychiatric Research* 28: 57 – 84.
- Worrall G, Angel J, Chaulk P, Clarke C & Robins M (1999): Effectiveness of an educational strategy to improve family physicians' detection and management of depression: a randomized controlled trial. *Canadian Medical Association Journal* 161:37-40.
- Zee KI van der, Sanderma R (1993) : Het meten van de algemene gezondheidstoestand met de RAND-36, een handleiding. Noordelijk Centrum voor gezondheidsvraagstukken, Rijksuniversiteit Groningen.

