

Predictors of perceived health status in patients after kidney transplantation

Predictors of perceived health status in patients after kidney transplantation. Rosenberger J, van Dijk JP, Nagyova I, Zezula I, Madarasova Geckova A, Roland R, van den Heuvel WJA, Groothoff JW. Transplantation 2006; 81: 1306-1310

Jaroslav Rosenberger ^{1,2,3}, Jitse P van Dijk ^{3,4}, Iveta Nagyova ³, Ivan Zezula ⁵, Andrea Madarasova Geckova ³, Robert Roland ^{1,2}, Wim JA van den Heuvel ^{6,7}, Johan W Groothoff ⁴

¹ Nephrology and Dialysis Center, Kosice, Slovak Republic

² Transplantation Department, University Hospital, Kosice, Slovak Republic

³ Kosice Institute for Society and Health, Institute of Social Sciences, Faculty of Science, University of PJ Safarik, Kosice, Slovak Republic

⁴ Department of Social Medicine, University Medical Center Groningen, University of Groningen, The Netherlands

⁵ Institute of Mathematics, Faculty of Science, University of PJ Safarik, Kosice, Slovak Republic

⁶ Institute for Rehabilitation Research, Hoensbroek, The Netherlands

⁷ Department of Medical Sociology, University of Maastricht, The Netherlands

Abstract

Background: Patients after kidney transplantation have decreased mortality, morbidity and better quality of life compared to people on dialysis. Major efforts are being directed towards research into graft and patient survival. Research into quality of life is less intensive. The aim of this study was to explore the predictors of perceived health status (PHS) in kidney transplant recipients.

Methods: Out of 218 patients after kidney transplantation 138 participated in the study. Linear regression analysis was performed to predict PHS, measured with the SF-36 questionnaire, in three age categories (<40, 40-59, ≥60 years). Independent variables included social support (measured with the Social Support List Discrepancies questionnaire), socio-demographic and medical variables, side-effects and compliance.

Results: Predictors of better PHS in patients <40 years were better social support ($p \leq 0.001$), lower creatinine ($p \leq 0.001$) and lower stress from adverse effects ($p \leq 0.001$). In the group of patients aged 40-59 years higher education ($p \leq 0.05$), increased housekeeping activities ($p \leq 0.01$) and lower stress from adverse effects ($p \leq 0.001$) predicted better PHS. In the last age group predictors of better PHS were lower rate of dialysis ($p \leq 0.05$) and post-transplant hospitalizations ($p \leq 0.01$), absence of diabetes mellitus ($p \leq 0.01$) and lower stress from adverse effects ($p \leq 0.05$).

Conclusions: Major differences exist in PHS among kidney transplant recipients depending on their age. Side-effects of therapy are the most important predictor of PHS for all age groups. PHS of young patients mostly depends on their renal function and their social support. Education and working activities are most important for middle-aged people whereas in older patients PHS is mostly affected by co-morbidity.

Introduction

At the present time increasing attention is being paid to quality of life evaluation. Together with mortality, morbidity and cost utilization it has become one of the major indicators of medical care ¹. Many factors are known to be predictors of morbidity and mortality in dialysed and transplanted patients. Less information is available about those variables that have an impact on quality of life.

Quality of life according to the WHO definition is a multidimensional construct comprising physical, mental, social and economic components ². Due to the complexity of each domain and depending on the method of evaluation there are many variables identified as predictors of good or bad quality of life, including demographic, psychosocial and medical parameters ³. Quality of life is an umbrella term and is often interchanged with perceived health status, functional status, self-rated health or health-related quality of life. This is the source of frequent problems in this field as definitions and measures for quality of life and the related terms are not clear. Many researchers therefore focus their research involving evaluation of the impact of disease and health on quality of life, and they use the term "health-related quality of life", while others refuse such simplification and prefer the term "perceived health status" ⁴. Both terms are interchanged frequently, meaning the same – they evaluate mainly the functional (physical) and mental status of patients as the reflection of their disease. Despite the uncertainty in definitions, health-related quality of life or perceived health status is not a mere construct devoid of clinical relevance. Recent research has shown that it is a very important predictor of other outcomes in patients with chronic renal disease ^{5,6}.

Research into perceived health status in patients after kidney transplantation is broad, but mostly limited to the description or analysis

of its determinants using univariate statistics. Studies with more proper analysis of predictive variables are unfortunately scarce. Searching the literature, we identified ten such studies⁷⁻¹⁶. Their results are heterogeneous, depending on study design, composition of the target population and explored variables. Age is the best explored variable, and the majority of studies found higher age to be the most important negative predictor of perceived health status^{8,11,13,14,16}. The results are less clear for gender, because only the study made by Wight et al., who compared cohorts of 292 dialysis and 228 transplanted patients, found female gender to be connected with worse physical functioning in kidney transplant recipients¹⁶. Race is not a frequently reported variable, as only a few multiethnic studies have been performed in this field, all showing white race associated with better results^{7,9,14}. Similarly, only a few studies have explored socio-economic variables. Hathaway et al. confirmed that higher education and employment status predict better results in the Sickness Impact Profile, Ferran's and Power's QoL Index and Adult Self-Image Scales questionnaires¹³. Crom et al. found higher education to be associated with better psychological score⁹. Griva et al. used the SF-36 questionnaire and found higher income to be a predictor of a better mental component¹¹. Two studies explored the impact of social support on perceived health status and found it to be an important predictor of psychological health status^{9,13}. Medical variables are at the center of attention and most research papers include them. Surprisingly, only two studies explored the impact of side effects of treatment and reported them to be predictors of poor health status^{12,15}. Fujisawa et al. found serum creatinine to be the only predictor of results in physical functioning, general health perceptions and vitality scales of the SF-36 questionnaire¹⁰. Many researchers have found co-morbid conditions, including anemia, diabetes mellitus, hypertension, joint and eye diseases, to be associated with poor health status^{9,11,14-16}. In addition, length of hospital stay, number of hospitalizations and time on dialysis, which reflect morbidity, have also been found to have impact on perceived health status^{9,11,13}.

The aim of this study is to explore predictors of perceived health status depending on age in patients after kidney transplantation. A wide range of variables is analyzed with the aim of merging the importance of medical and non-medical factors, including demographic, socio-economic variables, social support, dialysis and transplantation factors, side-effects of immunosuppression and compliance with the treatment. In addition, the authors discuss the clinical importance of medical and non-medical predictors of perceived health status and the consequences for possible interventions.

Materials and methods

Patients

Out of 218 patients with a functioning graft after kidney transplantation from two transplant centers in Slovakia, 208 agreed to participate in the study and 138 sufficiently completed the given questionnaires (effective response rate 63.3 %). Nonresponders did not differ significantly from the analysed group whether in age, gender, education or employment status. All patients after the third month and before 7 years post-transplantation were asked to participate, with the exception of those with severe dementia or mental retardation. The lower limit of three months was chosen as it is a common period for short-term evaluations^{8,11,13,15}. Two, five or ten years are usually used for long-term evaluations. As the aim of this study was to study approximately 200 patients, the upper limit was set at 7 years after transplantation after considering the number of patients in our transplant centres. All patients signed an informed consent before the interview. The local ethical committee approved the study.

Procedures and measures

Patients were interviewed by an independent observer in a structured interview, which focused on basic demographic information, education, employment status, house-keeping activities, family life, social activities, dialysis history, compliance with immunosuppressive treatment and adverse effects of immunosuppression.

Age, gender, education (elementary, secondary and university), employment status (employed full-time or part-time, not employed – disabled, retired or unemployed) and house-keeping activities (measured in hours per week spent shopping, cooking, cleaning, or caring for family members) were socio-demographic variables.

Patients completed the Short Form Health Survey (SF-36) and Social Support List Discrepancies questionnaire (SSL-D). The SF-36 is a 36-item questionnaire for assessment of perceived health status¹⁷. It consists of 8 sub-scales which can be combined as the physical summary component and the mental summary component. All sub-scales as well as the summary components are presented as scores between 0 and 100, with higher scores indicating better health status. The validity and reliability of SF-36 have been tested in patients with renal disease, including those after kidney transplantation^{6,10,16,18}. Skalska et al. validated the questionnaire in the Czech population¹⁹. The Cronbach α in the present sample was 0.95, while the Cronbach α for each sub-scale varied between 0.77 (for social functioning) and 0.91 (for physical functioning). For the purposes of this study only summary component scores were used.

SSL-D is an instrument designed for assessment of social support²⁰. The items in this questionnaire are grouped into 6 scales and they can

be computed into a summary score (higher score indicates better social support). The validity and reliability of this questionnaire have been previously tested in various patient populations²¹⁻²³. The Cronbach α of the questionnaire in the research we performed was 0.89. For the purposes of this study only the SSL-D sumscore was used.

Information about medical variables was mostly taken from patients' medical records. Dialysis variables were as follows: duration of dialysis period (in years), number of hospitalizations during dialysis period and number of operations during dialysis period. Transplantation variables were as follows: serum creatinine (SCr), time since transplantation (in months), number of hospitalizations after transplantation, number of operations after transplantation and type of immunosuppression protocol. Presence of diabetes mellitus was used as a measure of co-morbidity.

Patients were asked to select from the list of 16 various adverse effects of immunosuppression during the interview (malaise, pain, muscle weakness, weight gain, facial changes, depression, anxiety, sleep disorders, gingival hyperplasia, leg edemas, skin lesions, hair loss, facial edemas, sexual dysfunction, diarrhea, fragile skin). Stress from each of these adverse effects of immunosuppression was measured on a 5-point scale (0 – no stress, 1 – low stress, 2 – moderate stress, 3 – high stress, 4 – very high stress). For each patient a total score of all adverse effects was calculated as the sum of scores in all items.

Compliance with the immunosuppression therapy was measured on a 5-point scale: 1 – excellent, hardly ever modify the treatment (no more than once per last month); 2 – good, rarely modify the treatment (2-3x per last month); 3 – average, sometimes modify the treatment (once a week); 4 – fair, often modify the treatment (more than once a week), 5 – bad, always modify the treatment. Modification of treatment was explained as missing a dose, prolonging the intervals between doses by more than two hours or changing the dose of immunosuppressants. The nephrologist was interviewed about each patient's compliance with the immunosuppression therapy using the same scale as well. No specific single method was imposed on the nephrologist to identify noncompliance. Nephrologists mostly based their opinion on cyclosporin level variations or knowledge about prescribed and used immunosuppressants. Patients were considered to be compliant only if they declared their compliance by themselves as excellent, in accord with their physician's opinion.

Statistics

Stepwise multiple linear regression analysis was performed in order to find predictors of perceived health status. Physical and mental component summary scores of the SF-36 were entered as the dependent variables. The model of independent variables consisted of socio-demographic variables, social support (SSL-D sumscore), dialysis variables, transplantation

variables, co-morbidity, sum score of stress from side-effects of immunosuppression and compliance with the immunosuppressive medication. The analysis was performed with the sample divided into three age categories – patients aged less than 40 years, patients between 40 and 59 years and patients over 60 years. The cutoff values were selected based on data distribution and information from the literature. SPSS 10.0 was used for statistical analyses.

Results

Mean age was 48.4 ± 11.9 years and male/female ratio was 1.55. The majority of patients had a good graft function with median of serum creatinine $133 \mu\text{mol/l}$ (1.5 mg/dl). The predominant immunosuppressive protocol consisted of cyclosporin, mycophenolate mofetil and prednisolone. More detailed description of the patient sample is given in Tables 1 and 2.

The results of multiple linear regression analyses of predictors of perceived health status are presented in Tables 3-5. Only the models with the best explanation of variance in the mental and physical components of perceived health status are presented.

Better social support and fewer side-effects of immunosuppression predicted a better mental component of perceived health status in patients younger than 40 years. Better social support and lower SCr predicted a better physical component of perceived health status in this age group (Table 3).

Completely different results were found in the group of middle-aged patients (between 40 and 59 years). Higher education, increased housekeeping activities and lower stress from adverse effects of immunosuppression predicted better perceived health status (Table 4).

In the last age group (patients aged 60 years and older) the mental component of perceived health status was positively associated with lower post-transplant morbidity represented by lower post-transplant hospitalization rate, absence of diabetes mellitus and fewer side effects of immunosuppression. Similarly, the physical component was positively associated with lower dialysis hospitalization rate and fewer side effects of immunosuppression (Table 5).

Discussion

The aim of this study was to detect the most relevant bio-psycho-social predictors of perceived health status in kidney transplant recipients. Based on literature search and our previous experience²⁴, the presented models were created.

Age is the most important variable when considering perceived health status in kidney transplant recipients^{8,11,13,24-26}. We decided therefore

to analyse patients separately in three different age groups (less than 40, 40-59 and more than 60 years). In all groups both medical and non-medical variables play an important role.

Perceived health status in young patients is mostly influenced by their social support, renal function and side-effects of immunosuppression. The models give an excellent explanation of variance in perceived health status (60.7 % for the mental and 78.3 % for the physical component). The study by Hathaway et al. with 91 patients with mean age 39.2 years found social support to be the most important predictor of all measures in the Sickness Impact Profile questionnaire ¹³. Fujisawa et al. published their study based on 117 kidney transplant recipients with ages similar to our “young patients group”. They found SCr to be a significant predictor of better results in SF-36 subscales ¹⁰. Both of these papers support our results in this age category. It seems that for young people success of transplantation is one of the most important determinants of their health status. With good kidney function their reintegration into society is successful and they report excellent perceived health status.

The explanation of variance in perceived health status is less clear in middle-aged patients compared to the younger ones (47.4 % in the mental and 48.1 % in the physical component). Both groups also present major differences in predictors of perceived health status, as predicted by education, house-keeping activities and side effects of immunosuppression in middle-aged kidney transplant recipients. Surprisingly, lower education is a negative predictor of health status only in this age category. One possible explanation is that the sample contained only 10 university-educated persons, all middle-aged. Kidney function is not a predictor of health status in this age category, in contrast to younger patients. Adverse effects of immunosuppressive medication are the only significant medical variable. A possible explanation is that younger patients evaluate good kidney function as the most important prerequisite for long life without dialysis. Their major worries are connected with the fate of their graft ¹¹. On the other hand, older patients accept even lower kidney function as long as they feel the same benefit in terms of their health status. So they rate unpleasant effects of treatment ¹⁵ and barriers in their everyday life ¹² as more important.

The models of predictors of perceived health status among the oldest patients explained 48.2 % of variance in the mental component and 34.6 % in the physical component. Major variations in health status appear to exist in this age group and no variable is predominant. Despite the lower percentage of explained variance, perceived health status has a quite clear determinant: co-morbidity (number of pre-transplant and post-transplant hospitalizations, presence of diabetes, side-effects of immunosuppression). These data are similar to the results of the study by Siegal et al. ¹⁵.

Participants in this study were recruited from two major transplant centers in Slovakia. The sample is representative and therefore the results may be extrapolated to the whole Slovak transplant population. However, perceived health status might be influenced by many cultural, ethnic and national variables, so additional larger multicenter studies are required to verify the results and allow their extrapolation to other populations. Another limitation of the study is that it had a cross-sectional design, so the results must also be verified longitudinally.

The most important medical variable for all age groups is the presence of unpleasant side-effects of immunosuppressive treatment. The consequence of this for clinical practice is that the adverse symptoms should be constantly evaluated by the transplant team, and major efforts should aim at decreasing their severity. Another important medical variable is kidney function, but only in the group of younger patients. Worry about viability of the graft and duration of its function is the major stressor for these patients^{11,27}. The major concern in this age category lies therefore in optimizing the kidney function, but with careful balance in the treatment to minimize side-effects. On the other hand, kidney function is not the predictor of perceived health status in older patients at all, as they benefit even with „less successful transplantation“ with lower graft function. Instead, co-morbidity is more important for this age group, so the major implications for practice are connected with optimizing treatment and following up co-morbid diseases (mostly diabetes and cardiovascular morbidity). Of the non-medical variables, social support has a predominant effect on health status in younger patients, while ability to participate in everyday activities is the most important variable for older patients. These non-medical factors must be taken into account by the transplant team. Co-operation of medical staff with a psychologist, a social worker and the patient's family is necessary therefore in order to assure better quality of life as well as the patient's active rehabilitation and reintegration into society.

Acknowledgements

This work was supported by the Science and Technology Assistance Agency under contract No. APVT-20-028802.

Conflict of interests: none

References

1. Keown P. Improving quality of life-the new target for transplantation. *Transplantation* 2001 Dec;72(12 Suppl):S67-S74.

2. Kuyken W. The World Health Organization Quality of Life Assessment (WHOQOL): Position paper from the World Health Organization. 1995.
3. Edgell ET, Coons SJ, Carter WB, Kallich JD, Mapes D, Damush TM, Hays RD. A review of health-related quality-of-life measures used in end-stage renal disease. *Clin Ther* 1996 Sep;18(5):887-938.
4. Moons P. Why call it health-related quality of life when you mean perceived health status? *Eur J Cardiovasc Nurs* 2004 Dec;3(4):275-7.
5. Mapes DL, Lopes AA, Satayathum S, McCullough KP, Goodkin DA, Locatelli F, Fukuhara S, Young EW, Kurokawa K, Saito A, Bommer J, Wolfe RA, Held PJ, Port FK. Health-related quality of life as a predictor of mortality and hospitalization: the Dialysis Outcomes and Practice Patterns Study (DOPPS). *Kidney Int* 2003 Jul;64(1):339-49.
6. Phillips AL, Walker EL, Martin JE, First MR, Hanto DW, Whiting JF. Quality of life as a predictor of morbidity, mortality, and resource utilization after solid organ transplant. *Transplant Proc* 2001 Feb;33(1-2):1922.
7. Bakewell AB, Higgins RM, Edmunds ME. Does ethnicity influence perceived quality of life of patients on dialysis and following renal transplant? *Nephrol Dial Transplant* 2001 Jul;16(7):1395-401.
8. Chiu SF, Wong HS, Morad Z, Loo LH. Quality of life in cadaver and living-related renal transplant recipients in Kuala Lumpur hospital. *Transplant Proc* 2004 Sep;36(7):2030-1.
9. Crom DB, Hathaway DK, Tolley E. The use of stepwise multiple linear regression to predict quality of life after kidney transplantation. *J Transplant Coord* 1995;5(2):72-6.
10. Fujisawa M, Ichikawa Y, Yoshiya K, Isotani S, Higuchi A, Nagano S, Arakawa S, Hamami G, Matsumoto O, Kamidono S. Assessment of health-related quality of life in renal transplant and hemodialysis patients using the SF-36 health survey. *Urology* 2000 Aug;56(2):201-6.
11. Griva K, Ziegelmann JP, Thompson D, Jayasena D, Davenport A, Harrison M, Newman SP. Quality of life and emotional responses in cadaver and living related renal transplant recipients. *Nephrol Dial Transplant* 2002 Dec;17(12):2204-11.
12. Hathaway D, Winsett R, Prendergast M, Subaiya I. The first report from the patient outcomes registry for transplant effects on life (PORTEL): Differences in side-effects and quality of life by organ type, time since transplant and immunosuppressive regimens. *Clin Transplant* 2003;17(3):183-94.
13. Hathaway DK, Winsett RP, Johnson C, Tolley EA, Hartwig M, Milstead J, Wicks MN, Gaber AO. Post kidney transplant quality of life prediction models. *Clin Transplant* 1998 Jun;12(3):168-74.
14. Julius M, Hawthorne VM, Carpentier-Alting P, Kneisley J, Wolfe RA, Port FK. Independence in activities of daily living for end-stage

- renal disease patients: biomedical and demographic correlates. *Am J Kidney Dis* 1989 Jan;13(1):61-9.
15. Siegal B, Halbert RJ, McGuire MJ. Life satisfaction among kidney transplant recipients: demographic and biological factors. *Prog Transplant* 2002 Dec;12(4):293-8.
 16. Wight JP, Edwards L, Brazier J, Walters S, Payne JN, Brown CB. The SF36 as an outcome measure of services for end stage renal failure. *Qual Health Care* 1998 Dec;7(4):209-21.
 17. Ware JE, Jr., Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992 Jun;30(6):473-83.
 18. Franke GH, Reimer J, Philipp T, Heemann U. Aspects of quality of life through end-stage renal disease. *Qual Life Res* 2003 Mar;12(2):103-15.
 19. Skalska H, Sobotik Z, Jezberova D, Mares J. Use and evaluation of the Czech version of the SF-36 questionnaire self-reported health status of medical students. *Cent Eur J Public Health* 2000 May;8(2):88-93.
 20. van Sonderen FL. Measuring social support [dissertation]. Groningen: University of Groningen; 1991.
 21. Bridges KR, Sanderman R, van Sonderen E. An English language version of the social support list: preliminary reliability. *Psychol Rep* 2002 Jun;90(3 Pt 1):1055-8.
 22. Riemsma RP, Rasker JJ, Taal E, Griep EN, Wouters JM, Wiegman O. Fatigue in rheumatoid arthritis: the role of self-efficacy and problematic social support. *Br J Rheumatol* 1998 Oct;37(10):1042-6.
 23. Veenstra KM, Sprangers MA, van der Eyken JW, Taminiu AH. Quality of life in survivors with a Van Ness-Borggreve rotationplasty after bone tumour resection. *J Surg Oncol* 2000 Apr;73(4):192-7.
 24. Rosenberger J, van Dijk JP, Nagyova I, Roland R, Geckova AM, van den Heuvel WJ, Groothoff JW. Do dialysis- and transplantation-related medical factors affect perceived health status? *Nephrol Dial Transplant* 2005 Oct;20(10):2153-8.
 25. Gudex CM. Health-related quality of life in endstage renal failure. *Qual Life Res* 1995 Aug;4(4):359-66.
 26. Jofre R, Lopez-Gomez JM, Moreno F, Sanz-Guajardo D, Valderrabano F. Changes in quality of life after renal transplantation. *Am J Kidney Dis* 1998 Jul;32(1):93-100.
 27. White MJ, Ketefian S, Starr AJ, Voepel-Lewis T. Stress, coping, and quality of life in adult kidney transplant recipients. *ANNA J* 1990 Dec;17(6):421-4, 431.

Table 1 Demographic, social and social support characteristics of the patient sample (N=138)

variable		% or mean, SD (range)
gender	male	60.9 %
	female	39.1 %
age		48.4 ± 11.9 years (18-74)
	< 40 years	20.9 %
	40-59 years	60.4 %
	≥ 60 years	18.7 %
education	elementary	15.3 %
	secondary	77.4 %
	university	7.3 %
house-keeping activities		24.3 ± 18.5 hours/week
occupational status	full-time job	8.7 %
	part-time job	1.5 %
	disabled, retired or unemployed	89.8 %
family status	single	21.5 %
	married	67.4 %
	divorced	5.9 %
	widowed	5.2 %
social support (SSL-D questionnaire)	everyday emotional support	9.3 ± 2.1 (4-16)
	emotional support with problems	17.7 ± 3.8 (8-32)
	esteem support	13.0 ± 2.8 (6-24)
	social companionship	10.5 ± 2.7 (5-20)
	social support sumscore	50.4 ± 9.0 (23-92)

Table 2 Medical characteristics of the patient sample (N=138)

variable		% or mean, SD (range)
organ donor	cadaveric	96.3 %
	living	3.7 %
dialysis before transplantation	hemodialysis	74.3 %
	peritoneal dialysis	14.0 %
	both	11.7 %
time since transplantation		34.11 ± 24.6 months
primary kidney disease	glomerulonephritis	43.5 %
	tubulointerstitial nephritis	23.2 %
	polycystic kidneys	5.1 %
	diabetes mellitus	3.6 %
	congenital diseases	2.9 %
	systemic diseases, vasculitis	2.2 %
	other or unknown causes	19.6 %
immunosuppression protocol	Pred + CsA + Aza	12.8 %
	Pred + CsA	13.5 %
	Pred + CsA + MMF	40.6 %
	Pred + Tac + MMF	9.0 %
	CsA + Aza	3.8 %
	CsA + MMF	10.5 %
	Tac + MMF	0.8 %
	CsA	8.3 %
	Tac	0.8 %

Pred - prednisone, CsA - cyclosporin A, Aza - azathioprin, MMF - mycophenolate mofetil, Tac - tacrolimus

Table 3 Multiple linear regression analysis of predictors of perceived health status in age group < 40 years

predictors	R ²	mental component	R ²	physical component
		β, p		β, p
	60.7 %		78.3 %	
employment status				-0.123
social support		0.570***		0.412***
SCr				-0.876***
side effects of IS		-0.550***		

SCr - serum creatinine, IS - immunosuppression

R² - explained variance (adjusted), β - β coefficient, *** p≤0.001

Table 4 Multiple linear regression analysis of predictors of perceived health status in age group 40-59 years

predictors	R ²	mental component	R ²	physical component
		β, p		β, p
	47.4 %		48.1 %	
primary education		-0.274*		-0.300***
house keeping activities		0.238**		0.194*
side effects of IS		-0.584***		-0.605***

IS - immunosuppression

R² - explained variance (adjusted), β - β coefficient, * p≤0.05, ** p≤0.01, *** p≤0.001

Table 5 Multiple linear regression analysis of predictors of perceived health status in age group ≥ 60 years

predictors	mental component		physical component	
	R ²	β , p	R ²	β , p
	48.2 %		34.6 %	
male gender		-0.194		
elementary education		-0.369		-0.0296
house keeping activities				0.170
social support		-0.24		
D hospitalizations		-0.282		-0.453*
D operations		0.465		
SCr		-0.233		
Tx hospitalizations		-0.906**		
diabetes mellitus		-0.797**		
side effects of IS		-0.679*		-0.395*
compliance		0.485**		

SCr - serum creatinine, IS - immunosuppression, D hospitalizations - number of hospitalizations during dialysis period, D operations - number of operations during dialysis period, Tx hospitalization - number of hospitalizations after transplantation, R² - explained variance (adjusted), β - β coefficient, * $p \leq 0.05$, ** $p \leq 0.01$