

University of Groningen

## Application of a glutamate microsensor to brain tissue

Oldenziel, Weite Hendrik

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*  
2006

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Oldenziel, W. H. (2006). *Application of a glutamate microsensor to brain tissue*. s.n.

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## Chapter 9

*References*

- 1) Abbas AK, Jardemark K, Lehmann A, Weber SG, Sandberg M. Bicarbonate-sensitive cysteine induced elevation of extracellular aspartate and glutamate in rat hippocampus *in vitro*. *Neurochem. Int.* 1997; 30: 253-259.
- 2) Abel PU, Woedtke von T. Biosensors for *in vivo* glucose measurement: can we cross the experimental stage. *Biosens. Bioelectron.* 2002; 17: 1059-1070.
- 3) Alkahi KA, Tian LM. Veratradine-enhanced persistent sodium current induces bursting in CA1 pyramidal neurons. *Neurosci.* 1996; 71: 625-632
- 4) Amara SG, Fontana ACK. Excitatory amino acid transporters: keeping up with glutamate. *Neurochem. Int.* 2002; 41: 313-318.
- 5) Amine A, Mohammadi H, Bourais I, Palleschi G. Enzyme inhibition-based biosensors for food safety and environmental monitoring. *Biosens. Bioelectron.* 2006; 21: 1405-1423.
- 6) Anderson CM, Bridges RJ, Chamberlin AR, Shimamoto K, Yasuda-Kamatani Y, Swanson RA. Differing effects of substrate and non-substrate transport inhibitors on glutamate uptake reversal. *J. Neurochem.* 2001; 79: 1207-1216.
- 7) Angulo MC, Kozlov AS, Charpak S, Audinat E. Glutamate released from glial cells synchronizes neuronal activity in the hippocampus. *J. Neurosci.* 2004; 24: 6920-6927.
- 8) Anzai J, Takeshita H, Kobayashi Y, Osa T, Hoshi T. Layer-by-layer construction of enzyme multilayers on an electrode for the preparation of glucose and lactate sensors: elimination of ascorbate interference by means of an ascorbate multilayer. *Anal. Chem.* 1998; 70: 811-817.
- 9) Aoki A, Rajagopalan R and Heller A. Effect of quaternization on electron diffusion coefficients for redox hydrogels based on poly (4-vinylpyridine). *J. Phys. Chem.* 1995; 99: 5102-5110.
- 10) Araque A, Carmignoto G Haydon PG. Dynamic signalling between astrocytes and neurons. *Annu. Rev. Physiol.* 2001; 63: 795-813.
- 11) Ashton AC, Volynski KE, Lelianova VG, Orlova EV, Renterghem Van C, Caneparo M, Seagar M, Ushkaryov YA.  $\alpha$ -latrotoxin, acting via two  $Ca^{2+}$ -dependent pathways, triggers exocytosis of two pools of synaptic vesicles. *J Biol. Chem.* 2003; 276: 44695-44703
- 12) Avigliano L, Vecchini P, Sirianna P, Marcozzi G, Marchesini A, Mondovi B. A reinvestigation on the quaternary structure of ascorbate oxidase from *Cucurbita pepo medullosa*. *Mol. Cel. Biochem.* 1983; 56: 107-112.
- 13) Avshalumov MV, Chem BT, Marshall SP, Pena DM, Rice ME. Glutamate-dependent inhibition of dopamine release in striatum is mediated by a new diffusible messenger,  $H_2O_2$ . *J. Neurosci.* 2003; 23(7): 2744-2750.

- 14) Bahr BA. Long-term hippocampal slices: a model system for investigating synaptic mechanisms and pathologic processes. *J. Neurosci. Res.* 1995; 42: 294-305.
- 15) Baker A, Xi Z-X, Shen H, Swanson CJ and Kalivas PW. The origin and neuronal function of in-vivo nonsynaptic glutamate. *J. Neurosci.* 2002; 22: 9134-9141.
- 16) Baker DA, McFarland K, Lake RW, Shen H, Tang XC, Toda S, Kalivas PW. Neuroadaptions in cystine-glutamate exchange underlie cocaine relapse. *Nat. Neurosci.* 2003; 6: 743-749.
- 17) Bartlett PN, Birkin PR, Palmisano F, De Benedetto G. A study on the direct electrochemical communication between horseradish peroxidase and a poly(aniline) modified electrode. *J. Chem. Soc. Faraday Trans.* 1996; 92: 3123-3130.
- 18) Bartlett PN, Cooper JM. A review of the immobilization of enzymes in electropolymerized films. *J. Electroanal. Chem.* 1993; 362: 1-12.
- 19) Belay A, Collins A, Ruzgas T, Kissinger PT, Gorton L, Csöregi E. Redox hydrogel based bienzyme electrode for l-glutamate monitoring. *J. Pharm. Biomed. Anal.* 1999; 19: 93-105.
- 20) Belay A, Ruzgas T, Csöregi E, Moges G, Tessema M, Solomon T, Gorton L. LC-biosensor system for the determination of the neurotoxin  $\beta$ -N-oxalyl- $\alpha,\beta$ -diaminopropionic acid. *Anal. Chem.* 1997; 69: 3471-3475.
- 21) Benediktsson AM, Schachtele SJ, Green SH, Daily M. Ballistic labeling and dynamic imaging of astrocytes in organotypic hippocampal slice cultures. *J. Neurosci. Meth.* 2005; 141: 41-53.
- 22) Bensadoun A; Weinstein D. Assay of proteins in the presence of interfering materials. *Anal. Biochem.* 1976, 70 (1), 241-250
- 23) Benveniste H, Drejer J, Schousboe A, Diemer NH. Regional cerebral glucose phosphorylation and blood flow after insertion of a microdialysis fiber through the dorsal hippocampus in the rat. *J. Neurochem.* 1987; 49: 729-34.
- 24) Benveniste H, Hüttemeier PC. Microdialysis-theory and application. *Progr. Neurobiol.* 1990; 35: 195-215.
- 25) Bergles DE, Diamond JS, Jahr CE. Clearance of glutamate inside the synapse and beyond. *Curr. Opin. Neurobiol.* 1999; 9: 293-298.
- 26) Bergmeyer HU, Grassl M, Walter H.E. Methods of enzymatic analysis, 1983, 3rd ed, Vol. II, 157-158. On this article is based the "Quality control test procedure: Enzymatic assay of ascorbate oxidase", available at the website of the company Sigma.
- 27) Bezzi P, Carmignoto G, Pasti L, Vesce S, Rossi D, Rizzini BL, Pozzan T, Volterra A. Prostaglandins stimulate calcium-dependent glutamate release in astrocytes. *Nature* 1998; 391: 281-285.

- 28) Bezzi P, Gundersen V, Galbete JL, Seifert G, Steinhauser C, Pilate E, Volterra A. Astrocytes contain a vesicular compartment that is competent for regulated exocytosis of glutamate. *Nature Neurosci.* 2004; 7: 613-620.
- 29) Bezzi P, Volterra A. A neuron-glia signalling network in the active brain. *Curr. Opin. Neurobiol.* 2001; 11: 387-394.
- 30) Bianchi L, Colivicchi MA, Bolam JP, Della Corte L. The release of amino acids from rat neostriatum and substantia nigra *in vivo*: a dual microdialysis probe analysis. *Neurosci.* 1998; 87: 171-180.
- 31) Bianchi L, Della Corte L, Tipton KF. Simultaneous determination of basal and evoked output levels of aspartate, glutamate, taurine, and 4-aminobutyric acid during microdialysis and from superfused brain slices. *J. Chrom. B. Biomed. Sci. Appl.* 1999; 723: 47-59.
- 32) Bindra DS, Zhang Y, Wilson GS, Sternberg R, Thévenot DR, Moatti D, Reach G. Design and *in vitro* studies of a needle-type glucose sensor for subcutaneous monitoring. *Anal. Chem.* 1991; 63:1692-6.
- 33) Binns BC, Huang Y, Goettl VM, Hackshaw KV, Stephens jr. RL. Glutamate uptake is attenuated in spinal deep dorsal and ventral horn in the rat spinal nerve ligation model. *Brain Res.* 2005, 1041: 38-47.
- 34) Bo P, Soragna D, Specchia C, Chimento P, Favalli L. Quantified EEG analysis monitoring in a novel model of general anaesthesia in rats. *Brain Res. Protoc.* 2003; 11: 155-161.
- 35) Bonanno G, Raiteri M. Release-regulating presynaptic heterocarriers. *Prog. Neurobiol.* 1994; 44: 451-462.
- 36) Borland LM, Guoyue S, Yang H, Michael AC. Voltammetric study of extracellular dopamine near microdialysis probes acutely implanted in the striatum of the anesthetized rat. *J. Neurosci. Meth.* 2005; 146: 149-158.
- 37) Brahma B, Forman RE, Stewart EE, Nicholson C, Rice ME. Ascorbate inhibits edema in brain slices. *J. Neurochem.* 2000: 74: 1263-1270.
- 38) Bruno JP, Gerhardt GA, Gash CR, Zmarowski A, Martin B, Burmeister JJ, Huettl P, Pomerleau F. Second-to-second monitoring of cholinergic transmission. Monitoring Molecules in Neuroscience; 11<sup>th</sup> Int. Conf; Ed: Di Chiara G, Carboni E, Valentini V, Acquas E, Bassareo V, Cadoni C. 2006; 37-39.
- 39) Burmeister JJ, Moxon K, Gerhardt GA. Ceramic-based multisite microelectrodes for the detection and elimination of interferences from the measurement of l-glutamate and other analytes. *Anal. Chem.* 2000; 72: 187-192.

- 40) Burmeister JJ, Gerhardt GA. Self-referencing ceramic based multisite microelectrodes for the detection and elimination of interferences from the measurement of l-glutamate and other analytes. *Anal. Chem.* 2001; 73: 1037-1042.
- 41) Burmeister JJ, Pomerleau F, Palmer M, Day BK, Huettl P and Gerhardt GA. Improved ceramic-based multisite microelectrode for rapid measurement of l-glutamate in the CNS. *J. Neurosci. Meth.* 2002; 119: 163-171.
- 42) Burmeister JJ, Palmer M, Gerhardt GA. Ceramic-based multisite electrode array for rapid choline measures in brain tissue. *Anal. Chim. Acta.* 2003; 481: 65-74.
- 43) Burmeister JJ, Pomerleau F, Huettl P, Gerhardt GA. Simultaneous oxygen and glutamate measurements with ceramic-based microelectrode arrays in CNS. *Monitoring Molecules in Neuroscience*; 11<sup>th</sup> Int. Conf; Ed: Di Chiara G, Carboni E, Valentini V, Acquas E, Bassareo V, Cadoni C. 2006; 289-291.
- 44) Butcher SP, Lazarewicz JW, Hamberger A. *In vivo* microdialysis studies on the effects of decortication and excitotoxic lesions on kainic acid-induced calcium fluxes, and endogenous amino acid release, in the rat striatum. *J Neurochem.* 1987; 49: 1355-60.
- 45) Cahill PS, Wightman RM. Simultaneous amperometric measurement of ascorbate and catecholamine secretion from individual bovine adrenal medullary cells. *Anal. Chem.* 1995; 67: 2599-2605.
- 46) Calvo EJ, Etchenique R, Pietrasante L, Wolosiuk A. Layer-by-layer self-assembly of glucose oxidase and Os(Bpy)<sub>2</sub>CIPyCH<sub>2</sub>NH-poly(Allylamine) bioelectrode. *Anal. Chem.* 2001; 73: 1161-1168.
- 47) Casella L, Monzani E, Santagostini L, Gioia de L, Gullotti M, Fantucci P, Beringhelli T, Marchesini A. Inhibitor binding studies on ascorbate oxidase. *Coord. Chem. Rev.* 1999, 185-186: 619-628.
- 48) Castillo J, Gáspár S, Leth S, Niculescu M, Mortari A, Bontidean I, Soukharev v, Dorneanu SA, Ryabov AD, Csöregi E. Biosensors for life quality. Design, development and applications. *Sens. Actuators B* 2004; 102: 179-194.
- 49) Castillo J, Isik S, Blöchl A, Pereira-Rodrigues N, Bedioui F, Csöregi E, Schuhmann W, Oni J. Simultaneous detection of the release of glutamate and nitric oxide from adherently growing cells using an array of glutamate and nitric oxide selective electrodes. *Biosens. Bioelectron.* 2005; 20: 1559-65.
- 50) Cavalier P and Atwell D. Tonic release of glutamate by a DIDS-sensitive mechanism in rat hippocampal slices. *J. Physiol.* 2005; 564: 397-410.
- 51) Cavalier P, Hamann M, Rossi D, Mobbs P, Attwell D. Tonic excitation and inhibition of neurons: ambient transmitter sources and computational consequences. *Prog. Biophys. Mol. Biol.* 2005; 87: 3-16.

- 52) Chaubey A, Malhotra BD. Mediated biosensors. *Biosens. Bioelectron.* 2002; 17: 441-456.
- 53) Chen H, Sun D. The role of Na-K-Cl co-transporter in cerebral ischemia. *Neurol. Res.* 2005; 27(3): 280-286.
- 54) Chen T, Friedman KA and Heller A. *In situ* assembled mass-transport controlling micromembranes and their application in implanted amperometric glucose sensors. *Anal. Chem.* 2000; 72: 3757-3763.
- 55) Chen X, Matsumoto N, Hu Y, Wilson GS. Electrochemically mediated electrodeposition/electropolymerization to yield a glucose microbiosensor with improved characteristics. *Anal. Chem.* 2002; 74: 368-372.
- 56) Clapp-Lilly KL, Roberts RC, Duffy LK, Irons KP, Hu Y, Drew KL. An ultrastructural analysis of tissue surrounding a microdialysis probe. *J. Neurosci. Meth.* 1999; 90(2): 129-142.
- 57) Clark LC jr., Lyons C. Electrode systems for continuous monitoring in cardiovascular surgery. *Ann. N.Y. Acad. Sci.* 1962; 102: 29-45.
- 58) Clark LC Jr, Noyes LK, Spokane RB, Sudan R, Miller ML. Long-term implantation of voltammetric oxidase/peroxide glucose sensors in the rat peritoneum. *Meth. Enzymol.* 1988;137:68-89.
- 59) Coche-Guerente L, Cosnier S, Labbe L. Sol-Gel derived composite materials for the construction of oxidase/peroxidase mediatorless biosensors. *Chem. Mater.* 1997; 9: 1348-1352.
- 60) Conti F, Weinberg RJ. Shaping excitation at glutamergic synapses. *TINS* 1999; 22 (10): 451-458.
- 61) Cooper JR, Bloom FE, Roth RH. The biochemical basis of neuropharmacology. 1996, 7<sup>th</sup> ed., Oxford University Press.
- 62) Cosnier S. Biomolecule immobilization on the electrode surface by entrapment or attachment to electrochemically polymerized films. A review. *Biosens. Bioelectron.* 1999; 14: 443-456.
- 63) Coyle JT, Leski ML, Morrison JH. The diverse roles of glutamic acid in brain signal transduction, in: *Neuropsychopharmacology: The fifth generation of progress*, 2002, ed: Davis KL, Charney D, Coyle JT, Nemeroff C; on-line version: [www.acnp.org](http://www.acnp.org).
- 64) Coyle JT, Puttfarcken P. Oxidative stress, glutamate, and neurodegenerative disorders. *Science* 1993; 262: 689-95.
- 65) Csöregi E, Jönsson-Petterson G, Gorton L. Mediatorless electrocatalytic reduction of hydrogen peroxide at graphite electrodes chemically modified with peroxidases. *J. Biotechnol.* 1993; 30: 315-317.

- 66) Csöregi E, Schmidtke DW, Heller A. Design and optimization of a selective subcutaneously implantable glucose electrode based on "wired" glucose oxidase. *Anal. Chem.* 1995; 67: 1240-1244.
- 67) Cui J, Kulagina NV and Michael AC. Pharmacological evidence for the selectivity of *in vivo* signals obtained with enzyme-based electrochemical sensors. *J. Neurosci. Meth.* 2001; 104: 183-189.
- 68) D'Andrea G, Maccarrone M, Oratore A, Avigliano L, Messerschmidt A. Kinetic features of ascorbic acid oxidase after partial deglycation. *Biochem. J.* 1989, 264: 601-604.
- 69) Dale ND, Hatz S, Tian F, Llaudet E. Listening to the brain: microelectrode biosensors for neurochemicals. *Tr. Biotech.* 2005; 23: 420-428.
- 70) Dalton EF, SurrIDGE NA, Jernigan JC, Wilbourn KO, Facci JS, Murray RW. Charge transport in electroactive polymers consisting of fixed molecular redox sites. *Chem. Phys.* 1990; 141: 143-157.
- 71) Danbolt NC. Glutamate uptake. *Prog. Neurobiol.* 2001; 65: 1-105.
- 72) Danilowicz C, Cortón E, Battaglini F, Calvo EJ. An Os(bpy)<sub>2</sub>Cl PyCH<sub>2</sub> NH-poly(allylamine) hydrogel mediator for enzyme wiring at electrodes. *Electrochim. Acta* 1998; 43: 3525-3531.
- 73) Dawson CR, Strothkamp KG, Krul KG. Ascorbate oxidase and related copper proteins. *Ann. N.Y. Acad. Sci.* 1975; 258; 209–220.
- 74) Day BK, Pomerleau F, Burmeister JJ, Huettl P, Gerhardt GA. Microelectrode array studies of basal and potassium-evoked release of l-glutamate in the anesthetized rat brain. *J. Neurochem.* 2006; 96: 1625-35.
- 75) Dean JB, Mulkey DK, Garcia III AJ, Putnam RW, Henderson III R. Neuronal sensitivity to hyperoxia, hypercapnia, and inert gases at hyperbaric pressures. *J. Appl. Physiol.* 2003; 95: 883-909.
- 76) Degani Y, Heller A. Direct electrical communication between chemically modified enzymes and metal electrodes. I. Electron transfer from glucose oxidase to metal electrodes via electron relays, bound covalently to the enzyme. *J. Phys. Chem.* 1987; 91: 1285-1289.
- 77) Del Arco A, Segovia G, Fuxe K, Mora F. Changes in dialysate concentrations of glutamate and GABA in the brain: an index of volume transmission mediated actions. *J. Neurochem.* 2003; 85: 23-33.
- 78) Dequaire M, Heller A. Screen Printing of Nucleic Acid Detecting Carbon Electrodes. *Anal. Chem.* 2002; 74: 4370-4377.
- 79) Doherty AP, Stanley MA, Vos JG. Electrocatalytic oxidation of ascorbic acid at [osmium(2,2'-bipyridyl)<sub>2</sub>-poly-4-vinylpyridine)<sub>10</sub>Cl]Cl modified electrodes; Implications



- for the development of biosensors based on osmium-containing redox relays. *Analyst*. 1995; 120: 2371-2376.
- 80) Drew KL, Pehek EA, Rasley BT, Ma YL, Green TK. Sampling glutamate and GABA with microdialysis: suggestions on how to get the dialysis membrane closer to the synapse. *J Neurosci. Meth.* 2004; 140: 127-131.
- 81) Duan S, Anderson CM, Keung EC, Chen Y, Chen Y, Swanson RA. P2X7 receptor-mediated release of excitatory amino acids from astrocytes. *J. Neurosci.* 2003; 23: 1320-1328.
- 82) Elmgren M, Nordling M, Lindquist SE. The influence of flow rate on biosensors based on redox enzymes incorporated in a redox polymer mounted in a thin-layer flow cell. *Anal. Biochem.* 1993; 215: 261-265.
- 83) Enkvist MO, McCarthy KD. Astroglial gap junction communication is increased by treatment with either glutamate or high K<sup>+</sup> concentration. *J. Neurochem.* 1994; 62: 489-495.
- 84) Fei J, Wu Y, Ji X, Wang J, Hu S, Gao Z. An amperometric biosensor for glucose based on electrodeposited redox polymer/glucose oxidase film on a gold electrode. *Anal. Sci.* 2003; 19: 1259-1263.
- 85) Feldman B, Brazg R, Schwartz S, Weinstein R. A continuous glucose sensor based on wired enzyme technology – Results from a 3-day trial in patients with type 1 diabetes. *Diab. techn. therap.* 2003; 5 (5): 769-779.
- 86) Fellin T, Pascual O, Gobbo S, Porzan T, Haydon PG, Carmignoto G. Neuronal synchrony mediated by astrocytic glutamate through activation of extrasynaptic NMDA receptors. *Neuron* 2004; 43: 729-743
- 87) Fields RD, Stevens-Graham B. New insights into neuron-glia communication. *Science* 2001; 298: 556-562.
- 88) Fillenz M. *In vivo* neurochemical monitoring and the study of behaviour. *Neurosci. Biobeh. Rev.* 2005; 29: 949-962.
- 89) Forzani ES, Solis VM and Calvo EJ. Electrochemical behaviour of polyphenol oxidase immobilized in self-assembled structures layer by layer with cationic polyallylamine. *Anal. Chem.* 2000; 72: 5300-5307.
- 90) Foulds NC, Lowe CR. Immobilization of glucose oxidase in ferrocene-modified pyrrole polymers. *Anal. Chem.* 1988; 60: 2473-2478
- 91) Gaddum JH. Push-pull cannulae. *J. Physiol.* 1961; 155: 1P.
- 92) Gallo V, Chittajallu R. Unwrapping glial cells from the synapse: What lies inside? *Science* 2001; 292: 872-873.

- 93) Gao Z, Binyamin G, Kim H-H, Barton SC, Zhang Y and Heller A. Electrodeposition of redox-polymers and co-electrodeposition of enzymes by coordinative crosslinking. *Angew. Chem. Int. Ed.* 2002; 41: 810-813
- 94) Garguilo G and Michael AC. An enzyme-modified microelectrode that detects choline injected locally into brain tissue. *J. Am. Chem. Soc.* 1993a; 115: 12218-12219.
- 95) Garguilo MG, Huynh N, Proctor A, Michael AC. Amperometric sensors for peroxide, choline and acetylcholine based on electron transfer between horseradish peroxidase and a redox polymer. *Anal. Chem.* 1993b; 65: 523-528.
- 96) Garguilo G, Michael AC. Quantification of choline in the extracellular fluid of brain tissue with amperometric microsensors. *Anal. Chem.* 1994; 66: 2621-2629.
- 97) Garguilo G, Michael AC. Optimization of amperometric microsensors for monitoring choline in the extracellular fluid of brain tissue. *Anal. Chim. Acta* 1995; 307: 291-299.
- 98) Garguilo G and Michael AC. Amperometric microsensors for monitoring choline in the extracellular fluid of brain. *J. Neurosci. Meth.* 1996; 70: 73-82.
- 99) Garthwaite G, Williams GD, Garthwaite J. Glutamate toxicity: An experimental and theoretical analysis. *Eur. J. Neurosci.* 1992; 4: 353-360.
- 100) Gáspár S, Wang X, Suzuki H, Csöregi E. Amperometric biosensor-based flow-through microdetector for microdialysis applications. *Anal. Chim. Acta* 2004; 525: 75-82.
- 101) Geddes JW, Chang NG, Ackley DC, Sultanian NS, McGillis JP, Yokel RA. Postmortem elevation in extracellular glutamate in the rat hippocampus when brain temperature is maintained at physiological levels: implications for the use of human brain autopsy tissues. *Brain Res.* 1999; 831: 104-112.
- 102) Gegelashvili G, Robinson MB, Trotti D, Rauen T. Regulation of glutamate transporters in health and disease. *Prog. Brain Res.* 2001; 132: 267-286.
- 103) Gerhardt GA, Burmeister JJ. Voltammetry *in vivo* for chemical analysis of the nervous system. Encyclopedia of Analytical Chemistry. R.A. Meyers (ed.). Wiley & Sons, Chisester, 2000: 710-731.
- 104) Gerhardt GA, Hoffman AF. Effects of recording media composition on the responses of Nafion-coated carbon fiber microelectrodes measured using high-speed chronoamperometry. *J. Neurosci. Meth.* 2001; 109: 13-21.
- 105) Gilgun-Sherki Y, Melamed E, Offen D. Oxidative stress induced-neurodegenerative diseases: the need for antioxidants that penetrate the blood brain barrier. *Neuropharmacology* 2001; 40: 959-975.

- 106) Gorton L, Csöregi E, Domínguez E, Emnéus J, Jönsson-Pettersson G, Marko-Varga G, Persson B. Selective detection in flow analysis based on the combination of immobilized enzymes and chemically modified electrodes. *Anal. Chim. Acta* 1991; 250: 203-248.
- 107) Gorton L, Lindgren A, Larsson T, Munteanu FD, Ruzgas T, Gazaryan I. Direct electron transfer between heme-containing enzymes and electrodes as basis for third generation biosensors. *Anal. Chim. Acta* 1999; 400: 91-108.
- 108) Gough DA, Lucisano JY, Tse PHS. Two-dimensional enzyme electrode sensor for glucose. *Anal. Chem.* 1985; 57: 2351-2357.
- 109) Greengard P. The neurobiology of slow synaptic transmission. *Science* 2001; 294: 1024-1030.
- 110) Gregg BA, Heller A. Cross-linked redox gels containing glucose oxidase for amperometric biosensor applications. *Anal. Chem.* 1990; 62: 258-263.
- 111) Gregg BA, Heller A. Redox polymer films containing enzymes. 1. A redox-conducting epoxy cement: synthesis, characterization, and electrocatalytic oxidation of hydroquinone. *J. Phys. Chem.* 1991a; 95: 5970-5975.
- 112) Gregg BA, Heller A. Redox polymer films containing enzymes. 2. Glucose oxidase containing enzyme electrodes. *J. Phys. Chem.* 1991b; 95: 5976-5980.
- 113) Grünewald RA. Ascorbic acid in the brain. *Brain Res. Rev.* 1993; 18: 123-133.
- 114) Guyot LL, Diaz FG, O'Regan MH, McLeod S, Park H, Phillis JW. Real-time measurement of glutamate release from the ischemic penumbra of the rat cerebral cortex using a focal middle cerebral artery occlusion model. *Neurosci. Lett* 2001; 299: 37-40.
- 115) Habermüller K, Mosbach M, Schuhmann W. Electron-transfer mechanisms in amperometric biosensors. *Fres. J. Anal. Chem.* 2000; 366: 560-568.
- 116) Hama H, Hara C, Yamaguchi K, Miyawaki A. PKC signaling mediates global enhancement of excitatory synaptogenesis in neurons triggered by local contact with astrocytes. *Neuron* 2004; 41: 405-414.
- 117) Hanrahan G, Patil DG, Wang J. Electrochemical sensors for environmental monitoring: design, development and applications. *J. Environ. Monit.* 2004; 6: 657-664.
- 118) Hansson E, Muyderman H, Leonova J, Allansson L, Sinclair J, Blomstrand F, Thorlin T, Nilsson M, Rönnbäck L. Astroglia and glutamate in physiology and pathology: aspects on glutamate transport, glutamate-induced cell swelling and gap-junction communication. *Neurochem. Int.* 2000; 37: 317-329.

- 119) Hassinger TD, Guthrie PB, Atkinson PB, Bennett M, Kater SB. An extracellular signaling component in propagation of astrocytic calcium waves. *Proc. Natl. Acad. Sci.* 1996; 93: 13268-13273.
- 120) Hascup KN, Rutherford EC, Pomerleau F, Huettl P, Gerhardt GA. Second-to-second measures of l-glutamate using enzyme-based microelectrodes in the CNS of conscious freely moving mice. *Monitoring Molecules in Neuroscience*; 11<sup>th</sup> Int. Conf; Ed: Di Chiara G, Carboni E, Valentini V, Acquas E, Bassareo V, Cadoni C. 2006; 298-300.
- 121) Hayashi T. Effects of sodium glutamate on the nervous system. *Keio J. Med.* 1954; 3: 183-192.
- 122) Haydon PG. Glia: Listening and talking to the synapse. *Nat. Rev. Neurosci.* 2001; 2: 185-193.
- 123) Hazzard JT, Maritano S, Tollin G, Marchesini A. Laser flash photolysis experiments on the effects of freezing and salt addition on intramolecular electron transfer within one-electron reduced ascorbate oxidase. *Archiv. Biochem. Biophys.* 1997; 339: 24-32.
- 124) Heller A. Implanted electrochemical glucose sensors for the management of diabetes. *Annu. Rev. Biomed. Eng.* 1999; 01: 153-175.
- 125) Heller A. Chapter 1: Redox-hydrogel based electrochemical biosensors. Oxford University Press, web-edition ([www.oup.co.uk](http://www.oup.co.uk)), 2006.
- 126) Hertz L, Zielke R. Astrocytic control of glutamergic activity: astrocytes as stars of the show. *Tr. Neurosci.* 2004; 27 (12): 735-743.
- 127) Hirano A, Moridera N, Akashi M, Saito M, Sugawara M. Imaging of l-glutamate fluxes in mouse brain slices based on an enzyme-based membrane combined with a difference-image analysis. *Anal. Chem.* 2003; 75: 3775-3783.
- 128) Hirrlinger J, Hulsman S, Kirchhoff F. Astroglial processes show spontaneous motility at active synaptic terminals *in situ*. *Eur. J. Neurosci.* 2004; 20: 2235-2239.
- 129) Hlubek M, Tian D, Stuenkel EL. Mechanism of  $\alpha$ -latrotoxin action at nerve endings of neurohypophysis. *Brain Res.* 2003; 992: 30-42.
- 130) Hoshi T, Anzai J, Osa T. Controlled deposition of glucose oxidase on platinum electrode based on an avidin/biotin system for the regulation of output current of glucose sensors. *Anal. Chem.* 1995; 67: 770-774.
- 131) Hu Y, Mitchell KM, Albahadily FN, Michaelis EK and Wilson GS. Direct measurement of glutamate release in the brain using a dual enzyme-based electrochemical sensor. *Brain Res.* 1994; 659: 117-125.
- 132) Hu Y, Wilson GS. Rapid changes in local extracellular rat brain glucose observed with an *in vivo* glucose sensor. *J. Neurochem.* 1997; 68, no.4: 1745-1752.

- 133) Huettnner JE. Kainate receptors and synaptic transmission. *Progr. Neurobiol.* 2003; 70: 387-407.
- 134) Hutchinson PJ, O'Connell MT, Kirkpatrick PJ, Pickard JD. How can we measure substrate, metabolite and neurotransmitter concentrations in the human brain. *Physiol. Meas.* 2002; 23: 75-109.
- 135) Inagaki T, Skotheim TA, Okamoto Y. Syntheses and electrochemical properties of siloxane polymers containing ferrocene and dimethylferrocene. *J. Chem. Soc. Chem. Commun.* 1989: 1181-1183.
- 136) Jabaudon D, Shimamoto K, Yasuda-Kamatani Y, Scanziani M, Gähwiler BH, Gerber U. Inhibition of uptake unmasks rapid extracellular turnover of glutamate of nonvesicular origin. *Proc. Natl. Acad. Sci.* 1999; 96: 8733-8738.
- 137) JanákyR, Varga V, Hermann A, Saransaari P, Oja SS. Mechanisms of l-cysteine neurotoxicity. *Neurochem. Res.* 2000; 25: 1397-1405.
- 138) Javitt DC. Glutamate as a therapeutic target in psychiatric disorders. *Mol. Psych.* 2004; 9: 984-997.
- 139) Jernigan JC, Chidsey CED, Murray RW. Electrochemistry of polymer films not immersed in solution: electron transfer on an ion budget. *J. Am. Chem. Soc.* 1985; 107: 2824-2826.
- 140) Jiang C, Agulian S, Haddad GG. O<sub>2</sub> tension in adult and neonatal brain slices under several experimental conditions. *Brain Res.* 1991; 568: 159-164.
- 141) Jiménez C, Bartolí J, de Rooij NF, Koudelka-Hep M. Glucose sensor based on an amperometric microelectrode with a photopolymerizable enzyme membrane. *Sens. Actuators B* 1995; 27: 421-424.
- 142) Jönsson G, Gorton L, Petterson L. An electrochemical sensor for hydrogen peroxide based on peroxidase absorbed on a spectrographic graphite electrode. *Electroanal.* 1989; 1: 49-55.
- 143) Karyakin A.A., Karyakina E.E.; Gorton L. Amperometric biosensor for glutamate using prussian blue-based "artificial peroxidase" as a transducer for hydrogen peroxide. *Anal. Chem.* 2000, 72, 1720-1723.
- 144) Kasai N, Jimbo Y, Niwa O, Matsue T, Torimitsu K. Real-time multisite observation of glutamate release in rat hippocampal slices. *Neurosci. Lett.* 2001; 304: 112-116.
- 145) Kasai N, Jimbo Y, Torimitsu K. Electrochemical monitoring of glutamate release at multiple positions in a rat hippocampal slice. *Anal. Sci.* 2002; 18: 1325-1327.
- 146) Katsumori H, Baldwin RA, Wasterlain CG. Reverse transport of glutamate during depolarization in immature hippocampal slices. *Brain Res.* 1999; 819: 160-164.

- 147) Katz E, Riklin A, Helegshabtai V, Willner I, Bückmann AF. Glucose oxidase electrodes via reconstitution of the apo-enzyme: tailoring of novel glucose biosensors. *Anal. Chim. Acta* 1999; 385: 45-58.
- 148) Kawagoe KT, Zimmerman JB and Wightman RM. Principles of voltammetry and microelectrode surface states. *J. Neurosci. Meth.* 1993; 48: 225-240.
- 149) Kehr J. Determination of glutamate and aspartate in microdialysis samples by reversed-phase column liquid chromatography with fluorescence and electrochemical detection. *J. Chromatogr. B Biomed. Sci. Appl.* 1998; 708: 27-38.
- 150) Kenausis G, Chen Q, Heller A. Electrochemical glucose and lactate sensors based on wired thermostable soybean peroxidase operating continuously and stably at 37 °C. *Anal. Chem.* 1997; 69: 1054-1060.
- 151) Kennedy RT, Thompson JE, Vickroy TW. *In vivo* monitoring of amino acids by direct sampling of brain extracellular fluid at ultralow flow rates and capillary electrophoresis. *J. Neurosci. Meth.* 2002; 114: 39-49.
- 152) Kew JNC, Kemp JA. Ionotropic and metabotropic glutamate receptor structure and pharmacology. *Psychopharmacology* 2005; 179: 4-29.
- 153) Khan AS, Michael AC. Invasive consequences of using micro-electrodes and microdialysis probes in the brain. *Tr. Anal. Chem.* 2003; 22 (9): 503-508.
- 154) Kimelberg HK, Rutledge E, Goderie S, Charniga C. Astrocytic swelling due to hypotonic or high K<sup>+</sup> medium causes inhibition of glutamate and aspartate uptake and increases their release. *J Cereb. Blood Flow Metab.* 1995; 15: 409-416.
- 155) Kimelberg HK. Increased release of excitatory amino acids by the actions of ATP and peroxynitrite on volume-regulated anion channels (VRACs) in astrocytes. *Neurochem. Int.* 2004; 45: 511-519.
- 156) Koeneman BA, Lee KK, Singh A, He J, Raupp GB, Panitch A, Capco DG. An *ex vivo* method for evaluating the biocompatibility of neural electrodes in rat brain slice cultures. *J. Neurosci. Meth.* 2004; 137: 257-63.
- 157) Koopal CGJ, Bos AACM, Nolte RJM. Third-generation glucose biosensor incorporated in a conducting printing ink. *Sens. Actuators B* 1994; 18: 166-170.
- 158) Krebs HA. Metabolism of amino acids. IV. Synthesis of glutamine from glutamic acid and ammonia, and the enzymatic hydrolysis of glutamine in animal tissue. *Biochem. J.* 1935; 29: 1951-1969.
- 159) Kreft M, Stenovec M, Rupnik M, Grilc S, Krzan M, Potokar M, Pangrsic T, Haydon PG, Zorec R. Properties of Ca<sup>2+</sup>-dependent exocytosis in cultured astrocytes. *Glia* 2004; 46: 437-445.

- 160) Kulagina NV, Shankar L, Michael AC. Monitoring glutamate and ascorbate in the extracellular space of brain tissue with electrochemical microsensors. *Anal. Chem.* 1999; 71: 5093-5100.
- 161) Kulagina NV, Michael AC. Monitoring hydrogen peroxide in the extracellular space of the brain with amperometric microsensors. *Anal. Chem.* 2003; 75: 4875-4881.
- 162) Kurzawa C, Hengstenberg A, Schuhmann W. Immobilization method for the preparation of biosensors based on pH shift-induced deposition of biomolecule-containing polymer films. *Anal. Chem.* 2002; 74: 355-361.
- 163) Kusakabe H, Midorikawa Y, Fujishima T, Kuninaka A, Yoshino H. Occurrence of a new enzyme, l-glutamate oxidase in a wheat bran culture extract of streptomyces sp. X-119-6. *Agric. Biol. Chem.* 1983 ; 47 : 1323-1328.
- 164) Lada MW, Vickroy TW, Kennedy RT. Evidence for neuronal origin and metabotropic receptor-mediated regulation of extracellular glutamate and aspartate in rat striatum *in vivo* following electrical stimulation of the prefrontal cortex. *J. Neurochem.* 1998; 70(2): 617-625.
- 165) Lai YL, Shen EY, Pan WHT. Effects of ascorbate in microdialysis medium on the extracellular basal concentration of glutamate in rat's striatum. *Neurosci. Lett.* 2000: 279: 145-148.
- 166) Lamprecht R, LeDoux J. Structural plasticity and memory. *Nature Neurosci.* 2004; 5: 45-54.
- 167) Lay P.A.; Sargeson A.M.; Taube H. *cis*- bis(2,2'-bipyridine-*N,N'*) complexes of ruthenium (III)/(II) and osmium (III)/(II). *Inorganic Synthesis*, 24, ed. Shreeve H.J.M., Wiley, NY, 1986, 291-299.
- 168) Leis JA, Bekar LK, Walz W. Potassium homeostasis in the ischemic brain. *Glia* 2005; 50: 407-416.
- 169) Lerma J. Roles and Rules of kainate receptors in synaptic transmission. *Nat. Rev. Neurosci.* 2003; 4: 481-495.
- 170) Lomeli H, Sprengel R, Laurie DJ, Köhr G, Herb A, Seeburg PH, Wisden W. The rat delta-1 and delta-2 subunits extend the excitatory amino acid receptor family. *FEBS Lett.* 1993; 315: 318-322.
- 171) Lowry JP, Ryan MR, O'Neill RD. Behaviourally induced changes in extracellular levels of brain glutamate monitored at 1s resolution with an implanted biosensor. *Anal. Comm.* 1998; 35: 87-89.
- 172) Lowry JP, Ryan MR, O'Neill RD. Interference in biosensor detection of brain glutamate *in vivo*: possible role of endogeneous ECF hydrogen peroxide. Monitoring Molecules in Neuroscience; 9<sup>th</sup> Int. Conf; Ed: O'Conner WT, Lowry JP, O'Conner JJ, O'Neill RD. 2001; 70-71.

- 173) Lowry OH, Rosebrough NJ, Lewis Farr A, Randall RJ. Protein measurement with the Folin Phenol reagent. *J. Biol. Chem.* 1951, 193 (1), 265-275.
- 174) Lumley-Woodyear de T, Rocca P, Lindsay J, Dror Y, Freeman A, Heller A. Polyacrylamide-based redox polymer for connecting redox centers of enzymes to electrodes. *Anal. Chem.* 1995; 67: 1332-1338.
- 175) Madden DR. The structure and function of glutamate receptor ion channels. *Nature Neurosci.* 2002; 3: 91-101.
- 176) Maiden R, Heller A. Elimination of electrooxidizable interferant-produces currents in amperometric biosensors. *Anal. Chem.* 1992; 64: 2889-2896.
- 177) Mao F, Mano N, Heller A. Long tethers binding redox centers to polymer backbones enhance electron transport in enzyme "wiring" hydrogels. *J. Am. Chem. Soc.* 2003; 125: 4951-4957.
- 178) Marcus RA, Sutin N. Electron transfer in chemistry and biology. *Biochim Biophys Acta* 1985; 811: 265-322.
- 179) Marcus RA. Electron transfer reactions in chemistry: theory and experiment *Angew. Chem. Int. Ed. Engl.* 1993; 32: 1111-1121.
- 180) Marko-Varga G, Appelquist R, Gorton L. A glucose sensor based on glucose dehydrogenase absorbed on a modified carbon electrode. *Anal. Chim. Acta* 1986; 179: 371-379.
- 181) Masamoto K, Takizawa N, Kobayashi H, Oka K and Tanishita K. Dual responses of tissue partial pressure of oxygen after functional stimulation in rat somatosensory cortex. *Brain Res.* 2003; 979: 104-113.
- 182) Mayat E, Petralia RS, Wang YX, Wenthold RJ. Immunoprecipitation, immunoblotting, and immunocytochemistry studies suggest that glutamate receptor  $\delta$  subunits form novel postsynaptic receptor complexes. *J. Neurosci.* 1995; 15(3): 2533-2546.
- 183) Mayer ML. Glutamate receptor ion channels. *Curr. Opin. Neurobiol.* 2005; 15: 282-288.
- 184) Michael AC. personal communication, 2006.
- 185) Miele M, Berners M, BoutelleMG, Kusakabe H, Fillenz M. The determination of the extracellular concentration of brain glutamate using quantitative microdialysis. *Brain Res.* 1996a; 707: 131-133.
- 186) Miele M, Boutelle MG and Fillenz M. The source of physiologically stimulated glutamate efflux from the striatum of conscious rats. *J. Physiol.* 1996b; 497: 745-751.



- 187) Migneault I, Dartiguenave C, Bertrand MJ, Waldron KC. Glutaraldehyde: behavior in aqueous solution, reaction with proteins, and application to enzyme cross-linking. *Biotechniques* 2004; 37: 790-796, 798-802.
- 188) Mikeladze E, Schulte A, Mosbach M, Blöchl A, Csöregi E, Solomonias R. Redox hydrogel-based bienzyme microelectrodes for amperometric monitoring of L-glutamate. *Electroanal.* 2002; 14 (6): 393-399.
- 189) Millan MJ. N-Methyl-D-aspartate receptors as a target for improved antipsychotic agents: novel insights and clinical perspectives. *Psychopharmacology* 2005; 179: 30-53.
- 190) Miller G. Neurons get connected via glia. *Science* 2003; 312 (21): 1323.
- 191) Montiel T, Camacho A, Estrada-Sánchez AM, Massieu L. Differential effects of the substrate inhibitor L-trans-pyrrolidine-2,4-dicarboxylate (PDC) and the non-substrate inhibitor DL-threo-β-benzyloxyaspartate (DL-TBOA) of glutamate transporters on neuronal damage and extracellular amino acid levels in rat brain *in vivo*. *Neurosci.* 2005; 133: 667-678.
- 192) Moussy F, Harrison DJ. Prevention of the rapid degradation of subcutaneously implanted Ag/AgCl reference electrodes using polymer coatings. *Anal Chem.* 1994a; 66: 674-679.
- 193) Moussy F, Jakeway S, Harrison DJ, Rajotte RV. *In vitro* and *in vivo* performance and lifetime of perfluorinated ionomer-coated glucose sensors after high-temperature curing. *Anal Chem.* 1994b; 66: 3882-3888.
- 194) Mulligan SJ, MacVivar BA. Calcium transients in astrocyte endfeet cause cerebrovascular constrictions. *Nature* 2004; 431: 195-199.
- 195) Muzzolini A, Bregola G, Bianchi C, Beani L, Simonato M. Characterization of glutamate and [<sup>3</sup>H]D-aspartate outflow from various *in vitro* preparations of the rat hippocampus. *Neurochem. Int.* 1997; 31: 113-124.
- 196) Nakajima K, Yamagiwa T, Hirano A, Sugawara M. A glass capillary microelectrode based on capillarity and its application to the detection of L-glutamate release from mouse brain slices. *Anal. Sci.* 2003; 19: 55-60.
- 197) Nakamura N, Negishi K, Hirano A, Sugawara M. Real-time monitoring of L-glutamate release from mouse brain slices under ischemia with a glass capillary-based enzyme electrode. *Anal. Bioanal. Chem.* 2005; 383: 660-667.
- 198) Nakamura T, Kawamura Y, Miyakawa H. Optical bioimaging: From living tissue to a single molecule: optical detection of synaptically induced glutamate transporter activity in hippocampal slices. *J Pharmacol. Sci.* 2003; 93: 234-241
- 199) Nedergaard M, Takano T, Hansen J. Beyond the role of glutamate as a neurotransmitter. *Nat. Rev. Neurosci.* 2002; 3: 748-55.

- 200) Newman EA. Glial inhibition of neurons by release of ATP. *J. Neurosci.* 2003; 23: 1659-1666.
- 201) Ni JA, Ju HX, Chen HY and Leech D. Amperometric determination of epinephrine with an osmium complex and Nafion double-layer membrane modified electrode. *Anal. Chim. Acta.* 1999; 378: 151-157.
- 202) Nicholls DG. The glutamergic nerve terminal. *Eur. J. Biochem.* 1993; 212: 613-631.
- 203) Nickell J, Pomerleau F, Allen J, Gerhardt GA. Age-related changes in the dynamics of potassium-evoked L-glutamate release in the striatum of Fischer 344 rats. *J. Neural Transm.* 2005; 112: 87-96.
- 204) Nickell J, Salvatore M, Pomerleau F, Huettl P, Apparsundaram S, Gerhardt GA. Decreased glutamate uptake rate in aged rat striatum: Investigations coupling *in vivo*, enzyme-based microelectrode arrays with biotinylation. Monitoring Molecules in Neuroscience; 11<sup>th</sup> Int. Conf; Ed: Di Chiara G, Carboni E, Valentini V, Acquas E, Bassareo V, Cadoni C. 2006; 322-324.
- 205) Niwa O, Torimitsu K, Morita M, Osborne P, Yamamoto K. Concentration of extracellular l-glutamate released from cultured nerve cells measured with a small volume online sensor. *Anal. Chem.* 1996; 68: 1865-1870.
- 206) Noraberg J, Kristensen BW, Zimmer J. Markers for neuronal degeneration in organotypic slice cultures. *Brain Res. Protoc.* 1999; 3: 278-290.
- 207) O'Neill RD, Lowry JP and Mas M. Monitoring Brain Chemistry *in vivo*: Voltammetric techniques, Sensors, and Behavioural Applications. *Crit. Rev. in Neurobiol.* 1998; 12: 69-127.
- 208) Obrenovitch TP, Zilkha E. Microdialysis coupled to online enzymatic assays. *Methods* 2001; 23: 63-41
- 209) Ohara TJ, Rajagopalan R, Heller A. "Wired" enzyme electrodes for amperometric determination of glucose or lactate in the presence of interfering substances. *Anal. Chem.* 1994; 66: 2451-2457.
- 210) Ohara TJ, Rajagopalan R, Heller A. Glucose electrodes based on cross-linked [Os(bpy)<sub>2</sub>Cl]<sup>+2+</sup> complexed poly(1-vinylimidazole) films. *Anal. Chem.* 1993a; 65: 3512-3517.
- 211) Ohara TJ, Vreeke MS, Battaglini F, Heller A. Bienzyme sensors based on "electrically wired" peroxidase. *Electroanalysis* 1993b; 5: 825-831.
- 212) Oldenzienl WH, Beukema W, Westerink BHC. Improving the reproducibility of hydrogel-coated glutamate microsensors by using an automatic dipcoater. *J Neurosci. Meth.* 2004; 140: 117-126. (Chapter 2)
- 213) Oldenzienl WH, Westerink BHC. Improving glutamate microsensors by optimizing the composition of the redox hydrogel. *Anal. Chem.* 2005; 77: 5520-5528. (Chapter 3)

- 214) Oldenziel WH, Jong de LAA, Dijkstra G, Cremers TIFH, Westerink BHC. Improving the performance of glutamate microsensors by purification of ascorbate oxidase. *Anal. Chem.* 2006; 78: 2456-60. (Chapter 4)
- 215) Oldenziel WH, Dijkstra G, Cremers TIFH, Westerink BHC. Evaluation of hydrogel-coated glutamate microsensors. *Anal. Chem.* 2006; 78: 3366-78. (Chapter 5)
- 216) Oldenziel WH, Zeyden van der M, Dijkstra G, Cremers TIFH, Westerink BHC. Monitoring extracellular glutamate in hippocampal slices with a microsensor. *J. Neurosci. Meth.* 2006, in press. (Chapter 6)
- 217) Oldenziel WH, Dijkstra G, Cremers TIFH, Westerink BHC. *In vivo* monitoring of extracellular glutamate in the brain with a microsensor. *Brain Res.*, 2006, in press. (Chapter 7)
- 218) Oliet SH, Piet R, Poulain DA. Control of glutamate clearance and synaptic efficacy by glial coverage of neurons. *Science* 2001; 292: 923-926.
- 219) Orłowska-Madjak M. Effect of excitatory amino acids on activity of vasopressinergic and oxytocinergic neurons. *Endocrine Regulations* 2004; 38: 23-28.
- 220) Palmisano F, Zamboni PG, Centonze D. Amperometric biosensors based on electrosynthesised polymeric films. *Fres. J. Anal. Chem.* 2000; 366: 586-601.
- 221) Palucha A, Pilc A. The involvement of glutamate in the pathophysiology of depression. *Drug News Perspect.* 2005; 18(4): 262-268.
- 222) Parikh V, Pomerleau F, Huettl P, Gerhardt GA, Sarter M, Bruno P. Rapid measurements of *in vivo* cholinergic transmission by amperometric detection of changes in extracellular choline levels. *Eur. J. Neurosci.* 2004; 20: 1545-54.
- 223) Parpura V, Basarsky TA, Liu F, Jęftinija K, Jęftinija S, Haydon PG. Glutamate-mediated astrocyte-neuron signalling. *Nature* 1994; 369: 744-747.
- 224) Parpura V, Scemes E, Spray DC. Mechanisms of glutamate release from astrocytes: gap junction "hemichannels", purinergic receptors and exocytotic release. *Neurochem. Int.* 2004; 45: 193-205.
- 225) Patolsky F, Zayats M, Katz E, Willner I. Precipitation of an insoluble product on enzyme monolayer electrodes for biosensor applications: characterization by faradaic impedance spectroscopy, cyclic voltammetry, and microgravimetric quartz crystal microbalance analyses. *Anal. Chem.* 1999; 71:3171-3180.
- 226) Paxinos G, Watson C. The rat brain in stereotaxic coordinates. 1986. Academic Press, London.
- 227) Pearson JE, Gill A, Vagdama P. Analytical aspects of biosensors. *Ann. Clin. Biochem.* 2000; 37: 119-145.

- 228) Perea G, Araque A. Properties of synaptically evoked astrocyte calcium signal reveal synaptic information processing by astrocytes. *J. Neurosci.* 2005; 25: 2192-2203.
- 229) Peters JL, Yang H, Michael AC. Quantitative aspects of brain microdialysis. *Anal. Chim. Acta* 2000; 412: 1-12.
- 230) Pinnacle: see the website [www.pinnaclet.com](http://www.pinnaclet.com)
- 231) Pomerleau F, Day BK, Huettl P, Burmeister JJ, Gerhardt GA. Real time *in vivo* measures of l-glutamate in the rat central nervous system using ceramic-based multisite microelectrode arrays. *Ann. N.Y. Acad. Sci.* 2003; 1003: 454-457.
- 232) Pomerleau F, Huettl P, Burmeister JJ, Gerhardt GA. Second-to-second measurements of GABA in CNS using enzyme-based microelectrode arrays. *Monitoring Molecules in Neuroscience; 11<sup>th</sup> Int. Conf*; Ed: Di Chiara G, Carboni E, Valentini V, Acquas E, Bassareo V, Cadoni C. 2006; 46-48.
- 233) Qhobosheana M, Wu D, Gu Y, Tan W. A two-dimensional imaging biosensor to monitor enhanced brain glutamate release stimulated by nicotine. *J Neurosci. Meth.* 2004; 135: 71-78.
- 234) Quan Z, Liu YM. Capillary electrophoretic separation of glutamate enantiomers in neural samples. *Electrophoresis* 2003; 24: 1092-1096.
- 235) Rahman A, Kwon NH, Won MS, Choe ES, Shim YB. Functionalized conducting polymer as an enzyme-immobilizing substrate: An amperometric glutamate microbiosensor for *in vivo* measurements. *Anal. Chem.* 2005; 77: 4854-4860.
- 236) Raiteri L, Raiteri M, Bonanno G. Coexistence and function of different neurotransmitter transporters in the plasma membrane of CNS neurons. *Prog. Neurobiol.* 2002; 68: 287-309.
- 237) Rajagopalan R, Aoki A, Heller A. Effect of quarternization of the glucose oxidase "wiring" redox polymer on the maximum current densities of glucose electrodes. *J. Phys. Chem.* 1996; 100: 3719-3727.
- 238) Razumas V, Kazlauskaite J, Ruzgas T, Kulys J. Bioelectrochemistry of microperoxidases. *Bioelectrochem. Bionerg.* 1992; 28: 159-176.
- 239) Rea K, Cremers TIFH, Westerink BHC. HPLC conditions are critical for the detection of GABA by microdialysis. *J. Neurochem.* 2005; 94: 672-679.
- 240) Rebec GV, Witowski SR, Sandstrom MI, Rostand RD and Kennedy RT. Extracellular ascorbate modulates cortically evoked glutamate dynamics in rat striatum. *Neurosci. Lett.* 2005; 378(3): 166-170.
- 241) Reinhammer B, Aasa R, Vänngård T, Maritano S, Marchesini A. The type 2 copper of ascorbate oxidase. *Biochim. Biophys. Acta* 1997; 1337: 191-197.

- 242) Rhemrev-Boom MM, Korf J, Venema K, Urban G, Vadgama P. A versatile biosensor for continuous biomedical monitoring. *Biosens. Bioelectron.* 2001; 16: 839-847.
- 243) Rice ME. Ascorbate regulation and its neuroprotective role in the brain. *TINS* 2000; 23: 209-216.
- 244) Rice ME. Use of ascorbate in the preparation and maintenance of brain slices. *Methods: A Companion to methods in enzymology* 1999; 18: 144-149.
- 245) Robert F, Parisi L, Bert L, Renaud B, Stoppini L. Microdialysis monitoring of extracellular glutamate combined with the simultaneous recording of evoked field potentials in hippocampal organotypic slice cultures. *J. Neurosci. Meth.* 1997; 74: 65-76.
- 246) Roettger V, Lipton P. Mechanism of glutamate release from rat hippocampal slices during *in vitro* ischemia. *Neurosci.* 1996; 75: 677-85.
- 247) Rossel S, Gonzales LE, Hernández L. One-second time resolution brain microdialysis in fully awake rats. Protocol for the collection, separation and sorting of nanoliter dialysate volumes. *J. Chrom. B.* 2003; 784: 385-393.
- 248) Rozza A, Masoero E, Favalli L, Lanza E, Govoni S, Rizzo V, Montalbetti L. Influence of different anaesthetics on extracellular amino acids in rat brain. *J. Neurosci. Meth.* 2000; 101: 165-169
- 249) Ruan K, Xu C, Li T, Li J, Lange R, Balny C. The thermodynamic analysis of protein stabilization by sucrose and glycerol against pressure-induced unfolding. *Eur. J. Biochem.* 2003, 270: 1654-1661.
- 250) Rusling JF, Forster RJ. Electrochemical catalysis with redox polymer and polyion-protein films. *J. Coll. Inter. Sci.* 2003; 262: 1-15.
- 251) Rusling JF. Sensors for toxicity of chemicals and oxidative stress based on electrochemical catalytic DNA oxidation. *Biosens. Bioelectron.* 2004; 20: 1022-1028.
- 252) Rutherford EC, Pomerleau F, Huettl P, Strömbergl, Johnson KW, Gerhardt GA. Second-by second enzyme-based microelectrode recordings of basal l-glutamate in the prefrontal cortex of awake rats. Monitoring Molecules in Neuroscience; 11<sup>th</sup> Int. Conf; Ed: Di Chiara G, Carboni E, Valentini V, Acquas E, Bassareo V, Cadoni C. 2006; 334-336.
- 253) Ruzgas T, Csöregi E, Emneus J, Gorton L, Marko-Varga G. Peroxidase-modified electrodes: Fundamentals and application. *Anal. Chim. Acta* 1996; 330: 123-138.
- 254) Ryabov AD, Sukharev VS, Alexandrova L, Le Lagadec R, Pfeffer M. New synthesis and new bio-application of cyclometalated ruthenium (II) complexes for fast mediated electron transfer with peroxidase and glucose oxidase. *Inorg Chem.* 2001; 40: 6529-6532.

- 255) Ryabova ES, Goral VN, Csöregi E, Matthiasson B, Ryabov AD. Coordinative approach to mediate electron transfer: ruthenium complexed to native glucose oxidase. *Angew. Chem. Int. Ed. Engl.* 1999; 38: 804-807.
- 256) Ryan MR, Lowry JP, O'Neill D. Biosensor for neurotransmitter l-glutamic acid designed for efficient use of l-glutamate oxidase and effective rejection of interference. *Analyst* 1997; 122: 1419-1424.
- 257) Santiago M, Westerink BHC. Simultaneous recording of the release of nigral and striatal dopamine in the awake rat. *Neurochem. Int.* 1992; 20: 107-110.
- 258) Scholtmeijer K, Wessels JG, Wosten HA. Fungal hydrophobins in medical and technical applications. *Appl. Microbiol. Biotechnol.* 2001; 56:1-8.
- 259) Schoonen AJ, Schmidt FJ, Hasper H, Verbrugge DA, Tiessen RG, Lerk CF. Development of potentially wearable glucose sensor for patients with diabetes mellitus: Design and in-vitro evaluation. *Biosens. Bioelectron.* 1990; 5(1): 37- 46.
- 260) Schuhmann W, Ohara TJ, Schmidt HL, Heller A. Electron transfer between glucose oxidase and electrodes via redox mediators bound with flexible chains to the electrode surface. *J. Am. Chem. Soc.* 1991; 113: 1394-1397.
- 261) Schuhmann W, Wohlschläger H, Lammert R, Schmidt HL, Löffler U, Wiemhöfer AD, Göpel W. Pyrrole, a new possibility for covalent binding of oxidoreductases to electrode surfaces as a base for stable biosensors. *Sens. Actuators B* 1990; 1: 537-541.
- 262) Schuhmann W. Amperometric enzyme biosensors based on optimized electron-transfer pathways and non-manual immobilization procedures. *Rev. Mol. Biotech.* 2002; 82: 425-441.
- 263) Schuhmann W. Immobilisation using electrogenerated polymers. In: Cass T, Ligler FS (eds): *Immobilized biomolecules in analysis. A practical approach.* Oxford University Press, NY, 1998, 187-210.
- 264) Scott TA; Melvin EH. Determination of dextran with anthrone. *Anal. Chem.* 1953, 25, 1656-1661.
- 265) Sem'yanov AV. Diffusional extrasynaptic neurotransmission via glutamate and GABA. *Neurosci. Behav. Psych.* 2005; 35: 253-266.
- 266) Seri B, Garcia-Verdugo JM, McEwen BS, Alvarez-Buyllia A. Astrocytes give rise to new neurons in the adult mammalian hippocampus. *J. Neurosci.* 2001; 21: 7153-7160.
- 267) Shankar L, Garguilo G, Michael AC. Enzyme biosensors based on redox polymers. In: Mulchandani A, Rogers KR, editors. *Methods in Biotechnology, Enzymes and Microbial Biosensors: Techniques and Protocols*, vol. 6. Totowa, NJ: Humana Press, 1998: 121-32.

- 268) Shi G, Yamamoto K, Zhou T, Xu F, Kato T, Ji-ye J, Jin L. On-line biosensors for simultaneous determination of glucose, choline and glutamate integrated with a microseparation system. *Electrophoresis* 2003; 24: 3266-72.
- 269) Smolders I, Sarre S, Vanhaesendonck C, Ebinger G, Michotte Y. Extracellular striatal dopamine and glutamate after decortication and kainate receptor stimulation as measured by microdialysis. *J. Neurochem.* 1996; 66: 2373-2379.
- 270) Song H, Stevens CF, Gage FH. Astroglia induce neurogenesis from adult neural stem cells. *Nature* 2002; 417: 39-44.
- 271) Song JH, Shin SH, Chung IM. Effects of glutamate on dehydroascorbate uptake and its enhanced vulnerability to the peroxidation in cerebral cortical slices. *Exp. Mol. Med.* 2002; 34: 419-425.
- 272) Sonnewald U, Kondziella D. Neuronal glial interaction in different neurological diseases studied by ex vivo <sup>13</sup>C NMR spectroscopy. *NMR in Biomed.* 2003; 16: 424-429.
- 273) Sperlagh B, Kofalvi A, Deuchars J, Atkinson L, Milligan CJ, Buckley NJ, Vizi ES. Involvement of P2X7 receptors in the regulation of neurotransmitter release in the rat hippocampus. *J. Neurochem.* 2002; 81: 1196-11211.
- 274) Stamford JA, Palij P, Davidson C, Jorm CM and Phillips PEM. Fast cyclic voltammetry in brain slices. *Voltammetric Methods in Brain Systems*, ed. Boulton AA, Baker GB and Adams RN, Totowa, NJ: Humana Press 1995: 81-116
- 275) Stenken JA. Methods and issues in microdialysis calibration. *Anal. Chim. Acta* 1999; 379: 337-358.
- 276) Stephens ML, Quintero JE, Talauliker PM, Huettl P, Pomerleau F, Gerhardt GA. Second-by-second l-glutamate recordings in the CA1 and CA3 regions in the young rat hippocampus. *Monitoring Molecules in Neuroscience; 11<sup>th</sup> Int. Conf*; Ed: Di Chiara G, Carboni E, Valentini V, Acquas E, Bassareo V, Cadoni C. 2006; 337-339.
- 277) Stern JR, Eggleston LV, Hems R, Krebs HA. Accumulation of glutamic acid in isolated brain tissue. *Biochem. J.* 1949; 410-418.
- 278) Sternberg R, Barrau MB, Gangiotti L, Thévenot DR, Bindra DS, Wilson GS, Velho G, Froguel P, Reach G. Study and development of multilayer needle-type enzyme-based glucose microsensors. *Biosensors* 1989;4: 27-40.
- 279) Strike DJ, de Rooij NF, Koudelka-Hep M. Electrochemical techniques for the modification of microelectrodes. *Biosens. Bioelectron.* 1995; 10: 61-66.
- 280) Syková E. Extrasynaptic volume transmission and diffusion parameters of the extracellular space. *Neurosci.* 2004; 129: 861-876.
- 281) Syková E. Glia and volume transmission during physiological and pathological states. *J. Neural Tranm.* 2005; 112: 137-147.

- 282) Takano T, Kang J, Jaiswal JK, Simon SM, Lin JHC, Yu Y, Li Y, Yang J, Dienel G, Zielke HR, Nedergaard M. Receptor-mediated glutamate release from volume sensitive channels in astrocytes. *PNAS* 102 (45): 16466-16471.
- 283) Takashi M, Hashimoto M. Lowering extracellular Na<sup>+</sup> concentration causes NMDA receptor-mediated neuronal death in cultured rat hippocampal slices. *Brain. Res.* 1996; 735: 1-8.
- 284) Therasense: see the website [www.Therasense.com](http://www.Therasense.com). (automatic link to [www.abbottdiabetescare.com](http://www.abbottdiabetescare.com))
- 285) Thévenot DR, Toth K, Durst RA, Wilson GS. Electrochemical biosensors: Recommended definitions and classification. *Anal. Lett.* 2001; 34: 635-659.
- 286) Tikhonov DB, Zhorov BS. Sodium channel activators: Model of binding inside the pore and a possible mechanism of action. *FEBS lett.* 2005; 579: 4207-4212.
- 287) Timasheff SN. The control of protein stability and association by weak interactions with water: how do solvents affect these processes? *Annu. Rev. Biophys. Biomol. Struct.* 1993, 22: 67-97.
- 288) Timmerman W, Westerink BHC. Brain microdialysis of GABA and glutamate: What does it signify? *Synapse* 1997; 27: 242-261.
- 289) Trudeau LE. Glutamate co-transmission as an emerging concept in monoamine neuron function. *Rev. Psych. Neurosci.* 2004; 29: 296-310.
- 290) Turner APF, Karube I, Wilson GS (eds). *Biosensors: Fundamentals and Applications*. Oxford University Press, Oxford, 1987, p. 481-5
- 291) Turner APF. *Biosensors: past, present and future*. 1996; webpublication: [www.cranfield.ac.uk/biotech/chinap.htm](http://www.cranfield.ac.uk/biotech/chinap.htm).
- 292) Ullian EM, Sapperstein SK, Christopherson KS, Barres BA. Control of synapse number by glia. *Science* 2001; 294: 657-661.
- 293) Ungerstedt U. Introduction to microdialysis. In: Robinson TE, Justice JB Jr., editors. *Techniques in the behavioural and neural sciences: microdialysis in the neurosciences*, vol. 7. Amsterdam: Elsevier, 1991: 3-18.
- 294) Updike SJ, Hicks, GP. The enzyme electrode. *Nature* 1967; 214: 986-988.
- 295) Vaidya R, Wilkins E. Use of charged membranes to control interference by body chemicals in a glucose biosensor. *Med. Eng. Phys.* 1994; 16: 416-21.
- 296) Vitten H, Isaacson JS. Synaptic transmission: Exciting times for presynaptic receptors. *Curr. Opin Neurobiol.* 2001; 11: 695-697.
- 297) Volterra A, Meldolesi J. Astrocytes, from brain glue to communication elements: the revolution continues. *Nat. Rev. Neurosci.* 2005; 6: 626-640.



- 298) Waagpetersen HS, Qu H, Sonnewald U, Shimamoto K, Schousboe A. Role of glutamine and neuronal glutamate uptake in glutamate homeostasis and synthesis during vesicular release in cultured glutamergic neurons. *Neurochem. Int.* 2005; 47: 92-102.
- 299) Waldmeier PC, Wicki P, Feldtrauer JJ. Release of endogenous glutamate from rat cortical slices in presence of the glutamate uptake inhibitor L-trans-pyrrolidine-2,4-dicarboxylic acid. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 1993; 348: 478-485.
- 300) Waldmeier PC, Stocklin K, Feldtrauer JJ. Systemic administration of baclofen and the GABA<sub>B</sub> antagonist, CGP 35348, does not affect GABA, glutamate or aspartate in microdialysates of the striatum of conscious rats. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 1992; 345: 544-552.
- 301) Wang J, Lu F. J. Oxygen-rich oxidase enzyme electrodes for operation in oxygen free solutions. *J. Am Chem. Soc.* 1998; 120: 1048-1050.
- 302) Wang J, Lu M, Yang F, Zhang X, Baeyens WRG, Garcia-Campana AM. Microdialysis with on-line chemiluminescence detection for the study of nitric oxide release in rat brain following traumatic injury. *Anal. Chim. Acta* 2001; 428: 173-181.
- 303) Wang J, Li S, Mo JW, Porter J, Musameh MM, Dasgupta PK. Oxygen-independent poly(dimethylsiloxane)-based carbon-paste glucose biosensors. *Biosens. Bioelectron.* 2002; 17: 999-1003.
- 304) Wang X, Arcuino G, Takano T, Lin J, Peng WG, Wan P, Li P, Xu Q, Liu QS, Goldman SA, Nedergaard M. P2X7 receptor inhibition improves recovery after spinal cord injury. *Nat. Med.* 2004; 10: 821-827.
- 305) Wang X, Shi F, Wosten HA, Hektor H, Poolman B, Robillard GT. The SC<sub>3</sub> hydrophobin self-assembles into a membrane with distinct mass transfer properties. *Biophys. J.* 2005; 88: 3434-43.
- 306) Watkins LR, Milligan EE, Maier SF. Glial activation: a driving force for pathological pain. *Tr. Neurosci.* 2001; 24: 450-455.
- 307) Watkins JC. l-Glutamate as a central neurotransmitter: Looking back. *Biochem. Soc. Trans.* 2000; 28 (4): 297-310.
- 308) West AR, Moore H, Grace AA. Direct examination of local regulation of membrane activity in striatal and prefrontal cortical neurons *in vivo* using simultaneous intracellular recording and microdialysis. *J. Pharm. Exp. Therap.* 2002; 301: 867-877.
- 309) Westerink BHC. Brain microdialysis and its application for the study of animal behaviour. *Behav. Brain Res.* 1995; 70 (2): 103-124.
- 310) Westerink BHC and Timmerman W. Do neurotransmitters sampled by brain microdialysis reflect functional release. *Anal. Chim. Acta* 1999; 379: 263-274.

- 311) Westerink BHC, Damsma G, Rollema H, de Vries JB, Horn AS. Scope and limitations of *in vivo* brain dialysis: A comparison of its application to various neurotransmitter systems. *Life Sci.* 1987; 41: 1763-1776.
- 312) Westerink BHC, de Vries JB. A method to evaluate the diffusion rate of drugs from a microdialysis probe through the brain. *J. Neurosci. Meth.* 2001; 109: 53-58.
- 313) Willner I, Katz E. Integration of layered redox proteins and conductive supports for bioelectronic applications. *Angew. Chem. Int. Ed.* 2000; 39: 1180-1218.
- 314) Wilson GS, Gifford R. Biosensors for real-time *in vivo* measurements. *Biosens Bioelectr.* 2005; 20: 2388-2403.
- 315) Wilson JX, Peters CE, Sitar SM, Daoust P, Gelb AW. Glutamate stimulates ascorbate transport by astrocytes. *Brain Res.* 2000; 858: 61-66.
- 316) Wisniewski N, Moussy F, Reichert WM. Characterization of implantable biosensor membrane biofouling. *Fres. J. Anal. Chem.* 2000; 366: 611-621.
- 317) Wisniewski N, Reichert WM. Methods for reducing biosensor membrane biofouling. *Coll. and Surf. B: Biointerfaces* 2000; 18: 197-219.
- 318) Xi ZX, Shen H, Baker DA, Kalivas PW. Inhibition of non-vesicular glutamate release by group III metabotropic glutamate receptors in the nucleus accumbens. *J. Neurochem.* 2003; 87: 1204-1212.
- 319) Xiao Y, Patolsky F, Katz E, Hainfeld JF, Willner I. "Plugging into enzymes": Nanowiring of redox enzymes by a gold nanoparticle. *Science* 2003; 299: 1877-1881.
- 320) Yao T, Nanjyo Y, Nishino H. Micro-flow *in vivo* analysis of l-glutamate with an on-line enzyme amplifier based on substrate recycling. *Anal. Sci.* 2001; 17: 703-8.
- 321) Yusa T. Increased extracellular ascorbate release reflects glutamate re-uptake during the early stage of reperfusion after forebrain ischemia in rats. *Brain Res.* 2001; 897: 104-113.
- 322) Yuzaki M. The  $\delta 2$  glutamate receptor: 10 years later. *Neurosci. Res.* 2003; 46: 11-22.
- 323) Zhang FF, Wan Q, Li CX, Wang XL, Zhu ZQ, Xian YZ, Jin LT, Yamamoto K. Simultaneous assay of glucose, lactate, L-glutamate and hypoxanthine levels in a rat striatum using enzyme electrodes based on neutral red-doped silica nanoparticles. *Anal. Bioanal. Chem.* 2004a; 380: 637-42.
- 324) Zhang FF, Wan Q, Li CX, Wang XL, Zhu ZQ, Xian YZ, Jin LT, Yamamoto K. Simultaneous monitoring of glucose, lactate, L-glutamate and hypoxanthine levels in rat striatum by a flow-injection enzyme electrode array system with *in vivo* microdialysis sampling. *J. Electroanal. Chem.* 2005; 575: 1-7.

- 325) Zhang JM, Wang HK, Ye CQ, Ge W, Chen Y, Jiang ZL, Wu CP, Poo MM, Duan S. ATP released by astrocytes mediates glutamergic activity-dependent heterosynaptic transmission. *Neuron* 2003; 40: 971-982.
- 326) Zhang Q, Fukuda M, van Bockstaele E, Pascual O, Haydon PG. Fusion-related release of glutamate from astrocytes. *J. Biol. Chem.* 2004b; 279: 12724-12733.
- 327) Zhao H, Asai S, Kanematusu K, Kunimatsu T, Kohno T, Ishakawa K. Real time monitoring of the effects of normothermia and hypothermia on extracellular glutamate re-uptake in the rat following global brain ischemia. *Neuroreport* 1997; 8: 2389-2393.
- 328) Zhou F, Zhu X, Castellani RJ, Stimmelmayer R, Perry G, Smith MA, Drew KL. Hibernation, a model of neuroprotection. *Am. J. Pathol.* 2001;158: 2145-51.
- 329) Zhou F, Braddock JF, Hu Y, Zhu X, Castellani RJ, Smith MA, Drew KL. Microbial origin of glutamate, hibernation and tissue trauma: an *in vivo* microdialysis study. *J. Neurosci. Meth.* 2002; 119: 121-128.
- 330) Zilkha E, Obrenovitsch TP, Koshy A, Kusakabe H, Benetto HP. Extracellular glutamate: on-line monitoring using microdialysis coupled to enzyme-amperometric analysis. *J. Neurosci. Meth.* 1995; 60: 1-9.
- 331) Zimmer J, Kristensen BW, Jakobsen B, Noraberg J. Excitatory amino acid neurotoxicity and modulation of glutamate receptor expression in organotypic brain slice cultures. *Amino Acids* 2000; 19: 7-21.
- 332) Zimmermann H, Lindgren A, Schuhman W, Gorton L. Anisotropic orientation of horseradish peroxidase by reconstitution on a thiol-modified gold electrode. *Chem. Eur. J.* 2000; 6(4): 592-599.
- 333) Zonta M, Angulo MC, Gobbo S, Rosengarten B, Hossmann KA, Pozzan T, Carmignoto G. Neuron-to-astrocyte signalling is central to the dynamic control of brain microcirculation. *Nature Neurosci.* 2003; 6: 43-50.