

University of Groningen

Biochemical and functional characterization of Nudix hydrolase enzymes with novel regulatory roles in Gram positive methylotrophic bacteria

Kloosterman, Harmen

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2005

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kloosterman, H. (2005). *Biochemical and functional characterization of Nudix hydrolase enzymes with novel regulatory roles in Gram positive methylotrophic bacteria*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

References

- Abeygunawardana,C., Weber,D.J., Gittis,A.G., Frick,D.N., Lin,J., Miller,A.F. *et al.* (1995) Solution structure of the MutT enzyme, a nucleoside triphosphate pyrophosphohydrolase. *Biochemistry* **34**: 14997-15005.
- Abrhamova,Z., Patek,M., and Nesvera,J. (2002) Atypical location of double-strand origin of replication (nic site) on the plasmid pGA1 from *Corynebacterium glutamicum*. *Folia Microbiol (Praha)* **47**: 307-310.
- Akiyama,M., Horiuchi,T., and Sekiguchi,M. (1987) Molecular cloning and nucleotide sequence of the mutT mutator of *Escherichia coli* that causes A:T to C:G transversion. *Mol Gen Genet* **206**: 9-16.
- Altschul,S.F., Gish,W., Miller,W., Myers,E.W., and Lipman,D.J. (1990) Basic local alignment search tool. *J Mol Biol* **215**: 403-410.
- Alves,A.M., Meijer,W.G., Vrijbloed,J.W., and Dijkhuizen,L. (1996) Characterization and phylogeny of the *ppf* gene of *Ameycolatopsis methanolica* encoding PPI-dependent phosphofructokinase. *J Bacteriol* **178**: 149-155.
- Anthony,C. (1986) Bacterial oxidation of methane and methanol. *Adv Microb Physiol* **27**: 113-210.
- Anthony,C. (1982) The biochemistry of methylotrophs. Academic press, London.
- Anthony,C., Ghosh,M., and Blake,C.C. (1994) The structure and function of methanol dehydrogenase and related quinoproteins containing pyrrolo-quinoline quinone. *Biochem J* **304**: 665-674.
- Arakawa,K., Muller,R., Mahmud,T., Yu,T.W., and Floss,H.G. (2002) Characterization of the early stage aminoshikimate pathway in the formation of 3-amino-5-hydroxybenzoic acid: The RifN protein specifically converts kanosamine into kanosamine 6-phosphate. *J Am Chem Soc* **124**: 10644-10645.
- Archer,J.A., and Sinskey,A.J. (1993) The DNA sequence and minimal replicon of the *Corynebacterium glutamicum* plasmid pSR1: evidence of a common ancestry with plasmids from *C. diphtheriae*. *J Gen Microbiol* **139**: 1753-1759.
- Arfman,N., de Vries,K.J., Moezelaar,H.R., Attwood,M.M., Robinson,G.K., van Geel,M., and Dijkhuizen,L. (1992a) Environmental regulation of alcohol metabolism in thermotolerant methylotrophic *Bacillus* strains. *Arch Microbiol* **157**: 272-278.
- Arfman,N., Dijkhuizen,L., Kirchof,G., Ludwig,W., Schleifer,K.H., Bulygina,E.S.C.-K. *et al.* (1992b) *Bacillus methanolicus* sp. nov., a new species of thermotolerant, methanol-utilizing, endospore-forming bacteria. *Int J Syst Bacteriol* **42**: 439-445.
- Arfman,N., Hektor,H.J., Bystrykh,L.V., Govorukhina,N.I., Dijkhuizen,L., and Frank,J. (1997) Properties of an NAD(H)-containing methanol dehydrogenase and its activator protein from *Bacillus methanolicus*. *Eur J Biochem* **244**: 426-433.
- Arfman,N., Van Beeumen,J., de Vries,G.E., Harder,W., and Dijkhuizen,L. (1991) Purification and characterization of an activator protein for methanol dehydrogenase from thermotolerant *Bacillus* spp. *J Biol Chem* **266**: 3955-3960.
- Arfman,N., Watling,E.M., Clement,W., van Oosterwijk,R.J., de Vries,G.E., Harder,W., Attwood, M.M., and Dijkhuizen,L. (1989) Methanol metabolism in thermotolerant methylotrophic *Bacillus* strains involving a novel catabolic NAD-dependent methanol dehydrogenase as a key enzyme. *Arch Microbiol* **152**: 280-288.
- Armstrong,V.W., and Eckstein,F. (1976) Interaction of substrate analogues with *Escherichia coli* DNA-dependent RNA polymerase. *Eur J Biochem* **70**: 33-38.
- August,P.R., Tang,L., Yoon,Y.J., Ning,S., Muller,R., Yu,T.W. Taylor,M.; Hoffmann,D., Kim,C.G., Zhang,X., Hutchinson,C.R., and Floss,H.G. (1998) Biosynthesis of the ansamycin antibiotic rifamycin: deductions from the molecular analysis of the *rif* biosynthetic gene cluster of *Ameycolatopsis mediterranei* S699. *Chem Biol* **5**: 69-79.
- Bailey,S., Sedelnikova,S.E., Blackburn,G.M., Abdelghany,H.M., Baker,P.J., McLennan,A.G., and Rafferty,J.B. (2002) The crystal structure of diadenosine tetraphosphate hydrolase from *Caenorhabditis elegans* in free and binary complex forms. *Structure* **10**: 589-600.
- Bairoch,A. (1992) PROSITE: a dictionary of sites and patterns in proteins. *Nucleic Acids Res* **20**: 2013-2018.
- Bairoch,A., Bucher,P., and Hofmann,K. (1996) The PROSITE database, its status in 1995. *Nucleic Acids Res* **24**: 189-196.
- Bastide,A., Laget,M., Patte,J.C., and Dumenil,G. (1989) Methanol metabolism in *Corynebacterium* sp. XG, a facultatively methylotrophic strain. *J Gen Microbiol* **135**: 2869-2874.

References

- Belanger, A.E., and Hatfull, G.F.** (1999) Exponential-phase glycogen recycling is essential for growth of *Mycobacterium smegmatis*. *J Bacteriol* **181**: 6670-6678.
- Bentley, R.** (1990) The Shikimate Pathway. A metabolic tree with many branches. *Crit Rev Biochem Molec Biol* **25**: 307-384.
- Bentley, S.D., Chater, K.F., Cerdeno-Tarraga, A.M., Challis, G.L., Thomson, N.R., James, K.D., Harris, D.E., Quail, M.A., Kieser, H., Harper, D., Bateman, A., Brown, S., Chandra, G., Chen, C.W., Collins, M., Cronin, A., Fraser, A., Goble, A., Hidalgo, J., Hornsby, T., Howarth, S., Huang, C.H., Kieser, T., Larke, L., Murphy, L., Oliver, K., O'Neil, S., Rabinowitsch, E., Rajandream, M.A., Rutherford, K., Rutter, S., Seeger, K., Saunders, D., Sharp, S., Squares, R., Squares, S., Taylor, K., Warren, T., Wietzorrek, A., Woodward, J., Barrell, B.G., Parkhill, J., and Hopwood, D.A.** (2002) Complete genome sequence of the model actinomycete *Streptomyces coelicolor* A3(2). *Nature* **417**: 141-147.
- Bessman, M.J., Frick, D.N., and O'Handley, S.F.** (1996) The MutT proteins or "Nudix" hydrolases, a family of versatile, widely distributed, "housecleaning" enzymes. *J Biol Chem* **271**: 25059-25062.
- Bessman, M.J., Walsh, J.D., Dunn, C.A., Swaminathan, J., Weldon, J.E., and Shen, J.** (2001) The gene *ugdP*, associated with the invasiveness of *Escherichia coli* K1, designates a Nudix hydrolase, Orf176, active on adenosine (5')-pentaphospho-(5')-adenosine (Ap5A). *J Biol Chem* **276**: 37834-37838.
- Bhatnagar, S.K., and Bessman, M.J.** (1988) Studies on the mutator gene, *mutT* of *Escherichia coli*. Molecular cloning of the gene, purification of the gene product, and identification of a novel nucleoside triphosphatase. *J Biol Chem* **263**: 8953-8957.
- Bhatnagar, S.K., Bullions, L.C., and Bessman, M.J.** (1991) Characterization of the *mutT* nucleoside triphosphatase of *Escherichia coli*. *J Biol Chem* **266**: 9050-9054.
- Bibb, M.J., Freeman, R.F., and Hopwood, D.A.** (1977) Physical and genetical characterization of a second sex factor, SCP2, for *Streptomyces coelicolor* A3(2). *Mol Gen Genet* **154**: 155-166.
- Billington, S.J., Jost, B.H., and Songer, J.G.** (1998) The *Arcanobacterium (Actinomyces) pyogenes* plasmid pAPI is a member of the pIJ101/pJV1 family of rolling circle replication plasmids. *J Bacteriol* **180**: 3233-3236.
- Birnboim, H.C., and Doly, J.** (1979) A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* **7**: 1513-1523.
- Bradford, M.M.** (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**: 248-254.
- Branden, C., and Tooze, J.** (1991) Introduction to protein structure. Garland Publishing, Inc., New York and London.
- Brown, D.P., Chiang, S.J., Tuan, J.S., and Katz, L.** (1988) Site-specific integration in *Saccharopolyspora erythraea* and multisite integration in *Streptomyces lividans* of actinomycete plasmid pSE101. *J Bacteriol* **170**: 2287-2295.
- Bystrykh, L., Arfman, N., and Dijkhuizen, L.** (1993a) The methanol-oxidizing enzyme systems in Gram-positive methylotrophic bacteria. In *Microbial growth on C₁ compounds*. Murrell, J.C., and Kelly, D.P. (eds). Intercept Ltd, Andover., pp. 245-251.
- Bystrykh, L.V., Govorukhina, N.I., Dijkhuizen, L., and Duine, J.A.** (1997) Tetrazolium-dye-linked alcohol dehydrogenase of the methylotrophic actinomycete *Amycolatopsis methanolica* is a three-component complex. *Eur J Biochem* **247**: 280-287.
- Bystrykh, L.V., Govorukhina, N.I., van Ophem, P.W., Hektor, H.J., Dijkhuizen, L., and Duine, J.A.** (1993b) Formaldehyde dismutase activities in Gram-positive bacteria oxidizing methanol. *J Gen Microbiol* **139**: 1979-1985.
- Bystrykh, L.V., Vonck, J., van Bruggen, E.F., Van Beeumen, J., Samyn, B., Govorukhina, N.I., Arfman, N., Duine, J.A., and Dijkhuizen, L.** (1993c) Electron microscopic analysis and structural characterization of novel NADP(H)-containing methanol: N,N'-dimethyl-4-nitrosoaniline oxidoreductases from the gram-positive methylotrophic bacteria *Amycolatopsis methanolica* and *Mycobacterium gastri* MB19. *J Bacteriol* **175**: 1814-1822.
- Cabiscol, E., Aguilar, J., and Ros, J.** (1994) Metal-catalyzed oxidation of Fe²⁺ dehydrogenases. Consensus target sequence between propanediol oxidoreductase of *Escherichia coli* and alcohol dehydrogenase II of *Zymomonas mobilis*. *J Biol Chem* **269**: 6592-6597.
- Cai, J.P., Ishibashi, T., Takagi, Y., Hayakawa, H., and Sekiguchi, M.** (2003) Mouse MTH2 protein which prevents mutations caused by 8-oxoguanine nucleotides. *Biochem Biophys Res Commun* **305**: 1073-1077.

References

- Cai, J.P., Kakuma, T., Tsuzuki, T., and Sekiguchi, M. (1995) cDNA and genomic sequences for rat 8-oxo-dGTPase that prevents occurrence of spontaneous mutations due to oxidation of guanine nucleotides. *Carcinogenesis* **16**: 2343-2350.
- Cartwright, J.L., Britton, P., Minnick, M.F., and McLennan, A.G. (1999) The *IalA* invasion gene of *Bartonella bacilliformis* encodes a (de)nucleoside polyphosphate hydrolase of the MutT motif family and has homologs in other invasive bacteria. *Biochem Biophys Res Commun* **256**: 474-479.
- Cartwright, J.L., Gasmil, L., Spiller, D.G., and McLennan, A.G. (2000) The *Saccharomyces cerevisiae* PCD1 gene encodes a peroxisomal nudix hydrolase active toward coenzyme A and its derivatives. *J Biol Chem* **275**: 32925-32930.
- Cartwright, J.L., and Lennan, A.G. (1999) The *Saccharomyces cerevisiae* YOR163w gene encodes a diadenosine 5', 5'''-P1,P6-hexaphosphate (Ap6A) hydrolase member of the MutT motif (Nudix hydrolase) family. *J Biol Chem* **274**: 8604-8610.
- Cartwright, J.L., Safrany, S.T., Dixon, L.K., Darzynkiewicz, E., Stepinski, J., Burke, R., and McLennan, A.G. (2002) The g5R (D250) gene of African swine fever virus encodes a Nudix hydrolase that preferentially degrades diphosphoinositol polyphosphates. *J Virol* **76**: 1415-1421.
- Chen, S., von Bamberg, D., Hale, V., Breuer, M., Hardt, B., Muller, R. *et al.* (1999) Biosynthesis of ansatrienin (mycotrienin) and naphthomycin. Identification and analysis of two separate biosynthetic gene clusters in *Streptomyces collinus* Tu 1892. *Eur J Biochem* **261**: 98-107.
- Chen, Y.M., Lu, Z., and Lin, E.C. (1989) Constitutive activation of the *fucAO* operon and silencing of the divergently transcribed *fucPIK* operon by an IS5 element in *Escherichia coli* mutants selected for growth on L-1,2-propanediol. *J Bacteriol* **171**: 6097-6105.
- Cohen, A., Bar-Nir, D., Goedeke, M.E., and Parag, Y. (1985) The integrated and free states of *Streptomyces griseus* plasmid pSG1. *Plasmid* **13**: 41-50.
- Conway, T., Sewell, G.W., Osman, Y.A., and Ingram, L.O. (1987) Cloning and sequencing of the alcohol dehydrogenase II gene from *Zymomonas mobilis*. *J Bacteriol* **169**: 2591-2597.
- Conyers, G.B., and Bessman, M.J. (1999) The gene, *ialA*, associated with the invasion of human erythrocytes by *Bartonella bacilliformis*, designates a nudix hydrolase active on dinucleoside 5'-polyphosphates. *J Biol Chem* **274**: 1203-1206.
- Cue, D., Lam, H., Dillingham, R.L., Hanson, R.S., and Flickinger, M.C. (1997) Genetic manipulation of *Bacillus methanolicus*, a gram-positive, thermotolerant methylotroph. *Appl Environ Microbiol* **63**: 1406-1420.
- Daniel, R., Boenigk, R., and Gottschalk, G. (1995) Purification of 1,3-propanediol dehydrogenase from *Citrobacter freundii* and cloning, sequencing, and overexpression of the corresponding gene in *Escherichia coli*. *J Bacteriol* **177**: 2151-2156.
- Daubaras, D.L., Hershberger, C.D., Kitano, K., and Chakrabarty, A.M. (1995) Sequence analysis of a gene cluster involved in metabolism of 2,4,5-trichlorophenoxyacetic acid by *Burkholderia cepacia* AC1100. *Appl Environ Microbiol* **61**: 1279-1289.
- De Boer L., Dijkhuizen, L., Grobben, G., Goodfellow, M., Stackebrandt, E., Parlett, J.H., Whitehead, D., and Witt, D. (1990a) *Amycolatopsis methanolica* sp. nov., a facultatively methylotrophic actinomycete. *Int J Syst Bacteriol* **40**: 194-204.
- De Boer L., Grobben, G., Vrijbloed, J.W., and Dijkhuizen, L. (1990b) Biosynthesis of aromatic amino acids in *Nocardia* sp. 239: Effects of amino acid analogues on growth and regulatory enzymes. *Appl Microbiol Biotechnol* **33**: 183-189.
- De Boer, L., Harder, W., and Dijkhuizen, L. (1988) Phenylalanine and tyrosine metabolism in the facultative methylotroph *Nocardia* sp. 239. *Arch Microbiol* **149**: 459-465.
- De Boer, L., Vrijbloed, J.W., Grobben, G., and Dijkhuizen, L. (1989) Regulation of aromatic amino acid biosynthesis in the ribulose monophosphate cycle methylotroph *Nocardia* sp. 239. *Arch Microbiol* **151**: 325.
- De Hoop, M., Asgeirsdottir, S., Blaauw, M., Veenhuis, M., Cregg, J., Gleeson, M., and Geert, A.B. (1991) Mutations in the FAD-binding fold of alcohol oxidase from *Hansenula polymorpha*. *Protein Eng* **4**: 821-829.
- De Vries, G.E., Arfman, N., Terpstra, P., and Dijkhuizen, L. (1992) Cloning, expression, and sequence analysis of the *Bacillus methanolicus* C1 methanol dehydrogenase gene. *J Bacteriol* **174**: 5346-5353.
- Del Solar, G., Giraldo, R., Ruiz-Echevarria, M.J., Espinosa, M., and Diaz-Orejas, R. (1998) Replication and control of circular bacterial plasmids. *Microbiol Mol Biol Rev* **62**: 434-464.

References

- Dijkhuizen,L.** (1993) Methylo-trophs. In *Biotechnology*. Rehm,H.J., Reed,G., Puhler,A., and Stadler,P. (eds). VCH, Weinheim, pp. 265-284.
- Dijkhuizen,L., and Arfman,N.** (1990) Methanol metabolism in thermotolerant methylo-trophic *Bacillus* species. *FEMS Microbiol Rev* **87**: 215-220.
- Dijkhuizen,L., Arfman,N., Attwood,M.M., Brooke,A.G., Harder,W., and Watling,E.M.** (1988) Isolation and initial characterization of thermotolerant methylo-trophic *Bacillus* strains. *FEMS Microbiol Lett* **52**: 209-214.
- Dijkhuizen,L., Levering,P.R., and de Vries,G.E.** (1992) The physiology and biochemistry of aerobic methanol-utilizing Gram-negative and Gram-positive bacteria. In *The Biotechnology Handbook, The Methane and Methanol Utilizers*. Murrell,J.C., and Dalton,H. (eds). New York: Plenum Press, pp. 149-150.
- Dixon,M., and Webb,E.C.** (1979) *Enzymes*. Longman Group Ltd, London.
- Dopheide,T.A., Crewther,P., and Davidson,B.E.** (1972) Chorismate mutase-prephenate dehydratase from *Escherichia coli* K-12. II. Kinetic properties. *J Biol Chem* **247**: 4447-4452.
- Duine,J.A., and Frank,J., Jr.** (1980) The prosthetic group of methanol dehydrogenase. Purification and some of its properties. *Biochem J* **187**: 221-226.
- Duine,J.A., Frank,J., and Berkhout,M.P.** (1984) NAD-dependent, PQQ-containing methanol dehydrogenase: a bacterial dehydrogenase in a multienzyme complex. *FEBS Lett* **168**: 217-221.
- Duine,J.A., Frank,J., and Jongejan,J.A.** (1986) PQQ and quinoprotein enzymes in microbial oxidations. *FEMS Microbiol Rev* **32**: 165-178.
- Dunn,C.A., O'Handley,S.F., Frick,D.N., and Bessman,M.J.** (1999) Studies on the ADP-ribose pyrophosphatase subfamily of the nudix hydrolases and tentative identification of *trgB*, a gene associated with tellurite resistance. *J Biol Chem* **274**: 32318-32324.
- Dunn,M.F., and Bernhard,S.A.** (1971) Rapid kinetic evidence for adduct formation between the substrate analog p-nitroso-N,N-dimethylaniline and reduced nicotinamide-adenine dinucleotide during enzymic reduction. *Biochemistry* **10**: 4569-4575.
- Eshraghi,S., Hancock,I.C., and Williams,E.** (1990) Abstracts 6th Int. Symp. Microbial Growth on C1 Compounds, Göttingen. p. 105.
- Espinosa,J.C., Tercero,J.A., Rubio,M.A., and Jimenez,A.** (1999) The *pur7* gene from the puromycin biosynthetic pur cluster of *Streptomyces alboniger* encodes a nudix hydrolase. *J Bacteriol* **181**: 4914-4918.
- Euverink,G.J., Hessels,G.I., Franke,C., and Dijkhuizen,L.** (1995a) Chorismate mutase and 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase of the methylo-trophic actinomycete *Amycolatopsis methanolica*. *Appl Environ Microbiol* **61**: 3796-3803.
- Euverink,G.J., Vrijbloed,J.W., Hessels,G.I., and Dijkhuizen,L.** (1996) Isolation and analysis of mutants of the methylo-trophic actinomycete *Amycolatopsis methanolica* blocked in aromatic amino acid biosynthesis. *FEMS Microbiol Lett* **136**: 275-281.
- Euverink,G.J., Wolters,D.J., and Dijkhuizen,L.** (1995b) Prephenate dehydratase of the actinomycete *Amycolatopsis methanolica*: purification and characterization of wild-type and deregulated mutant proteins. *Biochem J* **308**: 313-320.
- Fisher,D.I., Safrany,S.T., Strike,P., McLennan,A.G., and Cartwright,J.L.** (2002) Nudix hydrolases that degrade dinucleoside and diphosphoinositol polyphosphates also have 5-phosphoribosyl 1-pyrophosphate (PRPP) pyrophosphatase activity that generates the glycolytic activator ribose 1,5-bisphosphate. *J Biol Chem* **277**: 47313-47317.
- Fowler,R.G., Erickson,J.A., and Isbell,R.J.** (1994) Activity of the *Escherichia coli* *mutT* mutator allele in an anaerobic environment. *J Bacteriol* **176**: 7727-7729.
- Frick,D.N., and Bessman,M.J.** (1995) Cloning, purification, and properties of a novel NADH pyrophosphatase. Evidence for a nucleotide pyrophosphatase catalytic domain in MutT-like enzymes. *J Biol Chem* **270**: 1529-1534.
- Frick,D.N., Townsend,B.D., and Bessman,M.J.** (1995a) A novel GDP-mannose mannosyl hydrolase shares homology with the MutT family of enzymes. *J Biol Chem* **270**: 24086-24091.
- Frick,D.N., Weber,D.J., Abeygunawardana,C., Gittis,A.G., Bessman,M.J., and Mildvan,A.S.** (1995b) NMR studies of the conformations and location of nucleotides bound to the *Escherichia coli* MutT enzyme. *Biochemistry* **34**: 5577-5586.

References

- Frick,D.N., Weber,D.J., Gillespie,J.R., Bessman,M.J., and Mildvan,A.S. (1994) Dual divalent cation requirement of the MutT dGTPase. Kinetic and magnetic resonance studies of the metal and substrate complexes. *J Biol Chem* **269**: 1794-1803.
- Fujikawa,K., Kamiya,H., Yakushiji,H., Fujii,Y., Nakabeppu,Y., and Kasai,H. (1999) The oxidized forms of dATP are substrates for the human MutT homologue, the hMTH1 protein. *J Biol Chem* **274**: 18201-18205.
- Fujikawa,K., and Kasai,H. (2002) The oxidized pyrimidine ribonucleotide, 5-hydroxy-CTP, is hydrolyzed efficiently by the *Escherichia coli* recombinant Orf135 protein. *DNA Repair (Amst)* **1**: 571-576.
- Gabelli,S.B., Bianchet,M.A., Ohnishi,Y., Ichikawa,Y., Bessman,M.J., and Amzel,L.M. (2002) Mechanism of the *Escherichia coli* ADP-ribose pyrophosphatase, a Nudix hydrolase. *Biochemistry* **41**: 9279-9285.
- Gaywee,J., Xu,W., Radulovic,S., Bessman,M.J., and Azad,A.F. (2002) The *Rickettsia prowazekii* invasion gene homolog (*invA*) encodes a Nudix hydrolase active on adenosine (5')-pentaphospho-(5')-adenosine. *Mol Cell Proteomics* **1**: 179-185.
- Glennon,M.C., and Shears,S.B. (1993) Turnover of inositol pentakisphosphates, inositol hexakisphosphate and diphosphoinositol polyphosphates in primary cultured hepatocytes. *Biochem J* **293**: 583-590.
- Goodlove,P.E., Cunningham,P.R., Parker,J., and Clark,D.P. (1989) Cloning and sequence analysis of the fermentative alcohol-dehydrogenase-encoding gene of *Escherichia coli*. *Gene* **85**: 209-214.
- Gosset,G., Bonner,C.A., and Jensen,R.A. (2001) Microbial origin of plant-type 2-keto-3-deoxy-D-arabino-heptulosonate 7-phosphate synthases, exemplified by the chorismate- and tryptophan-regulated enzyme from *Xanthomonas campestris*. *J Bacteriol* **183**: 4061-4070.
- Hagege,J., Pernodet,J.L., Friedmann,A., and Guerineau,M. (1993a) Mode and origin of replication of pSAM2, a conjugative integrating element of *Streptomyces ambofaciens*. *Mol Microbiol* **10**: 799-812.
- Hagege,J., Pernodet,J.L., Sezonov,G., Gerbaud,C., Friedmann,A., and Guerineau,M. (1993b) Transfer functions of the conjugative integrating element pSAM2 from *Streptomyces ambofaciens*: characterization of a *kil-kor* system associated with transfer. *J Bacteriol* **175**: 5529-5538.
- Harder,W., and Veenhuis,M. (1989) Metabolism of one-carbon compounds. In *The Yeasts*. Rose A.H., and Harrison,J.S. (eds). Academic Press, London, pp. 289-316.
- Hayakawa,H., Hofer,A., Thelander,L., Kitajima,S., Cai,Y., Oshiro,S., Yakushiji,H., Nakabeppu,Y., Kuwano,M., and Sekiguchi,M., (1999) Metabolic fate of oxidized guanine ribonucleotides in mammalian cells. *Biochemistry* **38**: 3610-3614.
- Hazeu,W., de Bruyn,J.C., and van Dijken,J.P. (1983) *Nocardia* sp. 239, a facultative methanol utilizer with the ribulose monophosphate pathway of formaldehyde fixation. *Arch Microbiol* **135**: 205-210.
- He,J., Magarvey,N., Pirace,M., and Vining,L.C. (2001) The gene cluster for chloramphenicol biosynthesis in *Streptomyces venezuelae* ISP5230 includes novel shikimate pathway homologues and a monomodular non-ribosomal peptide synthetase gene. *Microbiology* **147**: 2817-2829.
- Hektor,H.J. Physiology and biochemistry of primary alcohol oxidation in the Gram-positive bacteria *Amycolatopsis methanolica* and *Bacillus methanolicus*. Ph.D thesis. 1997. University of Groningen.
- Hektor,H.J., and Dijkhuizen,L. (1996) Mutational analysis of primary alcohol metabolism in the methylotrophic actinomycete *Amycolatopsis methanolica*. *FEMS Microbiol Lett* **144**: 73-79.
- Hektor,H.J., Kloosterman,H., and Dijkhuizen,L. (2000) Nicotinoprotein methanol dehydrogenase enzymes in Gram-positive methylotrophic bacteria. *J Mol Cat B Enzymatic* **8**: 103-109.
- Hektor,H.J., Kloosterman,H., and Dijkhuizen,L. (2002) Identification of a Magnesium-dependent NAD(P)(H)-binding Domain in the Nicotinoprotein Methanol Dehydrogenase from *Bacillus methanolicus*. *J Biol Chem* **277**: 46966-46973.
- Hensgens,C.M., Jansen,M., Nienhuis-Kuiper,M.E., Boekema,E.J., Van Beeumen,J.F.L., and Hansen,T.A. (1995) Purification and characterization of an alcohol dehydrogenase from 1,2-propanediol-grown *Desulfovibrio* strain HDv. *Arch Microbiol* **164**: 265-270.
- Hensgens,C.M., Vonck,J., Van Beeumen,J., van Bruggen,E.F., and Hansen,T.A. (1993) Purification and characterization of an oxygen-labile, NAD-dependent alcohol dehydrogenase from *Desulfovibrio gigas*. *J Bacteriol* **175**: 2859-2863.

References

- Hodgson,D.A.** (2000) Primary metabolism and its control in streptomycetes: a most unusual group of bacteria. *Adv Microb Physiol* **42**: 47-238.
- Hurtado,C., Ruiz,A., Sillero,A., and Sillero,M.A.** (1987) Specific magnesium-dependent diadenosine 5',5'''-P₁,P₃-triphosphate pyrophosphohydrolase in *Escherichia coli*. *J Bacteriol* **169**: 1718-1723.
- Ikeda,H., Ishikawa,J., Hanamoto,A., Shinose,M., Kikuchi,H., Shiba,T., Sakaki,Y., Hattori,M., and Omura,S.** (2003) Complete genome sequence and comparative analysis of the industrial microorganism *Streptomyces avermitilis*. *Nat Biotechnol* **21**: 526-531.
- Ilyina,T.V., and Koonin,E.V.** (1992) Conserved sequence motifs in the initiator proteins for rolling circle DNA replication encoded by diverse replicons from eubacteria, eucaryotes and archaeobacteria. *Nucleic Acids Res* **20**: 3279-3285.
- Ingram,S.W., Stratemann,S.A., and Barnes,L.D.** (1999) *Schizosaccharomyces pombe* Aps1, a diadenosine 5',5''-P₁,P₆-hexaphosphate hydrolase that is a member of the nudix (MutT) family of hydrolases: cloning of the gene and characterization of the purified enzyme. *Biochemistry* **38**: 3649-3655.
- Inoue,M., Kamiya,H., Fujikawa,K., Ootsuyama,Y., Murata,K.N., Osaki,T., Yasumoto,K., and Kasai,H.** (1998) Induction of chromosomal gene mutations in *Escherichia coli* by direct incorporation of oxidatively damaged nucleotides. New evaluation method for mutagenesis by damaged DNA precursors in vivo. *J Biol Chem* **273**: 11069-11074.
- Ishikawa,E., Ogushi,S., Ishikawa,T., and Uyeda,K.** (1990) Activation of mammalian phospho-fructokinases by ribose 1,5-bisphosphate. *J Biol Chem* **265**: 18875-18878.
- Jacoby,G.A., and Shapiro,J.A.** (1977) Plasmids studied in *Pseudomonas aeruginosa* and other Pseudomonads. In DNA insertion elements, Plasmids and Episomes. Bukhari,A.I., Shapiro,J.A., and Adhya,S. (eds). Cold Spring Harbor, NY.: Cold Spring Harbor Laboratory, pp. 639-656.
- Jakubowski,H.** (1986) Sporulation of the yeast *Saccharomyces cerevisiae* is accompanied by synthesis of adenosine 5'-tetraphosphate and adenosine 5'-pentaphosphate. *Proc Natl Acad Sci U S A* **83**: 2378-2382.
- Jornvall,H., Persson,B., and Jeffery,J.** (1987) Characteristics of alcohol/polyol dehydrogenases. The zinc-containing long-chain alcohol dehydrogenases. *Eur J Biochem* **167**: 195-201.
- Kakuma,T., Nishida,J., Tsuzuki,T., and Sekiguchi,M.** (1995) Mouse MTH1 protein with 8-oxo-7,8-dihydro-2'-deoxyguanosine 5'-triphosphatase activity that prevents transversion mutation. cDNA cloning and tissue distribution. *J Biol Chem* **270**: 25942-25948.
- Kamath,A.V., and Yanofsky,C.** (1993) Sequence and characterization of *mutT* from *Proteus vulgaris*. *Gene* **134**: 99-102.
- Kanagasundaram,V., and Scopes,R.K.** (1992) Cloning, sequence analysis, and expression of the structural gene encoding glucose-fructose oxidoreductase from *Zymomonas mobilis*. *J Bacteriol* **174**: 1439-1447.
- Kang,L.W., Gabelli,S.B., Bianchet,M.A., Xu,W.L., Bessman,M.J., and Amzel,L.M.** (2003) Structure of a coenzyme A pyrophosphatase from *Deinococcus radiodurans*: a member of the Nudix family. *J Bacteriol* **185**: 4110-4118.
- Kataoka,M., Seki,T., and Yoshida,T.** (1991) Regulation and function of the *Streptomyces* plasmid pSN22 genes involved in poek formation and inviability. *J Bacteriol* **173**: 7975-7981.
- Kato,N., Miyamoto,N., Shima,M., and Shakazawa,C.** (1988) 3-Hexulose phosphate synthase from a new facultative methylotroph. *Mycobacterium gastri* MB19. *Agric Biol Chem* **52**: 2659-2661.
- Kato,N., Tsui,K., Tani,Y., and Ogushi,S.** (1975) Utilization of methanol by an actinomycete. In Microbial growth on C1-compounds. Tokyo: The Society of Fermentation Technology, pp. 91-98.
- Kato,N., Yamagami,T., Shima,M., and Sakazawa,C.** (1986) Formaldehyde dismutase, a novel NAD-binding oxidoreductase from *Pseudomonas putida* F61. *Eur J Biochem* **156**: 59-64.
- Kendall,K.J., and Cohen,S.N.** (1987) Plasmid transfer in *Streptomyces lividans*: identification of a *kil-kor* system associated with the transfer region of pIJ101. *J Bacteriol* **169**: 4177-4183.
- Kendall,K.J., and Cohen,S.N.** (1988) Complete nucleotide sequence of the *Streptomyces lividans* plasmid pIJ101 and correlation of the sequence with genetic properties. *J Bacteriol* **170**: 4634-4651.
- Kieser,T., Bibb,M., Buttner,M.J., Chater,K.F., and Hopwood,D.A.** (2000) Practical *Streptomyces* Genetics. The John Innes Foundation, Norwich.

References

- Kim,C.G., Kirschning,A., Bergon,P., Zhou,P., Su,E., Sauerbrei,B., Ning,S., Ahn,Y., Breuer,M., Leistner,E., and Floss,H.G. (1996a) Biosynthesis of 3-amino-5-hydroxybenzoic acid, the precursor of mC(7)N units in ansamycin antibiotics. *J Am Chem Soc* **118**: 7486-7491.
- Kim,C.G., Kirschning,A., Bergon,P., Ahn,Y., Wang,J.J., Shibuya,M., and Floss,H.G. (1996b) Formation of 3-amino-5-hydroxybenzoic acid, the precursor of mC₇N Units in ansamycin antibiotics, by a new variant of the Shikimate pathway. *J Am Chem Soc* **114**: 4941-4943.
- Kingston,R.L., Scopes,R.K., and Baker,E.N. (1996) The structure of glucose-fructose oxidoreductase from *Zymomonas mobilis*: an osmoprotective periplasmic enzyme containing non-dissociable NADP. *Structure* **4**: 1413-1428.
- Kisselev,L.L., Justesen,J., Wolfson,A.D., and Frolova,L.Y. (1998) Diadenosine oligophosphates (Ap(n)A), a novel class of signalling molecules? *FEBS Lett* **427**: 157-163.
- Klaus,S.M., Wegkamp,A., Sybesma,W., Hugenholtz,J., Gregory,J.F., III, and Hanson,A.D. (2005) A nudix enzyme removes pyrophosphate from dihydroneopterin triphosphate in the folate synthesis pathway of bacteria and plants. *J Biol Chem* **280**: 5274-5280.
- Kloosterman,H., Vrijbloed,J.W., and Dijkhuizen,L. (2002) Molecular, biochemical and functional characterization of a Nudix hydrolase protein that stimulates activity of a nicotinoprotein alcohol dehydrogenase. *J Biol Chem* **277**: 34785-34792.
- Knaggs,A.R. (1999) The biosynthesis of shikimate metabolites. *Nat Prod Rep* **16**: 525-560.
- Koonin,E.V. (1993) A highly conserved sequence motif defining the family of MutT- related proteins from eubacteria, eukaryotes and viruses. *Nucleic Acids Res* **21**: 4847.
- Kovar,J., Simek,K., Kucera,L., and Matyska,L. (1984) Steady-state kinetics of horse-liver alcohol dehydrogenase with a covalently bound coenzyme analogue. *Eur J Biochem* **139**: 585-591.
- Kreutzer,D.A., and Essigmann,J.M. (1998) Oxidized, deaminated cytosines are a source of C-->T transitions in vivo. *Proc Natl Acad Sci U S A* **95**: 3578-3582.
- Laemmli,U.K., and Favre,K. (1973) Maturation of the head of bacteriophage T4. I. DNA packaging events. *J Mol Biol* **80**: 575-599.
- Lazzarini,A., Cavaletti,L., Toppo,G., and Marinelli,F. (2000) Rare genera of actinomycetes as potential producers of new antibiotics. *Antonie Van Leeuwenhoek* **78**: 399-405.
- Legler,P.M., Lee,H.C., Peisach,J., and Mildvan,A.S. (2002) Kinetic and magnetic resonance studies of the role of metal ions in the mechanism of *Escherichia coli* GDP-mannose mannosyl hydrolase, an unusual nudix enzyme. *Biochemistry* **41**: 4655-4668.
- Legler,P.M., Massiah,M.A., Bessman,M.J., and Mildvan,A.S. (2000) GDP-mannose mannosyl hydrolase catalyzes nucleophilic substitution at carbon, unlike all other Nudix hydrolases. *Biochemistry* **39**: 8603-8608.
- Li,D., and Stevenson,K.J. (1997) Purification and sequence analysis of a novel NADP(H)-dependent type III alcohol dehydrogenase from *Thermococcus* strain AN1. *J Bacteriol* **179**: 4433-4437.
- Lin,J., Abeygunawardana,C., Frick,D.N., Bessman,M.J., and Mildvan,A.S. (1997) Solution structure of the quaternary MutT-M2+-AMPCPP-M2+ complex and mechanism of its pyrophosphohydrolase action. *Biochemistry* **36**: 1199-1211.
- Lundin,A., Nilsson,C., Gerhard,M., Andersson,D.I., Krabbe,M., and Engstrand,L. (2003) The NudA protein in the gastric pathogen *Helicobacter pylori* is an ubiquitous and constitutively expressed dinucleoside polyphosphate hydrolase. *J Biol Chem* **278**: 12574-12578.
- Madon,J., and Hutter,R. (1991) Transformation system for *Amycolatopsis (Nocardia) mediterranei*: direct transformation of mycelium with plasmid DNA. *J Bacteriol* **173**: 6325-6331.
- Madon,J., Moretti,P., and Hutter,R. (1987) Site-specific integration and excision of pMEA100 in *Nocardia mediterranei*. *Mol Gen Genet* **209**: 257-264.
- Makarova,K.S., Aravind,L., Daly,M.J., and Koonin,E.V. (2000) Specific expansion of protein families in the radioresistant bacterium *Deinococcus radiodurans*. *Genetica* **108**: 25-34.
- Maki,H., and Sekiguchi,M. (1992) MutT protein specifically hydrolyses a potent mutagenic substrate for DNA synthesis. *Nature* **355**: 273-275.

References

- Maksel,D., Guranowski,A., Ilgoutz,S.C., Moir,A., Blackburn,M.G., and Gayler,K.R. (1998) Cloning and expression of diadenosine 5',5'''-P₁,P₄-tetraphosphate hydrolase from *Lupinus angustifolius* L. *Biochem J* **329**: 313-319.
- McCandliss,R.J., Poling,M.D., and Herrmann,K.M. (1978) 3-Deoxy-D-arabino-heptulosonate 7-phosphate synthase. Purification and molecular characterization of the phenylalanine- sensitive isoenzyme from *Escherichia coli*. *J Biol Chem* **253**: 4259-4265.
- Mejean,V., Salles,C., Bullions,L.C., Bessman,M.J., and Claverys,J.P. (1994) Characterization of the *mutX* gene of *Streptococcus pneumoniae* as a homologue of *Escherichia coli mutT*, and tentative definition of a catalytic domain of the dGTP pyrophosphohydrolases. *Mol Microbiol* **11**: 323-330.
- Michaels,M.L., and Miller,J.H. (1992) The GO system protects organisms from the mutagenic effect of the spontaneous lesion 8-hydroxyguanine (7,8-dihydro-8-oxoguanine). *J Bacteriol* **174**: 6321-6325.
- Mitchell,S.J., and Minnick,M.F. (1995) Characterization of a two-gene locus from *Bartonella bacilliformis* associated with the ability to invade human erythrocytes. *Infect Immun* **63**: 1552-1562.
- Mo,J.Y., Maki,H., and Sekiguchi,M. (1992) Hydrolytic elimination of a mutagenic nucleotide, 8-oxodGTP, by human 18-kilodalton protein: sanitization of nucleotide pool. *Proc Natl Acad Sci U S A* **89**: 11021-11025.
- Moreno-Bruna,B., Baroja-Fernandez,E., Munoz,F.J., Bastarrica-Berasategui,A., Zanduetta-Criado,A., Rodriguez-Lopez,M., Lasa,I., Akazawa,T., and Pozueta-Romero,J. (2001) Adenosine diphosphate sugar pyrophosphatase prevents glycogen biosynthesis in *Escherichia coli*. *Proc Natl Acad Sci U S A* **98**: 8128-8132.
- Murray,V. (1989) Improved double-stranded DNA sequencing using the linear polymerase chain reaction. *Nucleic Acids Res* **17**: 8889.
- Muth,G., Farr,M., Hartmann,V., and Wohlleben,W. (1995) *Streptomyces ghanaensis* plasmid pSG5: nucleotide sequence analysis of the self-transmissible minimal replicon and characterization of the replication mode. *Plasmid* **33**: 113-126.
- Nagy,I., Verheijen,S., De Schrijver,A., Van Damme,J., Proost,P., Schoofs,G. *et al.* (1995) Characterization of the *Rhodococcus* sp. NI86/21 gene encoding alcohol: N,N'-dimethyl-4-nitrosoaniline oxidoreductase inducible by atrazine and thiocarbamate herbicides. *Arch Microbiol* **163**: 439-446.
- Nair,R.V., Bennett,G.N., and Papoutsakis,E.T. (1994) Molecular characterization of an aldehyde/alcohol dehydrogenase gene from *Clostridium acetobutylicum* ATCC 824. *J Bacteriol* **176**: 871-885.
- Nelms,J., Edwards,R.M., Warwick,J., and Fotheringham,I. (1992) Novel mutations in the *pheA* gene of *Escherichia coli* K-12 which result in highly feedback inhibition-resistant variants of chorismate mutase/prephenate dehydratase. *Appl Environ Microbiol* **58**: 2592-2598.
- Nesvera,J., Patek,M., Hochmannova,J., Abrhamova,Z., Becvarova,V., Jelinkova,M., and Vohradsky,J. (1997) Plasmid pGA1 from *Corynebacterium glutamicum* codes for a gene product that positively influences plasmid copy number. *J Bacteriol* **179**: 1525-1532.
- Nesvera,J., Patek,M., Hochmannova,J., Chibisova,E., Serebrijski,I., Tsygankov,Y., and Netrusov,A. (1991) Transformation of a new Gram-positive methylotroph, *Brevibacterium methylicum*, by plasmid DNA. *Appl Microbiol Biotechnol* **35**: 777-780.
- O'Handley,S.F., Dunn,C.A., and Bessman,M.J. (2000) Orf135 from *Escherichia coli* is a Nudix hydrolase specific for CTP, dCTP, and 5-methyl-dCTP. *J Biol Chem* **276**: 5421-5426.
- O'Handley,S.F., Frick,D.N., Bullions,L.C., Mildvan,A.S., and Bessman,M.J. (1996) *Escherichia coli orf17* codes for a nucleoside triphosphate pyrophosphohydrolase member of the MutT family of proteins. Cloning, purification, and characterization of the enzyme. *J Biol Chem* **271**: 24649-24654.
- O'Handley,S.F., Frick,D.N., Dunn,C.A., and Bessman,M.J. (1998) Orf186 represents a new member of the Nudix hydrolases, active on adenosine(5')triphospho(5')adenosine, ADP-ribose and NADH. *J Biol Chem* **273**: 3192-3197.
- Oda,H., Nakabepu,Y., Furuichi,M., and Sekiguchi,M. (1997) Regulation of expression of the human MTH1 gene encoding 8-oxo-dGTPase. Alternative splicing of transcription products. *J Biol Chem* **272**: 17843-17850.
- Patel,N., Pierson,D.L., and Jensen,R.A. (1977) Dual enzymatic routes to L-tyrosine and L-phenylalanine via pretyrosine in *Pseudomonas aeruginosa*. *J Biol Chem* **252**: 5839-5846.
- Perkins,E.J., Gordon,M.P., Caceres,O., and Lurquin,P.F. (1990) Organization and sequence analysis of the 2,4-dichlorophenol hydroxylase and dichlorocatechol oxidative operons of plasmid pJP4. *J Bacteriol* **172**: 2351-2359.

References

- Perraud,A.L., Fleig,A., Dunn,C.A., Bagley,L.A., Launay,P., Schmitz,C., Stokes,A.J., Zhu,Q., Bessman,M.J., Penner,R., Kinet,J.P., and Scharenberg,A.M. (2001) ADP-ribose gating of the calcium-permeable LTRPC2 channel revealed by Nudix motif homology. *Nature* **411**: 595-599.
- Perraud,A.L., Takanishi,C.L., Shen,B., Kang,S., Smith,M.K., Schmitz,C., Knowles,H.M., Ferraris,D., Li,W., Zhang,J., Stoddard,B.L., and Scharenberg,A.M. (2005) Accumulation of free ADP-ribose from mitochondria mediates oxidative stress-induced gating of TRPM2 cation channels. *J Biol Chem* **280**: 6138-6148.
- Piersma,S.R. (1998) Structure and catalytic mechanism of nicotinoprotein alcohol dehydrogenases from *Amycolatopsis methanolica*, Ph.D thesis. University of Delft.
- Pierson,L.S., Gaffney,T., Lam,S., and Gong,F. (1995) Molecular analysis of genes encoding phenazine biosynthesis in the biological control bacterium. *Pseudomonas aureofaciens* 30-84. *FEMS Microbiol Lett* **134**: 299-307.
- Pohnert,G., Zhang,S., Husain,A., Wilson,D.B., and Ganem,B. (1999) Regulation of phenylalanine biosynthesis. Studies on the mechanism of phenylalanine binding and feedback inhibition in the *Escherichia coli* P-protein. *Biochemistry* **38**: 12212-12217.
- Possoz,C., Gagnat,J., Sezonov,G., Guerineau,M., and Pernodet,J.L. (2003) Conjugal immunity of *Streptomyces* strains carrying the integrative element pSAM2 is due to the *pif* gene (pSAM2 immunity factor). *Mol Microbiol* **47**: 1385-1393.
- Quan,S., and Dabbs,E.R. (1993) Nocardioform arsenic resistance plasmid characterization and improved *Rhodococcus* cloning vectors. *Plasmid* **29**: 74-79.
- Raffaelli,N., Lorenzi,T., Amici,A., Emanuelli,M., Ruggieri,S., and Magni,G. (1999) *Synechocystis* sp. Slr0787 protein is a novel bifunctional enzyme endowed with both nicotinamide mononucleotide adenylyltransferase and 'Nudix' hydrolase activities. *FEBS Lett* **444**: 222-226.
- Ray,J.M., and Bauerle,R. (1991) Purification and properties of tryptophan-sensitive 3-deoxy-D- arabino-heptulosonate-7-phosphate synthase from *Escherichia coli*. *J Bacteriol* **173**: 1894-1901.
- Raynal,A., Friedmann,A., Tophile,K., Guerineau,M., and Pernodet,J.L. (2002) Characterization of the *attP* site of the integrative element pSAM2 from *Streptomyces ambifaciens*. *Microbiology* **148**: 61-67.
- Read,T.D., Peterson,S.N., Tourasse,N., Baillie,L.W., Paulsen,I.T., Nelson,K.E., Tettelin,H., Fouts,D.E., Eisen,J.A., Gill,S.R., Holtzapple,E.K., Okstad,O.A., Helgason,E., Rilstone,J., Wu,M., Kolonay,J.F., Beanan,M.J., Dodson,R.J., Brinkac,L.M., Gwinn,M., DeBoy,R.T., Madpu,R., Daugherty,S.C., Durkin,A.S., Haft,D.H., Nelson,W.C., Peterson,J.D., Pop,M., Khouri,H.M., Radune,D., Benton,J.L., Mahamoud,Y., Jiang,L., Hance,I.R., Weidman,J.F., Berry,K.J., Plaut,R.D., Wolf,A.M., Watkins,K.L., Nierman,W.C., Hazen,A., Cline,R., Redmond,C., Thwaite,J.E., White,O., Salzberg,S.L., Thomason,B., Friedlander,A.M., Koehler,T.M., Hanna,P.C., Kolsto,A.B., and Fraser,C.M. (2003) The genome sequence of *Bacillus anthracis* Ames and comparison to closely related bacteria. *Nature* **423**: 81-86.
- Reid,M.F., and Fewson,C.A. (1994) Molecular characterization of microbial alcohol dehydrogenases. *Crit Rev Microbiol* **20**: 13-56.
- Ruzhenikov,S.N., Burke,J., Sedelnikova,S., Baker,P.J., Taylor,R., Bullough,P.A., Muir,N.M., Gore,M.G., and Rice,D.W. (2001) Glycerol dehydrogenase. structure, specificity, and mechanism of a family III polyol dehydrogenase. *Structure* **9**: 789-802.
- Safrany,S.T., Caffrey,J.J., Yang,X., Bembenek,M.E., Moyer,M.B., Burkhart,W.A., and Shears,S.B. (1998) A novel context for the 'MutT' module, a guardian of cell integrity, in a diphosphoinositol polyphosphate phosphohydrolase. *EMBO J* **17**: 6599-6607.
- Safrany,S.T., Ingram,S.W., Cartwright,J.L., Falck,J.R., McLennan,A.G., Barnes,L.D., and Shears,S.B. (1999) The diadenosine hexaphosphate hydrolases from *Schizosaccharomyces pombe* and *Saccharomyces cerevisiae* are homologues of the human diphosphoinositol polyphosphate phosphohydrolase. Overlapping substrate specificities in a MutT-type protein. *J Biol Chem* **274**: 21735-21740.
- Sakumi,K., Furuichi,M., Tsuzuki,T., Kakuma,T., Kawabata,S., Maki,H., and Sekiguchi,M. (1993) Cloning and expression of cDNA for a human enzyme that hydrolyzes 8-oxo-dGTP, a mutagenic substrate for DNA synthesis. *J Biol Chem* **268**: 23524-23530.
- Sambrook,J., Frisch,E.F., and Maniatis,T. (1989) Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.

References

- Sander,C., and Schneider,R. (1991) Database of homology-derived protein structures and the structural meaning of sequence alignment. *Proteins* **9**: 56-68.
- Schoner,R., and Herrmann,K.M. (1976) 3-Deoxy-D-arabino-heptulosonate 7-phosphate synthase. Purification, properties, and kinetics of the tyrosine-sensitive isoenzyme from *Escherichia coli*. *J Biol Chem* **251**: 5440-5447.
- Seery,L.T., Nolan,N.C., Sharp,P.M., and Devine,K.M. (1993) Comparative analysis of the pC194 group of rolling circle plasmids. *Plasmid* **30**: 185-196.
- Seifert,H.S., Chen,E.Y., So,M., and Heffron,F. (1986) Shuttle mutagenesis: a method of transposon mutagenesis for *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A* **83**: 735-739.
- Sekhar,V.C., and Plapp,B.V. (1990) Rate constants for a mechanism including intermediates in the interconversion of ternary complexes by horse liver alcohol dehydrogenase. *Biochemistry* **29**: 4289-4295.
- Servin-Gonzalez,L. (1993) Relationship between the replication functions of *Streptomyces* plasmids pJV1 and pIJ101. *Plasmid* **30**: 131-140.
- Sezonov,G., Possoz,C., Friedmann,A., Pernodet,J.L., and Guerineau,M. (2000) KorSA from the *Streptomyces* integrative element pSAM2 is a central transcriptional repressor: target genes and binding sites. *J Bacteriol* **182**: 1243-1250.
- Sharples,G.J., and Lloyd,R.G. (1991) Resolution of Holliday junctions in *Escherichia coli*: identification of the *ruvC* gene product as a 19-kilodalton protein. *J Bacteriol* **173**: 7711-7715.
- Shears,S.B., Ali,N., Craxton,A., and Bembenek,M.E. (1995) Synthesis and metabolism of bis-diphosphoinositol tetrakisphosphate in vitro and in vivo. *J Biol Chem* **270**: 10489-10497.
- Sheehan,M.C., Bailey,C.J., Dowds,B.C., and McConnell,D.J. (1988) A new alcohol dehydrogenase, reactive towards methanol, from *Bacillus stearothermophilus*. *Biochem J* **252**: 661-666.
- Sheikh,S., O'Handley,S.F., Dunn,C.A., and Bessman,M.J. (1998) Identification and characterization of the Nudix hydrolase from the Archaeon, *Methanococcus jannaschii*, as a highly specific ADP-ribose pyrophosphatase. *J Biol Chem* **273**: 20924-20928.
- Shultz,J., Hermodson,M.A., Garner,C.C., and Herrmann,K.M. (1984) The nucleotide sequence of the *aroF* gene of *Escherichia coli* and the amino acid sequence of the encoded protein, the tyrosine-sensitive 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase. *J Biol Chem* **259**: 9655-9661.
- Silakowski,B., Kunze,B., and Muller,R. (2000) *Stigmatella aurantiaca* Sg a15 carries genes encoding type I and type II 3-deoxy-D-arabino-heptulosonate-7-phosphate synthases: involvement of a type II synthase in aurachin biosynthesis. *Arch Microbiol* **173**: 403-411.
- Sofia,H.J., Burland,V., Daniels,D.L., Plunkett,G., III, and Blattner,F.R. (1994) Analysis of the *Escherichia coli* genome. V. DNA sequence of the region from 76.0 to 81.5 minutes. *Nucleic Acids Res* **22**: 2576-2586.
- Sohling,B., and Gottschalk,G. (1996) Molecular analysis of the anaerobic succinate degradation pathway in *Clostridium kluyveri*. *J Bacteriol* **178**: 871-880.
- Stephens,L., Radenberg,T., Thiel,U., Vogel,G., Khoo,K.H., Dell,A., Jackson,T.R., Hawkins,P.T., and Mayr,G.W. (1993) The detection, purification, structural characterization, and metabolism of diphosphoinositol pentakisphosphate(s) and bisdiphosphoinositol tetrakisphosphate(s). *J Biol Chem* **268**: 4009-4015.
- Stojiljkovic,I., Baumler,A.J., and Heffron,F. (1995) Ethanolamine utilization in *Salmonella typhimurium*: nucleotide sequence, protein expression, and mutational analysis of the *cchA cchB eutE eutJ eutG eutH* gene cluster. *J Bacteriol* **177**: 1357-1366.
- Sugimoto,S., and Shiio,I. (1980) Purification and properties of bifunctional 3-deoxy-D-arabino-heptulosonate 7-phosphate synthetase-chorismate mutase component A from *Brevibacterium flavum*. *J Biochem Tokyo* **87**: 881-890.
- Suzuki,I., Seki,T., and Yoshida,T. (1997) Nucleotide sequence of a nicking site of the *Streptomyces* plasmid pSN22 replicating by the rolling circle mechanism. *FEMS Microbiol Lett* **150**: 283-288.
- Taddei,F., Hayakawa,H., Bouton,M., Cirinesi,A., Matic,I., Sekiguchi,M., and Radman,M. (1997) Counteraction by MutT protein of transcriptional errors caused by oxidative damage. *Science* **278**: 128-130.
- Tamarit,J., Cabiscol,E., Aguilar,J., and Ros,J. (1997) Differential inactivation of alcohol dehydrogenase isoenzymes in *Zymomonas mobilis* by oxygen. *J Bacteriol* **179**: 1102-1104.

References

- Tauch,A., Bischoff,N., Brune,I., and Kalinowski,J. (2003) Insights into the genetic organization of the *Corynebacterium diphtheriae* erythromycin resistance plasmid pNG2 deduced from its complete nucleotide sequence. *Plasmid* **49**: 63-74.
- Tercero,J.A., Espinosa,J.C., Lacalle,R.A., and Jimenez,A. (1996) The biosynthetic pathway of the aminonucleoside antibiotic puromycin, as deduced from the molecular analysis of the *pur* cluster of *Streptomyces alboniger*. *J Biol Chem* **271**: 1579-1590.
- Treffers,H.P., Spinelli,V., and Belser,O. (1999) A factor (or mutator gene) influencing mutation rates in *Escherichia coli*. *Proc Natl Acad Sci U S A* **40**: 1064-1071.
- Tsuzuki,T., Egashira,A., Igarashi,H., Iwakuma,T., Nakatsuru,Y., Tominaga,Y., Kawate,H., Nakao,K., Nakamura,K., Ide,F., Kura,S., Nakabeppu,Y., Katsuki,M., Ishikawa,T., and Sekiguchi,N. (2001) Spontaneous tumorigenesis in mice defective in the MTH1 gene encoding 8-oxo-dGTPase. *Proc Natl Acad Sci U S A* **98**: 11456-11461.
- Valentin,H.E., Zwingmann,G., Schonebaum,A., and Steinbuechel,A. (1995) Metabolic pathway for biosynthesis of poly(3-hydroxybutyrate-co-4-hydroxybutyrate) from 4-hydroxybutyrate by *Alcaligenes eutrophus*. *Eur J Biochem* **227**: 43-60.
- Van der Geize,R., Hessels,G.I., van Gerwen,R., Vrijbloed,J.W., van der Meijden P., and Dijkhuizen,L. (2000) Targeted disruption of the *kstD* gene encoding a 3-ketosteroid delta(1)-dehydrogenase isoenzyme of *Rhodococcus erythropolis* strain SQ1. *Appl Environ Microbiol* **66**: 2029-2036.
- Van der Meer,J.R., Eggen,R.I., Zehnder,A.J., and de Vos,W.M. (1991) Sequence analysis of the *Pseudomonas* sp. strain P51 *tcb* gene cluster, which encodes metabolism of chlorinated catechols: evidence for specialization of catechol 1,2-dioxygenases for chlorinated substrates. *J Bacteriol* **173**: 2425-2434.
- Van Dijk,E., Cougot,N., Meyer,S., Babajko,S., Wahle,E., and Seraphin,B. (2002) Human Dep2: a catalytically active mRNA decapping enzyme located in specific cytoplasmic structures. *EMBO J* **21**: 6915-6924.
- Van Ophem,P., and Duine,J.A. (1990) Methanol dehydrogenase from *Nocardia* sp. 230. In PQQ and quinoproteins. Jongejan,J.A., and Duine,J.A. (eds). Kluwer Academic Publishers, Dordrecht, pp. 50-51.
- Van Ophem,P., Euverink,G.J., Dijkhuizen,L., and Duine,J.A. (1991) A novel dye-linked alcohol dehydrogenase activity present in some Gram-positive bacteria. *FEMS Microbiol Lett* **80**: 57-64.
- Van Ophem,P.W., and Duine,J.A. (1993) In *Enzymology and molecular biology of carbonyl metabolism* Vol. 4. Weiner,H., and Crabb,D.W. (eds). Plenum Press, New York. p. 605.
- Van Wageningen AM, Kirkpatrick,P.N., Williams,D.H., Harris,B.R., Kershaw,J.K., Lennard,N.J., Jones,M., Jones,S.J., and Solenberg,P.J. (1998) Sequencing and analysis of genes involved in the biosynthesis of a vancomycin group antibiotic. *Chem Biol* **5**: 155-162.
- Venkova-Canova,T., Patek,M., and Nesvera,J. (2003) Control of *rep* gene expression in plasmid pGA1 from *Corynebacterium glutamicum*. *J Bacteriol* **185**: 2402-2409.
- Vonck,J., Arfman,N., de Vries,G.E., Van Beeumen,J., van Bruggen,E.F., and Dijkhuizen,L. (1991) Electron microscopic analysis and biochemical characterization of a novel methanol dehydrogenase from the thermotolerant *Bacillus* sp. C1. *J Biol Chem* **266**: 3949-3954.
- Vrijbloed,J.W. (1996) Functional analysis of the integrative plasmid pMEA300 of the actinomycete *Amycolatopsis methanolica*. PhD-thesis. University of Groningen.
- Vrijbloed,J.W., Jelinkova,M., Hessels,G.I., and Dijkhuizen,L. (1995a) Identification of the minimal replicon of plasmid pMEA300 of the methylotrophic actinomycete *Amycolatopsis methanolica*. *Mol Microbiol* **18**: 21-31.
- Vrijbloed,J.W., Madon,J., and Dijkhuizen,L. (1994) A plasmid from the methylotrophic actinomycete *Amycolatopsis methanolica* capable of site-specific integration. *J Bacteriol* **176**: 7087-7090.
- Vrijbloed,J.W., Madon,J., and Dijkhuizen,L. (1995b) Transformation of the methylotrophic actinomycete *Amycolatopsis methanolica* with plasmid DNA: stimulatory effect of a pMEA300-encoded gene. *Plasmid* **34**: 96-104.
- Vrijbloed,J.W., van der Put N.M., and Dijkhuizen,L. (1995c) Identification and functional analysis of the transfer region of plasmid pMEA300 of the methylotrophic actinomycete *Amycolatopsis methanolica*. *J Bacteriol* **177**: 6499-6505.
- Vrijbloed,J.W., van Hylckama Vlieg,J., van der Put,N.M., Hessels,G.I., and Dijkhuizen,L. (1995d) Molecular cloning with a pMEA300-derived shuttle vector and characterization of the *Amycolatopsis methanolica* prephenate dehydratase gene. *J Bacteriol* **177**: 6666-6669.

References

- Walker,G.E., Dunbar,B., Hunter,I.S., Nimmo,H.G., and Coggins,J.R. (1996) Evidence for a novel class of microbial 3-deoxy-D-arabino- heptulosonate-7-phosphate synthase in *Streptomyces coelicolor* A3(2), *Streptomyces rimosus* and *Neurospora crassa*. *Microbiology* **142**: 1973-1982.
- Walter,K.A., Bennett,G.N., and Papoutsakis,E.T. (1992) Molecular characterization of two *Clostridium acetobutylicum* ATCC 824 butanol dehydrogenase isozyme genes. *J Bacteriol* **174**: 7149-7158.
- Wang,Z., Jiao,X., Carr-Schmid,A., and Kiledjian,M. (2002) The hDcp2 protein is a mammalian mRNA decapping enzyme. *Proc Natl Acad Sci U S A* **99**: 12663-12668.
- Weber,D.J., Abeygunawardana,C., Bessman,M.J., and Mildvan,A.S. (1993) Secondary structure of the MutT enzyme as determined by NMR. *Biochemistry* **32**: 13081-13088.
- Weber,D.J., Bhatnagar,S.K., Bullions,L.C., Bessman,M.J., and Mildvan,A.S. (1992) NMR and isotopic exchange studies of the site of bond cleavage in the MutT reaction. *J Biol Chem* **267**: 16939-16942.
- Wiegert,T., Sahn,H., and Sprenger,G.A. (1997) The substitution of a single amino acid residue (Ser-116 --> Asp) alters NADP-containing glucose-fructose oxidoreductase of *Zymomonas mobilis* into a glucose dehydrogenase with dual coenzyme specificity. *J Biol Chem* **272**: 13126-13133.
- Wierenga,R.K., Drenth,J., and Schulz,G.E. (1983) Comparison of the three-dimensional protein and nucleotide structure of the FAD-binding domain of p-hydroxybenzoate hydroxylase with the FAD- as well as NADPH-binding domains of glutathione reductase. *J Mol Biol* **167**: 725-739.
- Wierenga,R.K., Terpstra,P., and Hol,W.G. (1986) Prediction of the occurrence of the ADP-binding beta alpha beta-fold in proteins, using an amino acid sequence fingerprint. *J Mol Biol* **187**: 101-107.
- Williamson,V.M., and Paquin,C.E. (1987) Homology of *Saccharomyces cerevisiae* ADH4 to an iron-activated alcohol dehydrogenase from *Zymomonas mobilis*. *Mol Gen Genet* **209**: 374-381.
- Wong,J.T. (1975) Kinetics of enzyme mechanisms. Academic Press, London.
- Woodward,J.R. (1990) In Advances in autotrophic microbiology and one-carbon metabolism. Codd,G.A., Dijkhuizen,L., and Tabita,F.R. (eds). Kluwer Academic Publishers, Dordrecht. pp. 193-225.
- Xia,T., Zhao,G., and Jensen,R.A. (1992) Loss of allosteric control but retention of the bifunctional catalytic competence of a fusion protein formed by excision of 260 base pairs from the 3' terminus of pheA from *Erwinia herbicola*. *Appl Environ Microbiol* **58**: 2792-2798.
- Xu,W., Dunn,C.A., Jones,C.R., D'Souza,G., and Bessman,M.J. (2004) The 26 Nudix hydrolases of *Bacillus cereus*, a close relative of *Bacillus anthracis*. *J Biol Chem* **279**: 24861-24865.
- Xu,W., Gauss,P., Shen,J., Dunn,C.A., and Bessman,M.J. (2002) The gene e.1 (*nudE.1*) of T4 bacteriophage designates a new member of the Nudix hydrolase superfamily active on flavin adenine dinucleotide, adenosine 5'-triphospho-5'-adenosine, and ADP-ribose. *J Biol Chem* **277**: 23181-23185.
- Xu,W., Shen,J., Dunn,C.A., Desai,S., and Bessman,M.J. (2001) The Nudix hydrolases of *Deinococcus radiodurans*. *Mol Microbiol* **39**: 286-290.
- Yang,W., Li,E., Kairong,T., and Stanley,S.L., Jr. (1994) *Entamoeba histolytica* has an alcohol dehydrogenase homologous to the multifunctional *adhE* gene product of *Escherichia coli*. *Mol Biochem Parasitol* **64**: 253-260.
- Yanofsky,C., Cox,E.C., and Horn,V. (1966) The unusual mutagenic specificity of an *E. coli* mutator gene. *Proc Natl Acad Sci U S A* **55**: 274-281.
- Youngleson,J.S., Jones,W.A., Jones,D.T., and Woods,D.R. (1989) Molecular analysis and nucleotide sequence of the *adh1* gene encoding an NADPH-dependent butanol dehydrogenase in the Gram-positive anaerobe *Clostridium acetobutylicum*. *Gene* **78**: 355-364.
- Yu,T.W., Muller,R., Muller,M., Zhang,X., Draeger,G., Kim,C.G. *et al.* (2001) Mutational analysis and reconstituted expression of the biosynthetic genes involved in the formation of 3-amino-5-hydroxybenzoic acid, the starter unit of rifamycin biosynthesis in *Amycolatopsis mediterranei* S699. *J Biol Chem* **276**: 12546-12555.
- Zhang,S., Pohnert,G., Kongsaree,P., Wilson,D.B., Clardy,J., and Ganem,B. (1998) Chorismate mutase-prephenate dehydratase from *Escherichia coli*. Study of catalytic and regulatory domains using genetically engineered proteins. *J Biol Chem* **273**: 6248-6253.

References

- Zhang, Y., Praszker, J., Hodgson, A., and Pittard, A. J.** (1994) Molecular analysis and characterization of a broad-host-range plasmid, pEP2. *J Bacteriol* **176**: 5718-5728.
- Ziegler, M.** (2000) New functions of a long-known molecule. Emerging roles of NAD in cellular signaling. *Eur J Biochem* **267**: 1550-1564.

