

University of Groningen

Mechano- and electrophysiological studies on cochlear hair cells and lateral line cupulae

Dinklo, Theodorus

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2005

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Dinklo, T. (2005). *Mechano- and electrophysiological studies on cochlear hair cells and lateral line cupulae*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

-
- Alexandre, D. and Ghysen, A. (1999) Somatotopy of the lateral line projection in larval zebrafish. *Proc. Natl. Acad. Sci. U.S.A.* **96**, 7558-7562.
- Assad, J.A., Hacohen, N. and Corey, D.P. (1989) Voltage dependence of adaptation and active bundle movement in bullfrog saccular hair cells. *Proc. Natl. Acad. Sci. U.S.A.* **86**, 2918-2922.
- Assad, J.A. and Corey, D.P. (1992) An active motor model for adaptation by vertebrate hair cells. *J. Neurosci.* **12**, 3291-3309.
- Batchelor, G.K. (1967) *An introduction to fluid dynamics*. Cambridge University Press, Cambridge.
- Benser, M.E., Marquis, R.E. and Hudspeth, A.J. (1996) Rapid, active hair bundle movements in hair cells from the bullfrog's sacculus. *J. Neurosci.* **16**, 5629-5643.
- Bleckmann, H. (1980) Reaction time and stimulus frequency in prey localization in the surface-feeding fish *Aplocheilichthys lieatus*. *J. Comp. Physiol. A* **140**, 163-172.
- Bleckmann, H. (1993) Role of the lateral line in fish behaviour. In *Behaviour of Teleost Fishes*, ed. Pitcher, T.J., pp. 201-246. Chapman and Hall.
- Bosher, S.K. and Warren, R.L. (1978) Very low calcium content of cochlear endolymph, an extracellular fluid. *Nature* **273**, 377-378.
- Colquhoun, D. and Hawkes, A.G. (1995) The principles of the stochastic interpretation of ion-channel mechanisms. In *Single-Channel Recording*, eds. Sakmann, B. and Neher, E., pp. 397-482. Plenum Press, New York.
- Coombs, S., Janssen, J. and Webb, W.W. (1987) Diversity of lateral line systems: evolutionary and functional considerations. In *Sensory biology of aquatic animals*, eds. Atema, J., Fay, R.R., Popper, A.N. and Tavolga, W.N., pp. 553-593. Springer-Verlag, New York.
- Coombs, S. and Montgomery, J. (1994) Function and evolution of superficial neuromasts in an Antarctic notothenioid fish. *Brain Behav. Evol.* **44**, 287-298.
- Corey, D.P. and Hudspeth, A.J. (1979) Response latency of vertebrate hair cells. *Biophys. J.* **26**, 499-506.
- Corey, D.P. and Hudspeth, A.J. (1983) Kinetics of the receptor current in bullfrog saccular hair cells. *J. Neurosci.* **3**, 962-976.
- Corey, D.P., Garcia-Anoveros, J., Holt, J.R., Kwan, K.Y., Lin, S.Y., Vollrath, M.A., Amalfitano, A., Cheung, E.L., Derfler, B.H., Duggan, A., Geleoc, G.S., Gray, P.A., Hoffman, M.P., Rehm, H.L., Tamasauskas, D., & Zhang, D.S. (2004) TRPA1 is a candidate for the mechanosensitive transduction channel of vertebrate hair cells. *Nature* **432**, 723-730

- Crawford, A.C. and Fettiplace, R. (1983) Auditory nerve responses to imposed displacements of the turtle basilar membrane. *Hear. Res.* **12**, 199-208.
- Crawford, A.C. and Fettiplace, R. (1985) The mechanical properties of ciliary bundles of turtle cochlear hair cells. *J. Physiol* **364**, 359-379.
- Crawford, A.C., Evans, M.G. and Fettiplace, R. (1989) Activation and adaptation of transducer currents in turtle hair cells. *J. Physiol* **419**, 405-434.
- Crawford, A.C., Evans, M.G. and Fettiplace, R. (1991) The actions of calcium on the mechano-electrical transducer current of turtle hair cells. *J. Physiol.* **434**, 369-398.
- Ćurčić-Blake, B. and van Netten, S.M. (2005) Rapid responses of the cupula in the lateral line of ruffe (*Gymnocephalus cernuus*). *J. Comp. Physiol. A* (Accepted for publication)
- Dallos, P. and Fakler, B. (2002) Prestin, a new type of motor protein. *Nat. Rev. Mol. Cell Biol.* **3**, 104-111.
- Dallos, P., Pooper, A.N. and Fay, R.R. (1996) *The Cochlea*, Springer, New York.
- Denk, W. and Webb, W.W. (1992) Forward and reverse transduction at the limit of sensitivity studied by correlating electrical and mechanical fluctuations in frog saccular hair cells. *Hear. Res.* **60**, 89-102.
- Denk, W., Holt, J.R., Shepherd, G.M. and Corey, D.P. (1995) Calcium imaging of single stereocilia in hair cells: localization of transduction channels at both ends of tip links. *Neuron* **15**, 1311-1321.
- Dijkgraaf, S. (1963) The functioning and significance of the lateral line organs. *Biol. Rev.* **38**, 55-105.
- Dinklo, T., van Netten, S.M., Marcotti, W. and Kros, C.J. (2003) Signal processing by transducer channels in mammalian outer hair cells. In *Biophysics of the Cochlea: From Molecule to Model*. Ed. Gummer, A. W., pp. 73-79, World Scientific, Singapore.
- Eatock, R.A., Corey, D.P. and Hudspeth, A.J. (1987) Adaptation of mechano-electrical transduction in hair cells of the bullfrog's sacculus. *J. Neurosci.* **7**, 2821-2836.
- Eatock, R.A. (2000) Adaptation in hair cells. *Annu. Rev. Neurosci.* **23**, 285-314.
- Ehret, G. (1983) Development of hearing and response behavior to sound stimuli: behavioral studies. In *Development of the Auditory and Vestibular Systems*, ed. Romand, R., pp. 211-237. Academic Press, New York.
- Engelmann, J., Hanke, W., Mogdans, J. and Bleckmann, H. (2000) Hydrodynamic stimuli and the fish lateral line. *Nature* **408**, 51-52.

-
- Enger, P.S., Kalmijn, A.J. and Sand, O. (1989) Behavioral investigations on the functions of the lateral line and inner ear in predation. In *The mechanosensory lateral line: neurobiology and evolution*, eds. Coombs, S., Görner, P. and Münz, H., pp. 575-587. Springer-Verlag, New York.
- Ernstrom, G.G. and Chalfie, M. (2002) Genetics of sensory mechanotransduction. *Annu. Rev. Genet.* **36**, 411-453.
- Fante, R.L. (1986) *Signal analysis and estimation: An introduction* Wiley, New York.
- Fernandez, C. and Schmidt, R.S. (1963) Development of mammalian endocochlear potential. *J. Exp. Zool.* **153**, 227-235.
- Fettiplace, R., Ricci, A.J. and Hackney, C.M. (2001) Clues to the cochlear amplifier from the turtle ear. *Trends Neurosci.* **24**, 169-175.
- Flock, Å., Kimura, R., Lundquist, P.G. and Wersäll, J. (1962) Morphological basis of directional sensitivity of the outer hair cells in the organ of Corti. *J. Acoust. Soc. Am.* **34**, 1351-1355.
- Flock, Å. (1967) Ultrastructure and function in the lateral line organs. In *Lateral line detectors*, ed. Cahn, Ph.H., pp. 163-197. Indiana University Press.
- Flock, Å. and Cheung, H. C. (1977) Actin filaments in sensory hairs of inner ear receptor cells. *J. Cell Biol.* **75**, 339-343.
- Flock, Å. and Orman, S. (1983) Micromechanical properties of sensory hairs on receptor cells of the inner ear. *Hear. Res.* **11**, 249-260.
- Géléoc, G.S., Lennan, G.W., Richardson, G.P. and Kros, C.J. (1997) A quantitative comparison of mechano-electrical transduction in vestibular and auditory hair cells of neonatal mice. *Proc. R. Soc. Lond B Biol. Sci.* **264**, 611-621.
- Géléoc, G.S. and Holt, J.R. (2003) Developmental acquisition of sensory transduction in hair cells of the mouse inner ear. *Nat. Neurosci.* **6**, 1019-1020.
- Gillespie, P.G. and Walker, R.G. (2001) Molecular basis of mechanosensory transduction. *Nature* **413**, 194-202.
- Görner, P. (1963) Untersuchungen zur Morphologie und Elektrophysiologie des Seitenlinienorgans vom Krallenfrosch (*Xenopus laevis* Daudin). *Zeitschrift für vergleichende Physiologie* **47**, 316-338.
- Hacohen, N., Assad, J.A., Smith, W.J. and Corey, D.P. (1989) Regulation of tension on hair-cell transduction channels: displacement and calcium dependence. *J. Neurosci.* **9**, 3988-3997.

- Hassan, E.-S., Abdel-Latif, H. and Biebricher, R. (1992) Studies on the effects of Ca^{++} and Co^{++} on the swimming behavior of the blind Mexican cave fish. *J. Comp. Physiol. A* **171**, 413-419.
- Hill, J.K. and Gillespie, P.G. (2003) Seeing touch: moving closer to the worm mechanotransduction complex. *Curr. Biol.* **13**, R967-R969.
- Hille, B. (1992) *Ionic Channels of Excitable Membranes*, 2nd edition, Sinauer Associates Inc., Sunderland.
- Hodgkin, A.L. and Huxley, A.F. (1952) A quantitative description of membrane currents and its application to conduction and excitation in nerve. *J. Physiol.(Lond.)* **117**, 500-544.
- Holt, J.R. and Corey, D.P. (2000) Two mechanisms for transducer adaptation in vertebrate hair cells. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 11730-11735.
- Holt, J.R., Gillespie, S.K., Provance, D.W., Shah, K., Shokat, K.M., Corey, D.P., Mercer, J.A. and Gillespie, P.G. (2002) A chemical-genetic strategy implicates myosin-1c in adaptation by hair cells. *Cell* **108**, 371-381.
- Holton, T. and Hudspeth, A.J. (1986) The transduction channel of hair cells from the bullfrog characterized by noise analysis. *J. Physiol.* **375**, 195-227.
- Howard, J. and Ashmore, J.F. (1986) Stiffness of sensory hair bundles in the sacculus of the frog. *Hear. Res.* **23**, 93-104.
- Howard, J. and Hudspeth, A.J. (1987) Mechanical relaxation of the hair bundle mediates adaptation in mechano-electrical transduction by the bullfrog's saccular hair cell. *Proc. Natl. Acad. Sci. U.S.A.* **84**, 3064-3068.
- Howard, J. and Hudspeth, A.J. (1988) Compliance of the hair bundle associated with gating of mechano-electrical transduction channels in the bullfrog's saccular hair cell. *Neuron* **1**, 189-199.
- Hudspeth, A.J. (1982) Extracellular current flow and the site of transduction by vertebrate hair cells. *J. Neurosci.* **2**, 1-10.
- Hudspeth, A.J. (1989) How the ear's works work. *Nature* **341**, 397-404.
- Hudspeth, A.J. (1997) Mechanical amplification of stimuli by hair cells. *Curr. Opin. Neurobiol.* **7**, 480-486.
- Iwasa, K.H. and Ehrenstein, G. (2002) Cooperative interaction as the physical basis of the negative stiffness in hair cell stereocilia. *J. Acoust. Soc. Am.* **111**, 2208-2212.

-
- Janssen, J., Coombs, S., Hoekstra, D. and Platt, C. (1987) Anatomy and differential growth of the lateral line system of the mottled sculpin, *Cottus bairdi* (Scopaeniformes: Cottidae). *Brain Behav. Evol.* **30**, 210-229.
- Janssen, J., Jones, W.R., Whang, A. and Oshel, P.E. (1994) Use of the lateral line in particulate feeding in the dark by juvenile alewife (*Alosa pseudoharengus*). *Can. J. Fish. Aquat. Sci.* **52**, 358-363.
- Jaramillo, F. and Hudspeth, A.J. (1991) Localization of the hair cell's transduction channels at the hair bundle's top by iontophoretic application of a channel blocker. *Neuron* **7**, 409-420.
- Jaramillo, F. and Hudspeth, A.J. (1993) Displacement-clamp measurement of the forces exerted by gating springs in the hair bundle. *Proc. Natl. Acad. Sci. U.S.A.* **90**, 1330-1334.
- Jaramillo, F. and Wiesenfeld, K. (1998) Mechanoelectrical transduction assisted by Brownian motion: a role for noise in the auditory system. *Nat. Neurosci.* **1**, 384-388.
- Jielof, R., Spoor, A. and De Vries, H. (1952) The microphonic activity of the lateral line. *J. Physiol* **116**, 137-157.
- Jorgensen, F. and Kroese, A.B.A. (1995) Ca selectivity of the transduction channels in the hair cells of the frog sacculus. *Acta Physiol. Scand.* **155**, 363-376.
- Kachar, B., Parakkal, M., Kurc, M., Zhao, Y. and Gillespie, P.G. (2000) High-resolution structure of hair-cell tip links. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 13336-13341.
- Kalmijn, A.J. (1987) Hydrodynamic and acoustic field detection. In *Sensory biology of aquatic animals*, eds. Atema, J., Fay, R.R., Popper, A.N. and Tavolga, W.N., pp. 83-110. Springer-Verlag, New York.
- Kennedy, H.J., Evans, M.G., Crawford, A.C. and Fettiplace, R. (2003) Fast adaptation of mechanoelectrical transducer channels in mammalian cochlear hair cells. *Nat. Neurosci.* **6**, 832-836.
- Kim, J., Chung, Y.D., Park, D.Y., Choi, S., Shin, D.W., Soh, H., Lee, H.W., Son, W., Yim, J., Park, C.S., Kernan, M.J. and Kim, C. (2003) A TRPV family ion channel required for hearing in *Drosophila*. *Nature* **424**, 81-84.
- Kroese, A.B.A., Van der Zalm, J.M. and Van den Bercken, J. (1978) Frequency response of the lateral-line organ of *Xenopus laevis*. *Pflugers Arch.* **375**, 167-175.
- Kroese, A.B.A. (1979) *On the physiology of the sensory hair cells in the lateral-line organ of Xenopus laevis and the action of ototoxic antibiotics*. Thesis, University of Utrecht.
- Kroese, A.B.A., Van der Zalm, J.M. and Van den Bercken, J. (1980) Extracellular receptor potentials from the lateral-line organ of *Xenopus laevis*. *J. Exp. Biol.* **86**, 63-77.

- Kroese, A.B.A. and Schellart, N.A. (1987) Evidence for velocity- and acceleration sensitive units in the trunk lateral line of the trout. *J. Physiol.* **394**, 13p.
- Kroese, A.B.A. and Schellart, N.A. (1992) Velocity- and acceleration-sensitive units in the trunk lateral line of the trout. *J. Neurophysiol.* **68**, 2212-2221.
- Kros, C.J., Rüsçh, A. and Richardson, G.P. (1992) Mechano-electrical transducer currents in hair cells of the cultured neonatal mouse cochlea. *Proc. R. Soc. Lond B Biol. Sci.* **249**, 185-193.
- Kros, C.J. (1996) Physiology of mammalian hair cells. In *The Cochlea*, eds. Dallos, P., Popper, A.N. and Fay, R.R., pp. 199-208. Springer, New York.
- Kros, C.J., Ruppertsberg, J.P. and Rüsçh, A. (1998) Expression of a potassium current in inner hair cells during development of hearing in mice. *Nature* **394**, 281-284.
- Kros, C.J., Marcotti, W., van Netten, S.M., Self, T.J., Libby, R.T., Brown, S.D., Richardson, G.P. and Steel, K.P. (2002) Reduced climbing and increased slipping adaptation in cochlear hair cells of mice with Myo7a mutations. *Nat. Neurosci.* **5**, 41-47.
- Kuiper, J. W. (1956) *The microphonic effect of the lateral line organ*. Thesis, University of Groningen.
- Lamb, H. (1932) *Hydrodynamics*, 6th edition, Cambridge University Press.
- Landau, L.D. and Lifshitz, E.M. (1980) *Statistical Physics, Part 1*, 3rd edition, Pergamon Press, Oxford.
- Liberman, M.C., Gao, J., He, D.Z., Wu, X., Jia, S. and Zuo, J. (2002) Prestin is required for electromotility of the outer hair cell and for the cochlear amplifier. *Nature* **419**, 300-304.
- Lumpkin, E.A., Marquis, R.E. and Hudspeth, A.J. (1997) The selectivity of the hair cell's mechano-electrical-transduction channel promotes Ca^{2+} flux at low Ca^{2+} concentrations. *Proc. Natl. Acad. Sci. U.S.A.* **94**, 10997-11002.
- Marcotti, W. and Kros, C.J. (1999) Developmental expression of the potassium current $I_{K,n}$ contributes to maturation of mouse outer hair cells. *J. Physiol.* **520.3**, 653-660.
- Marcotti, W., Johnson, S.L., Holley, M.C. and Kros, C.J. (2003) Developmental changes in the expression of potassium currents of embryonic, neonatal and mature mouse inner hair cells. *J. Physiol.* **548**, 383-400.
- Markin, V.S. and Hudspeth, A.J. (1995) Gating-spring models of mechano-electrical transduction by hair cells of the internal ear. *Annu. Rev. Biophys. Biomol. Struct.* **24**, 59-83.

-
- Martin, P., Mehta, A.D. and Hudspeth, A.J. (2000) Negative hair-bundle stiffness betrays a mechanism for mechanical amplification by the hair cell. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 12026-12031.
- Metcalfé, W.K. (1985) Sensory neuron growth cones comigrate with posterior lateral line primordial cells in zebrafish. *J. Comp. Neurol.* **238**, 218-224.
- Metcalfé, W.K., Kimmel, C.B. and Schabtach, E. (1985) Anatomy of the posterior lateral line system in young larvae of the zebrafish. *J. Comp. Neurol.* **233**, 377-389.
- Metcalfé, W.K. (1989) Organization and development of the zebrafish posterior lateral line. In *The mechanosensory lateral line: neurobiology and evolution*, eds. Coombs, S., Görner, P. and Münz, H., pp. 147-159. Springer-Verlag, New York.
- Montgomery, J.C. and MacDonald, J.A. (1987) Sensory tuning of lateral line receptors in Antarctic fish to the movement of planktonic prey. *Science* **235**, 195-196.
- Montgomery, J.C., Baker, C.F. and Carton, A.G. (1997) The lateral line can mediate rheotaxis in fish. *Nature* **389**, 960-963.
- Münz, H. (1989) Functional organization of the lateral line periphery. In *The mechanosensory lateral line: neurobiology and evolution*, eds. Coombs, S., Görner, P. and Münz, H., pp. 285-297. Springer-Verlag, New York.
- Neugebauer, D.C. and Thurm, U. (1985) Interconnections between the stereovilli of the fish inner ear. *Cell Tissue Res.* **240**, 449-453.
- Nicolson, T., Rüsçh, A., Friedrich, R.W., Granato, M., Ruppertsberg, J.P. and Nüsslein-Volhard, C. (1998) Genetic analysis of vertebrate sensory hair cell mechanosensation: the zebrafish circler mutants. *Neuron* **20**, 271-283.
- Papoulis, A. (1991) *Probability, Random Variables and Stochastic Processes*, 3rd edition, McGraw-Hill, New York.
- Partridge, B.L. and Pitcher, T.J. (1980) The sensory basis of fish schools: relative role of lateral line and vision. *J. Comp. Physiol.* **135**, 315-325.
- Pickles, J.O., Comis, S.D. and Osborne, M.P. (1984) Cross-links between stereocilia in the guinea pig organ of Corti and their possible relation to sensory transduction. *Hear. Res.* **15**, 103-112.
- Pitcher, T.J., Partridge, B.L. and Wardle, C.S. (1976) A blind fish can school. *Science* **194**, 963-965.
- Pujol, R., Lavigne-Rebillard, M. and Lenoir, M. (1998) Development of sensory and neural structures in the mammalian cochlea. In *Development of the Auditory System*, eds. Rubel, E.W., Popper, A.N. and Fay, R.R., pp. 146-192. Springer, New York.

- Ricci, A., Wu, Y.C. and Fettiplace, R. (1998) The endogenous calcium buffer and the time course of transducer adaptation in auditory hair cells. *J. Neurosci.* **18**, 8261-8277.
- Ricci, A. (2002) Differences in mechano-transducer channel kinetics underlie tonotopic distribution of fast adaptation in auditory hair cells. *J. Neurophysiol.* **87**, 1738-1748.
- Ricci, A.J., Crawford, A.C. and Fettiplace, R. (2002) Mechanisms of active hair bundle motion in auditory hair cells. *J. Neurosci.* **22**, 44-52.
- Robles, L. and Ruggero, M.A. (2001) Mechanics of the mammalian cochlea. *Physiol. Rev.* **81**, 1305-1352.
- Rouse, G.W. and Pickles, J.O. (1991) Paired development of hair cells in neuromasts of the teleost lateral line. *Proc. R. Soc. Lond B Biol. Sci.* **246**, 123-128.
- Rüsch, A. and Thurm, U. (1990) In *The Mechanics and Biophysics of Hearing*, eds. Dallos, P., Geisler, C.D., Matthews, J.W., Ruggero, M.A. and Steel, C.R., pp. 42-49. Springer, Berlin.
- Rüsch, A., Kros, C.J. and Richardson, G.P. (1994) Block by amiloride and its derivatives of mechano-electrical transduction in outer hair cells of mouse cochlear cultures. *J. Physiol.* **474**, 75-86.
- Russell, I.J., Kossel, M. and Richardson, G.P. (1992) Nonlinear mechanical responses of mouse cochlear hair bundles. *Proc. R. Soc. Lond B Biol. Sci.* **250**, 217-227.
- Sadanaga, M. and Morimitsu, T. (1995) Development of endocochlear potential and its negative component in mouse cochlea. *Hear. Res.* **89**, 155-161.
- Saunders, J.C. and Szymko, Y.M. (1989) The design, calibration and use of a water microjet for stimulating hair cell sensory hair bundles. *J. Acoust. Soc. Am.* **86**, 1797-1804.
- Sidi, S., Friedrich, R.W. and Nicolson, T. (2003) NompC TRP channel required for vertebrate sensory hair cell mechanotransduction. *Science* **301**, 96-99.
- Steel, K.P. and Barkway, C. (1989) Another role for melanocytes: their importance for normal stria vascularis development in the mammalian inner ear. *Development* **107**, 453-463.
- Steel, K.P. and Kros, C.J. (2001) A genetic approach to understanding auditory function. *Nat. Genet.* **27**, 143-149.
- Stokes, G.G. (1851) On the Effect of the Internal Friction of Fluids on the Motion of Pendulums. *Cambridge Philos. Trans.* **9**, 8-106.
- Strelhoff, D. and Flock, Å. (1984) Stiffness of sensory-cell hair bundles in the isolated guinea pig cochlea. *Hear. Res.* **15**, 19-28.

Tilney, L.G., Derosier, D.J. and Mulroy, M.J. (1980) The organization of actin filaments in the stereocilia of cochlear hair cells. *J. Cell Biol.* **86**, 244-259.

van Netten, S.M. (1987) *Laser interferometric study of the mechanosensitivity of the fish lateral line*. Thesis, University of Groningen.

van Netten, S.M. and Kroese, A.B.A. (1987) Laser interferometric measurements on the dynamic behaviour of the cupula in the fish lateral line. *Hear. Res.* **29**, 55-61.

van Netten, S.M. (1988) Laser interferometer microscope for the measurement of nanometer vibrational displacements of a light-scattering microscopic object. *J. Acoust. Soc. Am.* **83**, 1667-1674.

van Netten, S. M. (1991) Hydrodynamics of the excitation of the cupula in the fish canal lateral line. *J. Acoust. Soc. Am.* **89**, 310-319.

van Netten, S.M. and Khanna, S.M. (1994) Stiffness changes of the cupula associated with the mechanics of hair cells in the fish lateral line. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 1549-1553.

van Netten, S.M. (1997) Hair cell mechano-transduction: its influence on the gross mechanical characteristics of a hair cell sense organ. *Biophys. Chem.* **68**, 43-52.

van Netten, S.M. and Kros, C.J. (2000) Gating energies and forces of the mammalian hair cell transducer channel and related hair bundle mechanics. *Proc. R. Soc. Lond. B Biol. Sci.* **267**, 1915-1923.

van Netten, S.M., Dinklo, T., Marcotti, W. and Kros, C.J. (2003) Channel gating forces govern accuracy of mechano-electrical transduction in hair cells. *Proc. Natl. Acad. Sci. U.S.A.* **100**, 15510-15515.

van Netten, S.M. (Submitted for publication) Hydrodynamic detection by cupulae in the lateral line canal: Functional relations between physics and physiology.

van Trees, H.L. (1968) *Detection, Estimation and Modulation Theory, Part 1* Wiley, New York.

von Campenhausen, C., Riess, I. and Weissert, R. (1981) Detection of stationary objects by the blind cave fish *Anoptichthys jordani* (Characidae). *J. Comp. Physiol.* **143**, 369-374.

Walker, Th.J. (1967) History, histological methods and details of the structure of the lateral line of the walleye surfperch. In *Lateral line detectors*, ed. Cahn, Ph.H., pp. 13-25. Indiana university press, Bloomington.

Weissert, R. and von Campenhausen, C. (1981) Discrimination between stationary objects by the blind cave fish *Anoptichtys jordani* (Characidae). *J. Comp. Physiol.* **143**, 375-382.

- Whitfield, T.T. (2003) from the Zebrafish Information Network (ZFIN), University of Oregon, URL: http://zfin.org/zf_info/anatomy/dict/lat_line/lat_line.html
- Wiersinga-Post, J.E.C. (1997) *Mechanophysiology of the cupulae and hair cells in the lateral line of fish and Pitch perception of complex sounds in humans*. Thesis, University of Groningen, The Netherlands.
- Williams, J.A. and Holder, N. (2000) Cell turnover in neuromasts of zebrafish larvae. *Hear. Res.* **143**, 171-181.
- Wu, Y.C., Ricci, A.J. and Fettiplace, R. (1999) Two components of transducer adaptation in auditory hair cells. *J. Neurophysiol.* **82**, 2171-2181.
- Yamasaki, M., Komune, S., Shimozono, M., Matsuda, K. and Haruta, A. (2000) Development of monovalent ions in the endolymph in mouse cochlea. *ORL J. Otorhinolaryngol. Relat. Spec.* **62**, 241-246.
- Yamoah, E.N. and Gillespie, P.G. (1996) Phosphate analogs block adaptation in hair cells by inhibiting adaptation-motor force production. *Neuron* **17**, 523-533.

