

University of Groningen

Structure-function relationships in type I signal peptidases of bacilli

Roosmalen, Maarten Leonardus van

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2001

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Roosmalen, M. L. V. (2001). *Structure-function relationships in type I signal peptidases of bacilli*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

REFERENCES

1. **Tjalsma, H., A. Bolhuis, J.D.H. Jongbloed, S. Bron, and J.M. van Dijl.** 2000. Signal peptide-dependent protein transport in *Bacillus subtilis*: a genome-based survey of the secretome. *Microbiol.Mol.Biol.Rev.* **64**:515-547.
2. **Dalbey, R.E. and A. Kuhn.** 2000. Evolutionarily related insertion pathways of bacterial, mitochondrial, and thylakoid membrane proteins. *Ann.Rev.Cell Dev.Biol.* **16**:51-87.
3. **Dalbey, R.E., M.Y. Chen, F.L. Jiang, and J.C. Samuelson.** 2000. Understanding the insertion of transporters and other membrane proteins. *Curr.Opin.Cell Biol.* **12**:435-442.
4. **Hartl, F.U., S. Lecker, E. Schiebel, J.P. Hendrick, and W. Wickner.** 1990. The binding cascade of SecB to SecA to SecY/E mediates preprotein targeting to the *Escherichia coli* plasma membrane. *Cell* **63**:269-280.
5. **Henry, R., M. Carrigan, M. McCaffery, X. Ma, and K. Cline.** 1997. Targeting determinants and proposed evolutionary basis for the Sec and the Delta pH protein transport systems in chloroplast thylakoid membranes. *J.Cell.Biol.* **136**:823-832.
6. **Andersson, H., E. Bakker, and G. von Heijne.** 1992. Different positively charged amino acids have similar effects on the topology of a polytopic transmembrane protein in *Escherichia coli*. *J.Biol.Chem.* **267**:1491-1495.
7. **Dalbey, R.E.** 1990. Positively charged residues are important determinants of membrane protein topology. *TIBS* **15**:253-257.
8. **Lamiet, A.A. and A. Pluckthun.** 1989. The precursor of beta-lactamase: purification, properties and folding kinetics. *EMBO J.* **8**:1469-1477.
9. **Liu, G., T.B. Topping, and L.L. Randall.** 1989. Physiological role during export for the retardation of folding by the leader peptide of maltose-binding protein. *Proc.Natl.Acad.Sci.USA* **86**:9213-9217.
10. **Park, S., G. Liu, T.B. Topping, W.H. Cover, and L.L. Randall.** 1988. Modulation of folding pathways of exported proteins by the leader sequence. *Science* **239**:1033-1035.
11. **von Heijne, G. and L. Abrahmsen.** 1989. Species-specific variation in signal peptide design. Implications for protein secretion in foreign hosts. *FEBS Lett.* **244** :439-446.
12. **Nielsen, H., J. Engelbrecht, S. Brunak, and G. von Heijne.** 1997. Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Protein Eng.* **10**:1-6.
13. **Nielsen, H., J. Engelbrecht, S. Brunak, and G. von Heijne.** 1997. A neural network method for identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Int.J.Neural Syst.* **8**:581-599.
14. **Briggs, M.S., D.G. Cornell, R.A. Dluhy, and L.M. Gierasch.** 1986. Conformations of signal peptides induced by lipids suggest initial steps in protein export. *Science* **233**:206-208.
15. **de Vrije, G.J., A.M. Batenburg, J.A. Killian, and B. de Kruijff.** 1990. Lipid involvement in protein translocation in *Escherichia coli*. *Mol.Microbiol.* **4**:143-150.
16. **von Heijne, G.** 1986. Net N-C charge imbalance may be important for signal sequence function in bacteria. *J.Mol.Biol.* **192**:287-290.
17. **von Heijne, G.** 1986. Towards a comparative anatomy of N-terminal topogenic protein sequences. *J.Mol.Biol.* **189**:239-242.
18. **von Heijne, G. and C. Manoil.** 1990. Membrane proteins: From sequence to structure. *Protein Eng.* **4**:109-112.
19. **von Heijne, G.** 1992. Membrane protein structure prediction. Hydrophobicity analysis and the positive-inside rule. *J.Mol.Biol.* **225**:487-494.
20. **Sipos, L. and G. von Heijne.** 1993. Predicting the topology of eukaryotic membrane proteins. *Eur.J.Biochem.* **213**:1333-1340.
21. **von Heijne, G.** 1994. Control of topology and mode of assembly of a polytopic membrane protein by positively charged residues. *Nature* **341**:456-458.
22. **Wallin, E. and G. von Heijne.** 1998. Genome-wide analysis of integral membrane proteins from eubacterial, archaean, and eukaryotic organisms. *Protein Sci.* **7**:1029-1038.
23. **van de Vossenberg, J.L.C.M., S.V. Albers, C. van der Does, A.J.M. Driessen, and W. van Klompenburg.** 1998. The positive inside rule is not determined by the polarity of the Delta psi. *Mol.Microbiol.* **29**:1125-1126.
24. **Gallusser, A. and A. Kuhn.** 1990. Initial steps in protein membrane insertion. Bacteriophage M13 procoat protein binds to the membrane surface by electrostatic interaction. *EMBO J.* **9**:2723-2729.
25. **Andersson, H. and G. von Heijne.** 1994. Membrane protein topology: Effects of $\Delta\mu_{H^+}$ on the translocation of charged residues explain the 'positive inside' rule. *EMBO J.* **13**:2267-2272.
26. **Schuenemann, T.A., V. Delgado-Nixon, and R.E. Dalbey.** 1999. Direct evidence that the proton motive force inhibits membrane translocation of positively charged residues within membrane proteins. *J.Biol.Chem.* **274**:6855-6864.
27. **Flewelling, R.F. and W.L. Hubbell.** 1986. The membrane dipole potential in a total membrane potential model. Applications to hydrophobic ion interactions with membranes. *Biophys.J.* **49**:541-552.
28. **van de Vossenberg, J.L., S.V. Albers, C. van der Does, A.J.M. Driessen, and W. van Klompenburg.** 1998. The positive inside rule is not determined by the polarity of the delta psi (transmembrane electrical potential). *Mol.Microbiol.* **29**:1125-1127.
29. **Paetzel, M., R.E. Dalbey, and N.-C.J. Strynadka.** 1998. Crystal structure of a bacterial signal peptidase in complex with a beta-lactam inhibitor. *Nature* **396**:186-190.

References

30. **Barkocy-Gallagher, G.A. and P.J.J. Bassford.** 1992. Synthesis of precursor maltose-binding protein with proline in the +1 position of the cleavage site interferes with the activity of *Escherichia coli* signal peptidase I *in vivo*. *J.Biol.Chem.* **267**:1231-1238.
31. **Nilsson, I. and G. von Heijne.** 1992. A signal peptide with a proline next to the cleavage site inhibits leader peptidase when present in a sec-independent protein. *FEBS Lett.* **299**:243-246.
32. **Watts, C., W. Wickner, and R. Zimmermann.** 1983. M13 procoat and a pre-immunoglobulin share processing specificity but use different membrane receptor mechanisms. *Proc.Natl.Acad.Sci.USA* **80**:2809-2813.
33. **Halpin, C., P.D. Elderfield, H.E. James, R. Zimmermann, B. Dunbar, and C. Robinson.** 1989. The reaction specificities of the thylakoidal processing peptidase and *Escherichia coli* leader peptidase are identical. *EMBO J.* **8**:3917-3921.
34. **Chou, K.-C.** 2001. Using subsite coupling to predict signal peptides. *Protein Eng.* **14**:75-79.
35. **Pohlschroder, M., W.A. Prinz, E. Hartmann, and J. Beckwith.** 1997. Protein translocation in the three domains of life: Variations on a theme. *Cell* **91**:563-566.
36. **Dalbey, R.E. and C. Robinson.** 1999. Protein translocation into and across the bacterial plasma membrane and the plant thylakoid membrane. *TIBS* **24**:17-22.
37. **Berks, B.C., F. Sargent, and T. Palmer.** 2000. The Tat protein export pathway. *Mol.Microbiol.* **35**:260-274.
38. **Robinson, C. and A. Bolhuis.** 2001. Protein targeting by the twin-arginine translocation pathway. *Nature Rev.Mol. Cell Biol.* **2**:350-356.
39. **Fekkes, P. and A.J.M. Driessen.** 1999. Protein targeting to the bacterial cytoplasmic membrane. *Microbiol.Mol.Biol.Rev.* **63**:161-173.
40. **Muller, M., H.G. Koch, K. Beck, and U. Schaefer.** 2000. Protein traffic in bacteria: Multiple routes from the ribosome to and across the membrane. *Prog.Nucleic Acid Res.Mol.Biol.* **66**:107-157.
41. **Driessen, A.J.M., E.H. Manting, and C. van der Does.** 2001. The structural basis of protein targeting and translocation in bacteria. *Nature Struct.Biol.* **8**:492-498.
42. **Lill, R., W. Dowhan, and W. Wickner.** 1990. The ATPase activity of SecA is regulated by acidic phospholipids, SecY, and the leader and mature domains of precursor proteins. *Cell* **60**:271-280.
43. **Breukink, E., R.A. Demel, G. de Korte Kool, and B. de Kruijff.** 1992. SecA insertion into phospholipids is stimulated by negatively charged lipids and inhibited by ATP: a monolayer study. *Biochemistry* **31**:1119-1124.
44. **Kumamoto, C.A. and J. Beckwith.** 1983. Mutations in a new gene, secB, cause defective protein localization in *Escherichia coli*. *J.Bacteriol.* **154**:253-260.
45. **Hendrick, J.P. and W. Wickner.** 1991. SecA protein needs both acidic phospholipids and SecY/E protein for functional high-affinity binding to the *Escherichia coli* plasma membrane. *J.Biol.Chem.* **266**:24596-24600.
46. **Tokuda, H., K. Shiozuka, and S. Mizushima.** 1990. Reconstitution of translocation activity for secretory proteins from solubilized components of *Escherichia coli*. *Eur.J.Biochem.* **192**:583-589.
47. **Ito, K., M. Wittekind, M. Nomura, K. Shiba, T. Yura, A. Miura, and H. Nashimoto.** 1983. A temperature-sensitive mutant of *E. coli* exhibiting slow processing of exported proteins. *Cell* **32**:789-797.
48. **Riggs, P.D., A.I. Derman, and J. Beckwith.** 1988. A mutation affecting the regulation of a secA-lacZ fusion defines a new sec gene. *Genetics* **118**:571-579.
49. **Douville, K., M. Leonard, L. Brundage, K. Nishiyama, H. Tokuda, S. Mizushima, and W. Wickner.** 1994. Band 1 subunit of *Escherichia coli* preprotein translocase and integral membrane export factor P12 are the same protein. *J.Biol.Chem.* **269**:18705-18707.
50. **Gardel, C., S. Benson, J. Hunt, S. Michaelis, and J. Beckwith.** 1987. secD, a new gene involved in protein export in *Escherichia coli*. *J.Bacteriol.* **169**:1286-1290.
51. **Gardel, C., K. Johnson, A. Jacq, and J. Beckwith.** 1990. The secD locus of *E.coli* codes for two membrane proteins required for protein export. *EMBO J.* **9**:3209-3216.
52. **Duong, F. and W. Wickner.** 1997. Distinct catalytic roles of the SecYE, SecG and SecDFyajC subunits of preprotein translocase holoenzyme. *EMBO J.* **16**:2756-2768.
53. **Scotti, P.A., M.L. Urbanus, J. Brunner, J.W.L. de Gier, G. von Heijne, C. van der Does, A.J.M. Driessen, B. Oudega, and J. Luirink.** 2000. YidC, the *Escherichia coli* homologue of mitochondrial Oxa1p, is a component of the Sec translocase. *EMBO J.* **19**:542-549.
54. **Bolhuis, A., C.P. Broekhuizen, A. Sorokin, M.L. van Roosmalen, G. Venema, S. Bron, W.J. Quax, and J.M. van Dijl.** 1998. SecDF of *Bacillus subtilis*, a molecular siamese twin required for the efficient secretion of proteins. *J.Biol.Chem.* **273**:21217-21224.
55. **Economou, A. and W. Wickner.** 1994. SecA promotes preprotein translocation by undergoing ATP-driven cycles of membrane insertion and deinsertion. *Cell* **78**:835-843.
56. **Economou, A.** 1998. Bacterial preprotein translocase: mechanism and conformational dynamics of a processive enzyme. *Mol.Microbiol.* **27**:511-518.
57. **Bernstein, H.D., M.A. Poritz, K. Strub, P.J. Hoben, S. Brenner, and P. Walter.** 1989. Model for signal sequence recognition from amino-acid sequence of 54K subunit of signal recognition particle. *Nature* **340**:482-486.
58. **de Gier, J.W.L., P. Mansournia, Q.A. Valent, G.J. Phillips, J. Luirink, and G. von Heijne.** 1996. Assembly of a cytoplasmic membrane protein in *Escherichia coli* is dependent on the signal recognition particle. *FEBS Lett.* **399**:307-309.
59. **Valent, Q.A., P.A. Scotti, S. High, J.W.L. de Gier, G. von Heijne, G. Lentzen, W. Wintermeyer, B. Oudega, and J. Luirink.** 1998. The *Escherichia coli* SRP and SecB targeting pathways converge at the translocon. *EMBO J.* **17**:2504-2512.

60. Berks, B.C. 1996. A common export pathway for proteins binding complex redox cofactors? *Mol.Microbiol.* **22**:393-404.
61. Brink, S., E.G. Bogsch, W.R. Edwards, P.J. Hynds, and C. Robinson. 1998. Targeting of thylakoid proteins by the DELTA_{pH}-driven twin-arginine translocation pathway requires a specific signal in the hydrophobic domain in conjunction with the twin-arginine motif. *FEBS Lett.* **434**:425-430.
62. Cristobal, S., J.W.L. de Gier, H. Nielsen, and G. von Heijne. 1999. Competition between Sec- and TAT-dependent protein translocation in *Escherichia coli*. *EMBO J.* **18**:2982-2990.
63. Mould, R.M. and C. Robinson. 1991. A proton gradient is required for the transport of two luminal oxygen-evolving proteins across the thylakoid membrane. *J.Biol.Chem.* **266**:12189-12193.
64. Robinson, C., D. Cai, A. Hulford, I.W. Brock, D. Michl, L. Hazell, I. Schmidt, R.G. Herrmann, and R.B. Klossgen. 1994. The presequence of a chimeric construct dictates which of two mechanisms are utilized for translocation across the thylakoid membrane: evidence for the existence of two distinct translocation systems. *EMBO J.* **13**:279-285.
65. Chaddock, A.M., A. Mant, I. Karnachov, S. Brink, R.G. Herrmann, R.B. Klossgen, and C. Robinson. 1995. A new type of signal peptide: Central role of twin-arginine motif in transfer signals for the DELTA-pH-dependent thylakoidal protein translocase. *EMBO J.* **14**:2715-2722.
66. Sargent, F., E.G. Bogsch, N.R. Stanley, M. Wexler, C. Robinson, B.C. Berks, and T. Palmer. 1998. Overlapping functions of components of a bacterial Sec-independent protein export pathway. *EMBO J.* **17**:3640-3650.
67. Bogsch, E.G., F. Sargent, N.R. Stanley, B.C. Berks, C. Robinson, and T. Palmer. 1998. An essential component of a novel bacterial protein export system with homologues in plastids and mitochondria. *J.Biol.Chem.* **273**:18003-18006.
68. Jongbloed, J.D.H., U. Martin, H. Antelmann, M. Hecker, H. Tjalsma, G. Venema, S. Bron, J.M. van Dijl, and J. Muller. 2000. TatC is a specificity determinant for protein secretion via the twin-arginine translocation pathway. *J.Biol.Chem.* **275**:41350-41357.
69. Thomas, J.D., R.A. Daniel, J. Errington, and C. Robinson. 2001. Export of active green fluorescent protein to the periplasm by the twin-arginine translocase (Tat) pathway in *Escherichia coli*. *Mol.Microbiol.* **39**:47-53.
70. Santini, C.L., A. Bernadac, M. Zhang, A. Chanal, B. Ize, C. Blanco, and L.F. Wu. 2001. Translocation of jellyfish green fluorescent protein via the Tat system of *Escherichia coli* and change of its periplasmic localization in response to osmotic up-shock. *J.Biol.Chem.* **276**:8159-8164.
71. Feilmeier, B.J., G. Iseminger, D. Schroeder, H. Webber, and G.J. Phillips. 2000. Green fluorescent protein functions as a reporter for protein localization in *Escherichia coli*. *J.Bacteriol.* **182**:4068-4076.
72. Wolfe, P.B., M. Rice, and W. Wickner. 1985. Effects of two sec genes on protein assembly into the plasma membrane of *Escherichia coli*. *J.Biol.Chem.* **260**:1836-1841.
73. Geller, B.L. and W. Wickner. 1985. M13 procoat inserts into liposomes in the absence of other membrane proteins. *J.Biol.Chem.* **260**:13281-13285.
74. Soekarjo, M., M. Eisenhauer, A. Kuhn, and H. Vogel. 1996. Thermodynamics of the membrane insertion process of the M13 procoat protein, a lipid bilayer traversing protein containing a leader sequence. *Biochemistry* **35**:1232-1241.
75. Kuhn, A., G. Kreil, and W. Wickner. 1986. Both hydrophobic domains of M13 procoat are required to initiate membrane insertion. *EMBO J.* **5**:3681-3685.
76. Cao, G., A. Kuhn, and R.E. Dalbey. 1995. The translocation of negatively charged residues across the membrane is driven by the electrochemical potential: Evidence for an electrophoresis-like membrane transfer mechanism. *EMBO J.* **14**:866-875.
77. Kuhn, A., H.Y. Zhu, and R.E. Dalbey. 1990. Efficient translocation of positively charged residues of M13 procoat protein across the membrane excludes electrophoresis as the primary force for membrane insertion. *EMBO J.* **9**:2385-2389.
78. Andersson, H. and G. von Heijne. 1993. Sec dependent and sec independent assembly of *E.coli* inner membrane proteins: The topological rules depend on chain length. *EMBO J.* **12**:683-691.
79. Kuhn, A. 1988. Alterations in the extracellular domain of M13 procoat protein make its membrane insertion dependent on secA and secY. *Eur.J.Biochem.* **177**:267-271.
80. Samuelson, J.C., M.Y. Chen, F.L. Jiang, I. Moller, M. Wiedmann, A. Kuhn, G.J. Phillips, and R.E. Dalbey. 2000. YidC mediates membrane protein insertion in bacteria. *Nature* **406**:637-641.
81. Hell, K., J. Herrmann, E. Pratje, W. Neupert, and R.A. Stuart. 1997. Oxa1p mediates the export of the N- and C-termini of pCoxII from the mitochondrial matrix to the intermembrane space. *FEBS Lett.* **418**:367-370.
82. Hell, K., J.M. Herrmann, E. Pratje, W. Neupert, and R.A. Stuart. 1998. Oxa1p, an essential component of the N-tail protein export machinery in mitochondria. *Proc.Natl.Acad.Sci.USA* **95**:2250-2255.
83. Moore, M., M.S. Harrison, E.C. Peterson, and R. Henry. 2000. Chloroplast Oxa1p homolog albino3 is required for post-translational integration of the light harvesting chlorophyll-binding protein into thylakoid membranes. *J.Biol.Chem.* **275**:1529-1532.
84. Zwizinski, C., T. Date, and W. Wickner. 1981. Leader peptidase is found in both the inner and outer membranes of *Escherichia coli*. *J.Biol.Chem.* **256**:3593-3597.
85. Wolfe, P.B., W. Wickner, and J.M. Goodman. 1983. Sequence of the leader peptidase gene of *Escherichia coli* and the orientation of leader peptidase in the bacterial envelope. *J.Biol.Chem.* **258**:12073-12080.
86. Dalbey, R.E. and W. Wickner. 1985. Leader peptidase catalyzes the release of exported proteins from the outer surface of the *Escherichia coli* plasma membrane. *J.Biol.Chem.* **260**:15925-15931.
87. Bairl, A. and P. Muller. 1998. A second gene for type I signal peptidase in *Bradyrhizobium japonicum*, sipF is located near genes involved in RNA processing and cell division. *Mol.Gen.Genet.* **260**:346-356.

References

88. Muller, P., K. Ahrens, T. Keller, and A. Klaucke. 1995. A TnpA insertion within the *Bradyrhizobium japonicum sipS* gene, homologous to prokaryotic signal peptidases, results in extensive changes in the expression of PBM-specific nodulins of infected soybean (*Glycine max*) cells. *Mol. Microbiol.* **18**:831-840.
89. Dalbey, R.E., M.O. Lively, S. Bron, and J.M. van Dijl. 1997. The chemistry and enzymology of the type I signal peptidases. *Protein Sci.* **6**:1129-1138.
90. Moore, K.E. and S. Miura. 1987. A small hydrophobic domain anchors leader peptidase to the cytoplasmic membrane of *Escherichia coli*. *J. Biol. Chem.* **262**:8806-8813.
91. San-Millan, J.L., D. Boyd, R. Dalbey, W. Wickner, and J. Beckwith. 1989. Use of phoA fusions to study the topology of the *Escherichia coli* inner membrane protein leader peptidase. *J. Bacteriol.* **171**:5536-5541.
92. van Klompenburg, W., P. Whitley, R. Diemel, G. von Heijne, and B. de Kruijff. 1995. A quantitative assay to determine the amount of signal peptidase I in *E. coli* and the orientation of membrane vesicles. *Mol. Membr. Biol.* **12**:349-353.
93. Zwizinski, C. and W. Wickner. 1980. Purification and characterization of leader (signal) peptidase from *Escherichia coli*. *J. Biol. Chem.* **255**:7973-7977.
94. Wolfe, P.B., P. Silver, and W. Wickner. 1982. The isolation of homogeneous leader peptidase from a strain of *Escherichia coli* which overproduces the enzyme. *J. Biol. Chem.* **257**:7898-7902.
95. Wolfe, P.B., C. Zwizinski, and W. Wickner. 1983. Purification and characterization of leader peptidase from *Escherichia coli*. *Methods Enzymol.* **97**:40-57.
96. Tschantz, W.R. and R.E. Dalbey. 1994. Bacterial leader peptidase I. *Methods Enzymol.* **244**:285-301.
97. Talarico, T.L., I.K. Dev, P.J.Jr. Bassford, and P.H. Ray. 1991. Inter-molecular degradation of signal peptidase I *in vitro*. *Biochem. Biophys. Res. Com.* **181**:650-656.
98. Kuo, D.W., H.K. Chan, C.J. Wilson, P.R. Griffin, H. Williams, and W.B. Knight. 1993. *Escherichia coli* leader peptidase: Production of an active form lacking a requirement for detergent and development of peptide substrates. *Arch. Biochem. Biophys.* **303**:274-280.
99. Paetzel, M., N.-C.J. Strynadka, W.R. Tschantz, R. Casareno, P.R. Bullinger, and R.E. Dalbey. 1997. Use of site-directed chemical modification to study an essential lysine in *Escherichia coli* leader peptidase. *J. Biol. Chem.* **272**:9994-10003.
100. Bilgin, N., J.I. Lee, H.Y. Zhu, R.E. Dalbey, and G. von Heijne. 1990. Mapping of catalytically important domains in *Escherichia coli* leader peptidase. *EMBO J.* **9**:2717-2722.
101. Tschantz, W.R., M. Paetzel, G. Cao, D. Suci, M. Inouye, and R.E. Dalbey. 1995. Characterization of a Soluble, Catalytically Active Form of *Escherichia coli* Leader Peptidase: Requirement of Detergent or Phospholipid for Optimal Activity. *Biochemistry* **34**:3935-3941.
102. van Klompenburg, W., M. Paetzel, J.M. de Jong, R.E. Dalbey, R.A. Demel, G. von Heijne, and B. de Kruijff. 1998. Phosphatidylethanolamine mediates insertion of the catalytic domain of leader peptidase in membranes. *FEBS Lett.* **431**:79
103. van Dijl, J.M., A. de Jong, J. Vehmaanperä, G. Venema, and S. Bron. 1992. Signal peptidase I of *Bacillus subtilis*: patterns of conserved amino acids in prokaryotic and eukaryotic type I signal peptidases. *EMBO J.* **11**:2819-2828.
104. Bolhuis, A., A. Sorokin, V. Azevedo, S.D. Ehrlich, P.G. Braun, A. de Jong, G. Venema, S. Bron, and J.M. van Dijl. 1996. *Bacillus subtilis* can modulate its capacity and specificity for protein secretion through temporally controlled expression of the *sipS* gene for signal peptidase I. *Mol. Microbiol.* **22**:605-618.
105. Tjalsma, H., M.A. Noback, S. Bron, G. Venema, K. Yamane, and J.M. van Dijl. 1997. *Bacillus subtilis* contains four closely related type I signal peptidases with overlapping substrate specificities. Constitutive and temporally controlled expression of different *sip* genes. *J. Biol. Chem.* **272**:25983-25992.
106. Tjalsma, H., A. Bolhuis, M.L. van Roosmalen, T. Wiegert, W. Schumann, C.P. Broekhuizen, W.J. Quax, G. Venema, S. Bron, and J.M. van Dijl. 1998. Functional analysis of the secretory precursor processing machinery of *Bacillus subtilis*: Identification of a eubacterial homolog of archaeal and eukaryotic signal peptidases. *Genes Dev.* **12**:2318-2331.
107. Meijer, W.J.J., A. de Jong, G.B.A. Wisman, H. Tjalsma, G. Venema, S. Bron, and J.M. van Dijl. 1995. The endogenous *Bacillus subtilis* (natto) plasmids pTA1015 and pTA1040 contain signal peptidase-encoding genes: Identification of a new structural module on cryptic plasmids. *Mol. Microbiol.* **17**:621-631.
108. Tjalsma, H., J. van den Dolder, W.J.J. Meijer, G. Venema, S. Bron, and J.M. van Dijl. 1999. The plasmid-encoded signal peptidase SipP can functionally replace the major signal peptidases SipS and SipT of *Bacillus subtilis*. *J. Bacteriol.* **181**:2448-2454.
109. Bron, S., A. Bolhuis, H. Tjalsma, S. Holsappel, G. Venema, and J.M. van Dijl. 1998. Protein secretion and possible roles for multiple signal peptidases for precursor processing in Bacilli. *J. Biotech.* **64**:1-13.
110. Tjalsma, H., A.G. Stöver, A. Driks, G. Venema, S. Bron, and J.M. van Dijl. 2000. Conserved serine and histidine residues are critical for activity of the ER-type signal peptidase SipW of *Bacillus subtilis*. *J. Biol. Chem.* **275**:25102-25108.
111. Black, M.T. 1993. Evidence that the catalytic activity of prokaryote leader peptidase depends upon the operation of a serine-lysine catalytic dyad. *J. Bacteriol.* **175**:4957-4961.
112. Tschantz, W.R., M. Sung, V.M. Delgado-Partin, and R.E. Dalbey. 1993. A serine and a lysine residue implicated in the catalytic mechanism of the *Escherichia coli* leader peptidase. *J. Biol. Chem.* **268**:27349-27354.
113. van Dijl, J.M., A. de Jong, G. Venema, and S. Bron. 1995. Identification of the potential active site of the signal peptidase SipS of *Bacillus subtilis*: Structural and functional similarities with LexA-like proteases. *J. Biol. Chem.* **270**:3611-3618.

114. **Dalbey, R.E. and G. von Heijne.** 1992. Signal peptidases in prokaryotes and eukaryotes - a new protease family. *TIBS* **17**:474-478.
115. **van Valkenburgh, C., X.M. Chen, C. Mullins, H. Fang, and N. Green.** 1999. The catalytic mechanism of endoplasmic reticulum signal peptidase appears to be distinct from most eubacterial signal peptidases. *J.Biol.Chem.* **274**:11519-11525.
116. **Stöver, A.G. and A. Driks.** 1999. Secretion, localization, and antibacterial activity of TasA, a *Bacillus subtilis* spore-associated protein. *J.Bacteriol.* **181**:1664-1672.
117. **Stöver, A.G. and A. Driks.** 1999. Control of synthesis and secretion of the *Bacillus subtilis* protein YqxM. *J.Bacteriol.* **181**:7065-7069.
118. **van Roosmalen, M.L., J.D.H. Jongbloed, A. de Jong, J. van Eerden, G. Venema, S. Bron, and J.M. van Dijl.** 2001. Detergent-independent *in vitro* activity of a truncated *Bacillus* signal peptidase. *Microbiology* **147**:909-917.
119. **Tjalsma, H., G. Zanen, G. Venema, S. Bron, and J.M. van Dijl.** 1999. The potential active site of the lipoprotein-specific (type II) signal peptidase of *Bacillus subtilis*. *J.Biol.Chem.* **274**:28191-28197.
120. **Tokunaga, M., J.M. Loranger, S.Y. Chang, M. Regue, S. Chang, and H.C. Wu.** 1985. Identification of prolipoprotein signal peptidase and genomic organization of the *lsp* gene in *Escherichia coli*. *J.Biol.Chem.* **260**:5610-5615.
121. **van Roosmalen, M.L., J.D.H. Jongbloed, A. Kuipers, G. Venema, S. Bron, and J.M. van Dijl.** 2000. A truncated soluble *Bacillus* signal peptidase produced in *Escherichia coli* is subject to self-cleavage at its active site. *J.Bacteriol.* **182**:5765-5770.
122. **Vehmaanperä, J., A. Görner, G. Venema, S. Bron, and J.M. van Dijl.** 1993. *In vitro* assay for the *Bacillus subtilis* signal peptidase SipS : Systems for efficient *in vitro* transcription-translation and processing of precursors of secreted proteins. *FEMS Microbiol.Lett.* **114**:207-214.
123. **Jackson, R.C. and G. Blobel.** 1980. Post-translational processing of full-length presecretory proteins with canine pancreatic signal peptidase. *Ann.NY Acad.Sci.* **343**:391-404.
124. **Sung, M. and R.E. Dalbey.** 1992. Identification of potential active-site residues in the *Escherichia coli* leader peptidase. *J.Biol.Chem.* **267**:13154-13159.
125. **Black, M.T., J.G. Munn, and A.E. Allsop.** 1992. On the catalytic mechanism of prokaryotic leader peptidase I. *Biochem.J.* **282**:539-543.
126. **Stryer, L.** 1988. Introduction to enzymes, L. Stryer (ed.), Biochemistry, 3 ed. W.H. Freeman and Company, New York.
127. **Dao-pin, S., D.E. Anderson, W.A. Baase, F.W. Dahlquist, and B.W. Matthews.** 1991. Structural and thermodynamic consequences of burying a charged residue within the hydrophobic core of T4 lysozyme. *Biochemistry* **30**:11521-11529.
128. **Wickner, W., K. Moore, N. Dibb, D. Geissert, and M. Rice.** 1987. Inhibition of purified *Escherichia coli* leader peptidase by the leader (signal) peptide of bacteriophage M13 procoat. *J.Bacteriol.* **169**:3821-3822.
129. **Kuo, D.W., W. Jeffrey, P.R. Griffin, S.K. Shah, and W.B. Knight.** 1994. Determination of the kinetic parameters of *Escherichia coli* leader peptidase activity using a continuous assay: the pH dependence and time-dependent inhibition by β -lactams are consistent with a novel serine protease mechanism. *Biochemistry* **33**:8347-8354.
130. **Allsop, A.E., G. Brooks, G. Bruton, S. Coulton, P.D. Edwards, I.K. Hatton, A.C. Kaura, S.D. McLean, N.D. Pearson, T.C. Smale, and R. Southgate.** 1995. Penem inhibitors of bacterial signal peptidase. *Bioorg.Med.Chem.Lett.* **5**:443-448.
131. **Allsop, A., G. Brooks, P.D. Edwards, A.C. Kaura, and R. Southgate.** 1996. Inhibitors of bacterial signal peptidase: a series of 6-(substituted oxyethyl)penems. *J.Antibiot.Tokyo* **49**:921-928.
132. **Paetzel, M., R.E. Dalbey, and N.C.J. Strynadka.** 2000. The structure and mechanism of bacterial type I signal peptidases - A novel antibiotic target. *Pharmacol.Ther.* **87**:27-49.
133. **Black, M.T. and G. Bruton.** 1998. Inhibitors of bacterial signal peptidases. *Curr.Pharm.Des.* **4**:133-154.
134. **Chatterjee, S., D. Suci, R.E. Dalbey, P.C. Kahn, and M. Inouye.** 1995. Determination of K_m and K_{cat} for signal peptidase I using a full length secretory precursor, pro-OmpA-nuclease A. *J.Mol.Biol.* **245**:311-314.
135. **Suci, D., S. Chatterjee, and M. Inouye.** 1997. Catalytic efficiency of signal peptidase I of *Escherichia coli* is comparable to that of members of the serine protease family. *Protein Eng.* **10**:1057-1060.
136. **Smith, H., S. Bron, J. van Ee, and G. Venema.** 1987. Construction and use of signal sequence selection vectors in *Escherichia coli* and *Bacillus subtilis*. *J.Bacteriol.* **169**:3321-3328.
137. **Caulfield, M.P., L.T. Duong, R. O'Brien, J.A. Majzoub, and M. Rosenblatt.** 1988. A chemically synthesized radiolabeled signal peptide: design, preparation, and biological evaluation of an iodinated analog of preproparathyroid hormone. *Mol.Endocrinol.* **2**:452-458.
138. **Caulfield, M.P., L.T. Duong, R.K. Baker, M. Rosenblatt, and M.O. Lively.** 1989. Synthetic substrate for eukaryotic signal peptidase. Cleavage of a synthetic peptide analog of the precursor region of preproparathyroid hormone. *J.Biol.Chem.* **264**:15813-15817.
139. **Dev, I.K., P.H. Ray, and P. Novak.** 1990. Minimum substrate sequence for signal peptidase I of *Escherichia coli*. *J.Biol.Chem.* **265**:20069-20072.
140. **Zhong, W. and S.J. Benkovic.** 1998. Development of an internally quenched fluorescent substrate for *Escherichia coli* leader peptidase. *Anal.Biochem.* **255**:66-73.
141. **Little, J.W., B. Kim, K.L. Roland, M.H. Smith, L.L. Lin, and S.N. Slilaty.** 1994. Cleavage of LexA repressor. *Methods Enzymol.* **244**:266-284.
142. **Peat, T.S., E.G. Frank, R. Woodgate, and W.A. Hendrickson.** 1996. Production and crystallization of a selenomethionyl variant of UmuD', an *Escherichia coli* SOS response protein. *Proteins* **25**:506-509.
143. **Keiler, K.C. and R.T. Sauer.** 1996. Identification of active site residues of the Tsp protease. *J.Biol.Chem.* **270**:28864-28868.

References

144. Strynadka, N.C., H. Adachi, S.E. Jensen, K. Johns, A. Sielecki, C. Betzel, K. Sutoh, and M.N. James. 1992. Molecular structure of the acyl-enzyme intermediate in beta-lactam hydrolysis at 1.7 Å resolution. *Nature* **359**:700-705.
145. Paetzel, M., F. Danel, L. de Castro, S.C. Mosimann, M.G.P. Page, and N.C.J. Strynadka. 2000. Crystal structure of the class D beta-lactamase OXA-10. *Nature Struct.Biol.* **7**:918-925.
146. Kelly, J.A., J.R. Knox, H. Zhao, J.M. Frere, and J.M. Ghaysen. 1989. Crystallographic mapping of beta-lactams bound to a D-alanyl-D-alanine peptidase target enzyme. *J.Mol.Biol.* **209**:281-295.
147. Haase, J. and E. Lanka. 1997. A specific protease encoded by the conjugative DNA transfer systems of IncP and Ti plasmids is essential for pilus synthesis. *J.Bacteriol.* **179**:5728-5735.
148. Patricelli, M.P. and B.F. Cravatt. 1999. Fatty acid amide hydrolase competitively degrades bioactive amides and esters through a nonconventional catalytic mechanism. *Biochemistry* **38**:14125-14130.
149. Birghan, C., E. Mundt, and A.E. Gorbalenya. 2000. A non-canonical Lon proteinase lacking the ATPase domain employs the Ser-Lys catalytic dyad to exercise broad control over the life cycle of a double-stranded RNA virus. *EMBO J.* **19**:114-123.
150. Lejal, N., B. Da Costa, J.C. Huet, and B. Delmas. 2000. Role of Ser-652 and Lys-692 in the protease activity of infectious bursal disease virus VP4 and identification of its substrate cleavage sites. *J.Gen.Virol.* **81**:983-992.
151. Yamamoto, Y., N. Inagaki, and K. Satoh. 2001. Overexpression and characterization of carboxyl-terminal processing protease for precursor D1 protein - Regulation of enzyme-substrate interaction by molecular environments. *J.Biol.Chem.* **276**:7518-7525.
152. Miller, M., J.-K.M. Rao, A. Wlodawer, and M.R. Gribskov. 1993. A left-handed crossover involved in amidohydrolase catalysis: Crystal structure of *Erwinia chrysanthemi* L-asparaginase with bound L-aspartate. *FEBS Lett.* **328**:vol178
153. Rawlings, N.D. and A.J. Barrett. 1994. Families of serine peptidases. *Methods Enzymol.* **244**:19-61.
154. Paetzel, M., M. Chernaia, N.-C.J. Strynadka, W.R. Tschantz, C. Cao, R.E. Dalbey, and M.-N.G. James. 1995. Crystallization of a soluble, catalytically active form of *Escherichia coli* leader peptidase. *Prot.Struct.Funct.Gen.* **23**:122-125.
155. Carlos, J.L., P.A. Klenotic, M. Paetzel, N.C.J. Strynadka, and R.E. Dalbey. 2000. Mutational evidence of transition state stabilization by Serine 88 in *Escherichia coli* type I signal peptidase. *Biochemistry* **39**:7276-7283.
156. Izard, J.W. and D.A. Kendall. 1994. Signal peptides: exquisitely designed transport promoters. *Mol.Microbiol.* **13**:765-773.
157. Karamyshev, A.L., Z.N. Karamysheva, A.V. Kajava, V.N. Ksenzenko, and M.A. Nesmeyanova. 1998. Processing of *Escherichia coli* alkaline phosphatase: Role of the primary structure of the signal peptide cleavage region. *J.Mol.Biol.* **277**:859-870.
158. Bolhuis, A., H. Tjalsma, K. Stephenson, C.R. Harwood, G. Venema, S. Bron, and J.M. van Dijl. 1999. Different mechanisms for thermal inactivation of *Bacillus subtilis* signal peptidase mutants. *J.Biol.Chem.* **274**:15865-15868.
159. Pugsley, A.P. 1993. The complete general secretory pathway in gram-negative bacteria. *Microbiol.Rev.* **57**:50-108.
160. Smith, H., A. de Jong, S. Bron, and G. Venema. 1988. Characterization of signal-sequence-coding regions selected from the *Bacillus subtilis* chromosome. *Gene* **70**:351-361.
161. Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. Molecular cloning: A laboratory manual., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.,
162. Thompson, J.D., D.G. Higgins, and T.J. Gibson. 1994. CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nuc.Acids Res.* **22**:4673-4680.
163. Wu, X.C., W. Lee, L. Tran, and S.-L. Wong. 1991. Engineering a *Bacillus subtilis* expression-secretion system with a strain deficient in six extracellular proteases. *J.Bacteriol.* **173**:4952-4958.
164. van Dijl, J.M., A. de Jong, H. Smith, S. Bron, and G. Venema. 1991. Signal peptidase I overproduction results in increased efficiencies of export and maturation of hybrid secretory proteins in *Escherichia coli*. *Mol.Gen.Genet.* **227**:40-48.
165. van Dijl, J.M., A. de Jong, H. Smith, S. Bron, and G. Venema. 1991. Non-functional expression of *Escherichia coli* signal peptidase I in *Bacillus subtilis*. *J.Gen.Microbiol.* **137**:2073-2084.
166. Welsh, K.M., K.A. Trach, C. Folger, and J.A. Hoch. 1994. Biochemical characterization of the essential GTP-binding protein Obg of *Bacillus subtilis*. *J.Bacteriol.* **176**:7161-7168.
167. Studier, F.W., A.H. Rosenberg, J.J. Dunn, and J.W. Dubendorff. 1990. Use of T7 RNA polymerase to direct expression of cloned genes. *Methods Enzymol.* **6**:60-89.
168. Bolhuis, A., H. Tjalsma, H. Smith, A. de Jong, R. Meima, G. Venema, S. Bron, and J.M. van Dijl. 1999. Evaluation of bottlenecks in the late stages of protein secretion in *Bacillus subtilis*. *Appl.Environ.Microbiol.* **65** :2934-2941.
169. Paetzel, M. and R.E. Dalbey. 1997. Catalytic hydroxyl/amine dyads within serine proteases. *TIBS* **22**:28-31.
170. van Wely, K.H.M., J. Swaving, and A.J.M. Driessen. 1998. Translocation of the precursor of alpha-amylase into *Bacillus subtilis* membrane vesicles. *Eur.J.Biochem.* **255**:690-697.
171. Bolhuis, A., G. Venema, W.J. Quax, S. Bron, and J.M. van Dijl. 1999. Functional analysis of paralogous thiol-disulfide oxidoreductases in *Bacillus subtilis*. *J.Biol.Chem.* **274**:24531-24538.
172. Cregg, K.M., E.I. Wilding, and M.T. Black. 1996. Molecular cloning and expression of the *spsB* gene encoding an essential type I signal peptidase from *Staphylococcus aureus*. *J.Bacteriol.* **178**:5712-5718.
173. Wertman, K.F., A.R. Wyman, and D. Botstein. 1986. Host/vector interactions which affect the viability of recombinant phage lambda clones. *Gene* **49**:253-262.
174. Kunst, F., N. Ogasawara, I. Moszer, A.M. Albertini, G. Alloni, V. Azevedo, M.G. Bertero, P. Bessières et al. 1997. The complete genome sequence of the Gram-positive bacterium *Bacillus subtilis*. *Nature* **390**:249-256.

175. von Heijne, G. 1985. Signal sequences. The limits of variation. *J.Mol.Biol.* **184**:99-105.
176. von Heijne, G. 1994. Sec-independent protein insertion into the inner *E. coli* membrane. *FEBS Lett.* **346**:69-72.
177. von Heijne, G. 1997. Getting greasy: how transmembrane polypeptide segments integrate into the lipid bilayer. *Mol.Microbiol.* **24**:249-253.
178. von Heijne, G. 1983. Patterns of amino acids near signal-sequence cleavage sites. *Eur.J.Biochem.* **133**:17-21.
179. von Heijne, G. 1984. How signal sequences maintain cleavage specificity. *J.Mol.Biol.* **173**:243-251.
180. Shen, L.M., J.I. Lee, S. Cheng, H.-J.A. Kuhn, and R.E. Dalbey. 1991. Use of site-directed mutagenesis to define the limits of sequence variation tolerated for processing of the M13 procoat protein by the *Escherichia coli* leader peptidase. *Biochemistry* **30**:11775-11781.
181. Folz, R.J., S.F. Nothwehr, and J.I. Gordon. 1988. Substrate specificity of eukaryotic signal peptidase. Site-saturation mutagenesis at position -1 regulates cleavage between multiple sites in human pre (delta pro) apolipoprotein A-II. *J.Biol. Chem.* **263**:2070-2078.
182. Fikes, J.D., G.A. Barkocy-Gallagher, D.G. Klapper, and P.J.J. Bassford. 1990. Maturation of *Escherichia coli* maltose-binding protein by signal peptidase I *in vivo*. Sequence requirements for efficient processing and demonstration of an alternate cleavage site. *J.Biol. Chem.* **265**:3417-3423.
183. Schanstra, J.P., R. Rink, F. Pries, and D.B. Janssen. 1993. Construction of an expression and site-directed mutagenesis system of haloalkane dehalogenase in *Escherichia coli*. *Prot. Expr. Pur.* **4**:479-489.
184. Kyhse-Andersen, J. 1984. Electrophoretic transfer of multiple gels: a simple apparatus without buffer tank for rapid transfer of proteins from polyacrylamide to nitrocellulose. *J.Biochem.Biophys.Methods* **10**:203-209.
185. Blattner, F.R., G. Plunkett, C.A. Bloch, N.T. Perna, V. Burland, M. Riley, J. Collado-Vides, J.D. Glasner, C.K. Rode, and G.F. Mayhew. 1997. The complete genome sequence of *Escherichia coli* K-12. *Science* **277**:1453-1462.
186. Fleischmann, R.D., M.D. Adams, O. White, R.A. Clayton, E.F. Kirkness, A.R. Kesteven, C.J. Bult, J.F. Tomb, B.A. Dougherty, and J.M. Merrick. 1995. Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science* **269**:496-512.
187. Tomb, J.F., O. White, A.R. Kerlavage, R.A. Clayton, G.C. Sutton, R.D. Fleischmann, K.A. Ketchum, H.P. Klenk, S. Gill, and B.A. Dougherty. 1997. The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* **377**:539-547.
188. Philipp, W.J., S. Poulet, K. Eiglmeir, S. Pascopella, V. Balasubramanian, B. Heym, S. Bergh, B.R. Bloom, W.R. Jacobs, and S.T. Cole. 1996. An integrated map of the genome of the tubercle Bacillus, *Mycobacterium tuberculosis* H37Rv, and comparison with *Mycobacterium leprae*. *Proc.Natl.Acad.Sci.USA* **93**:3132-3137.
189. Bult, C.J., C.O. White, G.J. Olsen, L. Zhou, R.D. Fleischmann, G.C. Sutton, J.A. Blake, T.M. Fitzgerald, R.A. Clayton, and J.D. Gocayne. 1996. Complete genome sequence of the methanogenic archaeon, *Methanococcus jannaschii*. *Science* **273**:1058-1073.
190. Smith, D.R., L.A. Doucette-Stamm, C. Deloughery, H. Lee, J. Dubois, T. Aldrege, R. Bashirzadeh, D. Blakely, R. Cook, and K. Gilbert. 1997. Complete genome sequence of *Methanobacterium thermoautotrophicum* deltaH: functional analysis and comparative genomics. *J.Bacteriol.* **179**:7135-7155.
191. Goffeau, A., R. Aert, M.L. Agostini-Carbone, A. Ahmed, M. Aigle, L. Alberghina, K. Albermann, M. Albers, M. Aldea, and D. Alexandraki. 1997. The yeast genome directory. *Nature* **387**:5-105.
192. van Dijk, J.M., H. Smith, S. Bron, and G. Venema. 1988. Synthesis and processing of *Escherichia coli* TEM-beta-lactamase and *Bacillus licheniformis* alpha-amylase in *E. coli*: the role of signal peptidase I. *Mol.Gen.Genet.* **214**:55-61.
193. Bohni, P.C., R.J. Deshaies, and R.W. Schekman. 1988. Sec11 is required for signal peptide processing and yeast cell growth. *J.Cell.Biol.* **106**:1035-1042.
194. Kaneko, T., S. Sato, H. Kotani, A. Tanaka, E. Asamizu, Y. Nakamura, N. Miyajima, M. Hirose, M. Sugiura, and S. Sasamoto. 1996. Sequence analysis of the genome of the unicellular cyanobacterium *Synechocystis* sp. strain PCC6803. II Sequence determination of the entire genome and assignment of potential protein-coding regions. *DNA Res.* **30**:109-136.
195. Behrens, M., G. Michaelis, and E. Pratje. 1991. Mitochondrial inner membrane protease 1 of *Saccharomyces cerevisiae* shows sequence similarity to the *Escherichia coli* leader peptidase. *Mol.Gen.Genet.* **228**:167-176.
196. Nunnari, J., T.D. Fox, and P. Walter. 1993. A mitochondrial protease with two catalytic subunits of nonoverlapping specificities. *Science* **262**:1997-2004.
197. Klenk, H.P., R.A. Clayton, J.F. Tomb, O. White, K.E. Nelson, K.A. Ketchum, R.J. Dodson, M. Gwinn, E.K. Hickey, and J.D. Peterson. 1997. The complete genome sequence of the hyperthermophilic, sulphate-reducing archaeon *Archaeoglobus fulgidus*. *Nature* **390**:364-370.
198. Altschul, S.F., T.L. Madden, A.A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D.J. Lipman. 1997. Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nuc.Acids Res.* **25**:3389-3402.
199. Cronan, J.E. 1990. Biotinylation of proteins *in vivo*. *J.Biol. Chem.* **265**:10327-10333.
200. Miller, J.H. 1982. Experiments in molecular genetics, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
201. Palva, I. 1982. Molecular cloning of alpha-amylase gene from *Bacillus amyloliquefaciens* and its expression in *Bacillus subtilis*. *Gene* **19**:81-87.
202. Kim, L., A. Mogk, and W. Schumann. 1996. A xylose-inducible *Bacillus subtilis* integration vector and its application. *Gene* **181**:71-76.
203. van Sinderen, D., S. Withoff, H. Boels, and G. Venema. 1990. Isolation and characterisation of *comL*, a transcriptional unit involved in competence development of *Bacillus subtilis*. *Mol.Gen.Genet.* **224**:396-404.
204. Vieria, J. and J. Messing. 1990. New pUC-derived cloning vectors with different selectable markers and DNA replication origins. *Gene* **100**:253-262.

References

205. **Bron, S. and G. Venema.** 1972. Ultraviolet inactivation and excision-repair in *Bacillus subtilis*. IV. Integration and repair of ultraviolet-inactivated transforming DNA. *Mutat.Res.* **15**:395-409.
206. **Shelness, G.S. and G. Blobel.** 2001. Two subunits of the canine signal peptidase complex are homologous to yeast Sec11 protein. *J.Biol.Chem.* **265**:9512-9519.
207. **Greenburg, G., G.S. Shelness, and G. Blobel.** 1989. A subunit of mammalian signal peptidase is homologous to yeast Sec11 protein. *J.Biol.Chem.* **264**:15762-15765.
208. **Signs, S.A. and R. Jacquet.** 1994. Induction of ethanol dependence increases signal peptidase mRNA levels in rat brain. *Mol.Cell.Biochem.* **139**:21-26.
209. **Kontinen, V.P. and M. Sarvas.** 1988. Mutants of *B. subtilis* defective in protein export. *J.Gen.Microbiol.* **134**:2333-2344.
210. **Jander, G., J.E. Cronan, and J. Beckwith.** 1996. Biotinylation *in vivo* as a sensitive indicator of protein secretion and membrane protein insertion. *J.Bacteriol.* **178**:3049-3058.
211. **Ferrari, E., A.S. Jarnagin, and B.F. Schmidt.** 1993. p.917-937. In A.L. Sonenshein, J.A. Hoch, and R. Losick (ed.), *Bacillus subtilis* and other Gram-positive bacteria, American society for Microbiology, Washington, D.C.
212. **Msadek, T., F. Kunst, and G. Rapoport.** 1993. p.729-745. In A.L. Sonenshein, J.A. Hoch, and G. Rapoport (ed.), *Bacillus subtilis* and other Gram-positive bacteria, American society for Microbiology, Washington, D.C.
213. **Jacobs, M.F.** 1995. Expression of the subtilisin Carlsberg-encoding gene in *Bacillus licheniformis* and *Bacillus subtilis*. *Gene* **152**:69-74.
214. **Hoang, V. and J. Hofemeister.** 1995. *Bacillus amyloliquefaciens* possesses a second type I signal peptidase with extensive sequence similarity to other *Bacillus* SPases. *Biochim.Biophys.Acta* **1269**:64-68.
215. **Newsome, A.L., J.W. McLean, and M.O. Lively.** 1992. Molecular cloning of a cDNA encoding the glycoprotein of hen oviduct microsomal signal peptidase. *Biochem.J.* **282**:447-452.
216. **Fang, H., C. Mullins, and N. Green.** 1997. In addition to SEC11, a newly identified gene, SPC3, is essential for signal peptidase activity in the yeast endoplasmic reticulum. *J.Biol.Chem.* **272**:13152-13158.
217. **Meyer, H.A. and E. Hartmann.** 1997. The yeast SPC22/23 homolog Spc3p is essential for signal peptidase activity. *J.Biol.Chem.* **272**:13159-13164.
218. **Thompson, J.D., T.J. Gibson, F. Plewniak, F. Jeanmougin, and D.G. Higgins.** 1997. The CLUSTAL-X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nuc.Acids Res.* **25**:4876-4882.
219. **Strimmer, K. and A. Von Haeseler.** 1996. Quartet puzzling: A quartet maximum-likelihood method for reconstructing tree topologies. *Mol.Biol.Evol.* **13**:964-969.
220. **Vriend, G.** 1990. WHAT IF: A molecular modeling and drug design program. *J.Mol.Graph.* **8**:52-56.
221. **Carlos, J.L., M. Paetzel, G. Brubaker, A. Karla, C.M. Ashwell, M.O. Lively, G.Q. Cao, P. Bullinger, and R.E. Dalbey.** 2000. The role of the membrane-spanning domain of type I signal peptidases in substrate cleavage site selection. *J.Biol.Chem.* **275**:38813-38822.
222. **van Roosmalen, M.L., J.D.H. Jongbloed, J.-Y.F. Dubois, G. Venema, S. Bron, and J.M. van Dijl.** 2001. Distinction between Major and Minor *Bacillus* Signal Peptidases Based on Phylogenetic and Structural Criteria. *J.Biol.Chem.* **276**:25230-25235.
223. **Mall, S., R. Broadbridge, R.P. Sharma, A.G. Lee, and J.M. East.** 2000. Effects of aromatic residues at the ends of transmembrane alpha-helices on helix interactions with lipid bilayers. *Biochemistry* **39**:2071-2078.
224. **Monne, M., I. Nilsson, M. Johansson, N. Elmhed, and G. von Heijne.** 1998. Positively and negatively charged residues have different effects on the position in the membrane of a model transmembrane helix. *J.Mol.Biol.* **284**:1177-1183.
225. **Kim, Y.T., T. Muramatsu, and K. Takahashi.** 1995. Identification of Trp300 as an important residue for *Escherichia coli* leader peptidase activity. *Eur.J.Biochem.* **234**:358-362.
226. **Dalbey, R.E. and W. Wickner.** 1987. Leader peptidase of *Escherichia coli*: critical role of a small domain in membrane assembly. *Science* **235**:783-787.
227. **Andersson, H. and G. von Heijne.** 1993. Position-specific Asp-Lys pairing can affect signal sequence function and membrane protein topology. *J.Biol.Chem.* **268**:21389-21393.
228. **Dalbey, R.E., A. Kuhn, and W. Wickner.** 1987. The internal signal sequence of *Escherichia coli* leader peptidase is necessary, but not sufficient, for its rapid membrane assembly. *J.Biol.Chem.* **262**:13241-13245.
229. **Dalbey, R.E.** 1991. Leader peptidase. *Mol.Microbiol.* **5**:2855-2860.