

University of Groningen

Transport and sorting of sphingolipids in polarized cells

IJzendoorn, Sven Christian David van

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1999

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

IJzendoorn, S. C. D. V. (1999). *Transport and sorting of sphingolipids in polarized cells: the involvement of the sub-apical compartment*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

- Ahmed, S.N., Brown, D.A. and E. London.** (1997). On the origin of sphingolipid/cholesterol-rich detergent-insoluble cell membranes: physiological concentrations of cholesterol and sphingolipid induce formation of a detergent-insoluble, liquid-ordered lipid phase in model membranes. *Biochem.* 36: 10944-10953.
- Ali, N. and Evans, W.H.** 1990. Priority targeting of glycosyl-phosphatidylinositol-anchored proteins to the bile-canalicular (apical) plasma membrane of hepatocytes. Involvement of 'late' endosomes. *Biochem. J.* 271:193-1990.
- Alonso, M.A., Fan, L. and Alarcon, B.** (1997). Multiple sorting signals determine apical localization of a nonglycosylated integral membrane protein. *J. Biol. Chem.* 272:30748-30752.
- Ammala, C., Ashcroft, F.M. and Rorsman, P.** (1993). Calcium-independent potentiation of insulin release by cyclic AMP in single beta-cells. *Nature* 363, 356-358.
- Apodaca, G., M. Bomsel, J.Arden, P.P. Breitfeld, K. Tang and K.E. Mostov.** (1991). The polymeric immunoglobulin receptor: a model protein to study transcytosis. *J. Clin. Invest.* 87:1877-1882.
- Apodaca, G., Enrich, C. and Mostov, K.E.** (1994). The calmodulin antagonist, W-13, alters transcytosis, recycling, and the morphology of the endocytic pathway in Madin-Darby canine kidney cells. *J. Biol. Chem.* 269:19005-19013.
- Apodaca, G., Katz, L.A. and Mostov, K.E.** (1994). Receptor-mediated transcytosis of IgA in MDCK cells is via apical recycling endosomes. *J. Cell Biol.* 125:67-86.
- Arni, S., Keilbaugh, S.A., Ostermeyer, A.G. and Brown, D.A.** (1998). Association of GAP-43 with detergent-resistant membranes requires two palmitoylated cysteine residues. *J. Biol. Chem.* 273:28478-28485.
- Aroeti, B. and Mostov, K.E.** (1994). Polarized sorting of the polymeric immunoglobulin receptor in the exocytotic and endocytotic pathways is controlled by the same amino acids. *EMBO J.* 13:2297-2304.
- Aroeti, B., Okhrimenko, H., Reich, V. and Orzech, E.** (1998). Polarized trafficking of plasma membrane proteins: emerging roles for coats, SNAREs, GTPases and their link to the cytoskeleton. *Biochim. Biophys. Acta* 1376:57-90.
- Ayscough, K.R., Stryker, J., Pokala, N., Sanders, M., Crews, P. and Drubin, D.G.** (1997). High rates of actin filament turnover in budding yeast and roles for actin in establishment and maintenance of cell polarity revealed using the actin inhibitor latrunculin-A. *J. Cell Biol.* 137:399-416.
- Babia, T., Kok, J.W. and Hoekstra, D.** (1994). The use of fluorescent lipid analogues to study endocytosis of glycosphingolipids. In Receptor Research Methods. B. Greenstein, editor. Harwood Acad. Publ./London, UK. 155-174.
- Barroso, M. and Sztul, E.S.** (1994). Basolateral to apical transcytosis in polarized cells is indirect and involves BFA and trimeric G protein sensitive passage through the apical endosome. *J. Cell Biol.* 124:83-100.
- Barr, V.A., L.J. Scott and A.L. Hubbard.** (1995). Immunoabsorption of hepatic vesicles carrying newly synthesized dipeptidyl peptidase IV and polymeric IgA receptor. *J. Biol. Chem.* 270:27834-27844.
- Barr, V.A. and A.L. Hubbard.** (1993). Newly synthesized hepatocyte plasma membrane proteins are transported in transcytotic vesicles in the bile duct-ligated rat. *Gastroenterology* 105:44-571.
- Bartles, J.R., Feracci, H.M., Stieger, B. and Hubbard, A.L.** (1987). Biogenesis of the rat hepatocyte plasma membrane in vivo: comparison of the pathways taken by apical and basolateral proteins using subcellular fractionation. *J. Cell Biol.* 105:1241-1251.
- Bender, V., Buschlen, S. and Cassio, D.** (1998). Expression and localization of hepatocyte domain-specific plasma membrane proteins in hepatoma x fibroblast hybrids and in hepatoma dedifferentiated variants. *J. Cell Sci.* 111, 3437-50.
- Bligh, E.G. and Dyer, W.J.** (1959). A rapid method of total lipid extraction and purification. *Can. J. Biochem. Biophys.* 37:911-017.

- Boyer, J.L. and Soroka, C.J.** (1995). Vesicle targeting to the apical domain regulates bile excretory function in isolated rat hepatocyte couplets. *Gastroenterology* 109:1600-1611.
- Bomsel, M., Prydz, K., Parton, R.G., Gruenberg, J. AND Simons K** (1989). Endocytosis in filter-grown Madin-Darby canine kidney cells. *J. Cell Biol.* 109:3243-3258.
- Bomsel, M. and Mostov, K.** (1993). Possible role of both the α and $\beta\gamma$ subunits of the heterotrimeric G protein, G_s , in transcytosis of the polymeric immunoglobulin receptor. *J. Biol. Chem.* 268, 25824-25835.
- Bonilha, V.L., Marmorstein, A.D., Cohen-Gould, L. and Rodriguez-Boulan, E.** (1997). Apical sorting of influenza hemagglutinin by transcytosis in retinal pigment epithelium. *J. Cell Sci.* 110:1717-1727.
- Breitfeld, P.P., J.M. Harris and K.E. Mostov.** (1989). Postendocytic sorting of the ligand for the polymeric immunoglobulin receptor in Madin-Darby canine kidney cells. *J. Cell Biol.* 109:475-486.
- Brignoni, M., Pignataro, O.P., Rodriguez, M.L., Vega-Salas, D.E., Rodriguez-Boulan, E. and Salas, P.J.I.** (1995). Cyclic AMP modulates the rate of 'constitutive' exocytosis of apical membrane proteins in Madin-Darby canine kidney cells. *J. Cell Sci.* 108:1931-1943.
- Brown, D.A. and London, E.** (1998). Functions of lipid rafts in biological membranes. *Annu. Rev. Cell Dev. Biol.* 14:111-136.
- Brown, D.A. and J.K. Rose.** (1992). Sorting of GPI-anchored proteins to glycolipid-enriched membrane subdomains during transport to the apical surface. *Cell* 68:533-544.
- Calhoun, B.C. and Goldenring, J.R.** (1997). Two Rab proteins, vesicle-associated membrane protein 2 (VAMP-2) and secretory carrier membrane proteins (SCAMPs), are present on immunoisolated parietal cell tubulovesicles. *Biochem. J.* 325:559-564.
- Cariappa, R. and M.S. Kilberg.** (1992). Plasma membrane domain localization and transcytosis of the glucagon-induced hepatic system A carrier. *Am. J. Physiol.* 263:E1021-E1028.
- Cardone, M.H., Smith, B.L., Song, W., Mochly-Rosen, D. and Mostov, K.E.** (1994). Phorbol Myristate Acetate-mediated stimulation of transcytosis and apical recycling in MDCK cells. *J. Cell Biol.* 124:717-727.
- Casanova, J.E., Breitfeld, P.P., Ross, S.A. and Mostov, K.E.** (1990). Phosphorylation of the polymeric immunoglobulin receptor required for its efficient transcytosis. *Science* 248:742-745.
- Casanova, J.E., Wang, X., Kumar, R., Bhartur, S.G., Navarre, J., Woodrum, J.E., Altschuler, Y., Ray, G.S. and Goldenring, J.R.** (1999). Association of rab25 and rab11a with the apical recycling system of polarized madin-darby canine kidney cells. *Mol. Biol. Cell.* 10:47-61.
- Casu, A. and Camogliano, L.** (1990). Glycerophospholipids and cholesterol composition of bile in bile-fistula rats treated with monensin. *Biochim. Biophys. Acta* 1043:113-115.
- Chapin, S.J., Enrich, C., Aroeti, B., Havel, R.J. and Mostov, K.E.** (1996). Calmodulin binds to the basolateral targeting signal of the polymeric immunoglobulin receptor. *J. Biol. Chem.* 271:1336-1342.
- Chen, W., Feng, Y, Chen, D. and Wandinger-Ness, A.** (1998). Rab11 is required fro trans-Golgi network-to-plasma membrane transport and a preferential target for GDP dissociation inhibitor. *Mol. Biol. Cell.* 9: 3241-3257.
- Chiu, J., Hu, C., Lui, W., Lo, S. and Chang, C.** (1990). The formation of bile canaliculi in hepatoma cell lines. *Hepatology* 11:834-842.
- Chuang, J.Z. and Sung, C.H.** (1998). The cytoplasmic tail of rhodopsin acts as a novel apical sorting signal in polarized MDCK cells. *J. Cell Biol.* 142:1245-1256.
- Clague, M.J.** (1998). Molecular aspects of the endocytic pathway. *Biochem. J.* 336:271-282.
- Daro, E., van der Sluijs, P., Galli, T. and Mellman, I.** (1996). Rab4 and cellubrevin define different early endosome populations on the pathway of transferrin receptor recycling. *Proc. Natl. Acad. Sci. U. S. A.* 93:9559-9564.
- de Vries, H., Schrage, C. and Hoekstra D.** (1998). An apical-type trafficking pathway is present in cultured oligodendrocytes but the sphingolipid-enriched myelin membrane is the target of a basolateral-type pathway. *Mol. Biol. Cell* 9:599-609.

- D'Souza-Schorey, C., van Donselaar, E., Hsu, V.W., Yang, C., Stahl, P.D. and Peters, P.J.** (1998). ARF6 targets recycling vesicles to the plasma membrane: insights from an ultrastructural investigation. *J. Cell Biol.* 140:603-616.
- Enrich, C., Jackle, S. and Havel, R.J.** (1996). The polymeric immunoglobulin receptor is the major calmodulin-binding protein in an endosome fraction from rat liver enriched in recycling receptors. *Hepatology* 24:226-232.
- Enrich, C., Bachs, O., Evans, W.H.** (1988). A 115 kDa calmodulin-binding protein is located in rat liver endosome fractions. *Biochem. J.* 255:999-1005.
- Farquhar, M.G.** (1985). Progress in unraveling pathways of Golgi traffick. *Annu. Rev. Cell Biol.* 1:447-488.
- Forte, J.G. and Yao, D.X.** (1996). The membrane recruitment-and-recycling hypothesis of gastric HCl secretion. *Trends in Cell Biol.*6:45-48.
- Fushimi, K., Sasaki, S. and Marumo, F.** (1997). Phosphorylation of serine 256 is required for cAMP-dependent regulatory exocytosis of the aquaporin-2 water channel. *J. Biol. Chem.* 272, 14800-14804.
- Futter, C.E., Connolly, C.N., Cutler, D.F. and Hopkins, C.R.** (1995). Newly synthesized transferrin receptors can be detected in the endosome before they appear on the cell surface. *J. Biol. Chem.* 270:10999-11003.
- Futter, C.E., Gibson, A., Allchin, E.H., Maxwell, S., Ruddock, L.J., Odorizzi, G., Domingo, D., Trowbridge, I.S. and Hopkins, C.R.** (1998). In polarized MDCK cells basolateral vesicles arise from clathrin-gamma-adaptin-coated domains on endosomal tubules. *J. Cell Biol.* 141,611-623.
- Geuze, H.J., Slot, J.W., Strous, G.J., Peppard, J., von Figura, K., Haslik, A. and Schwartz, A.L.** (1984). Intracellular receptor sorting during endocytosis: comparative immunoelectron microscopy of multiple receptors in rat liver. *Cell* 37:195-204.
- Geli, M.I., Wesp, A. and Riezman, H.** (1998). Distinct functions of calmodulin are required for the uptake step of receptor-mediated endocytosis in yeast: the type I myosin Myo5p is one of the calmodulin targets. *EMBO J.* 17:635-647.
- Gibson, A., Futter, C.E., Maxwell, S., Allchin, E.H., Shipman, M., Kraehenbuhl, J.P., Domingo, D., Odorizzi, G., Trowbridge, I.S. and Hopkins, C.R.** (1998). Sorting mechanisms regulating membrane protein traffic in the apical transcytotic pathway of polarized MDCK cells. *J. Cell Biol.* 143, 81-94.
- Goda S., Kobayashi T and I Goto.** (1987). Hydrolysis of galactosylsphingosine and lactosylsphingosine by beta-galactosidases in human brain and cultured fibroblasts. *Biochim Biophys Acta* 920:259-265.
- Goldenring, J.R., Shen, K.R., Vaughan, H.D. and Modlin, I.M.** (1993). Identification of a small GTP-binding protein, Rab25, expressed in the gastrointestinal mucosa, kidney, and lung. *J. Biol. Chem.* 268:18419-18422.
- Goldenring, J.R., Smith, J., Vaughan, H.D., Cameron, P., Hawkins, W. and Navarre, J.** (1996). Rab11 is an apically located small GTP-binding protein in epithelial tissues. *Am. J. Physiol.* 270:G515-G525.
- Gottlieb, T.A., Ivanov, I.E., Adesnik, M. and Sabatini, D.D.** (1993). Actin microfilaments play a critical role in endocytosis at the apical but not the basolateral surface of polarized epithelial cells. *J. Cell Biol.* 120:695-710.
- Green, E.G., Ramm, E., Riley, N.M., Spiro, D.J., Goldenring, J.R. and Wessling-Resnick, M.** (1997). Rab11 is associated with transferrin-containing recycling compartments in K562 cells. *Biochem. Biophys. Res. Commun.* 239:612-616.
- Gruenberg, J., G. Griffiths and K.E. Howell.** (1989). Characterization of the early endosome and putative endocytic carrier vesicles in vivo and with an assay of vesicle fusion in vitro. *J. Cell Biol.* 108:1301-1316.
- Gruenberg, J. and F.R. Maxfield.** (1995). Membrane transport in the endocytic pathway. *Curr. Opin. Cell Biol.* 7:552-563.

- Gut, A., Balda, M.S. and Matter, K.** (1998). The cytoplasmic domains of a beta1 integrin mediate polarization in Madin-Darby canine kidney cells by selective basolateral stabilization. *J. Biol. Chem.* 273:29381-29388.
- Hansen, S.H. and Casanova, J.E.** (1994). Gsa stimulates transcytosis and apical secretion in MDCK cells through cAMP and protein kinase A. *J. Cell Biol.* 126:677-687.
- Hansen, G.H., Niels-Christiansen, L.L., Immerdal, L., Hunziker, W., Kenny, A.J. and Michael Danielsen, E.** (1999). Transcytosis of Immunoglobulin A in the Mouse Enterocyte Occurs Through Glycolipid Raft- and Rab17-Containing Compartments. *Gastroenterology* 116:610-622.
- Hauri, H.P. and Matter, K.** (1991). Protein traffic in intestinal epithelial cells. *Semin. Cell Biol.* 2:355-364.
- Heilker, R., Manning-Krieg, U., Zuber, J.F. and Spiess, M.** (1996). In vitro binding of clathrin adaptors to sorting signals correlates with endocytosis and basolateral sorting. *EMBO J.* 15:2893-2899.
- Hemery, I., Durand-Schneider, A., Feldmann, G., Vaerman, J. and M. Maurice.** (1996). The transcytotic pathway of an apical plasma membrane protein (B10) in hepatocytes is similar to that of IgA and occurs via a tubular pericentriolar compartment. *J. Cell Sci.* 109:1215-1227.
- Hirschberg, K., Miller, C.M., Ellenberg, J., Presley, J.F., Siggia, E.D., Phair, R.D. and Lippincott-Schwartz, J.** (1998). Kinetic analysis of secretory protein traffic and characterization of golgi to plasma membrane transport intermediates in living cells. *J. Cell Biol.* 143:1485-1503.
- Hirst, J. and Robinson, M.S.** (1998). Clathrin and adaptors. *Biochim. Biophys. Acta.* 1404:173-193.
- Hoekstra, D. and Kok, J.W.** (1992). Trafficking of glycosphingolipids in eukaryotic cells; sorting and recycling of lipids. *Biochim. Biophys. Acta* 1113:277-294.
- Hoppe, C.A., Connolly, T.P. and Hubbard, A.L.** (1985). Transcellular transport of polymeric IgA in the rat hepatocyte: biochemical and morphological characterization of the transport pathway. *J. Cell Biol.* 101:2113-2123.
- Hopkins, C.R., Gibson, A., Shipman, M., Strickland, D.K. and Trowbridge, I.S.** (1994). In migrating fibroblasts, recycling receptors are concentrated in narrow tubules in the pericentriolar area, and then routed to the plasma membrane of the leading lamella. *J. Cell Biol.* 125:1265-1274.
- Houliston, E., Pickering, S.J. and Maro, B.** (1987). Redistribution of microtubules and pericentriolar material during the development of polarity in mouse blastomeres. *J. Cell Biol.* 104:1299-1308.
- Hughson, E.J. and C.R. Hopkins.** (1990). Endocytic pathways in polarized Caco-2 cells: identification of an endosomal compartment accessible from both apical and basolateral surfaces. *J. Cell Biol.* 110:337-348.
- Hunziker, W., Harter, C., Matter, K., and Mellman, I.** (1991). Basolateral sorting in MDCK cells requires a distinct cytoplasmic domain determinant. *Cell* 66:907-920.
- Hunziker, W., Male, P. and Mellman, I.** (1990). Differential microtubule requirements for transcytosis in MDCK cells. *EMBO J.* 9:3515-3525.
- Hunziker, W. and I. Mellman.** (1989). Expression of macrophage-lymphocyte Fc receptors in Madin-Darby canine kidney cells: polarity and transcytosis differ for isoforms with or without coated pit localization domains. *J. Cell Biol.* 109:3291-3302.
- Hunziker, W. and Peters, P.J.** (1998). Rab17 localizes to recycling endosomes and regulates receptor-mediated transcytosis in epithelial cells. *J. Biol. Chem.* 273, 15734-15741.
- Ihrke, G., Martin, G.V., Shanks, M.R., Schrader, M., Schroer, T.A. and A. L. Hubbard.** (1998). Apical plasma membrane proteins and endolyn-78 travel through a sub-apical compartment in polarized WIF-B hepatocytes. *J. Cell Biol.* 141:115-133.

- Iwabuchi, K., Handa, K. and Hakomori, S.** (1998). Separation of "glycosphingolipid signaling domain" from caveolin-containing membrane fraction in mouse melanoma B16 cells and its role in cell adhesion coupled with signaling. *J. Biol. Chem.* 273:33766-33773.
- Jackman, M.R., Shurety, W., Ellis, J.A. and Luzio, J.P.** (1994). Inhibition of apical but not basolateral endocytosis of ricin and folate in Caco-2 cells by cytochalasin D. *J. Cell Sci.* 107:2547-2456.
- Jones AL, G.T. Hradek, D.L. Schmucker and B.J. Underdown.** (1984). The fate of polymeric and secretory immunoglobulin A after retrograde infusion into the common bile duct in rats. *Hepatology* 4:1173-1183.
- Katsura, T., Verbavatz, J-M., Farinas, J., Ma, T., Ausiello, D.A., Verkman, A.S. and Brown, D.** (1997). Constitutive and regulatory membrane expression of aquaporin 1 and aquaporin 2 water channels in stably transfected LLC-PK1 epithelial cells. *Proc. Natl. Acad. Sci. U.S.A.* 92, 7212-7216.
- Keller, P. and Simons, K.** (1998). Cholesterol is required for surface transport of influenza virus hemagglutinin. *J. Cell Biol.* 140:1357-1367.
- Kipp, H. Sai, Y. and Arias, I.M.** (1998). Direct targeting of newly synthesized ABC-transporters from Golgi to the apical membrane in rat hepatocytes and GFP-mdr1 transfected Wif-B9 cells. *Mol. Biol. Cell* 9:209a
- Kirchhausen, T., Bonifacino, J.S. and Riezman, H.** (1997). Linking cargo to vesicle formation: receptor tail interactions with coat proteins. *Curr. Opin. Cell Biol.* 9:488-495.
- Kishimoto, Y.** (1975). A facile synthesis of ceramides. *Chem. Phys. Lip.* 15:33-36.
- Knight, A., Hughson, E., Hopkins, C.R. and Cutler, D.F.** (1995). Membrane protein trafficking through the common apical endosome compartment of polarized Caco-2 cells. *Mol. Biol. Cell* 6:597-610.
- Kok, J.W., Babia, T. and Hoekstra, D.** (1991). Sorting of sphingolipids in the endocytic pathway of HT29 cells. *J. Cell Biol.* 114:231-239.
- Kok, J.W., K. Hoekstra, S. Eskelinen and D. Hoekstra.** (1992). Recycling pathways of glucosylceramide in BHK cells: distinct involvement of early and late endosomes. *J Cell Sci.* 103:1139-1152.
- Kok, J.W. and Hoekstra, D.** (1994). Glycosphingolipid trafficking in the endocytic pathway. *Curr. topics in membranes* 40:503-537.
- Kramer, E.M., Koch, T., Niehaus, A. and Trotter, J.** (1997). Oligodendrocytes direct glycosyl phosphatidylinositol-anchored proteins to the myelin sheath in glycosphingolipid-rich complexes. *J. Biol. Chem.* 272, 8937-8945.
- Kundu, A., Avalos, R.T., Sanderson, C.M. and Nayak, D.P.** (1996). Transmembrane domain of influenza virus neuraminidase, a type II protein, possesses an apical sorting signal in polarized MDCK cells. *J. Virol.* 70:6508-6515.
- Lafont, F., Lecat, S., Verkade, P. and Simons, K.** (1998). Annexin XIIIb associates with lipid microdomains to function in apical delivery. *J. Cell Biol.* 142:1413-1427.
- Lamaze, C., Fujimoto, L.M., Yin, H.L. and Schmid, S.L.** (1997). The actin cytoskeleton is required for receptor-mediated endocytosis in mammalian cells. *J. Biol. Chem.* 272:20332-20335.
- Le Borgne, R. and Hoflack, B.** (1998). Mechanisms of protein sorting and coat assembly: insights from the clathrin-coated vesicle pathway. *Curr. Opin. Cell Biol.* 10:499-503.
- Leitinger, B., Hille-Rehfeld, A. and Spiess, M.** (1995). Biosynthetic transport of the asialoglycoprotein receptor H1 to the cell surface occurs via endosomes. *Proc. Natl. Acad. Sci. U. S. A.* 92:10109-10113.
- Lichtenstein, Y., Desnos, C., Faundez, V., Kelly, R.B. and Clift-O'Grady, L.** (1998). Vesiculation and sorting from PC12-derived endosomes in vitro. *Proc. Natl. Acad. Sci. U. S. A.* 95:11223-11228.
- Lin, S., Naim, H.Y., Rodriguez, A.C. and Roth, M.G.** (1998). Mutations in the middle of the transmembrane domain reverse the polarity of transport of the influenza virus hemagglutinin in MDCK epithelial cells. *J. Cell Biol.* 142:51-57.
- Lipsky, N.G. and Pagano, R.E.** (1985). Intracellular translocation of fluorescent sphingolipids in cultured fibroblasts: endogenously synthesized sphingomyelin and glucocerebroside analogues pass through the Golgi

apparatus en route to the plasma membrane. *J. Cell Biol.* 100:27-34.

Lisanti, M.P. and Rodriguez-Boulan, E. (1990). Glycosphingolipid membrane anchoring provides clues to the mechanism of protein sorting in polarized epithelial cells. *Trends Biochem. Sci.* 15:13-118.

Lisanti, M.P., Caras, I.W., Davitz, M.A. and Rodriguez-Boulan, E. (1989). A glycosphingolipid membrane anchor acts as an apical targeting signal in polarized epithelial cells. *J. Cell Biol.* 109:2145-2156.

Lochter, A. and Schachner, M. (1997) Inhibitors of protein kinases abolish ECM-mediated promotion of neuronal polarity. *Exp. Cell Res.* 235, 124-129.

Low, S.H., Chapin, S.J., Weimbs, T., Komuves, L.G., Bennett, M.K. and Mostov, K.E. (1996). Differential localization of syntaxin isoforms in polarized Madin-Darby canine kidney cells. *Mol. Biol. Cell* 7:2007-2018.

Lutcke, A., Jansson, S., Parton, R.G., Chavrier, P., Valencia, A., Huber, L.A., Lehtonen, E. and Zerial, M. (1993). Rab17, a novel small GTPase, is specific for epithelial cells and is induced during cell polarization. *J. Cell Biol.* 121:553-564.

Luton, F., Cardone, M.H., Zhang, M. and Mostov, K.E. (1998). Role of tyrosine phosphorylation in ligand-induced regulation of transcytosis of the polymeric Ig receptor. *Mol. Biol. Cell* 9:1787-1802.

Maric, K., Oksche, A. and Rosenthal, W. (1998). Aquaporin-2 expression in primary cultured rat inner medullary collecting duct cells. *Am. J. Physiol.* 275, F796-F801.

Matlin, K.S. and K. Simons. (1984). Sorting of an apical plasma membrane glycoprotein occurs before it reaches the cell surface in cultured epithelial cells. *J. Cell Biol.* 99:2131-2139.

Maratos-Flier, E., C.Y.Y. Kao, E.M. Verdin and G.L. King. (1987). Receptor-mediated vectorial transcytosis of epidermal growth factor by Madin-Darby canine kidney cells. *J. Cell Biol.* 105:1595-1601.

Marmorstein, A.D., Gan, Y.C., Bonilha, V.L., Finnemann, S.C., Csaky, K.G. and Rodriguez-Boulan, E. (1998). Apical polarity of N-CAM and EMMPRIN in retinal pigment epithelium resulting from suppression of basolateral signal recognition. *J. Cell Biol.* 142:697-710.

Matsuoka, K., Orci, L., Amherdt, M., Bednarek, S.Y., Hamamoto, S., Schekman, R. and Yeung, T. (1998). COPII-coated vesicle formation reconstituted with purified coat proteins and chemically defined liposomes. *Cell* 17:263-275.

Matter, K., K. Bucher and H.P. Hauri. (1990). Microtubule perturbation retards both the direct and the indirect apical pathway but does not affect sorting of plasma membrane proteins in intestinal epithelial cells (Caco-2). *EMBO J.* 9:3163-3170.

Matter, K. and Mellman, I. (1994). Mechanisms of cell polarity: sorting and transport in epithelial cells. *Curr. Opin. Cell Biol.* 6:545-554.

Matter, K., Yamamoto, E.M. and Mellman, I. (1994). Structural requirements and sequence motifs for polarized sorting and endocytosis of LDL and Fc receptors in MDCK cells. *J. Cell Biol.* 126:991-1004.

Mayor, S., Sabharanjak, S. and Maxfield, F.R. (1998). Cholesterol-dependent retention of GPI-anchored proteins in endosomes. *EMBO J.* 17:4626-4638.

McIntyre, J.C. and Sleight, R.G. (1991). Fluorescence assay for phospholipid membrane asymmetry. *Biochemistry* 30:11819-11827.

Mellman, I. (1996). Membranes and sorting. *Curr. Opin. Cell Biol.* 8:497-498.

Merisko, E.M. (1985). Evidence for the interaction of alpha-actinin and calmodulin with the clathrin heavy chain. *Eur. J. Cell Biol.* 39:167-172.

Merisko, E.M., Welch, J.K., Chen, T.Y., Chen, M. (1988). Alpha-actinin and calmodulin interact with distinct sites on the arms of the clathrin trimer. *J. Biol. Chem.* 263:15705-15712.

Misek, D.E., E. Bard and E. Rodriguez-Boulan. (1984). Biogenesis of epithelial cell polarity: intracellular sorting and vectorial exocytosis of an apical plasma membrane glycoprotein. *Cell* 39:537-546.

- Mostov, K.E. and Cardone, M.H.** (1995). Regulation of protein traffic in polarized epithelial cells. *Bioessays* 17:129-138.
- Mostov, K.E. and D.L. Deitcher.** (1986). Polymeric IgA receptor expressed in MDCK cells transcytoses IgA. *Cell* 46:613-621.
- Mostov, K.E., Y. Altschuler, S. Chapin, C. Enrich, S-H. Low, F. Luton, J. Richman-Eisenstat, K. Singer, K. Tang and T. Weimbs.** (1995). Regulation of protein traffic in polarized epithelial cells: the polymeric immunoglobulin receptor model. *Cold Spring Harbor Symp. Quant. Biol.* 60:775-781.
- Mukherjee, S., Ghosh, R.N. and Maxfield, F.R.** (1997). Endocytosis. *Physiol. Rev.* 77:759-803.
- Muniz, M., Martin, M.E., Hidalgo, J. and Velasco, A.** (1997). Protein kinase A activity is required for the budding of constitutive transport vesicles from the trans-Golgi network. *Proc. Natl. Acad. Sci. U. S. A.* 94:14461-14466.
- Nichols, G.E., Shiraishi, T., Allietta, M., Tillack, T.W. and Young, W.W. Jr.** (1987). Polarity of the Forssman glycolipid in MDCK epithelial cells. *Biochim. Biophys. Acta.* 930:154-166.
- Novick, P. and Zerial, M.** (1997). The diversity of Rab proteins in vesicle transport. *Curr. Opin. Cell Biol.* 9:496-504.
- Odorizzi, G. Pearse, A. Domingo, D., Trowbridge, I.S. and Hopkins, C.R.** (1996). Apical and basolateral endosomes of MDCK cells are interconnected and contain a polarized sorting mechanism. *J. Cell Biol.* 135:139-152.
- Odorizzi, G. and Trowbridge, I.S.** (1997). Structural requirements for basolateral sorting of the human transferrin receptor in the biosynthetic and endocytic pathways of Madin-Darby canine kidney cells. *J. Cell Biol.* 137:1255-1264.
- Okamoto, C.T., Karam, S.M., Jeng, Y.Y, Forte, J.G. and Goldenring, J.R.** (1998). Identification of clathrin and clathrin adaptors on tubulovesicles of gastric acid secretory (oxyntic) cells. *Am. J. Physiol.* 274:C1017-C1029.
- Okamoto, C.T. and Jeng, Y.Y.** (1998b). An immunologically distinct β -adaptin on tubulovesicles of gastric oxyntic cells. *Am. J. Physiol.* 275:C1323-C1329.
- Orzech, E., Schlessinger, K., Weiss, A., Okamoto, C.T. and Aroeti, B.** (1999). Interactions of the AP-1 Golgi adaptor with the polymeric immunoglobulin receptor and their possible role in mediating brefeldin A-sensitive basolateral targeting from the trans-Golgi network. *J. Biol. Chem.* 274:2201-2215.
- Pagano, R.E. and Sleight, R.G.** (1985). Defining lipid transport pathways in animal cells. *Science* 229:1051-1057.
- Parton, R.G., K. Prydz, M. Bomsel, K. Simons and G. Griffiths.** (1989). Meeting of the apical and basolateral endocytic pathways of the Madin-Darby canine kidney cell in late endosomes. *J. Cell Biol.* 109:3259-3272.
- Pearse, B.M. and Robinson, M.S.** (1990). Clathrin, adaptors, and sorting. *Annu. Rev. Cell Biol.* 6:151-171.
- Pelham, H.R.** (1996). The dynamic organisation of the secretory pathway. *Cell Struct. Funct.* 21, 413-419.
- Pfeiffer, S, S.C. Fuller and K. Simons.** (1985). Intracellular sorting and basolateral appearance of the G protein of vesicular stomatitis virus in Madin-Darby canine kidney cells. *J. Cell Biol.* 101:470-476.
- Pimplikar, S.W. and Simons, K.** (1993). Role of heterotrimeric G proteins in polarized membrane transport. *J. Cell Sci. Suppl.* 17P:27-32.
- Pimplikar, S.W. and Simons, K.** (1994). Activators of protein kinase A simulate apical but not basolateral transport in epithelial Madin-Darby canine kidney cells. *J. Biol. Chem.* 269:19054-19059.
- Pomorski, T., Herrmann, A., Zimmermann, B., Zachowski, A. and Mueller, P.** (1995). An improved

assay for measuring the transverse redistribution of fluorescent phospholipids in plasma membranes. *Chem. Phys. Lip.* 77:139-146.

Presley, J.F., Mayor, S., Dunn, K.W., Johnson, L.S., McGraw, T.E. and Maxfield, F.R. (1993). The End2 mutation in CHO cells slows the exit of transferrin receptors from the recycling compartment but bulk membrane recycling is unaffected. *J. Cell Biol.* 122:1231-1241.

Putz, U. and Schwarzmann, G. (1995). Golgi staining by two fluorescent ceramide analogues in cultured fibroblasts requires metabolism. *Eur. J. Cell Biol.* 68:113-121.

Radhakrishna, H. and Donaldson, J.G. (1997). ADP-ribosylation factor 6 regulates a novel plasma membrane recycling pathway. *J. Cell Biol.* 139:49-61.

Rahkila, P., Luukela, V., Vaananen, K. and Metsikko, K. (1998). Differential targeting of vesicular stomatitis virus G protein and influenza virus hemagglutinin appears during myogenesis of L6 muscle cells. *J. Cell Biol.* 140:1101-1111.

Rapoport, I., Miyazaki, M., Boll, W., Duckworth, B., Cantley, L.C., Shoelson, S. and Kirchhausen, T. (1997). Regulatory interactions in the recognition of endocytic sorting signals by AP-2 complexes. *EMBO J.* 16:2240-2250.

Ren, M., Xu, G., Zeng, J., De Lemos-Chiarandini, C., Adesnik, M. and Sabatini, D.D. (1998). Hydrolysis of GTP on rab11 is required for the direct delivery of transferrin from the pericentriolar recycling compartment to the cell surface but not from sorting endosomes. *Proc. Natl. Acad. Sci. U. S. A.* 95:6187-6192.

Roelofsen, H., Soroka, C.J., Keppler, D. and Boyer, J.L. (1998). Cyclic AMP stimulates sorting of the canalicular organic anion transporter (Mrp2/cMoat) to the apical domain in hepatocyte couplets. *J. Cell Sci.* 111:1137-1145.

Saha, C., Nigam, S.K. and Denker, B.M. (1998). Involvement of Galphai2 in the maintenance and biogenesis of epithelial cell tight junctions. *J. Biol. Chem.* 273, 21629-21633.

Salisbury, J.L., Condeelis, J.S. and Satir, P. (1980). Role of coated vesicles, microfilaments, and calmodulin in receptor-mediated endocytosis by cultured B lymphoblastoid cells. *J. Cell Biol.* 87:132-141.

Sariola, M., Saraste, J. and Kuismanen, E. (1995). Communication of post-Golgi elements with early endocytic pathway: regulation of endoproteolytic cleavage of Semliki Forest virus p62 precursor. *J. Cell Sci.* 108:2465-2475.

Scheiffele, P., Verkade, P., Fra, A.M., Virta, H., Simons, K. and Ikonen, E. (1998). Caveolin-1 and -2 in the exocytic pathway of MDCK cells. *J. Cell Biol.* 140:795-806.

Schell, M.J., Maurice, M., Stieger, B. and Hubbard, A.L. (1992). 5'nucleotidase is sorted to the apical domain of hepatocytes via an indirect route. *J. Cell Biol.* 119:1173-1182.
29:10865-10871.

Schroeder, R., London, E. and Brown, D. (1994). Interactions between saturated acyl chains confer detergent resistance on lipids and glycosylphosphatidylinositol (GPI)-anchored proteins: GPI-anchored proteins in liposomes and cells show similar behavior. *Proc. Natl. Acad. Sci. U. S. A.* 91:12130-12134.

Schwartz, G.J., Barasch, J. and Al-Awqati, Q. (1985). Plasticity of functional epithelial polarity. *Nature* 318:368-371.

Song, J., Khachikian, Z., Radhakrishna, H. and Donaldson, J.G. (1998). Localization of endogenous ARF6 to sites of cortical actin rearrangement and involvement of ARF6 in cell spreading. *J. Cell Sci.* 111:2257-2267.

Sormunen, R., Eskelinen, S. and Lehto, V. (1993). Bile canaliculus formation in cultured HepG2 cells. *Lab. Invest.* 68:652-662.

Simons, K. and S.D. Fuller. (1985). Cell surface polarity in epithelia. *Annu. Rev. Cell Biol.* 1:243-288.

Simons, K. and G. van Meer. (1988). Lipid sorting in epithelial cells. *Biochemistry* 27:6197-6202.

Simons, K. and A. Wandinger-Ness. (1990). Polarized sorting in epithelia. *Cell* 62:207-210.

- Simons, K. and E. Ikonen.** (1997) Functional rafts in cell membranes. *Nature* 387:569-572.
- Spang, A., Matsuoka, K., Hamamoto, S., Schekman, R. and Orci, L.** (1998). Coatamer, Arf1p, and nucleotide are required to bud coat protein complex I-coated vesicles from large synthetic liposomes. *Proc. Natl. Acad. Sci. U. S. A.* 95:11199-11204.
- Stephens, D.J. and Banting, G.** (1998). Specificity of interaction between adaptor-complex medium chains and the tyrosine-based sorting motifs of TGN38 and Igp120. *Biochem. J.* 335:567-572.
- Takei, K., Haucke, V., Slepnev, V., Farsad, K., Salazar, M., Chen, H. and De Camilli, P.** (1998). Generation of coated intermediates of clathrin-mediated endocytosis on protein-free liposomes. *Cell* 94:131-141.
- Teter, K., Chandy, G., Quinones, B., Pereyra, K., Machen, T. and Moore, H.P.** (1998). Cellubrevin-targeted fluorescence uncovers heterogeneity in the recycling endosomes. *J. Biol. Chem.* 273:19625-19633.
- Tomana, M., R. Kulhavy and J. Mestecky.** (1988). Receptor-mediated binding and uptake of immunoglobulin A by human liver. *Gastroenterology* 94:762-770.
- Toomre, D., Keller, P., White, J., Olivo, J.C. and Simons, K.** (1998). Dual-color visualization of trans-golgi network to plasma membrane traffic along microtubules in living cells. *J. Cell Sci.* 112:21-33.
- Traub, L.M. and Kornfeld, S.** (1997). The trans-Golgi network: a late secretory sorting station. *Curr. Opin. Cell Biol.* 9:527-533.
- Trinchera, M., Carrettoni, D. and Ghidoni, R.** (1991). A part of glucosylceramide formed from exogenous lactosylceramide is not degraded to ceramide but re-cycled and glycosylated in the Golgi apparatus. *J. Biol. Chem.* 266:9093-9099.
- Turner, J.R. and A.M. Tartakoff.** (1989). The response of the Golgi complex to microtubule alterations: the roles of metabolic energy and membrane traffic in Golgi complex organization. *J. Cell Biol.* 109:2081-2088.
- Ullrich, O., Reinsch, S., Urbe, S., Zerial, M. and Parton, R.G.** (1996). Rab11 regulates recycling through the pericentriolar recycling endosome. *J. Cell Biol.* 135:913-924.
- Valtorta, F. and Meldolesi, J.** (1994). The presynaptic compartment: signals and targets. *Semin. Cell Biol.* 5, 211-219.
- van Adelsberg, J., Edwards, J.C., Takito, J., Kiss, B. and al-Awqati, Q.** (1994). An induced extracellular matrix protein reverses the polarity of band 3 in intercalated epithelial cells. *Cell* 76, 1053-1061.
- van den Berg, C.W., Cinek, T., Hallett, M.B., Horejsi, V. and Morgan, B.P.** (1995). Exogenous glycosyl phosphatidylinositol-anchored CD59 associates with kinases in membrane clusters on U937 cells and becomes Ca(2+)-signaling competent. *J. Cell Biol.* 131:669-677.
- van der Woude I, A. Wagenaar, A.A. Meekel, M.B.A. ter Beest, M.H. Ruiters, J.B. Engberts and D. Hoekstra.** (1997). Novel pyridinium surfactants for efficient, nontoxic in vitro gene delivery. *Proc. Natl. Acad. Sci. U. S. A.* 94:1160-1165.
- van der Bijl, P., Lopes-Cardozo, M. and van Meer, G.** (1996). Sorting of newly synthesized galactosylphingolipids to the two surface domains of epithelial cells. *J. Cell Biol.* 132:813-821.
- Van Genderen, I. and van Meer, G.** (1995). Differential targeting of glucosylceramide and galactosylceramide analogues after synthesis but not during transcytosis in Madin-Darby canine kidney cells. *J. Cell Biol.* 131:645-654.
- van Helvoort, A., A.J. Smit, H. Sprong, I. Fritzsche, A.H. Schinkel, P. Borst and G. van Meer.** (1996). MDR1 P-Glycoprotein is a lipid translocase of broad specificity, while MDR3 P-Glycoprotein specifically translocates phosphatidylcholine. *Cell* 87:507-517.
- van IJzendoorn, S.C.D. and Hoekstra, D.** (1998). (Glyco)sphingolipids are sorted in sub-apical compartments in HepG2 cells: A role for non-Golgi-related intracellular sites in the polarized distribution of (Glyco)sphingolipids. *J. Cell Biol.* 142:683-696
- van IJzendoorn, S.C.D. and Hoekstra, D.** (1999). The subapical compartment: a novel sorting center? *Trends Cell Biol.* 9:144-149.

- van IJzendoorn, S.C.D., M.M.P. Zegers, J.W. Kok and D. Hoekstra.** (1997). Segregation of glucosylceramide and sphingomyelin occurs in the apical to basolateral transcytotic route in HepG2 cells. *J. Cell Biol.* 137:347-357.
- Van Meer, G.** (1993). Transport and sorting of membrane lipids. *Curr. Opin. Cell Biol.* 5:661-673.
- Van Meer, G. and Simons, K.** (1988). Lipid polarity and sorting in epithelial cells. *J. Cell Biochem.* 36:51-58.
- Van Meer, G., Stelzer, E.H.K., Wijnaendts-van-Resandt, R.W. and Simons, K.** (1987). Sorting of sphingolipids in epithelial (Madin-Darby canine kidney cells. *J. Cell Biol.* 105:1623-1635.
- Van 't Hof, W. and Van Meer, G.** (1990). Generation of lipid polarity in intestinal epithelial (Caco-2) cells: sphingolipid synthesis in the Golgi complex and sorting before vesicular traffic to the plasma membrane. *J. Cell Biol.* 111:977-986.
- Wang, L.H., Rothberg, K.G. and Anderson, R.G.** (1993). Mis-assembly of clathrin lattices on endosomes reveals a regulatory switch for coated pit formation. *J. Cell Biol.* 123:1107-1117.
- Wei, M.L., Bonzelius, F., Scully, R.M., Kelly, R.B. and Herman, G.A.** (1998). GLUT4 and transferrin receptor are differentially sorted along the endocytic pathway in CHO cells. *J. Cell Biol.* 140:565-575.
- Weimbs, T., S.H. Low, S.J. Chapin and K.E. Mostov.** (1997). Apical targeting in polarized epithelial cells: there's more afloat than rafts. *Trends Cell Biology* 7:393-399.
- West, A.E., Neve, R.L. and Buckley, K.M.** (1997). Identification of a somatodendritic targeting signal in the cytoplasmic domain of the transferrin receptor. *J. Neurosci.* 17, 6038-6047.
- Whitney, J.A., Gomez, M., Sheff, D., Kreis, T.E. and Mellman, I.** (1995). Cytoplasmic coat proteins involved in endosome function. *Cell* 83:703-713.
- Wilson, W.M. and Colton, T.L.** (1997). Targeting of an intestinal apical endosomal protein to endosomes in nonpolarized cells. *J. Cell Biol.* 136:319-330.
- Zaal, K.J.M., Kok, J., Sormunen, R., Eskelinen, S. and Hoekstra, D.** (1993). Intracellular sites involved in the biogenesis of bile canaliculi in hepatic cells. *Eur. J. Cell Biol.* 63:10-19.
- Zaal, K.J.M., Kok, J.W., Kuipers, F. and Hoekstra, D.** (1994). Lipid trafficking in hepatocytes: relevance to biliary lipid secretion. *Adv. in Molec. and Cell Biol.* 8:133-150.
- Yoshimori, T., Keller, P., Roth, M.G. and Simons, K.** (1996). Different biosynthetic transport routes to the plasma membrane in BHK and CHO cells. *J. Cell Biol.* 133:247-256.
- Yeaman, C., Grindstaff, K.K. and Nelson, W.J.** (1999). New perspectives on mechanisms involved in generating epithelial cell polarity. *Physiol. Rev.* 79:73-98.
- Zacchi, P., Stenmark, H., Parton, R.G., Orioli, D., Lim, F., Giner, A., Mellman, I., Zerial, M and C. Murphy.** (1998). Rab17 regulates membrane traffic through apical recycling endosomes in polarized epithelial cells. *J. Cell Biol.* 140:1039-1053.
- Zegers, M.M.P. and D. Hoekstra.** (1997). Sphingolipid transport to the apical plasma membrane in human hepatoma cells is controlled by PKC and PKA activity: a correlation with cell polarity in HepG2 cells. *J. Cell Biol.* 138:307-321.
- Zegers, M.M.P., Zaal, K.J.M., van IJzendoorn, S.C.D., Klappe, K. and Hoekstra, D.** (1998). Actin filaments and microtubules are involved in different membrane traffic pathways that transport sphingolipids to the apical surface of polarized HepG2 cells. *Mol. Biol. Cell* 9:1939-1949.
- Zegers, M.M.P. and Hoekstra, D.** (1998). Mechanisms and functional features of polarized membrane traffic in epithelial and hepatic cells. *Biochem. J.* 336:257-269.
- Zheng, X., Lu, D. and Sadler, J.E.** (1999). Apical sorting of bovine enteropeptidase does not involve detergent-resistant association with sphingolipid-cholesterol rafts. *J. Biol. Chem.* 274:1596-1605.
- Zurzolo, C., Le Bivic, A., Quaroni, A., Nitsch, L. and Rodriguez-Boulan, E.** (1992). Modulation of transcytotic and direct targeting pathways in a polarized thyroid cell line. *EMBO J.* 11, 2337-2344.

Zurzolo, C., van't Hof, W., van Meer, G. and Rodriguez-Boulan, E. (1994). VIP21/caveolin, glycosphingolipid clusters and the sorting of glycosylphosphatidylinositol-anchored proteins in epithelial cells. *EMBO J.* 13:42-53.

Zurzolo, C., Lisanti, M.P., Caras, I.W., Nitsch, L. and Rodriguez-Boulan, E. (1993). Glycosylphosphatidylinositol-anchored proteins are preferentially targeted to the basolateral surface in Fischer rat thyroid epithelial cells. *J. Cell Biol.* 121:1031-1039.