

University of Groningen

Studies on the bacterial protein conducting pore

Does, Christiaan van der

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2000

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Does, C. V. D. (2000). *Studies on the bacterial protein conducting pore: the SecYEG complex*. [S.n.].

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter 10

List of publications

- Manting, E.H., **Van der Does, C.**, Remigy, H., Engel, A., and Driessen, A.J.M. (2000) SecYEG assembles into a tetramer to form the active protein translocation channel. *EMBO J.* *19*, 852-861.
- Scotti, P.A., Urbanus, M.L., Brunner, J., de Gier, J.W., von Heijne, G., **Van der Does, C.**, Driessen, A.J.M., Oudega, B., and Luirink, J. (2000) YidC, the *Escherichia coli* homologue of mitochondrial Oxa1p, is a component of the Sec translocase. *EMBO J.* *19*, 542-549.
- Van der Does, C.**, Swaving, J., van Klompenburg, W., and Driessen, A.J.M. (2000) Non-bilayer lipids stimulate the activity of the reconstituted bacterial protein translocase. *J.Biol.Chem.* *275*, 2472-2478.
- Den Blaauwen, T., van der Wolk, J.P.W., **Van der Does, C.**, van Wely, K.H.M., and Driessen, A.J.M. (1999). Thermodynamics of nucleotide binding to NBS-I of the *Bacillus subtilis* preprotein translocase subunit SecA. *FEBS Lett.* *458*, 145-150.
- Kaufmann, A., Manting, E.H., Veenendaal, A.K.J., Driessen, A.J.M., and **Van der Does, C.** (1999). Cysteine-directed cross-linking demonstrates that helix 3 of SecE is close to helix 2 of SecY and helix 3 of a neighboring secE. *Biochemistry* *38*, 9115-9125.
- Manting, E.H., Kaufmann, A., **Van der Does, C.**, and Driessen, A.J.M. (1999). A Single Amino Acid Substitution in SecY Stabilizes the Interaction with SecA. *J.Biol.Chem.* *274*, 23868-23874.
- Van de Vossenbergh, J.L., Albers, S.-V., **Van der Does, C.**, Driessen, A.J.M., Van Klompenburg, W. (1998) The positive inside rule is not determined by the polarity of the delta psi. *Mol Microbiol.* *29*(4):1125-7.
- Van der Does, C.**, Manting, E.H., Kaufmann, A., Lutz, M., and Driessen, A.J.M. (1998). Interaction between SecA and SecYEG in micellar solution and formation of the membrane-inserted state. *Biochemistry* *37*, 201-210.
- van Voorst, F., **Van der Does, C.**, Brunner, J., Driessen, A.J.M., and de Kruijff, B. (1998). Translocase-bound SecA is largely shielded from the phospholipid acyl chains. *Biochemistry* *37*, 12261-12268.
- Fekkes, P., **Van der Does, C.**, and Driessen, A.J.M. (1997). The molecular chaperone SecB is released from the carboxy- terminus of SecA during initiation of precursor protein translocation. *EMBO J.* *16*, 6105-6113.
- Manting, E.H., **Van der Does, C.**, and Driessen, A.J.M. (1997). In vivo cross-linking of the SecA and SecY subunits of the *Escherichia coli* preprotein translocase. *J.Bacteriol.* *179*, 5699-5704.
- Den Blaauwen, T., de Wit, J.G., Gosker, H., **Van der Does, C.**, Breukink, E.J., de Leij, L., and Driessen, A.J.M. (1997). Inhibition of preprotein translocation and reversion of the membrane inserted state of SecA by a carboxyl terminus binding mAb. *Biochemistry* *36*, 9159-9168.

- Van der Does, C.**, den Blaauwen, T., de Wit, J.G., Manting, E.H., Groot, N.A., Fekkes, P., and Driessen, A.J.M. (1996). SecA is an intrinsic subunit of the Escherichia coli preprotein translocase and exposes its carboxyl terminus to the periplasm. *Mol.Microbiol.* *22*, 619-629.
- Poolman, B., Knol, J., **Van der Does, C.**, Henderson, P.J., Liang, W.J., Leblanc, G., Pourcher, T., and Mus-Veteau, I. (1996). Cation and sugar selectivity determinants in a novel family of transport proteins. *Mol.Microbiol.* *19*, 911-922.
- Driessen, A.J.M., de Wit, J.G., Kuiper, W., Van der Wolk, J.P., Fekkes, P., **Van der Does, C.**, van Wely, K.H.M., Manting, E.H., and Den Blaauwen, T. (1995). SecA, a novel ATPase that converts chemical energy into a mechanical force to drive precursor protein translocation. *Biochem.Soc.Trans.* *23*, 981-985.
- Balandin, T., **Van der Does, C.**, Albert, J.M., Bol, J.F., and Linthorst, H.J. (1995). Structure and induction pattern of a novel proteinase inhibitor class II gene of tobacco. *Plant Mol.Biol.* *27*, 1197-1204.
- Linthorst, H.J., **Van der Does, C.**, Brederode, F.T., and Bol, J.F. (1993). Circadian expression and induction by wounding of tobacco genes for cysteine proteinase. *Plant Mol.Biol.* *21*, 685-694.
- Linthorst, H.J., **Van der Does, C.**, van Kan, J.A., and Bol, J.F. (1993). Nucleotide sequence of a cDNA clone encoding tomato (*Lycopersicon esculentum*) cysteine proteinase. *Plant Physiol.* *101*, 705-706.
- Linthorst, H.J., Brederode, F.T., **Van der Does, C.**, and Bol, J.F. (1993). Tobacco proteinase inhibitor I genes are locally, but not systemically induced by stress. *Plant Mol.Biol.* *21*, 985-992.

