

University of Groningen

## The development of depression in children and adolescents with ADHD

Roy, Arunima

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2016

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Roy, A. (2016). *The development of depression in children and adolescents with ADHD*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## **Peer dislike and victimisation in pathways from ADHD symptoms to depression**

*Roy, A., Hartman, C.A.H., Veenstra, R., Oldehinkel, A.J.*

Based on:

Roy, A., Hartman, C. A., Veenstra, R., & Oldehinkel, A. J. (2015). Peer dislike and victimisation in pathways from ADHD symptoms to depression. *European child & adolescent psychiatry*, 24(8), 887-895.

ABSTRACT

Background: The following hypotheses were tested in a longitudinal, population-based study: 1) ADHD symptoms are associated with peer dislike and victimisation; 2) Peer dislike and victimisation increase the risk for subsequent depression; and 3) The effect of ADHD symptoms on depression is partly mediated through peer dislike and victimisation. Gender differences in mediating pathways through peer dislike and victimisation to depression were additionally explored.

Method: The Child Behaviour Checklist (CBCL), Youth Self Report (YSR) and Teacher's Checklist of Pathology (TCP) assessed ADHD symptoms in 728 adolescents. Peer nominations were used to assess peer dislike and victimisation. The Composite International Diagnostic Interview (CIDI) was used to assess depression. Effects of peer dislike, victimisation, and ADHD symptoms on depression were modelled using Cox regression.

Results: ADHD symptoms were associated with peer dislike ( $rs=.17, p<.001$ ) and victimisation ( $rs=.11, p=.001$ ). Dislike, victimisation, and ADHD symptoms increased risk for depression. Risk for depression associated with victimisation and ADHD symptoms reduced with time. Dislike and victimisation mediated 7% of the effect of ADHD symptoms on depression. Pathways through dislike and victimisation were present in girls but not in boys.

Conclusion: Peer dislike and victimisation explain, to a limited extent, the prospective association between ADHD and depression, particularly in girls.

Keywords: Attention Deficit Hyperactivity Disorder; Depression; Peer Dislike; Victimization; Prospective Study

## INTRODUCTION

Attention deficit hyperactivity disorder (ADHD), a common neurodevelopmental disorder (American Psychiatric Association, 2013), is often comorbid with depression (Connor et al., 2003). Depression not only leads to morbidity by itself, but when associated with ADHD also increases the severity and duration of ADHD and worsens prognosis (Jensen, Shervette, Xenakis, & Richters, 1993). Consequently, it is of importance to prevent the development of depression in ADHD. A better understanding of the comorbid association of ADHD and depression is required as a first step in order to develop preventive measures and interventions. Since ADHD manifests at an early age (Kessler et al., 2007), while depressive disorders typically have an onset in adolescence and young adulthood (Oldehinkel, Wittchen, & Schuster, 1999), it could be said that the onset of depression generally follows the onset of ADHD. Depression may arise in ADHD either directly or through the development of other, intermediate problems. Being disliked and being a victim of bullying by peers may be two such intermediate problems.

Adolescents with ADHD have problems in regulating their behaviours, and exhibiting behaviours deemed aberrant by peers could lead to being disliked (Diamantopoulou, Henricsson, & Rydell, 2005). Impairments in the development of age-appropriate social skills in ADHD can further compound the peer dislike (Murray-Close et al., 2010). Being disliked may in turn spiral into further difficulties in forming and maintaining relationships with peers (Oldehinkel, Rosmalen, Veenstra, Dijkstra, & Ormel, 2007). The resulting social disconnection can increase feelings of stress, anxiety, and loneliness, setting the stage for development of depression (Lansford et al., 2007; Oldehinkel et al., 2007).

Due to difficulties in controlling behaviours, adolescents with ADHD often respond aggressively to bullying, making them not only easy but also attractive victims for bullies (Perry, Kusel, & Perry, 1988). Victimisation involves infliction of purposeful hurtful actions by bullies, which ostracize and demoralize the victims (Olweus, 1994). Being victimised may intensify the spiral of isolation and rejection (Dake, Price, & Telljohann, 2003; Rigby & Slee, 1997), providing an increased risk for depression.

The above-mentioned influences of dislike and victimisation on the risk for future depression may act differently in adolescent boys and girls with ADHD. Boys with ADHD face greater social impairment at school (de Boo & Prins, 2007), which in turn may make them more likely to be disliked or victimised. Girls, meanwhile, tend to be more sensitive to peers' opinions,

and thus may be affected more by peer dislike and victimisation (Bakker, Ormel, Verhulst, & Oldehinkel, 2010; Oldehinkel et al., 2007; Oldehinkel, Verhulst, & Ormel, 2011). The net outcome of these contrasting predispositions might lead to a roughly similar influence of dislike and victimisation on pathways to depression in the two genders. On the other hand, the higher likelihood of social impairment in boys and the greater sensitivity to peer problems in girls might as well produce unexpected differences in these pathways between boys and girls.

Existing studies on the influences of peer problems in the ADHD-depression relationship have shown mixed results. In a short-term prospective study, Mikami and Hinshaw (2003) found peer nominated rejection measures to be associated with depressed-anxious behaviour in girls diagnosed with ADHD (Mikami & Hinshaw, 2003). Humphreys et al. (2013) in a population-based cohort of individuals followed-up from birth to 20 years of age, found that teacher- and self-reported peer rejection and popularity mediated the pathways from ADHD symptoms to depressive symptoms (Humphreys et al., 2013). In another retrospective study, Humphrey, Storch, and Geffken (2007), found a correlation between victimisation and depression in children diagnosed with ADHD (Humphrey, Storch, & Geffken, 2007). Finally, Mrug et al. (2012) found no evidence to suggest that peer rejection specifically predicted depression in boys and girls diagnosed with ADHD (Mrug et al., 2012).

Previously, we demonstrated that not only a full diagnosis of ADHD, but subthreshold ADHD too increased the risk for depression (Roy, Oldehinkel, Verhulst, Ormel, & Hartman, 2014). Relatively mild symptoms of ADHD in adolescents may therefore be sufficient to increase the risk for being disliked by peers or being victimised, which in turn could increase the risk for depression. The present study was set up to better understand pathways from ADHD symptoms to depression through peer dislike and victimisation. Using a prospective general population sample of adolescents, we tested the hypotheses that: 1) ADHD symptoms are associated with peer dislike and victimisation; 2) Peer dislike and victimisation increase the risk for subsequent depression; and 3) The effect of ADHD symptoms on depression is partly mediated through peer dislike and victimisation. In addition, we explored gender differences in mediating pathways through peer dislike and victimisation to depression.

## METHODS

The data were collected as part of the TRacking Adolescents' Individual Lives Survey (TRAILS), a Dutch prospective cohort study focusing on psychosocial development and mental health of adolescents from the general population. TRAILS involves bi- or triennial measurements from age 11 to at least age 25 (de Winter et al., 2005; Huisman et al., 2008). The methodology has been described in detail elsewhere (Ormel et al., 2012) and is summarized briefly below. The study was approved by the Dutch Central Committee on Research Involving Human Subjects (CCMO). Participants were treated in accordance with the Declaration of Helsinki, and all measurements were carried out with their adequate understanding and written consent.

### **Cohort**

Children were recruited from five municipalities in the north of The Netherlands, including both urban and rural areas. Primary school participation was a requisite for inclusion. Of the 2935 children who met these criteria, 2230 (76.0%) provided informed consent from both parent and child to participate in the study. Four assessment waves have been completed to date.

The present study used data from the second (T2) and fourth wave (T4) which ran from September 2003 to December 2004 (T2), and October 2008 to September 2010 (T4) respectively. The response rate at T2 was 96.4% (N=2149, mean age 13.6, SD=0.53, 51.0% girls) of whom 47% (N= 1007) completed the peer assessments. The response rate at T4 was 83.4% (N=1881, mean age 19.1, SD=0.60, 52.3% girls), of whom 84.2% (N=1584) completed the below described diagnostic interview. Adolescents without peer nomination data at T2 (N=827) were excluded. Adolescents who had developed depression before T2 (N=29) were also excluded, to allow only participants who either did not develop depression, or developed depression between T2 and T4. These led to a total of 728 participants (54% girls) being included in this study. Figure 5.1 presents a flowchart of participants included at each wave. The study was approved by the Dutch Central Committee on Research Involving Human Subjects (CCMO). Participants were treated in accordance with the Declaration of Helsinki, and all measurements were carried out with their adequate understanding and written consent.

### **Measures**

ADHD symptoms were measured using the Child Behaviour Checklist (CBCL) filled out by parents, the Youth Self Report (YSR) filled out by the children, and the Teacher's Checklist of Psychopathology (TCP) filled out by teachers in the second measurement wave (T2) (Achenbach, 1991b; Achenbach, 1991c). The CBCL and YSR are internationally validated questionnaires for child emotional and behavioural problems at ages 4-18 (Achenbach, Dumenci, & Rescorla, 2003). We used the attention problem scale, which consists of ten items in the CBCL ( $M = .32$ ,  $SD = .30$ ) and nine items in the YSR ( $M = .55$ ,  $SD = .33$ ). The TCP was developed by the TRAILS team and is a shorter version of the Teacher's Report Form (TRF) (Achenbach, 1991c). The TCP requires a shorter time than the TRF to be completed and therefore is easier to use in classrooms with many pupils. It contains nine descriptions of behaviours with response options for these behaviours ranging from 1 = 'not applicable' to 5 = 'very clearly applicable'. Information from teacher-reported attention and hyperactivity/impulsivity problems (corresponding to the CBCL-YSR attention problems scales) were used in this study ( $M = -.001$ ,  $SD = .91$ ). Multi-informant scores have been reported to give better estimates for ADHD than single-informant scores (Loeber, Green, Lahey, & Stouthamer-Loeber, 1989; Tripp, Schaughency, & Clarke, 2006). We used mean ADHD symptom scores, calculated from standardized parent, adolescent and teacher ratings ( $M = .005$ ,  $SD = .14$ ).

Information on peer dislike and victimisation was obtained through peer nominations at the second measurement wave (T2) (Kupersmidt & Coie, 1990). Peer nominations were collected in classes with at least 3 TRAILS participants. This led to the inclusion of 3798 adolescents (54.1% girls) from 150 classes in 34 schools, of whom 1078 were TRAILS participants (Dijkstra, Cillessen, Lindenberg, & Veenstra, 2010). Participants received a list of all classmates and were asked to indicate, among other things, whom in their classroom they disliked and who they bullied. The number of nominations participants could make ranged from none to all classmates. All participants completed the peer nominations independently. Variables used in this study reflect the percentage of classmates nominating the TRAILS participants as being disliked ( $M = 11$ ,  $SD = 0.13$ ) and victimised ( $M = 2$ ,  $SD = 0.05$ ). Percentage nominations were used to offset differences in the number of participants per classroom.

During the fourth assessment wave, psychiatric disorders were assessed by means of the World Health Organization Composite International Diagnostic Interview (CIDI), version 3.0

(N=1584). The CIDI is a structured diagnostic interview, which yields lifetime and current diagnoses according to the definitions and criteria of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV). The CIDI has been used in a large number of surveys worldwide (Kessler et al., 2004), and has shown to have a good concordance with clinical diagnoses (Haro et al., 2006; Kessler et al., 2009). In addition to the occurrence of psychiatric disorders, the CIDI yields their age of onset. In the present study data on Major Depressive Episode (MDE) was used and depression was operationalized as MDE, either with or without (hypo)manic symptoms. Age of onset refers to the age depression emerged for the first time.

Participants with and without completed CIDI interviews differed significantly in terms of dislike ( $t = 2.8, p = .005$ ), victimisation ( $t = 3.2, p = .002$ ) and ADHD symptom ( $t = 4.1, p < .001$ ) scores at T2. ADHD symptoms scores differed significantly between (excluded) participants with an MDE prior to T2 ( $N = 29, M = .63, SD = 1.2$ ) and included participants ( $N = 728, M = -.02, SD = 1.0$ ) ( $t = -3.5, p = .001$ ). The two groups had comparable dislike ( $t = -1.6, = .11$ ) and victimisation ( $t = .40, p = .69$ ) scores.

### **Statistical analysis**

Spearman rank correlations were used to estimate associations between ADHD symptoms, and peer dislike and victimisation. ADHD symptom, peer dislike and victimisation variables were non-normally distributed for which Spearman correlations are better suited. Cox proportional hazards regression was used to estimate the effects of peer dislike and victimisation on the probability to develop an MDE. Next, the effect of ADHD symptoms on the probability to develop an MDE was examined by means of Cox regression both with and without adjusting for dislike and victimisation. The percentage change (reduction) in the hazard ratio (HR) before and after adjusting for dislike and victimisation was computed to assess mediation by these two variables. Lastly, Cox regression models estimating mediation through dislike and victimisation were stratified by gender to explore gender differences.

Continuous variables (i.e., ADHD symptoms, peer dislike, victimisation) were standardised to mean 0 and SD 1 before analysis to ease interpretation of the regression coefficients. The proportional hazards assumption was tested using Schoenfeld residuals. For significant residuals, time-dependent variables were additionally included in the regression



models. Time-dependant variables were computed by multiplying the continuous variables with survival-time. Survival-time was the number of years between assessment of peer data (T2) and onset of depression. Peer dislike and victimisation were included both separately and simultaneously in the above described regression models to assess their individual and combined effects. Except for the gender-stratified analysis, all models were adjusted for gender. Analyses were performed using SPSS v. 20.0 (IBM Corp., Armonk, NY). All tests were two-tailed and a p-value of  $<.05$  was considered statistically significant.

## RESULTS

In the sample, 12% ( $n = 90$ ) of the adolescents eventually developed a depressive episode between T2 and T4, of whom 76% ( $n = 68$ ) were girls. While 66% ( $n = 483$ ) of the adolescents received one or more nominations for being disliked, only 21% ( $n = 149$ ) received at least one nomination for being victimised. Nominations (at least one) for both peer dislike and victimisation were received by 18% ( $n = 130$ ) of the adolescents.

Spearman's rho revealed weak but statistically significant relationships between ADHD symptoms and peer dislike ( $r_s = .17, p < .001$ ), and between ADHD symptoms and victimisation ( $r_s = .11, p = .001$ ).

Table 5.1 presents the effects of peer dislike and victimisation on major depression onset. With each unit rise in the standardized peer dislike score, the risk for depression increased by 1.4 times. Victimization but not peer dislike was found to have a time-varying effect (Schoenfeld residual =  $-0.25; p = .018$ ), and a time-dependent victimisation variable was additionally included in the model. The initial (i.e. at T2) effect of victimisation was stronger, that is, a 1.6 times increased depression risk per unit change in standardized victimisation score which gradually decreased over time and disappeared after about 2 years. When included simultaneously, the effects of dislike and victimisation were roughly similar to their effects on including them separately in the model, suggesting that the effects were non-overlapping.

Table 5.2 presents the effect of ADHD symptoms on major depression onset before and after adjusting for a low peer status and victimisation. ADHD symptoms violated the proportional hazards assumption (Schoenfeld residual =  $-0.30; p = .005$ ), and an additional time-dependent ADHD symptom variable was included in the model. Peer dislike mediated 4%,

while victimisation mediated 3% of the effect of ADHD symptoms on depression. When included simultaneously, the peer variables explained 7% of the effect of ADHD symptoms on depression.

The time-varying effect of ADHD symptoms on the risk for depression, before and after adjusting for dislike and victimisation, is illustrated in Table 5.3. While ADHD symptoms were strongly associated with an increased risk for depression at T2, the estimated risk declined over time and after 6 years, no excess risk remained.

In order to explore gender differences in pathways from ADHD symptoms to depression, gender stratified analyses were carried out. Table 5.4 presents results of these exploratory analyses. In boys, peer dislike and victimisation did not mediate the effect of ADHD symptoms on depression. For girls, peer dislike mediated 7% and victimisation mediated 3% of the effect of ADHD symptoms on depression. Please note that the estimated effect of victimisation on depression in girls was not significant when adjusted for ADHD symptoms.

In a previous study, we showed that disruptive behaviours and anxiety problems mediated the pathways from ADHD to depression (Roy et al., 2014). We carried out post-hoc analyses to assess if effects of peer dislike and victimization on depression were confounded by disruptive behaviours and anxiety problems. Inclusion of these additional covariates in the models did not change the essence of the results (please see supplementary material), suggesting that effects of peer dislike and victimization on depression are largely independent of disruptive behaviour and anxiety problems.

## DISCUSSION

Through this study, we aimed to further our understanding of the paths to depression in adolescents with symptoms of ADHD. We found that ADHD symptoms were associated with peer dislike and victimisation. Peer dislike, victimisation, and symptoms of ADHD were associated with an increased risk for depressive outcomes. Peer problems of dislike and victimisation mediated to a limited extent the paths to depression. Lastly, exploratory gender analyses showed mediation through peer dislike and victimisation only in girls and not in boys with ADHD symptoms.

Studies on clinical samples have shown a high risk for dislike and victimisation in children with ADHD as compared to other classmates (Humphrey et al., 2007; Murray-Close et al., 2010). We found ADHD symptoms to be only weakly associated with dislike and victimisation. These weak associations could be related to the use of a population sample and a continuous measure of ADHD in our study. Moreover, some peers may admire the disruptive and aggressive behaviours of adolescents with ADHD (Glass, Flory, & Hankin, 2012; Sandstrom & Coie, 1999), and peer support from a few classmates can protect against dislike and victimisation (Sainio, Veenstra, Huitsing, & Salmivalli, 2011). Consequently, not all adolescents with ADHD symptoms may be susceptible to face dislike or victimisation, explaining the weak associations of ADHD with these problems.

In accordance with previous literature, we found that peer dislike, victimisation, and ADHD symptoms independently increase the risk for depression (Angold, Costello, & Erkanli, 1999; Bukowski, Laursen, & Hoza, 2010; Kaltiala-Heino, Rimpela, Marttunen, Rimpela, & Rantanen, 1999; Rajendran, O'Neill, & Halperin, 2013). In this study, we found that pathways from ADHD symptoms to depression were mediated by peer dislike and victimisation, but to only a limited extent. The limited mediation may be attributable to positive illusory bias (Hoza, Pelham, Milich, Pillow, & McBride, 1993). Adolescents with ADHD tend to make inaccurate and overtly positive evaluations of their social skills and classroom peer status (Hoza, Murray-Close, Arnold, Hinshaw, & Hechtman, 2010), which, even though erroneous, may protect against negative outcomes such as depression. Alternatively, the use of a population sample and a continuous measure for ADHD symptoms may have contributed to the lack of substantial mediating effects. It is also possible that peer relationship difficulties indeed play a little role in the development of depression in ADHD, and other factors such as comorbid psychiatric illnesses, poor parental relationships, and traumatic life events mediate these paths to a greater extent (Daviss, Diler, & Birmaher, 2009).

The strength of the association between ADHD symptoms and depression decreased during the course of adolescence. In part, this decrease may be due to transitory state effects at the time ADHD symptoms were assessed. Yet another part may be explained by the fact that adolescents who are most vulnerable to the depressogenic effects of ADHD symptoms are likely to develop a depression first, thereby gradually leaving a more and more resilient group

(Hillegers et al., 2004; Surtees & Wainwright, 1999). The reduced risk for depression could also be related to changes in peer perceptions of problem behaviours with age such that, from childhood to adolescence, the symptoms of ADHD become more and more acceptable to peers (Moffitt, 1993; Sandstrom & Coie, 1999). Reduction in social impact of ADHD symptoms with age may consequently reduce the risk for depression.

Exploratory analyses of gender differences showed that dislike and victimisation mediated the paths to depression in girls but not in boys. This finding could be explained on the basis of girls' higher sensitivity to peer opinions (Bakker et al., 2010). In addition, symptoms and behaviours of ADHD in girls are considered gender inappropriate and are less well tolerated by peers than in boys (de Boo & Prins, 2007). This may give rise to more negative appraisals by peers and in turn increase risk for depressive outcomes in girls. Alternatively, the lack of mediation in boys could be related to a lack of power. Since our gender-stratified analyses were conducted in a relatively small sample of boys, our findings have to be corroborated by future research.

The main limitation of our study was that we did not follow up on the peer relationships of the participants. Thus, we could not assess how ADHD symptoms might have affected peer relationships over a period of time, and if this contributed to changes in depressive outcomes. It was also not known whether participants utilised any interventions for their ADHD symptoms. Interventions may lead to cessation of ADHD symptoms, reducing the risk of developing depression. Furthermore, stimulant medications used in the treatment of ADHD may reduce symptoms of depression in ADHD (Gurkan et al., 2010). Treatment can thus alter the occurrence of depression in ADHD and lack of this information may bias interpretations.

The strengths of our study include the use of multi informant ratings for assessment of ADHD, and peer nominations for assessment of peer relationships. We also used a longitudinal, population-based sample, which allowed us to follow up the effects of ADHD on depression in time. Population based samples also have the benefit of providing more accurate estimates of prevalences (Goodman et al., 1997), and a balanced representation of genders compared to clinical and referred samples (Gaub & Carlson, 1997). Lastly, in light of findings suggesting that dimensional approaches depict ADHD more accurately than categorical ones (Hyman, 2010; Levy, Hay, McStephen, Wood, & Waldman, 1997), our study, which used ADHD symptom

scores, may provide a better picture of this mental health problem than studies based on diagnoses.

Peer problems not only prove to be a possible hindrance to development of adolescents with ADHD but also, as we found in our study, increase the risk for depression. Therefore, interventions aimed at improving peer relationships in ADHD may be of importance. However, such interventions have failed in many instances to modify or improve peer problems (McQuade & Hoza, 2008; Nijmeijer et al., 2008). In addition, negative perceptions of peers and teachers may make it difficult for adolescents with ADHD to interact and form friendships (Atkinson, Robinson, & Shute, 1997; Chi & Hinshaw, 2002; Martin, Pescosolido, Olafsdottir, & McLeod, 2007). This points to the need for developing newer modalities of interventions, which focus on development of a sensitive environment in schools and in the community, responsive to the symptoms of ADHD. Peer problems in ADHD are resistant to change, such that adolescents who cease to show symptoms of ADHD may yet continue to face peer problems (Bagwell, Molina, Pelham, & Hoza, 2001). Thus, it is difficult for adolescents to ease into social situations and establish a normal life even with effective treatment and control of ADHD symptoms. In short, interventions aimed at making school and community environments more conducive for adolescents with ADHD may have a better effect in the long term than attempts at 'integrating' these adolescents into non-favourable and incompatible environments. A recent study aimed at increasing peers' social inclusion of children with ADHD has shown promising results in this direction (Mikami et al., 2013).

To conclude, this study elaborates the role of problems in peer relationships as partial mediators of the pathways from ADHD to depression. Though these may not be at work singly and may be complicated by presence of other factors, they provide an insight into the underlying mechanisms leading to depression. Knowledge of the possible precursors to depression in ADHD could help in the development of effective interventions, prevention of depression, and an improvement in the quality of life.

Table 5.1 Cox regression estimates of the effect of peer dislike and victimisation on depression onset controlling for gender

<b>Covariate</b>	<b>B (SE)</b>	<b><i>p</i></b>	<b>Wald <math>\chi^2</math></b>	<b>Hazard Ratio</b>	<b>95% CI</b>
<b>Model 1</b>					
Peer dislike	0.32 (0.09)	0.001	12.10	1.37	1.14 to 1.64
<b>Model 2</b>					
Victimisation	0.47 (0.17)	0.005	7.79	1.60	1.15 to 2.22
Victimisation*time	-0.21 (0.10)	0.033	4.56	0.82	0.68 to 0.98
<b>Model 3</b>					
Peer dislike	0.35 (0.10)	0.001	11.56	1.41	1.16 to 1.73
Victimisation	0.30 (0.19)	0.112	2.53	1.35	0.93 to 1.96
Victimisation*time	-0.22 (0.10)	0.029	4.76	0.80	0.66 to 0.98

Table 5.2 Cox regression estimates of the effect of ADHD symptoms on depression onset before and after adjusting for peer dislike and victimisation and controlling for gender

<b>Covariate</b>	<b>B (SE)</b>	<b><i>p</i></b>	<b>Wald <math>\chi^2</math></b>	<b>Hazard Ratio</b>	<b>95% CI</b>
<b>Model 1</b>					
ADHD	0.77 (0.18)	<0.001	18.05	2.16	1.52 to 3.09
ADHD*time	-0.15 (0.06)	0.021	5.36	0.86	0.76 to 0.98
<b>Model 2</b>					
ADHD	0.72 (0.18)	<0.001	16.08	2.06	1.45 to 2.93
ADHD*time	-0.14 (0.06)	0.024	5.11	0.87	0.77 to 0.98
Peer dislike	0.29 (0.10)	0.003	9.01	1.33	1.10 to 1.60
<b>Model 3</b>					
ADHD	0.74 (0.18)	<0.001	16.01	2.09	1.46 to 2.99
ADHD*time	-0.13 (0.06)	0.042	4.15	0.88	0.77 to 1.00
Victimisation	0.39 (0.18)	0.026	4.96	1.48	1.05 to 2.09
Victimisation*time	-0.20 (0.10)	0.047	3.96	0.82	0.68 to 1.00
<b>Model 4</b>					
ADHD	0.69 (0.18)	<0.001	14.71	1.99	1.40 to 2.85
ADHD*time	-0.13 (0.06)	0.047	3.94	0.88	0.78 to 1.00
Peer dislike	0.33 (0.11)	0.002	9.61	1.38	1.13 to 1.70
Victimisation	0.25 (0.20)	0.200	1.64	1.28	0.88 to 1.88
Victimisation*time	-0.21 (0.10)	0.041	4.19	0.81	0.66 to 0.99

Table 5.3 Time varying effects of ADHD symptoms on the risk for depression with and without controlling for dislike and victimisation

Time <sup>#</sup>	ADHD symptoms <sup>*</sup>	
	<i>B</i> (HR)	
	I	II
0	0.77 (2.16)	0.69 (1.99)
1	0.62 (1.86)	0.56 (1.75)
2	0.47 (1.60)	0.43 (1.54)
3	0.32 (1.38)	0.30 (1.35)
4	0.17 (1.19)	0.17 (1.19)
5	0.02 (1.02)	0.04 (1.04)
6	-0.12 (0.89)	-0.09 (0.91)

<sup>#</sup> Time in years after T2

<sup>\*</sup> I = Not controlling for dislike or victimisation; II = Controlling for both dislike and victimisation



Table 5.4 Gender differences in Cox regression estimates of the effect of ADHD symptoms on depression onset before and after adjusting for peer dislike and victimisation

Covariate	Girls			Boys		
	B (SE)	p	HR	B (SE)	p	HR
<b>Model 1</b>						
ADHD	0.80 (0.19)	<0.001	2.22	0.64 (0.42)	0.12	1.90
ADHD*time	-0.18 (0.07)	0.013	0.84	-0.08 (0.13)	0.51	0.92
<b>Model 2</b>						
ADHD	0.72 (0.19)	<0.001	2.06	0.64 (0.38)	0.13	1.90
ADHD*time	-0.17 (0.07)	0.01	0.84	-0.09 (0.12)	0.50	0.92
Peer dislike	0.30 (0.10)	0.004	1.34	0.20 (0.17)	0.29	1.23
<b>Model 3</b>						
ADHD	0.77 (0.19)	<0.001	2.15	0.68 (0.45)	0.13	1.97
ADHD*time	-0.17 (0.07)	0.02	0.84	-0.09 (0.13)	0.48	0.91
Victimisation	0.17 (0.24)	0.49	1.18	0.69 (0.26)	0.009	1.95
Victimisation*time	-0.07 (0.10)	0.52	0.94	-0.41 (0.21)	0.05	0.67
<b>Model 4</b>						
ADHD	0.72 (0.19)	<0.001	2.05	0.68 (0.45)	0.13	1.97
ADHD*time	-0.16 (0.07)	0.02	0.85	-0.10 (0.14)	0.45	0.90
Peer dislike	0.33 (0.11)	0.003	1.38	0.25 (0.21)	0.24	1.28
Victimisation	0.02 (0.26)	0.95	1.02	0.58 (0.28)	0.04	1.79
Victimisation*time	-0.07 (0.11)	0.51	0.93	-0.43 (0.21)	0.04	0.65

Figure 5.1 Flowchart of participants at each wave with mean ages (in years)

