

University of Groningen

Neuroanatomical changes in patients with loss of visual function

Prins, Doety

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:
2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Prins, D. (2016). *Neuroanatomical changes in patients with loss of visual function*. [Thesis fully internal (DIV), University of Groningen]. Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

Alcauter S, Barrio FA, Diaz R, and Fernandez-Ruiz J. Gray and white matter alterations in spinocerebellar ataxia type 7: an in vivo DTI and VBM study. *Neuroimage* 2011; 55 (1):1-7.

Alexander AL, Lee JE, Lazar M, Field AS. Diffusion Tensor Imaging of the brain. *Neurotherapeutics* 2007; 4:316-329.

Arden GB. Age-related macular degeneration. *Journal of the British Menopause Society* 2006; 12 (2):64-70.

Ashburner J and Friston KJ. Voxel-based morphometry--the methods. *Neuroimage* 2000; 11 (6 Pt 1):805-821.

Ashburner J and Friston KJ. Why voxel-based morphometry should be used. *Neuroimage* 2001; 14:1238-1243.

Ashburner J. A fast diffeomorphic image registration algorithm. *Neuroimage* 2007; 38 (1):95-113.

Augood CA, Vingerling JR, De Jong PT, Chakravarthy U, Seland J, Soubrane G, et al. Prevalence of age-related maculopathy in older Europeans: the European Eye Study (EUREYE). *Archives of Ophthalmology* 2006; 124 (4):529-535.

Backus BT, Fleet DJ, Parker AJ, Heeger DJ. Human cortical activity correlates with stereoscopic depth perception. *Journal of Neurophysiology* 2001; 86(4):2054–2068.

Baecke S, Lützkendorf R, Tempelmann C, Müller C, Adolf D, Scholz M, and Bernarding J. Event-related functional magnetic resonance imaging (efMRI) of depth-by-disparity perception: additional evidence for right-hemispheric lateralization. *Experimental Brain Research* 2009; 196(3):453–458.

Baker CI, Peli E, Knouf N, and Kanwisher N. Reorganization of visual processing in macular degeneration. *Journal of Neuroscience* 2005; 25 (3):614-618.

Baker CI, Dilks DD, Peli E, and Kanwisher N. Reorganization of visual processing in macular degeneration: replication and clues about the role of foveal loss. *Vision Research* 2008; 48:1910-1919.

Baker CI, Dilks DD, Peli E, and Kanwisher N. Reorganization of visual processing in macular degeneration: replication and clues about the role of foveal loss. *Vision Research* 2008; 48 (18):1910-1919.

- Balk LJ, Twisk JWR, Steenwijk MD, Daams M, Tewarie P, Killestein J, et al. A dam for retrograde axonal degeneration in multiple sclerosis? *Journal of Neurology, Neurosurgery & Psychiatry* 2014; 85: 782–9.
- Ban H, Preston TJ, Meeson A, and Welchman AE. The integration of motion and disparity cues to depth in dorsal visual cortex. *Nature Neuroscience* 2012; 15(4):636–643.
- Barnes GR, Li X, Thompson B, Singh KD, Dumoulin SO, Hess RF. Decreased gray matter concentration in the lateral geniculate nuclei in human amblyopes. *Investigative Ophthalmology & Visual Science* 2010; 51:1432-1438.
- Baseler HA, Gouws A, Crossland MD, Leung C, Tufail A, Rubin GS, et al. Objective visual assessment of antiangiogenic treatment for wet age-related macular degeneration. *Optometry & Vision Science* 2011; 88: 1255–61.
- Baseler HA, Gouws A, Haak KV, Racey C, Crossland MD, Tufail A, et al. Large-scale remapping of visual cortex is absent in adult humans with macular degeneration. *Nature Neuroscience* 2011; 14 (5):649-655.
- Basser PJ, Pajevic S, Pierpaoli C, Duda J, Aldroubi A. In vivo fiber tractography using DT-MRI data. *Magnetic Resonance in Medicine* 2000; 44(4):625-632.
- Bayer AU, Ferrari F, Erb C. High occurrence rate of glaucoma among patients with Alzheimer's disease. *European Journal of Neurology* 2002; 47(3):165-168.
- Beatty RM, Sadun AA, Smith L, Vonsattel JP, and Richardson EP. Direct demonstration of transsynaptic degeneration in the human visual system: a comparison of retrograde and anterograde changes. *Journal of Neurology, Neurosurgery, and Psychiatry*. 1982; 45(2):143–146.
- Biarnés M, Monés J, Alonso J, and Arias L. Update on geographic atrophy in age-related macular degeneration. *Optometry & Vision Science* 2011; 88 (7):881-889.
- Bookstein F. "Voxel-based morphometry" should not be used with imperfectly registered images. *Neuroimage* 2001; 14:1454-1462.
- Boucard CC, Hernowo AT, Maguire RP, Jansonius NM, Roerdink JB, Hooymans JM, and Cornelissen FW. Changes in cortical grey matter density associated with long-standing retinal visual field defects. *Brain* 2009; 132 (Pt 7):1898-1906.

Bridge H, von dem Hagen E, Davies G, Chambers C, Gouws A, Hoffmann M, et al. Changes in brain morphology in albinism reflect reduced visual acuity. *Cortex* 2012; 56:64-72.

Briggs F and Usrey WM. Corticogeniculate feedback and visual processing in the primate. *Journal of Physiology* 2011; 589 (pt1):33-40.

Brouwer GJ, van Ee R, and Schwarzbach J. Activation in visual cortex correlates with the awareness of stereoscopic depth. *The Journal of Neuroscience* 2005; 25(45):10403–10413.

Bürgel U, Amunts K, Hoemke L, Mohlberg H, Gilsbach JM, and Zilles K White matter fiber tracts of the human brain: three-dimensional mapping at microscopic resolution, topography and intersubject variability. *NeuroImage* 2006; 29(4):1092–1105.

Bürgel U, Schormann T, Schleicher A, and Zilles K. Mapping of histologically identified long fiber tracts in human cerebral hemispheres to the MRI volume of a reference brain: position and spatial variability of the optic radiation. *Neuroimage* 1999; 10 (5):489-499.

Bürgel U, Amunts K, Hoemke L, Mohlberg H, Gilsbach J, and Coenen VA. White matter fiber tracts of the human brain: three-dimensional mapping at microscopic resolution, topography and intersubject variability. *Neuroimage* 2006; 29 (4):1092-1105.

Chang EE, Goldberg JL. Glaucoma 2.0: Neuroprotection, neuroregeneration, neuroenhancement. *Ophthalmology* 2012; 119:979-986.

Chang ST, Xu J, Trinkaus K, Pekmezci M, Arthur SN, Song S-K, et al. Optic nerve Diffusion Tensor Imaging parameters and their correlation with optic disc topography and disease severity in adult glaucoma patients and controls. *Journal of Glaucoma* 2014; 23(8):513-520.

Chen WW, Wang N, Cai S, Fang Z, Yu M, Wu Q, et al. Structural brain abnormalities in patients with primary open-angle glaucoma: a study with 3T MR imaging. *Investigative Ophthalmology & Visual Science* 2013; 54: 545-54.

Chen Z, Lin F, Wang J, Li Z, Dai H, Mu K, et al. Diffusion tensor magnetic resonance imaging reveals visual pathway damage that correlates with clinical severity in glaucoma. *Clinical & Experimental Ophthalmology* 2013; 41:43-49.

Cheung CM, Tai ES, Kawasaki R, Tay WT, Lee JL, Hamzah H, and Wongh TY. Prevalence of and Risk Factors for Age-Related Macular Degeneration in a Multiethnic Asian Cohort. *Archives of Ophthalmology* 2012; 130(4):480-486.

- Clarke S. Modular organization of human extrastriate visual cortex: evidence from cytochrome oxidase pattern in normal and macular degeneration cases. *European Journal of Neuroscience* 1994; 6 (5):725-736.
- Congdon N, O'Colmain B, Klaver CCW, Klein R, Muñoz B, et al. Causes and prevalence of visual impairment among adults in the United States. *Archives of Ophthalmology* 2004; 122:477-485.
- Culham JC, and Kanwisher NG. Neuroimaging of cognitive functions in human parietal cortex. *Current Opinion in Neurobiology* 2001; 11(2):157-163.
- Cumurcu T, Dorak F, Cumurcu BE, Erbay LG, Ozsoy E. Is there any relation between pseudoexfoliation syndrome and Alzheimer's type dementia? *Seminars in Ophthalmology* 2013; 28:224-229.
- Da Cruz L, Coley BF, Dorn J, Merlini F, Filley E, Christopher P, et al. The Argus II epiretinal prosthesis system allows letter and word reading and long-term function in patients with profound vision loss. *The British Journal of Ophthalmology* 2013; 97(5):632-636.
- Dai H, Yin D, Hu C, Morelli JN, Hu S, Yan X, et al. Whole-brain voxel-based analysis of diffusion tensor MRI parameters in patients with primary open angle glaucoma and correlation with clinical glaucoma stage. *Neurology* 2013; 55(2):233-243.
- Dale AM, Fisch B, Sereno MI. Cortical surface-based analysis. I. Segmentation and surface reconstruction. *Neuroimage* 1999; 9:179-194.
- Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, et al. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage* 2006; 31 (3):968-980.
- Dilks DD, Baker CI, Peli E, and Kanwisher N. Reorganization of visual processing in macular degeneration is not specific to the "preferred retinal locus". *Journal of Neuroscience* 2009; 29 (9):2768-2773.
- Doerfler A, Waerntges S, Michelson G, Otto M, Struffert T, El-Rafei et al. Changes of radial diffusivity and fractional anisotropy in the optic nerve and optic radiation of glaucoma patients. *The Scientific World Journal* 2012; 2012:1-5.
- Eckhardt C, Eckhardt U. Macular translocation in nonexudative age-related macular degeneration. *Retina* 2002; 22:786-794.

El-Rafei A, Engelhorn T, Wärtges S, Dörfler A, Hornegger J, Michelson G. Glaucoma classification based on visual pathway analysis using diffusion tensor imaging. *Magnetic Resonance Imaging* 2013; 31:1081-1091.

Fechtner RD, Weinreb RN. Mechanisms of optic nerve damage in primary open angle glaucoma. *Survey of Ophthalmology* 1994; 39:23-42.

Fields RD. Neuroscience. Change in the brain's white matter. *Science* 2010; 330(6005):768–769.

Finger RP, Fimmers R, Holz FG, and Scholl HP. Prevalence and causes of registered blindness in the largest federal state of Germany. *British Journal of Ophthalmology* 2011; 95 (8):1061-1067.

Fischl B, van der Kouwe A, Destrieux C, Halgren E, Ségonne F, et al. Automatically parcellating the human cerebral cortex. *Cerebral Cortex* 2004; 14:11-22.

Fischl B, Liu A, Dale AM. Automated manifold surgery: constructing geometrically accurate and topologically correct models of the human cerebral cortex. *IEEE Transactions on Medical Imaging* 2001; 20:70-80.

Fischl B, Sereno MI, Dale AM. Cortical surface-based analysis. II: Inflation, flattening, and a surface-based coordinate system. *Neuroimage* 1999a; 9:195-207.

Fischl B, Sereno MI, Tootell RB, Dale AM. High-resolution intersubject averaging and a coordinate system for the cortical surface. *Human Brain Mapping* 1999b; 8:272-284.

Forrester JV, Xu H, Kuffová L, Dick AD, and McMenamin PG. Dendritic cell physiology and function in the eye. *Immunological Reviews* 2010; 234 (1):282-304.

Frazier JA, Chiu S, Breeze JL, Makris N, Lange N, Kennedy DN, et al. Structural brain magnetic resonance imaging of limbic and thalamic volumes in pediatric bipolar disorder. *American Journal of Psychiatry* 2005; 162 (7):1256-1265.

Gabilondo I, Martínez-Lapiscina EH, Martínez-Heras E, Fraga-Pumar E, Llufríu S, Ortiz S, et al. Trans-synaptic axonal degeneration in the visual pathway in multiple sclerosis. *Annals of Neurology* 2014; 75: 98–107.

Garaci FG, Bolacchi F, Cerulli A, Melis M, Spanò A, Cedrone C, et al. Optic nerve and optic radiation neurodegeneration in patients with glaucoma: in vivo analysis with 3-T diffusion-tensor MR imaging. *Radiology* 2009; 252:496-501.

Gehrs KM, Anderson DH, Johnson LV, and Hageman GS. Age-related macular degeneration--emerging pathogenetic and therapeutic concepts. *Annals of Medicine* 2006; 38 (7):450-471.

Ghiso JA, Doudevski I, Ritch R, Rostagno AA. Alzheimer's disease and glaucoma: mechanistic similarities and differences. *Journal of Glaucoma* 2013; 22 suppl 5:S36-38.

Von Glehn F, Jarius S, Cavalcanti Lira RP, Alves Ferreira MC, Von Glehn FHR, et al. Structural brain abnormalities are related to retinal nerve fiber layer thinning and disease duration in neuromyelitis optica spectrum disorders. *Multiple Sclerosis Journal* 2014; 20(9):1189-1197.

Göbel AP, Fleckenstein M, Schmitz-Valckenberg S, Brinkmann CK, and Holz FG. Imaging Geographic Atrophy in Age-Related Macular Degeneration. *Ophthalmologica* 2011; 226 (4):182-190.

Goldstein JM, Seidman LJ, Makris N, Ahern T, O'Brien LM, Caviness VS Jr, et al. Hypothalamic abnormalities in schizophrenia: sex effects and genetic vulnerability. *Biological Psychiatry* 2007; 61 (8):935-945.

Good C, Johnsrude I, Ashburner J, Henson RN, Friston KJ, Frackowiak RS. A voxel-based morphometric study of ageing in 465 normal adult human brains. *Neuroimage* 2001; 14:21-36.

Goncalves NR, Ban H, Sánchez-Panchuelo RM, Francis ST, Schluppeck D, and Welchman AE. 7 tesla fMRI reveals systematic functional organization for binocular disparity in dorsal visual cortex. *The Journal of Neuroscience* 2015; 35(7):3056-3072.

Guillery RW, Okoro AN, Witkop CJ. Abnormal visual pathways in the brain of a human albino. *Brain Research* 1975; 96:373-377.

Gupta N, Ang L-C, Noël de Tilly L, Bidaisee L, Yücel YH. Human glaucoma and neural degeneration in intracranial optic nerve, lateral geniculate nucleus, and visual cortex. *British Journal of Ophthalmology* 2006; 90:674-678.

Gupta N, Greenberg G, de Tilly LN, Gray B, Polemidiotis M, Yücel YH. Atrophy of the lateral geniculate nucleus in human glaucoma detected by magnetic resonance imaging. *British Journal of Ophthalmology* 2009; 93:56-60.

Gupta N, Yücel YH. What changes can we expect in the brain of glaucoma patients? *Survey of Ophthalmology* 2007; 52 suppl 2:S122-126.

Haak KV, Langers DRM, Renken R, van Dijk P, Borgstein J, Cornelissen FW. Abnormal visual field maps in human cortex: A mini-review and a case report. *Cortex* 2014; 56:14-25.

Von dem Hagen EAH, Houston GC, Hoffmann MB, Jeffery G, Morland AB. Retinal abnormalities in human albinism translate into a reduction of grey matter in the occipital cortex. *European Journal of Neuroscience* 2005; 22:2475-2480.

Hernowo AT, Boucard CC, Jansonius NM, Hooymans JM, and Cornelissen FW. Automated Morphometry of The Visual Pathway in Primary Open-Angle Glaucoma. *Investigative Ophthalmology & Visual Science* 2011; 52 (5):2758-2766.

Hernowo AT, Prins D, Baseler HA, Plank T, Gouws AD, Hooymans JMM, et al. Morphometric analyses of the visual pathways in macular degeneration. *Cortex* 2014; 56:99-110.

Holz FG, Pauleikhoff D, Klein R, and Bird AC. Pathogenesis of lesions in late age-related macular disease. *American Journal of Ophthalmology* 2004; 137 (3):504-510.

Hutton C, Draganski B, Ashburner J, Weiskopf N. A comparison between voxel-based cortical thickness and voxel-based morphometry in normal aging. *Neuroimage* 2009; 48:371-380.

Ikram MK, Cheung CY, Wong TY, Chen CPLH. Retinal pathology as biomarker for cognitive impairment and Alzheimer's disease. *Journal of Neurology, Neurosurgery & Psychiatry* 2012; 83:917-922.

Inoue T, Kawaji T, Tanihara H. Elevated levels of multiple biomarkers of Alzheimer's disease in the aqueous humor of eyes with open-angle glaucoma. *Investigative Ophthalmology & Visual Science* 2013; 54:5353-5358.

Ip IB, Minini L, Dow J, Parker AJ, and Bridge H Responses to interocular disparity correlation in the human cerebral cortex. *Ophthalmic & Physiological Optics* 2014; 34(2):186-198.

Iwata F, Patronas NJ, Caruso RC, Podgor MJ, Remaley NA, Kupfer C, et al. Association of visual field, cup-disc ratio, and magnetic resonance imaging of optic chiasm. *Archives of Ophthalmology* 1997:729-732.

- Jack CR Jr, Bernstein MA, Fox NC, Thompson P, Alexander G, Hervey D, et al. The Alzheimer's Disease Neuroimaging Initiative (ADNI): MRI methods. *Journal of Magnetic Resonance Imaging* 2008; 27 (4):685-691.
- Janssen SF, Gorgels TGMF, Ramdas WD, Klaver CCW, van Duijn CM, Jansonius NM, et al. The vast complexity of primary open angle glaucoma: disease genes, risks, molecular mechanisms and pathobiology. *Progress in Retinal and Eye Research* 2013; 37:31-67.
- Jenchitr W, Ruamviboonsuk P, Sanmee A, and Pokawattana N. Prevalence of age-related macular degeneration in Thailand. *Ophthalmic Epidemiology* 2011; 18 (1):48-52.
- Jenkinson M, Bannister P, Brady M, and Smith S. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *NeuroImage* 2002; 17(2):825-841.
- Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, and Smith SM. FSL. *NeuroImage* 2002; 62(2):782-790.
- Jenkinson M, and Smith S. A global optimisation method for robust affine registration of brain images. *Medical Image Analysis* 2001; 5(2):143-156.
- Jones DK, Horsfield MA, Simmons A. Optimal strategies for measuring diffusion in anisotropic systems by magnetic resonance imaging. *Magnetic Resonance in Medicine* 1999a; 42(3):515-525.
- Jones DK, Simmons A, Williams SC, Horsfield MA. Non-invasive assessment of axonal fiber connectivity in the human brain via diffusion tensor MRI. *Magnetic Resonance in Medicine* 1999b; 42(1):37-41
- Jiang J, Zhu W, Shi F, Liu Y, Li J, Qin W, et al. Thick visual cortex in the early blind. *The Journal of Neuroscience* 2009; 29(7):2205-2011.
- Jindahra P, Hedges TR, Mendoza-Santiesteban CE, Plant GT. Optical coherence tomography of the retina: applications in neurology. *Current Opinion in Neurology* 2010; 23: 16-23.
- Johansson BB. Brain plasticity in health and disease. *Keio Journal of Medicine* 2004; 53 (4):231-246.

Jonas JB, Wang NL, Wang YX, You QS, Xie XB, Yang DY, et al. Estimated trans-lamina cribrosa pressure difference versus intraocular pressure as biomarker for open-angle glaucoma. The Beijing Eye Study 2011. *Acta Ophthalmologica* 2015; 93(1):e7-e13.

Jonasson F, Arnarsson A, Eiríksdóttir G, Harris TB, Launer LJ, Meuer SM, et al. Prevalence of age-related macular degeneration in old persons: Age, Gene/environment Susceptibility Reykjavik Study. *Ophthalmology* 2011; 118 (5):825-830.

Kashiwagi K, Okubo T, Tsukahara S. Association of magnetic resonance imaging of anterior optic pathway with glaucomatous visual field damage and optic disc cupping. *Journal of Glaucoma* 2004; 13:189-195.

Keane PA and Sadda SR. Imaging chorioretinal vascular disease. *Eye* 2010; 24 (3):422-427.

Keenan TD, Goldacre R, and Goldacre MJ. Associations between age-related macular degeneration, Alzheimer disease, and dementia: record linkage study of hospital admissions. *JAMA Ophthalmology* 2014; 132(1):63-68.

Kessing LV, Lopez AG, Andersen PK, Kessing SV. No increased risk of developing Alzheimer disease in patients with glaucoma. *Journal of Glaucoma* 2007; 16:47-51.

Kim SY, Sadda S, Humayun MS, de Juan E Jr, Melia BM, and Green WR. Morphometric analysis of the macula in eyes with geographic atrophy due to age-related macular degeneration. *Retina* 2002; 22 (4):464-470.

Kinney PE, Jay B, Witkop CJ. Albinism. *Survey of Ophthalmology* 1988; 30:75-101.

Kirbas S, Turkyilmaz K, Tufekci A, Durmus M. Retinal nerve fiber layer thickness in Parkinson disease. *Journal of Neuroophthalmology* 2013; 33: 62–5.

Kirby E, Bandelow S, Hogervorst E. Visual impairment in Alzheimer's disease: a critical review. *Journa of Alzheimer's disease* 2010; 21:15-34.

Kitajima M, Korogi Y, Hirai T, Hamatake S, Ikushima I, Sugahara T, et al. MR changes in the calcarine area resulting from retinal degeneration. *American Journal of Neuroradiology* 1997; 18 (7):1291-1295.

Klaver CCW, Ott A, Hofman A, Assink JJM, Breteler MMB and De Jong PTVM. Is age-related maculopathy associated with Alzheimer's disease? *American Journal of Epidemiology* 1999; 150:963-8.

- Klaver CC, Wolfs RC, Vingerling JR, Hofman A, De Jong PT. Age-specific prevalence and causes of blindness and visual impairment in an older population: the Rotterdam Study. *Archives of Ophthalmology* 1998; 116:653-658.
- Klein A, Andersson J, Ardekani BA, Ashburner J, Avants B, Chiang MC, et al. Evaluation of 14 nonlinear deformation algorithms applied to human brain MRI registration. *Neuroimage* 2009; 46 (3):786-802.
- Klein R, Chou CF, Klein BE, Zhang X, Meuer SM, and Saaddine JB. Prevalence of age-related macular degeneration in the US population. *Archives of Ophthalmology* 2011, 129 (1):75-80.
- Klistorner A, Sriram P, Vootakuru N, Wang C, Barnett MH, Garrick R, et al. Axonal loss of retinal neurons in multiple sclerosis associated with optic radiation lesions. *Neurology* 2014; 82: 2165–72.
- Kocur I and Resnikoff S. Visual impairment and blindness in Europe and their prevention. *British Journal of Ophthalmology* 2002; 86 (7):716-722.
- Leporé N, Voss P, Lepore F, Chou Y-Y, Fortin M, Gougoux F, et al. Brain structure changes visualized in early- and late-onset blind subjects. *NeuroImage* 2010; 49(1):134–140.
- Levin N, Dumoulin SO, Winawer J, Dougherty RF, and Wandell BA. Cortical maps and white matter tracts following long period of visual deprivation and retinal image restoration. *Neuron* 2010 ; 65(1):21–31.
- Li C, Cai P, Shi L, Lin Y, Zhang J, Liu S, et al. Voxel-based morphometry of the visual-related cortex in primary open angle glaucoma. *Current Eye Research* 2012; 37:794-802.
- Li J, Liu Y, Qin W, Jiang J, Qiu Z, Xu J, et al. Age of onset of blindness affects brain anatomical networks constructed using diffusion tensor tractography. *Cerebral Cortex* 2013b; 23(3):542–551.
- Li K, Lu C, Huang Y, Yuan L, Zeng D, Wu K. Alteration of fractional anisotropy and mean diffusivity in glaucoma: novel results of a meta-analysis of diffusion tensor imaging studies. *PLoS One* 2014; 9:e97445
- Li Q, Jiang Q, Guo M, Li Q, Cai C, Yin X. Grey and white matter changes in children with monocular amblyopia: voxel-based morphometry and diffusion tensor imaging study. *British Journal of Ophthalmology* 2013a; 97:524-529.

Liepert J, Bauder H, Wolfgang HR, Miltner WH, Taub E, and Weiller C. Treatment-induced cortical reorganization after stroke in humans. *Stroke* 2000; 31(6):1210-1216.

Liu T, Cheung SH, Schuchard RA, Glielmi CB, Hu X, He S, and Legge GE. Incomplete cortical reorganization in macular degeneration. *Investigative Ophthalmology & Visual Science* 2010; 51 (12):6826-6834.

Liu T, Shi L, Wang J, Li C, Du H, Lu P, et al. Reduced white matter integrity in primary open-angle glaucoma: A DTI study using tract-based spatial statistics. *Journal of Neuroradiology* 2012; 40:89-93.

Lou AR, Madsen KH, Julian HO, Toft PB, Kjaer TW, Paulson OB, et al. Postoperative increase in grey matter volume in visual cortex after unilateral cataract surgery. *Acta Ophthalmologica* 2013; 91: 58–65.

Lv B, He H, Li X, Zhang Z, Huang W, Li M, et al. Structural and functional deficits in human amblyopia. *Neuroscience Letters* 2008; 437:5-9.

Lyoo IK, Sung YH, Dager SR, Friedman SD, Lee J-Y, et al. Regional cerebral cortical thinning in bipolar disorder. *Bipolar disorders* 2006; 8:65-74.

MacLaren RE, Groppe M, Barnard AR, Cottrill CL, Tolmachova T, Seymour L, et al. Retinal gene therapy in patients with choroideremia: initial findings from a phase 1/2 clinical trial. *Lancet* 2014; 383(9923):1129–1137.

Makris N, Goldstein JM, Kennedy D, Hodge SM, Caviness VS, Faraone SV, et al. Decreased volume of left and total anterior insular lobule in schizophrenia. *Schizophrenia Research* 2006; 83 (2-3):155-171.

Margalit E and Sadda SR. Retinal and optic nerve diseases. *Artificial Organs* 2003; 27 (11):963-974.

Masuda Y, Dumoulin SO, Nakadomari S, and Wandell BA. V1 projection zone signals in human macular degeneration depend on task, not stimulus. *Cerebral Cortex* 2008; 18 (11):2483-2493.

Medeiros NE and Curcio CA. Preservation of ganglion cell layer neurons in age-related macular degeneration. *Investigative Ophthalmology & Visual Science* 2001; 42 (3):795-803.

- Mendola JD, Conner IP, Roy A, Chan S-T, Schwartz TL, Odom JV, et al. Voxel-based analysis of MRI detects abnormal visual cortex in children and adults with amblyopia. *Human Brain Mapping* 2005; 25:222-236.
- Menke MN, Dabov S, Knecht P, and Sturm V. Reproducibility of retinal thickness measurements in patients with age-related macular degeneration using 3D Fourier-domain optical coherence tomography (OCT) (Topcon 3D-OCT 1000). *Acta Ophthalmologica* 2011; 89 (4):346-351.
- Merzenich MM, Nelson RJ, Stryker MP, Cynader MS, Schoppmann A, and Zook JM. Somatosensory cortical map changes following digit amputation in adult monkeys. *Journal of Comparative Neurology* 1984; 224 (4):591-605.
- Michelson G, Engelhorn T, Warntges S, El Rafei A, Hornegger J, Doerfler A. DTI parameters of axonal integrity and demyelination of the optic radiation correlate with glaucoma indices. *Graefe's Archives for Clinical and Experimental Ophthalmology* 2013; 251:243-253.
- Minini L, Parker AJ, and Bridge H. Neural modulation by binocular disparity greatest in human dorsal visual stream. *Journal of Neurophysiology* 2010; 104(1):169–178.
- Muoz B, West SK, Rubin GS, Schein OD, Quigley HA, et al. Causes of blindness and visual impairment in a population of older Americans: The Salisbury Eye Evaluation Study. *Archives of Ophthalmology* 118:819-825.
- Murai H, Suzuki Y, Kiyosawa M, Tokumaru AM, Ishii K, Mochizuki M. Positive correlation between the degree of visual field defect and optic radiation damage in glaucoma patients. *Japanese Journal of Ophthalmology* 2013; 57:257-262.
- Nangia V, Jonas JB, Kulkarni M, and Matin A. Prevalence of age-related macular degeneration in rural central India: the Central India Eye and Medical Study. *Retina* 2011; 31 (6):1179-1185.
- Neri P, Bridge H, and Heeger DJ. Stereoscopic processing of absolute and relative disparity in human visual cortex. *Journal of Neurophysiology* 2004; 92(3):1880–1891.
- Neveu MM, von dem Hagen E, Morland AB, Jeffery G. The fovea regulates symmetrical development of the visual cortex. *Journal of Comparative Neurology* 2008; 506:791-800.

Ngai LY, Stocks N, Sparrow JM, Patel R, Rumley A, Lowe G, et al. The prevalence and analysis of risk factors for age-related macular degeneration: 18-year follow-up data from the Speedwell eye study, United Kingdom. *Eye* 2011; 25 (6):784-793.

Nguyen TH, Stievenart JL, Saucet JC, Le Gargasson JF, Cohen YS, Pelegrini-Issac M, et al. [Cortical response to age-related macular degeneration (Part II). Functional MRI study]. *Journal Français d'Ophthalmologie* 2004; 27 (9 Pt 2):3S72-3S86.

Nickells RW. Retinal ganglion cell death in glaucoma: the how, the why, and the maybe. *Journal of Glaucoma* 1996; 5:345-356.

Noppeney U. The effects of visual deprivation on functional and structural organization of the human brain. *Neuroscience and Biobehavioral Reviews* 2007; 31(8):1169–1180.

Noppeney U, Friston KJ, Ashburner J, Frackowiak R, and Price CJ. Early visual deprivation induces structural plasticity in gray and white matter. *Current Biology* 2005; 15(13):R488–490.

Nucci C, Strouthidis NG, Khaw PT. Neuroprotection and other novel therapies for glaucoma. *Current Opinion in Pharmacology* 2013; 13:1-4.

Ohno-Matsui K. Parallel findings in age-related macular degeneration and Alzheimer's disease. *Progress in Retinal and Eye Research* 2011; 30:217-238.

Osborne NN. Recent clinical findings with memantine should not mean that the idea of neuroprotection in glaucoma is abandoned. *Acta Ophthalmology* 2009; 87:450-454.

Ou Y, Grossman DS, Lee PP, Sloan FA. Glaucoma, Alzheimer disease and other dementia: a longitudinal analysis. *Ophthalmic Epidemiology* 2012; 19:285-292.

Palaniyappan L, Liddle PF. Differential effects of surface area, gyrification and cortical thickness on voxel-based morphometric deficits in schizophrenia. *Neuroimage* 2008; 43:665-675.

Pan W-J, Wu G, Li C-X, Lin F, Sun J, and Lei H. Progressive atrophy in the optic pathway and visual cortex of early blind Chinese adults: A voxel-based morphometry magnetic resonance imaging study. *NeuroImage* 2007; 37(1):212–220.

Park H-J, Jeong S-O, Kim EY, Kim J Il, Park H, Oh M-K, et al. Reorganization of neural circuits in the blind on diffusion direction analysis. *Neuroreport* 2007; 18(17), 1757–1760.

- Park H-J, Lee JD, Kim EY, Park B, Oh M-K, Lee S, and Kim J-J. Morphological alterations in the congenital blind based on the analysis of cortical thickness and surface area. *NeuroImage* 2009; 47(1):98–106.
- Pascale A, Drago F, Govoni S. Protecting the retinal neurons from glaucoma: Lowering ocular pressure is not enough. *Pharmacological Research* 2012; 66:19-32.
- Pascual-Leone A, Amedi A, Fregni F, Merabet LB. The plastic human brain cortex. *Annual Review of Neuroscience* 2005; 28:377-401
- Pham TQ, Kifley A, Mitchell P and Wang JJ. Relation of age-related macular degeneration and cognitive impairment in an older population. *Gerontology* 2006; 52:353-358.
- Pi LH, Chen L, Liu Q, Ke N, Fang J, Zhang S, et al. Prevalence of eye diseases and causes of visual impairment in school-aged children in Western China. *Journal of Epidemiology* 2012; 22 (1):37-44.
- Plank T, Frolo J, Brandl-Rühle S, Renner AB, Hufendiek K, Helbig H, and Greenlee MW. Gray matter alterations in visual cortex of patients with loss of central vision due to hereditary retinal dystrophies. *Neuroimage* 2011; 56 (3):1556-1565.
- Prins D, Hanekamp S, and Cornelissen FW. Structural brain MRI studies in eye diseases: are they clinically relevant? A review of current findings. *Acta Ophthalmologica* 2016; 94(2):113–121.
- Prins D, Plank T, Baseler HA, Gouws AD, Beer A, Morland AB, et al. Surface-Based Analyses of Anatomical Properties of the Visual Cortex in Macular Degeneration. *PloS One* 2016; 11(1): e0146684.
- Proitsi P, Lupton MK, Dudbridge F, Tsolaki M, Hamilton G, Daniilidou M, et al. Alzheimer's disease and age-related macular degeneration have different genetic models for complement gene variation. *Neurobiology of Aging* 2012; 33(8):1843.e9-17.
- Przedborski S, Vila M, Jackson-Lewis V. Neurodegeneration: What is it and where are we? *Journal of Clinical Investigation* 2003; 111:3-10.
- Ptito M, Schneider FCG, Paulson OB, and Kupers R. Alterations of the visual pathways in congenital blindness. *Experimental Brain Research* 2008; 187(1):41–49.

Pu M, Xu L, Zhang H. Visual response properties of retinal ganglion cells in the Royal College of Surgeons dystrophic rat. *Investigative Ophthalmology & Visual Science* 2006; 47:3579.

Rahmani B, Tielsch JM, Katz J, Gottsch J, Quigley H, et al. The cause-specific prevalence of visual impairment in an urban population. The Baltimore Eye Survey. *Ophthalmology* 1996; 103:1721-1726.

Raz N, Lindenberger U, Rodrigue KM, Kennedy KM, Head D, Williamson A, et al. Regional brain changes in aging healthy adults: general trends, individual differences and modifiers. *Cerebral Cortex* 2005; 15 (11):1676-1689.

Resnikoff S, Pascolini D, Etya'ale D, Kocur I, Pararajasegaram R, Pokharel GP, and Mariotti SP. Global data on visual impairment in the year 2002. *Bulletin of the World Health Organization* 2004; 82 (11):844-851.

Rosengarth K, Keck I, Brandl-Rühle S, Frolo J, Hufendiek K, Greenlee MW, et al. Functional and structural brain modifications induced by oculomotor training in patients with age-related macular degeneration. *Frontiers in Psychology* 2013; 4: 428.

Rutschmann RM, and Greenlee MW. BOLD response in dorsal areas varies with relative disparity level. *Neuroreport* 2004; 15(4):615–619.

Schlegel AA, Rudelson JJ, and Tse PU. White matter structure changes as adults learn a second language. *Journal of Cognitive Neuroscience* 2012; 24(8):1664–1670.

Schmitz B, Schaefer T, Krick CM, Reith W, Backens M, Käsmann-Kellner B. Configuration of the optic chiasm in humans with albinism as revealed by magnetic resonance imaging. *Investigative Ophthalmology & Visual Science* 2003; 44:16-21.

Scholz J, Klein MC, Behrens TEJ, and Johansen-Berg H. Training induces changes in white-matter architecture. *Nature Neuroscience* 2009; 12(11):1370–1371.

Schoth F, Burgel U, Dorsch R, Reinges MHT, and Krings T. Diffusion tensor imaging in acquired blind humans. *Neuroscience Letters* 2006; 398(3):178–182.

Schotz F, Burgel U, Dorsch R, Reinges MHT, Krings T. Diffusion tensor imaging in acquired blind humans. *Neuroscience Letters* 2006; 398:178-182.

- Schumacher EH, Jacko JA, Primo SA, Main KL, Molomey KP, Kinzel EN, and Ginn J. Reorganization of visual processing is related to eccentric viewing in patients with macular degeneration. *Restorative Neurology and Neuroscience* 2008; 26 (4-5):391-402.
- Ségonne F, Dale AM, Busa E, Glessner M, Salat D, et al. A hybrid approach to the skull stripping problem in MRI. *Neuroimage* 2004; 22:1060-1075.
- Ségonne F, Pacheco J, Fischl B. Geometrically accurate topology-correction of cortical surfaces using nonseparating loops. *IEEE Transactions on Medical Imaging* 2007; 26:518-529.
- Shimony JS, Burton H, Epstein AA, McLaren DG, Sun SW, and Snyder AZ. Diffusion tensor imaging reveals white matter reorganization in early blind humans. *Cerebral Cortex* 2006; 16(11):1653–1661.
- Shu N, Li J, Li K, Yu C, and Jiang T. Abnormal diffusion of cerebral white matter in early blindness. *Human Brain Mapping* 2009; 30(1):220–227.
- Sivak JM. The aging eye: common degenerative mechanisms between the Alzheimer's brain and retinal disease. *Investigative Ophthalmology & Visual Science* 2013; 54:871-880.
- Sled JG, Zijdenbos AP, Evans AC. A nonparametric method for automatic correction of intensity nonuniformity in MRI data. *IEEE Transactions on Medical Imaging* 1998; 17:87-97.
- Smirnakis SM, Brewer AA, Schmid MC, Tolia AS, Schüz A, Augath M, et al. Lack of long-term cortical reorganization after macaque retinal lesions. *Nature* 2005; 435 (7040):300-307.
- Smith SM. Fast robust automated brain extraction. *Human Brain Mapping* 2002; 17(3):143–155.
- Smith SM and Brady JM. SUSAN - a new approach to low level image processing. *International Journal of Computer Vision* 1997; 23 (1):45-78.
- Smith SM and Nichols TE. Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localization in cluster inference. *Neuroimage* 2009; 44 (1):83-98.
- Spanish Eyes Epidemiological (SEE) study group. Prevalence of age-related macular degeneration in Spain. *British Journal of Ophthalmology* 2011; 95 (7):931-936.

Stasheff SF. Emergence of sustained spontaneous hyperactivity and temporary preservation of OFF responses in ganglion cells of the retinal degeneration (rd1) mouse. *Journal of Neurophysiology* 2008; 99:1408-1421.

Steeves JKE, González EG, and Steinbach MJ. Vision with one eye: a review of visual function following unilateral enucleation. *Spatial Vision* 2008; 21(6):509–529.

Stein JD, Vanderbeek BL, Talwar N, Nan B, Musch DC, and Zachs DN. Rates of nonexudative and exudative age-related macular degeneration among Asian American ethnic groups. *Investigative Ophthalmology & Visual Science* 2011; 52 (9):6842-6848.

Stingl K, Greppmaier U, Wilhelm B, Zrenner E. Subretinal visual implants. *Klinische Monatsblätter für Augenheilkunde* 2010; 227:940-945.

Sunness JS, Liu T, and Yantis S. Retinotopic mapping of the visual cortex using functional magnetic resonance imaging in a patient with central scotomas from atrophic macular degeneration. *Ophthalmology* 2004; 111 (8):1595-1598.

Szlyk JP, Little DM. An fMRI study of word-level recognition and processing in patients with age-related macular degeneration. *Investigative Ophthalmology & Visual Science* 2009; 50:4487-4495.

Tamura H, Kawakami H, Kanamoto T, Kato T, Yokoyama T, Sasaki K, et al. High frequency of open-angle glaucoma in Japanese patients with Alzheimer's disease. *Journal of the Neurological Sciences* 2006; 246:79-83.

Tanito M, Ohira A. Hemianopic inner retinal thinning after stroke. *Acta Ophthalmologica* 2013; 91: e237–8.

VanNewkirk MR, Weih L, McCarty CA, Taylor HR. Cause-specific prevalence of bilateral visual impairment in Victoria, Australia: the Visual impairment project. *Ophthalmology* 2001; 108:960-967.

Voets NL, Hough MG, Douaud G, Matthews PM, James A, et al. Evidence for abnormalities of cortical development in adolescent-onset schizophrenia. *Neuroimage* 2008; 43:665-675.

Wandell BA, Smirnakis SM. Plasticity and stability of visual field maps in adult primary visual cortex. *Nature Reviews Neuroscience* 2009; 10:873-884.

- Wang D, Qin W, Liu Y, Zhang Y, Jiang T, and Yu C. Altered white matter integrity in the congenital and late blind people. *Neural Plasticity* 2013; 128236.
- Wang M-Y, Wu K, Xu J-M, Dai J, Qin W, Liu J, et al. Quantitative 3-T diffusion tensor imaging in detecting optic nerve degeneration in patients with glaucoma: association with retinal nerve fiber layer thickness and clinical severity. *Neuroradiology* 2013; 55, 493–498.
- Wang YX, Xu L, Lu W, Liu FJ, Qu YZ, Wang J, et al. Parapapillary atrophy in patients with intracranial tumours. *Acta Ophthalmologica* 2013; 91:521-525.
- Welchman AE, Deubelius A, Conrad V, Bühlhoff HH, and Kourtzi Z. 3D shape perception from combined depth cues in human visual cortex. *Nature Neuroscience* 2005; 8(6):820–827.
- Whitwell JL, Tosakulwong N, Weigand SD, Senjem ML, Lowe VJ, et al. Does amyloid deposition produce a specific atrophic signature in cognitively normal subjects? *Neuroimage Clinical* 2013; 2:249-257.
- Williams AL, Lackey J, Wizov SS, Chia TMT, Gatla S, Moster ML, et al. Evidence for widespread structural brain changes in glaucoma: a preliminary voxel-based MRI study. *Investigative Ophthalmology & Visual Science* 2013; 54:5880-5887.
- Winkler AM, Ridgway GR, Webster MA, Smith SM, and Nichols TE. Permutation inference for the general linear model. *NeuroImage* 2014 92:381–397.
- Woo SJ, Park KH, Ahn J, Choe JY, Jeong H, Han JW, et al. Cognitive impairment in age-related macular degeneration and geographic atrophy. *Ophthalmology* 2012; 119 (10):2094-2101.
- Woolrich MW, Jbabdi S, Patenaude B, Chappell M, Makni S, Behrens T, et al. Bayesian analysis of neuroimaging data in FSL. *NeuroImage* 2009; 45(1 Suppl):S173–186.
- Wostyn P, Audenaert K, De Deyn PP. Alzheimer's disease: cerebral glaucoma? *Medical Hypotheses* 2010; 74:973-977.
- Wostyn P, De Groot V, Van Dam D, Audenaert K, De Deyn PP. Intracranial pressure fluctuations: a potential risk factor for glaucoma? *Acta Ophthalmologica* 2015; 93(1):e83-84.

Xiao JX, Xie S, Ye JT, Liu HH, Gan XL, Gong GL, et al. Detection of abnormal visual cortex in children with amblyopia by voxel-based morphometry. *American Journal of Ophthalmology* 2007; 143(3):489-493.

Xie S, Gong GL, Xiao JX, Ye JT, Liu HH, Gan XL, et al. Underdevelopment of optic radiation in children with amblyopia: a tractography study. *American Journal of Ophthalmology* 2007; 143(4):642-646.

Ye JH, Goo YS. The slow wave component of retinal activity in rd/rd mice recorded with a multi-electrode array. *Physiological Measurement* 2007; 28:1079-1088.

Yeatman JD, Weiner KS, Pestilli F, Rokem A, Mezer A, and Wandell BA. The vertical occipital fasciculus: A century of controversy resolved by in vivo measurements. *Proceedings of the National Academy of Sciences of the United States of America* 2014; 111(48):e5214-5223.

Yoon KC, Mun GH, Kim SD, Kim SH, Kim CY, Park KH, et al. Prevalence of eye diseases in South Korea: data from the Korea National Health and Nutrition Examination Survey 2008-2009. *Korean Journal of Ophthalmology* 2011; 25 (6):421-33.

Zarbin MA. Current concepts in the pathogenesis of age-related macular degeneration. *Archives of Ophthalmology* 2004; 122 (4):598-614.

Van Zeeburg EJT, Maaijwee KJM, Missotten TOAR, Heimann H, Van Meurs JC. A free retinal pigment epitheliumchoroid graft in patients with exsudative age-related macular degeneration: Results up to 7 years. *American Journal of Ophthalmology* 2012; 153(1):120-127.

Zhang Y, Brady M, and Smith S. Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Transactions on Medical Imaging* 2001; 20(1):45-57.

Zhang YQ, Li J, Xu L, Zhang L, Wang ZC, Yang H, et al. Anterior visual pathway assessment by magnetic resonance imaging in normal-pressure glaucoma. *Acta Ophthalmologica* 2012a; 90:e295-302.

Zhang Y, Wan S, Ge J, and Zhang X. Diffusion tensor imaging reveals normal geniculocalcarine-tract integrity in acquired blindness. *Brain Research* 2012b; 1458:34-39.

Zhang Z, Wang X, Jonas JB, Wang H, Zhang X, Peng X, et al. Intracranial pressure fluctuations: a potential risk factor for glaucoma? *Acta Ophthalmologica* 2015; 93(1):e84-85.

Zhang Z, Wang X, Jonas JB, Wang H, Zhang X, Peng X, et al. Valsalva manoeuvre, intra-ocular pressure, cerebrospinal fluid pressure, optic disc topography: Beijing intracranial and intra-ocular pressure study. *Acta Ophthalmologica* 2014; 92:e475;480.

Zikou AK, Kitsos G, Tzarouchi LC, Astrakas L, Alexiou GA, Argyropoulou MI. Voxel-based morphometry and diffusion tensor imaging of the optic pathway in primary open-angle glaucoma: a preliminary study. *American Journal of Neuroradiology* 2012; 33:128-134.

Zrenner E. Will retinal implants restore vision? *Science* 2002; 295 (5557):1022-1025.

