

University of Groningen

The educational gradient in self-rated health in Europe

Präg, Patrick; Wittek, Rafael; Mills, Melinda C.

Published in:
Acta Sociologica

DOI:
[10.1177/0001699316670715](https://doi.org/10.1177/0001699316670715)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Präg, P., Wittek, R., & Mills, M. C. (2017). The educational gradient in self-rated health in Europe: Does the doctor-patient relationship make a difference? . *Acta Sociologica*, 60(4), 325-341.
<https://doi.org/10.1177/0001699316670715>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

The educational gradient in self-rated health in Europe: Does the doctor–patient relationship make a difference?

Acta Sociologica
1–17

© The Author(s) 2016

Reprints and permissions:

sagepub.co.uk/journalsPermissions.nav

DOI: 10.1177/0001699316670715

asj.sagepub.com



Patrick Präg

Department of Sociology and Nuffield College, University of Oxford, UK

Rafael Wittek

Department of Sociology, Interuniversity Center for Social Science Theory and Methodology (ICS), University of Groningen, Netherlands

Melinda C Mills

Department of Sociology and Nuffield College, University of Oxford, UK

Abstract

Research suggests that doctor–patient relations have evolved from a doctor-centered, paternalistic approach towards a more patient-centered, egalitarian model of interactions between physicians and their patients. Given the long-running debate on the positive relationship between education and health, the question arises how this development in doctor–patient relations affects social inequalities in health. First, we test to what extent egalitarian (e.g. discussing treatment decisions with patients) doctor–patient relations are underlying the education–self-reported health association. Second, we test whether egalitarian and paternalistic (e.g. withholding some information from patients) doctor–patient relations show differential effects on self-reported health across educational groups. Analyses of the European Social Survey (ESS) 2004/2005 for 24 countries demonstrate that a more egalitarian doctor–patient relationship does not substantially reduce educational inequalities in self-reported health. However, some direct positive effects of egalitarian and direct negative effects of paternalistic doctor–patient relations on health ratings can be found. Finally, results show how the health status of the lower educated can improve with a more egalitarian and less paternalistic doctor–patient relationship.

Keywords

Physician–patient relations, health inequalities, patient-centered care, health care

Corresponding Author:

Patrick Präg, Department of Sociology, University of Oxford, Manor Road, Oxford OX1 3UQ, UK.

Email: patrick.prag@sociology.ox.ac.uk

Introduction

Social inequality in health and disease is a pervasive phenomenon, both over time and across societies. The mechanisms underlying it, however, are not fully understood (Elo, 2009). Earlier research has suggested that one important cause of social inequality in health is differences in quality of health care (Beckfield et al., 2015; Fiscella et al., 2000). Quality of health care comprises both the effectiveness of and access to health care (Campbell et al., 2000). Focusing on a crucial aspect of access to health care, namely the doctor–patient relationship, we examine the effects of different models of doctor–patient interactions on educational inequalities in self-reported health.

Education is a crucial indicator of socioeconomic status and low education is an important risk factor for poor health (Gesthuizen et al., 2012; Mirowsky and Ross, 2003). The lower educated suffer from poorer health compared to their better educated counterparts; for instance, with respect to self-rated health or mortality (Mackenbach et al., 2008; Präg et al., 2014). While remaining largely stable over the life course, education is a major predictor of future life chances. It is strongly connected to access to resources important for health, such as material resources and social connections (Ellwardt et al., 2014).

The aim of this study is to examine whether some aspects of doctor–patient relations exacerbate or compress health inequalities. Existing research points in this direction. Firstly, it has been shown that doctor–patient communication is affected by patient’s SES in the sense that doctors convey less information and directions to patients from the lower social strata of a society, deploy a more directive consulting style, and give patients fewer positive socio-emotional responses (Willems et al., 2005). Secondly, aspects of doctor–patient relationships have been shown to be related to outcomes such as patient satisfaction, coping with disease, and recall and understanding of information received from the physician (Ong et al., 1995), as well as to self-reported and objective health measures (Griffin et al., 2004; Stewart, 1995a). However, there has been little research connecting the two issues of the SES–health gradient and SES differences in the doctor–patient relationship. Our study fills this gap by explicitly addressing the role of doctor–patient relations in relation to educational inequalities in self-rated health.

The current drastic transformations in the doctor–patient relationship make it a particularly interesting factor in the study of inequalities in health. There are strong tendencies towards a more egalitarian relationship between physicians and their patients, in part due to the growing awareness of the crucial role of doctor–patient relations for health outcomes. The traditional ‘doctor-centered’ model (Byrne and Long, 1976), with its dominance of the doctor’s role, is gradually being replaced with the ‘patient-centered’ model that gives greater weight to patients’ needs and preferences, and aims at a partnership between the doctor and patient. For our analyses, we rely on survey data from 24 European countries. As there are no empirical data on doctor–patient relations over a longer period of time, these cross-national data allow us to increase the available variation in doctor–patient relations and allow for additional validity checks of our findings.

By addressing the role of patient–doctor relations in a model of the social gradient in health, we are extending existing research on health inequalities. Our central argument is that the differences in the doctor–patient relationship could affect the social gradient in health in two ways. First, a more egalitarian relationship between doctors and patients might be accompanied by a more egalitarian distribution of health outcomes. Thus, an egalitarian doctor–patient relationship could potentially compress the social gradient in health. The doctor–patient relationship could hence be a pathway factor that *mediates* the relationship between education and self-rated health.

Second, differences in the doctor–patient relationship could potentially affect the health of different educational groups in different ways. Compared to the lower educated, higher-educated individuals might command greater psychosocial resources and have greater experience negotiating with high-status actors such as doctors, irrespective of the doctor–patient relationship. The lower educated might thus benefit to a greater extent from a more egalitarian doctor–patient relationship. We will test whether aspects of the doctor–patient relationship *moderate* the relationship between education and health,

i.e. testing whether educational groups are differently affected by different forms of the doctor–patient relationship.

Theoretical background

The doctor–patient relationship

The relationship between doctors and patients is a complex phenomenon. It involves communication about issues of vital importance, but is often emotionally laden, sometimes involuntary, requires close cooperation, and is based upon interaction between individuals in unequal positions (Ong et al., 1995). Traditionally, the doctor–patient relationship is an asymmetric one, with the doctor being advantaged in several ways. First, doctors command superior knowledge with regard to health and illness. Second, doctors have a gatekeeper function with respect to many aspects of health and illness, both in a formal (e.g. sickness certifications for employers) and an informal way (e.g. the sick role (Parsons, 1951)). Third, doctors have great power in steering interactions with patients, e.g. in terms of waiting times and the content of meetings.

There is a long-standing tradition in sociology of research on the doctor–patient relationship (Henderson, 1935; Heritage and Maynard, 2011) and some classical works of sociology deal with the diffusion of innovations among physicians (Coleman et al., 1957) or interactions between physicians and patients (Parsons, 1951). However, there are relatively few studies which focus on the relationship between the doctor–patient relationship and health outcomes. A number of studies have shown that aspects of doctor–patient relationships can impact patient wellbeing and behavior, such as the ability to understand medical information, compliance with treatment regimes, coping with illness, satisfaction with care, overall quality of life, and health status (Griffin et al., 2004; O’Connor et al., 2009; Ong et al., 1995; Stewart, 1995a; Stewart, 1995b; Weiner et al., 2013; Williams et al., 2000).

The existing literature identified different pathways along which aspects of the doctor–patient relationship can impact health. One potential pathway is the quality of health care. A review of 139 studies revealed that the doctor–patient relationship is the most important predictor of patient-perceived quality of health care (Crow et al., 2003). Another possible pathway is adherence to doctors’ recommendations. Stavropoulou (2011) shows that individuals’ reports of adherence to medication recommendations depend on their relationships to their doctors, corroborating results of earlier studies (Bultman and Svarstad, 2000; Stavropoulou, 2012).

Paternalistic and egalitarian models

The asymmetry between doctors and patients is fundamental for different models of the doctor–patient relationship. Two different models are treated here, namely, the ‘paternalistic’ or ‘doctor-centered’ model and the ‘egalitarian’ or ‘patient-centered’ one. The paternalistic model actively builds upon the asymmetry between doctor and patient, as the doctor’s role as an expert is crucial for this model to function. The general idea of the model is straightforward: Based on their superior medical knowledge, doctors give advice to their patients, who, in turn, comply with this advice. Driven by a desire to appear competent (White, 1959), i.e. to make correct decisions, and aiming at behavioral confirmation (Ormel et al., 1997), patients defer to the doctors’ authority (Cialdini and Griskevicius, 2010). This authority is based on the doctors’ expert status and their superior skills and knowledge. As the model greatly draws upon characteristics of the doctor while the patient holds a largely passive role, this model is sometimes called the ‘doctor-centered model’ (Byrne and Long, 1976).

In the last decades, however, the doctor-centered model has given way to the egalitarian model (Kaba and Sooriakumaran, 2007; Peck and Conner, 2011; Steinhart, 2002). This model tries to bridge the asymmetry between doctor and patient by also conceiving the patient as an expert, namely an expert for their own situation (Department of Health, 2001), and assigning them a more active role. This

development has been fostered by campaigns encouraging patients to ask their physicians questions (Judson et al., 2013). Reasons for this shift of paradigm are numerous, with the most important ones being the crisis of the expert role in society (Beck, 1992), the greater availability of medical information for patients in self-help groups and internet sites (Neuberger, 2000), the increasing role of chronic incurable diseases (Charmaz and Rosenfeld, 2010), and medical progress that has created more treatment options and, thus, more opportunities for decision-making in the cure process. The general idea of the model consists of multiple aspects: Both the doctor and patient are seen as experts—the doctor for conveying medical information, and the patient for their own preferences and living situation. The doctor's task is seen less as providing the patient with binding advice, but more to communicate medical information about diseases and treatments to the patient and to discuss the information with them, such as benefits and risks of different treatment options. If decisions are to be made, they will be made jointly (Barry and Edgman-Levitan, 2012), taking into account not only the medical facts, but also the patient's preferences.

The mechanism at the core of the patient-centered model is also different from the one underlying the doctor-centered one. Instead of satisfying the needs for competence and behavioral confirmation via complying with doctors' orders, the patient-centered model fulfills the desires for patient competence and autonomy by informing the patient and treating them as capable of deciding for themselves. According to self-determination theory (Djundeva et al., 2015; Ryan and Deci, 2000), a combination of competence and autonomy will induce intrinsic motivation in patients. Feelings of competence can be achieved via the provision of information, but these feelings of competence need to be accompanied with a sense of autonomy to result in intrinsic motivation (Deci et al., 1999). Intrinsic motivation is potentially of crucial importance in the cure process, as individuals with intrinsic motivation have more interest, confidence, and self-esteem, and perform better in comparison to individuals who are externally controlled (e.g. merely following orders).

Socioeconomic status and doctor–patient relations

There are two ways doctor–patient relations can operate in the context of the educational gradient in self-rated health. Firstly, there might be an educational gradient in doctor–patient relations. Previous research has shown that doctor–patient relations vary with patients' SES. Doctor–patient communication is affected by patients' SES in the sense that doctors give lower-SES patients fewer positive socio-emotional responses, convey less information and directions, and make use of a more directive consulting style with less involvement in treatment decisions (Willems et al., 2005). Furthermore, doctors are able to identify the SES of their patients: Psychologists demonstrated that individuals can reliably identify the approximate income and SES of others within seconds (Kraus and Keltner, 2009). These differences appear to be driven both by patients' characteristics as well as the perceptions of doctors: High-SES patients communicate more actively and are able to elicit more information from their physicians, whereas doctors often misperceive the need for information of low-SES patients (Willems et al., 2005). Given the evidence that higher SES might be associated with a different, more egalitarian model of doctor–patient relations (Willems et al., 2005) and the evidence suggesting that a more egalitarian model of doctor–patient relations is associated with better wellbeing and health outcomes (Griffin et al., 2004; O'Connor et al., 2009; Ong et al., 1995; Weiner et al., 2013; Williams et al., 2000; McMillan et al., 2013; Rathert et al., 2013), the relationship between education and self-rated health might be at least partially due to differences in the doctor–patient relationship. In line with this reasoning, we formulate the following expectation:

H1: An egalitarian doctor–patient relationship reduces the size of the positive association between education and self-rated health.

Secondly, the models of the doctor–patient relationship might have differing effects on different educational groups. On the one hand, the lower-educated might particularly benefit from a more patient-

centered approach that takes their living conditions into account (e.g. Lutfey and Freese, 2005). In turn, the paternalistic approach might be particularly detrimental for the lower educated, as it does not pay attention to special needs they might have. Furthermore, if doctors give fewer positive socio-emotional responses, convey less information and directions to lower-educated patients than to higher-educated patients (Willems et al., 2005), then it follows by implication that lower-educated individuals benefit more from a more egalitarian style than higher-educated individuals. On the other hand, the higher educated can be expected to be less affected by different types of the doctor–patient relationship: The higher educated know better how to deal with directive high-status actors, as they have more self-confidence and other important psychological resources (Schnittker and McLeod, 2005) and more experience in negotiating and asking, irrespective of the style of the other party. Based on this, we further hypothesize:

H2: The doctor–patient relationship moderates the education–health relationship as follows: (a) An egalitarian doctor–patient relationship is associated with higher self-rated health, and that this is more strongly the case for the lower educated than for the higher educated, and (b) A paternalistic doctor–patient relationship is associated with lower self-rated health, and that is more strongly the case for the lower educated than for the higher educated.

Data and methods

Data

This study exploits the second round of the European Social Survey (ESS) (Jowell et al., 2007), a large-scale survey conducted in 26 countries in 2004/2005. This survey does not only comprise information on education and health, it also includes items on the doctor–patient relationship, making it a unique resource for our endeavor. We include 24 countries in our study, namely Austria, Belgium, Switzerland, the Czech Republic, Germany, Denmark, Estonia, Spain, Finland, Great Britain, Greece, Hungary, Italy, Ireland, Iceland, Luxembourg, The Netherlands, Norway, Poland, Portugal, Sweden, and Slovenia. Turkey and the Ukraine were excluded from the analysis, as first results revealed substantial cultural and socioeconomic differences to the other countries in the sample. The quality of the ESS data collection process is generally considered to be high (Koch et al., 2009), as in all countries random probability samples are drawn and face-to-face interviews are conducted, following standardized procedures and strict quality controls to ensure representativeness and comparability of the data. Response rates are on average 61.8%, ranging from 43.6% (France) to 78.8% (Greece). More detailed descriptions of the ESS, including detailed information on data collection and response rates for all countries are available on the ESS web site (www.europeansocialsurvey.org/).

Our study excludes respondents 25 years old or younger, as a substantial number of them has not yet finished education, which is one of our focal predictors in this study. Our sample size is further reduced by respondents who have missing values on any of the study variables. Descriptive statistics of the final sample of 31,189 respondents are provided in Table 1.

Health outcome: self-rated health. The outcome variable of this study is self-rated health. Self-rated health was measured on a five-point scale ranging from ‘very bad’ (0) to ‘very good’ (4). Self-rated health is a general assessment of one’s health status, not connected to any specific illness, but covering mental, physical, and social aspects of health (Idler et al., 1999). It has been shown to predict mortality and morbidity and has high test–retest reliability in a number of studies (Idler and Benyamini, 1997). Furthermore, this variable has been recommended by the World Health Organization for comparative research (De Bruin et al., 1996) and a large number of researchers have followed this advice (Huijts and Kraaykamp, 2012; Präg et al., 2016). Research has also shown that the associations between objective health indicators and self-perceived health are largely similar across countries (Bardage et al., 2005).

Table 1. Descriptive statistics (uncentered variables), $N = 31,198$.

	Mean	Std. dev.	Min.	Max.
Self-rated health	2.73	.91	0	4
<i>Education</i>				
Lower education	.34	—	0	1
Medium education	.42	—	0	1
High education	.24	—	0	1
<i>Egalitarian model</i>				
Easy to understand	3.30	1.15	1	5
Treat patients as equal	3.20	1.26	1	5
Discuss decisions	3.33	1.24	1	5
<i>Paternalistic model</i>				
Patients reluctant to ask questions	2.56	1.14	1	5
Keep truth from patients	2.26	1.07	1	5
Doctors deny mistakes	4.16	1.07	1	5
Age	50.83	15.50	26	101
Female	.54	—	0	1
Married/cohabiting	.70	—	0	1
Migrant	.08	—	0	1
Single mother	.07	—	0	1
<i>Children in the household</i>				
None	.54	—	0	1
One child	.20	—	0	1
Two children	.18	—	0	1
Three+ children	.08	—	0	1

Predictor variables

Doctor–patient relationship. The ESS questionnaire contains six items covering different aspects of the doctor–patient relationship. Respondents were asked how often statements applied to doctors in general. These six statements largely correspond to the two models of doctor–patient relations described previously and are reproduced in Table 2.

All answers are recorded on a five-point scale, ranging from ‘never or almost never’ (1) to ‘always or almost always’ (5). Zero-order correlations between the six items are low to moderate (.05 to .41, see Table 2). The items of the egalitarian model and the paternalistic model are positively correlated among one another and negatively correlated with the items of the other model, yet a factor analysis (shown in the online supplement) does not yield a clear two-dimensional structure. This points to the fact that the items capture distinct features of the doctor–patient relationship, but does not, however, allow for the creation of composite scores that would be more reliable than the single-item indicators we use for our analyses. Table 2 also shows that the bivariate relationships between the indicators of the doctor–patient relationship and education are substantively small (yet mostly statistically significant). For the items of the egalitarian model, the higher educated are more likely to agree with the statements that doctors are not difficult to understand and that doctors treat their patients as equal, yet there is no statistically significant difference between the educational groups for the statement that doctors discuss treatment decisions with their patients. The direction of these associations, however, is not always straightforward: For the items of the paternalistic model, agreement with the statement that doctors keep the whole truth from their patients is lower among the higher educated, yet for the statement that they are not willing to admit their mistakes, agreement is higher among the higher educated.

Table 2. Indicators of doctor–patient relationship by educational group and correlations between indicators.

	Average by educational group			F-test	Item correlations				
	Lower educated	Medium educated	Higher educated		1)	2)	3)	4)	5)
<i>Egalitarian model:</i>									
1) Doctors use words or phrases that their patients do not find difficult to understand (item reversed)	3.25	3.26	3.45	$F = 82.46$, $p = 0.01$	1.00				
2) Doctors treat patients as their equals	3.20	3.15	3.30	$F = 32.94$, $p = 0.03$.15	1.00			
3) Before doctors decide on a treatment, they discuss it with their patient	3.34	3.34	3.31	$F = 1.60$, $p = 0.46$.11	.41	1.00		
<i>Paternalistic model:</i>									
4) Patients are reluctant to ask their doctor all the questions they would like to ask	2.53	2.59	2.54	$F = 8.24$, $p = 0.11$	-.26	-.13	-.09	1.00	
5) Doctors keep the whole truth from their patients	2.31	2.28	2.14	$F = 62.45$, $p = 0.02$	-.18	-.11	-.08	.18	
6) Doctors are not willing to admit their mistakes to their patients (item reversed)	4.08	4.21	4.14	$F = 43.63$, $p = 0.02$	-.09	-.22	-.22	.05	.05

Notes: F-tests are based on OLS regression models that regress the respective doctor–patient relationship item on two education dummies to assess the correlation between education and the item. For all F-tests, the degrees of freedom are $N-1 = 31,195$ and $K-1 = 2$. Item correlations are Pearson product-moment correlation coefficients.

Education. We rely on education as our focal indicator of socioeconomic status. Based on the International Standard Classification of Education (ISCED), we distinguish between three educational groups: low (ISCED 0–2), medium (ISCED 3–4), and the higher educated (ISCED 5–6). There are several reasons for relying on education for our purposes. First, education reflects both individuals' material and non-material resources and their social status in a broad fashion. Second, the ISCED, with its high degree of cross-national standardization, allows meaningful comparison of educational groups across countries. Third, educational attainment is usually completed in early adulthood and remains stable across the life course, thus reducing the chance of reverse causation: unlike occupational status, educational degrees do not change even when an adult experiences a health shock (Elstad, 2004). Fourth, as the response rate for questions about education is usually much higher than for those about income, education is a comparatively straightforward indicator of SES. Finally, education is a meaningful measure for the SES of both men and women (Lahelma, 2010).

Control variables. As individual-level control variables, the models contain marital status/cohabitation (0 = not married or cohabiting with a partner, 1 = married/cohabiting with a partner), sex (0 = male, 1 = female), age, migrant status (1 = yes, 0 = no), being a lone mother (1 = yes, 0 = no), and number of children (ranging from no children to three and more children, as a set of dummy variables).

Method and modeling strategy. For analyzing a clustered dataset, multilevel (random effects) modeling is often recommended (Snijders and Boskers, 2012). Given the relatively low number of countries (Bryan and Jenkins, 2016; Mills and Präg, 2016) and our prime interest in the relationship between variables at the individual level, we present results as obtained from ordinary least squares (OLS) regression models. These models account for the country variance by including country fixed effects, and correct for the clustering in the data by reporting cluster-robust standard errors. We have centered our predictor variables at their respective country means, allowing us to obtain unbiased estimates of the within-country regressions (Allison, 2009). Alternative estimation strategies (which lead to the same substantive conclusions) are reported in the online supplement.

Table 3. Self-rated health regressed on education and doctor–patient relationship, OLS regression.

	(1)	(2)	(3)
<i>Education (ref. low education)</i>			
Medium education	0.192*** (0.019)	0.187*** (0.019)	0.191*** (0.020)
High education	0.358*** (0.021)	0.349*** (0.021)	0.353*** (0.022)
<i>Egalitarian model</i>			
Easy to understand		0.022*** (0.005)	
Treat patients as equal		0.026*** (0.005)	
Discuss decisions		-0.002 (0.004)	
<i>Paternalistic model</i>			
Patients reluctant to ask questions			-0.004 (0.004)
Keep truth from patients			-0.026*** (0.006)
Doctors deny mistakes			-0.013 (0.007)
Age	-0.018*** (0.001)	-0.018*** (0.001)	-0.017*** (0.001)
Female (ref. male)	-0.055** (0.018)	-0.054** (0.019)	-0.055** (0.018)
Married/cohabiting (ref. not m./c.)	0.065*** (0.017)	0.063*** (0.017)	0.065*** (0.017)
Single mother (ref. not s.m.)	-0.033 (0.032)	-0.032 (0.032)	-0.034 (0.033)
Migrant (ref. not migrant)	-0.063* (0.025)	-0.059* (0.026)	-0.061* (0.026)
<i>Children in household (ref. none)</i>			
One child in household	0.023 (0.013)	0.023 (0.013)	0.024 (0.013)
Two children in household	0.087*** (0.014)	0.088*** (0.013)	0.088*** (0.013)
Three+ children in household	0.069** (0.025)	0.071*** (0.025)	0.070** (0.025)
Intercept	2.865*** (0.014)	2.869*** (0.015)	2.868*** (0.014)
Country dummies included	YES	YES	YES
N countries	24	24	24
N individuals	31,198	31,198	31,198

Robust standard errors in parentheses. Coefficients for country dummies not displayed.

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Results

Model 1 reported in Table 3 confirms the presence of an educational gradient in health. Respondents with medium and high education report significantly better health than lower educated respondents. A




Figure 1. Educational gradient in health by country, before and after adjusting for egalitarian doctor–patient relationship, based on country-specific ordinary least squares (OLS) regression models. Note: All models control for age, sex, marital status, migrant status, single motherhood, and number of children.

Wald test further confirms that the coefficients for medium and highly educated respondents differ from one another ($F(1, 23) = 196.82, p = .000$), indicating that there is not a single dividing line between the lower educated and anyone with higher education, but that the health benefits of education increment gradually. The control variables in the model behave as can be expected from the literature: Older respondents and females report worse health, and those with a partner report better health. A comparison of the coefficients shows that apart from age, education is the most important predictor of health.

The indicators of an egalitarian doctor–patient relationship are entered in Model 2 of Table 3. Reporting that doctors are easy to understand and that they treat their patients as equals is associated with significantly better health, whereas the third indicator—whether doctors discuss their decisions with patients—is unrelated to self-reported health. Hypothesis 1, which had posited that a more egalitarian doctor–patient relationship attenuates the educational gradient in health, has to be rejected. When comparing the coefficients for education as obtained from Model 1 and Model 2, barely any difference can be detected, indicating that by and large, more egalitarian doctor–patient relations do not reduce educational differences in self-rated health. A reason for this is that, as shown before, the educational differences in the doctor–patient relationship are comparatively small.

In order to assess differences between countries for this finding, we have run separate regression models by country and report the education coefficients before and after adjusting for the egalitarian doctor–patient relationship in Figure 1. The size of the educational gradient in health varies considerably across countries, but closer inspection reveals that differences between the adjusted and unadjusted coefficients

are negligible, indicating that the finding of no meaningful reduction in the educational gradient in health in case of a more egalitarian doctor–patient relationship holds for all countries under study.

For the indicators of a paternalistic doctor–patient relationship, Model 3 of Table 3 indicates direct negative associations with self-rated health for one of the three indicators. When patients report that doctors keep the truth from patients, it goes along with statistically significant poorer self-rated health. Reporting about a reluctance to ask questions and about doctors denying mistakes have negative effects that are not statistically different from zero. The size of the educational gradient in health is, however, unaffected by including indicators of a paternalistic doctor–patient relationship, as can be seen from comparing the coefficients for education between Models 1 and 3.

Table 4 reports the interactions between indicators of the doctor–patient relationship and education. For two of the six indicators, we can detect a statistically significant interaction with education, namely when respondents report doctors to be easy to understand (Model 4 of Table 4) and when patients are reluctant to ask questions (Model 7 of Table 4). To facilitate interpretation, we have plotted the interactions in Figure 2. The upper panel of Figure 2 plots the interaction of Model 4 from Table 4. It shows that whether doctors are easy to understand plays virtually no role for the self-rated health of the higher and medium educated; the slopes of their predicted lines are virtually flat over the continuum of how easy doctors are to understand. For the group of the lower educated, however, we can see a marked increase in self-rated health as the reports of doctors being easy to understand increase. The plot provides evidence that it is the lower educated who benefit from a more egalitarian doctor–patient relationship, giving support for Hypothesis 2a. The lower panel of Figure 2 plots the interaction of Model 7 of Table 4. Again, for the higher educated and for those with medium education, it shows that there is virtually no association of self-rated health with reporting that patients are reluctant to ask questions, as the slopes of the predictions are flat. However, for the lower educated, we can see a falling slope as the agreement to the statement that patients are reluctant to ask questions gets stronger. Substantively, this means that a reluctance to answer questions is irrelevant for those with educational degrees classified as medium and high, but for the lower educated, such a reported reluctance is associated with worse self-rated health. This gives support for Hypothesis 2b.

Sensitivity analyses. The online supplement to this article shows the following robustness checks: We were able to replicate our findings from pooled OLS models with country dummies and robust standard errors by using random effects models without obtaining substantially different results. We also ran analyses making use of the weights calculated by the data providers, leading to substantively identical findings. We furthermore replicated our analyses dropping one country at a time or dropping all Eastern European countries altogether, yielding substantively similar findings.

Discussion

How doctor–patient relations affect the educational gradient in health is largely unknown. Drawing on a large European general population survey, our study generated several new insights into this field.

First, our analyses showed that two different models of the doctor–patient relationship, namely the traditional paternalistic model and the more egalitarian ‘patient-centered’ model, are directly associated with self-rated health. Reports that doctors are easy to understand and treat their patients as equals go along with better self-rated health. Conversely, reports that doctors do not tell the whole truth are associated with worse self-ratings of health. This confirms findings from previous studies, which have suggested that a more egalitarian doctor–patient relationship is associated with better wellbeing and health outcomes (Griffin et al., 2004; O’Connor et al., 2009; Ong et al., 1995; Stewart, 1995a; Stewart, 1995b; Weiner et al., 2013; Williams et al., 2000).

Second, the different models of the doctor–patient relationship do not mediate the educational gradient in self-rated health. Despite the evidence for differential treatment of patients by their doctors according to their SES in the literature (Willems et al., 2005), our data show that differences in

Table 4. Self-rated health regressed on education and doctor–patient relationship including interaction terms, OLS regression.

	(4)	(5)	(6)	(7)	(8)	(9)
<i>Education (ref. lower)</i>						
Medium education	0.186*** (0.020)	0.187*** (0.020)	0.188*** (0.019)	0.191*** (0.020)	0.190*** (0.020)	0.190*** (0.020)
High education	0.349*** (0.022)	0.349*** (0.021)	0.349*** (0.022)	0.354*** (0.022)	0.353*** (0.021)	0.354*** (0.022)
<i>Egalitarian model</i>						
Easy to understand	0.039*** (0.010)	0.022*** (0.005)	0.022*** (0.005)			
Medium ed. X Easy to understand	-0.024* (0.011)					
High ed. X Easy to understand	-0.036** (0.012)					
Treats patients as equal	0.026*** (0.005)	0.032** (0.009)	0.026*** (0.005)			
Medium ed. X Treats patients as equal		-0.009 (0.008)				
High ed. X Treats patients as equal		-0.011 (0.011)				
Doctor discuss decisions	-0.002 (0.004)	-0.002 (0.004)	0.002 (0.008)			
Medium ed. X Discuss decisions			-0.005 (0.011)			
High ed. X Discuss Decisions			-0.008 (0.010)			
<i>Paternalistic model</i>						
Patients reluctant to ask questions				-0.020** (0.006)	-0.004 (0.004)	-0.004 (0.004)
Medium ed. X Patients reluctant to ask questions				0.023* (0.008)		
High ed. X Patients reluctant to ask questions				0.029** (0.009)		
Keep truth from patients				-0.026*** (0.005)	-0.031** (0.011)	-0.026*** (0.005)
Medium ed. X Keep truth from patients					0.005 (0.013)	
High ed. X Keep truth from from patients					0.011 (0.0105)	
Doctors deny mistakes				-0.013 (0.007)	-0.013 (0.007)	-0.011 (0.010)
Medium ed. X Doctors deny mistakes						0.003 (0.012)
High ed. X Doctors deny mistakes						-0.011 (0.014)
Age	-0.018*** (0.001)	-0.018*** (0.0014)	-0.018*** (0.001)	-0.018*** (0.001)	-0.018*** (0.001)	-0.018*** (0.001)
Female (ref. male)	-0.054** (0.019)	-0.054** (0.019)	-0.054** (0.019)	-0.056** (0.018)	-0.055** (0.018)	-0.055** (0.018)

(continued)

Table 4. (continued)

	(4)	(5)	(6)	(7)	(8)	(9)
Married/cohabiting (ref. not m./c.)	0.063*** (0.017)	0.063*** (0.017)	0.063*** (0.017)	0.065*** (0.017)	0.064*** (0.017)	0.065*** (0.017)
Single mother (ref. not single mother)	-0.031 (0.032)	-0.032 (0.032)	-0.032 (0.032)	-0.033 (0.032)	-0.034 (0.033)	-0.034 (0.033)
Migrant (ref. not migrant)	-0.059* (0.027)	-0.059* (0.025)	-0.059* (0.026)	-0.061* (0.027)	-0.061* (0.026)	-0.061* (0.026)
Children in household (ref. none)						
One child in household	0.023 (0.013)	0.023 (0.013)	0.023 (0.013)	0.024 (0.013)	0.023 (0.013)	0.024 (0.013)
Two children in household	0.088*** (0.013)	0.088*** (0.013)	0.088*** (0.013)	0.087*** (0.013)	0.088*** (0.013)	0.088*** (0.013)
Three+ children in household	0.071*** (0.019)	0.071*** (0.019)	0.071*** (0.019)	0.070** (0.020)	0.070** (0.019)	0.070** (0.019)
Constant	2.872*** (0.015)	2.870*** (0.015)	2.869*** (0.015)	2.867*** (0.014)	2.869*** (0.014)	2.868*** (0.014)
Country dummies included	YES	YES	YES	YES	YES	YES
N countries	24	24	24	24	24	24
N individuals	31,198	31,198	31,198	31,198	31,198	31,198

Robust standard errors in parentheses.

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

doctor–patient relations are substantively small. Accounting for those differences does not decrease the education–self-rated health association.

Third, when doctors are more paternalistic and less egalitarian, educational differences in self-rated health can indeed be larger. We found a number of interactions between education and indicators of the different models of doctor–patient relations. For the lower educated, reporting that doctors are easy to understand—an indicator of the egalitarian model of doctor–patient relations—is associated with better self-rated health, while for those with medium and high levels of education this does not translate into better health. Also, reports that patients are reluctant to ask questions—an indicator of the paternalistic model of the doctor–patient relationship—is negatively associated with self-rated health for the lower educated, but for the medium and higher educated there is no negative association. This finding has important policy implications, as measures to improve doctor–patient relations can have a positive impact on overall population health, and can be expected to compress the social gradient in health.

As has been argued prominently by Marmot (2015), doctors should engage with the social background of their patients. Marmot encourages doctors to recognize (and improve) the social context (e.g. homelessness) in which poor health develops when treating patients, and suggests that doctors should cooperate with other community actors such as social workers, and engage in advocacy to improve the living conditions of their patients. Yet rather than expanding the responsibilities of doctors, our findings build, however, on the core business of physicians and show how doctors' bedside manners go along with particular health effects and the educational gradient in health. We believe that this can have an easier and greater impact on population health and health inequalities rather than overburdening doctors with new tasks and new arenas for action.

Furthermore, future research should engage with the effects different health care systems have on doctor–patient relations. While our study focused on the micro-level processes between doctors and patients that govern the educational gradient in health, considering differences between health care systems would be a promising next step (Blom et al., 2016; Wendt, 2009), which, however, needs




Figure 2. Upper panel: Plot of ‘doctors easy to understand’ by education interaction. Lower panel: Plot of ‘Patients reluctant to ask questions’ by education interaction.

Note: Calculations based on Model 4 of Table 4 (upper panel) and Model of Table 4 (lower panel).

careful theorizing to disentangle the mechanisms that govern the macro–micro link between health care systems and doctor–patient relations. As a first step, our study already documented important country differences in the strength of the education–self-rated health relationship, yet future research is necessary. Exploiting differences in health care system financing would also allow further investigation of any effects of a marketization of health care.

We acknowledge that our conclusions are based on a cross-sectional, observational study. Our results are associational and cannot make any claims about causality. Despite these caveats, we believe that our study gives important impulses for future research under improved methodological conditions. Drawing on longitudinal data would help to rule out causality issues and could yield greater insights as to how the change in doctor–patient relations occurred over time. Also, using validated and more comprehensive indicators of the different models of the doctor–patient relationship could strengthen the case for the arguments laid out in this article. Additional health outcomes that do not rely on self-reports would help overcome any biases that might stem from cross-national or individual (e.g. education-related)

heterogeneity (Jylhä, 2009; Manderbacka, 1998). However, our results proved to be robust to a multitude of model specifications and were not sensitive to the selection of countries, making it unlikely that institutional or cultural specificities (Hall and Lamont, 2009; Wendt et al., 2009) are driving our findings. Notwithstanding these limitations, our study was the first to show that in a large European population sample the move towards a more egalitarian model of doctor–patient relations can have positive effects on reducing social inequality in health, which opens up avenues for further research on the social gradient in health.

Acknowledgements

The authors wish to thank the four anonymous reviewers as well as the editors for helpful comments.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the European Union's Seventh Framework Program (FP7/2007–13, grant number 320116) and the European Research Council (Consolidator Grant 'Sociogenome', grant number 615603).

Supplemental material

The online supplement is available at <http://asj.sagepub.com/supplemental>

References

- Allison PD (2009) *Fixed Effects Regression Models*. Thousand Oaks, CA: Sage.
- Bardage C, Pluijm SMF, Pedersen NL, et al. (2005) Self-rated health among older adults. A cross-national comparison. *European Journal of Ageing* 2(2): 149–158.
- Barry MJ and Edgman-Levitan S (2012) Shared decision making: The pinnacle of patient-centered care. *New England Journal of Medicine* 366(9): 780–781.
- Beck U (1992) *Risk Society: Towards a New Modernity*. London: Sage.
- Beckfield J, Bambra C, Eikemo TA, et al. (2015) An institutional theory of welfare state effects on the distribution of population health. *Social Theory and Health* 13(3–4): 227–244.
- Blom N, Huijts T and Kraaykamp G (2016) Ethnic health inequalities in Europe: The moderating and amplifying role of healthcare system characteristics. *Social Science and Medicine* 158: 43–51.
- Bryan ML and Jenkins SP (2016) Multilevel modeling of country effects: A cautionary tale. *European Sociological Review* 32(1): 3–22.
- Bultman DC and Svarstad BL (2000) Effects of physician communication style on client medication beliefs and adherence with antidepressant treatment. *Patient Education and Counseling* 40(2): 173–185.
- Byrne PS and Long BEL (1976) *Doctors Talking to Patients: A Study of the Verbal Behaviour of General Practitioners Consulting in Their Surgeries*. London: HMSO.
- Campbell SM, Roland MO and Buetow SA (2000) Defining Quality of Care. *Social Science and Medicine* 51(11): 1611–1625.
- Charmaz K and Rosenfeld D (2010) Chronic illness. In: Cockerham WC (ed) *The New Blackwell Companion to Medical Sociology*. Malden, MA: Wiley-Blackwell, 312–333.
- Cialdini RB and Giskevicius V (2010) Social influence. In: Baumeister RF and Finkel EJ (eds) *Advanced Social Psychology: The State of the Science*. New York: Oxford University Press, 385–417.
- Coleman J, Katz E and Menzel H (1957) The diffusion of an innovation among physicians. *Sociometry* 20(4): 253–270.

- Crow R, Gage H, Hampson S, et al. (2003) The measurement of satisfaction with healthcare: Implications for practice from a systematic review of the literature. *Health Technology Assessment* 32(6): 1–244.
- De Bruin A, Picavet HSJ and Nossikov A (1996) *Health Interview Surveys: Towards International Harmonization of Methods and Instruments*. Copenhagen: WHO Regional Publications.
- Deci EL, Koestner R and Ryan RM (1999) A meta-analytic review of experiments examining the effects of extrinsic rewards on intrinsic motivation. *Psychological Bulletin* 125(6): 627–668.
- Department of Health (2001) *The Expert Patient: A New Approach to Chronic Disease Management for the 21st Century*. London: HMSO.
- Djundeva M, Mills M, Wittek R, et al. (2015) Receiving instrumental support in late parent–child relationships and parental depression. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences* 70(6): 981–994.
- Ellwardt L, Peter S, Präg P, et al. (2014) Social contacts of older people in 27 European countries: The role of welfare spending and economic inequality. *European Sociological Review* 30(4): 413–430.
- Elo IT (2009) Social class differentials in health and mortality: Patterns and explanations in comparative perspective. *Annual Review of Sociology* 35: 553–572.
- Elstad JI (2004) Health and status attainment: Effects of health on occupational achievement among employed Norwegian men. *Acta Sociologica* 47(2): 127–140.
- Fiscella K, Franks P, Gold MR, et al. (2000) Inequality in quality: Addressing socioeconomic, racial, and ethnic disparities in health care. *Journal of the American Medical Association* 283(19): 2579–2584.
- Gesthuizen M, Huijts T and Kraaykamp G (2012) Explaining health marginalisation of the lower educated The role of cross-national variations in health expenditure and labour market conditions. *Sociology of Health and Illness* 34(4): 591–607.
- Griffin SJ, Kinmonth A-L, Veltman MWM, et al. (2004) Effect on health-related outcomes of interventions to alter the interaction between patients and practitioners: A systematic review of trials. *Annals of Family Medicine* 2(6): 595–608.
- Hall PA and Lamont M (2009) *Successful Societies: How Institutions and Culture Affect Health*. Cambridge: Cambridge University Press.
- Henderson LJ (1935) Physician and patient as a social system. *New England Journal of Medicine* 212(18): 819–823.
- Heritage J and Maynard DW (2011) After 30 years, problems and prospects in the study of doctor–patient interaction. In: Pescosolido BA, Martin JK, McLeod JD, et al. (eds) *Handbook of the Sociology of Health, Illness, and Healing: A Blueprint for the 21st Century*. New York: Springer, 323–342.
- Huijts T and Kraaykamp G (2012) Formal and informal social capital and self-rated health in Europe: A new test of accumulation and compensation mechanisms using a multi-level perspective. *Acta Sociologica* 55(2): 143–158.
- Idler EL and Benyamini Y (1997) Self-rated health and mortality: A review of twenty-seven community studies. *Journal of Health and Social Behavior* 38(1): 21–37.
- Idler EL, Hudson SV and Leventhal H (1999) The meanings of self-ratings of health: A qualitative and quantitative approach. *Research on Aging* 21(3): 458–476.
- Jowell R, Roberts C, Fitzgerald R, et al. (2007) *Measuring Attitudes Cross-Nationally: Lessons from the European Social Survey*. Los Angeles, CA: Sage.
- Judson TJ, Detsky AS and Press MJ (2013) Encouraging patients to ask questions: How to overcome “white-coat silence”. *Journal of the American Medical Association* 309(22): 2325–2326.
- Jylhä M (2009) What is self-rated health and why does it predict mortality? Towards a unified conceptual model. *Social Science and Medicine* 69(3): 307–316.

- Kaba R and Sooriakumaran P (2007) The evolution of the doctor–patient relationship. *International Journal of Surgery* 5(1): 57–65.
- Koch A, Blom AG, Stoop I, et al. (2009) Data collection quality assurance in cross-national surveys: The example of the ESS. *Methods Data Analyses* 3(2): 219–247.
- Kraus MW and Keltner D (2009) Signs of socioeconomic status: A thin-slicing approach. *Psychological Science* 20(1): 99–106.
- Lahelma E (2010) Health and social stratification. In: Cockerham WC (ed) *The New Blackwell Companion to Medical Sociology*. Hoboken, NJ: Wiley-Blackwell, 69–96.
- Lutfey K and Freese J (2005) Toward some fundamentals of fundamental causality: Socioeconomic status and health in the routine clinic visit for diabetes. *American Journal of Sociology* 110(5): 1326–1372.
- Mackenbach JP, Stirbu I, Roskam A-JR, et al. (2008) Socioeconomic inequalities in health in 22 European countries. *New England Journal of Medicine* 358(23): 2468–2481.
- Manderbacka K (1998) Examining what self-rated health question is understood to mean by respondents. *Scandinavian Journal of Public Health* 26(2): 145–153.
- Marmot MG (2015) *The Health Gap: The Challenge of an Unequal World*. London: Bloomsbury.
- McMillan SS, Kendall E, Sav A, et al. (2013) Patient-centered approaches to health care: A systematic review of randomized controlled trials. *Medical Care Research and Review* 70(6): 567–596.
- Mills MC and Präg P (2016) Methodological advances in cross-national research: Multilevel challenges and solutions. *European Sociological Review* 32(1): 1–2.
- Mirowsky J and Ross CE (2003) *Education, Social Status, and Health*. New York: de Gruyter.
- Neuberger J (2000) The educated patient: New challenges for the medical profession. *Journal of Internal Medicine* 247(1): 6–10.
- O'Connor AM, Bennett CL, Stacey D, et al. (2009) *Decision Aids for People Facing Health Treatment or Screening Decisions (Review)*, London: Cochrane Collaboration.
- Ong LML, De Haes HCJM, Hoos AM, et al. (1995) Doctor–patient communication: a review of the literature. *Social Science and Medicine* 40(7): 903–918.
- Ormel J, Lindenberg S, Steverink N, et al. (1997) Quality of life and social production functions: A framework for understanding health effects. *Social Science and Medicine* 45(7): 1051–1063.
- Parsons T (1951) *The Social System*. New York: Free Press.
- Peck BM and Conner S (2011) Talking with me or talking at me? The impact of status characteristics on doctor–patient interaction. *Sociological Perspectives* 54(4): 547–567.
- Präg P, Mills MC and Wittek R (2014) Income and income inequality as social determinants of health: Do social comparisons play a role? *European Sociological Review* 30(2): 218–229.
- Präg P, Mills MC and Wittek R (2016) Subjective socioeconomic status and health in cross-national comparison. *Social Science and Medicine* 149: 84–92.
- Rathert C, Wyrwich MD and Boren SA (2013) Patient-centered care and outcomes: A systematic review of the literature. *Medical Care Research and Review* 70(4): 351–379.
- Ryan RM and Deci EL (2000) Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. *American Psychologist* 55(1): 68–78.
- Schnittker J and McLeod JD (2005) The social psychology of health disparities. *Annual Review of Sociology* 31: 75–103.
- Snijders T and Boskers R (2012) *Multilevel Analysis. An Introduction to Basic and Advanced Multilevel Modeling*. London: Sage.
- Stavropoulou C (2011) Non-adherence to medication and doctor–patient relationship. Evidence from a European survey. *Patient Education and Counseling* 83(1): 7–13.

- Stavropoulou C (2012) The doctor–patient relationship: A review of the theory and policy implications. In: McGuire A and Costa-Font J (eds) *The London School of Economics Companion To Health Policy*. Cheltenham, UK: Edward Elgar, 314–326.
- Steinhart B (2002) Patient autonomy. Evolution of the doctor–patient relationship. *Haemophilia* 8(3): 441–446.
- Stewart MA (1995a) Effective physician–patient communication and health outcomes: A review. *Canadian Medical Association Journal* 152(9): 1423–1433.
- Stewart MA (1995b) Studies of health outcome and patient-centered communication. In: Stewart MA, Brown JB, Weston WW, et al. (eds) *Patient-Centered Medicine: Transforming the Clinical Method*. Thousand Oaks, CA: Sage, 185–190.
- Weiner SJ, Schwartz A, Sharma G, et al. (2013) Patient-centered decision making and health care outcomes: An observational study. *Annals of Internal Medicine* 158(8): 573–580.
- Wendt C (2009) Mapping European healthcare systems: A comparative analysis of financing, service provision, and access to healthcare. *Journal of European Social Policy* 19(5): 432–445.
- Wendt C, Frisina L and Rothgang H (2009) Healthcare system types. A conceptual framework for comparison. *Social Policy and Administration* 43(1): 70–90. doi: 10.1111/j.1467-9515.2008.00647.x
- White RW (1959) Motivation reconsidered: The concept of competence. *Psychological Review* 66(5): 297–333.
- Willems S, De Maesschalck S, Deveugele M, et al. (2005) Socio-economic status of the patient and doctor–patient communication: Does it make a difference? *Patient Education and Counseling* 56(2): 139–146.
- Williams GC, Frankel RM, Campbell TL, et al. (2000) Research on relationship-centered care and healthcare outcomes from the Rochester biopsychosocial program: A self-determination theory integration. *Families, Systems, and Health* 18(1): 79–90.

Author biographies

Patrick Präg is a postdoctoral researcher at the University of Oxford and a Research Fellow at Nuffield College. He obtained his PhD from the Interuniversity Center for Social Science Theory and Methodology, University of Groningen. His research interests comprise health and wellbeing, social stratification, and work and family reconciliation. Recent articles have appeared in *Social Science and Medicine*, *Journal of Vocational Behavior*, and *European Sociological Review*.

Rafael Wittek is Professor of Theoretical Sociology at the University of Groningen and the Interuniversity Center for Social Science Theory and Methodology (ICS). His main research areas are social theory, social networks, the sociology of organizations, and the sociology of aging. Recent articles have appeared in *Social Science Research*, *Social Networks*, and *European Sociological Review*.

Melinda Mills is Nuffield Professor of Sociology at the University of Oxford and Professorial Fellow at Nuffield College. Her research focuses on family formation, fertility and partnerships, the life course, flexible work arrangements, globalization, and event history methods. She currently leads a large-scale sociogenetics project linking sociological and genetic research on the family. Recent articles have appeared in *Nature Genetics*, *Social Forces*, and *Journal of Marriage and Family*.