

University of Groningen

Parental and endosymbiont effects on sex determination in haplodiploid wasps

Geuverink, Elzemiek

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Geuverink, E. (2017). *Parental and endosymbiont effects on sex determination in haplodiploid wasps: Who is in control?* [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Bibliography

- Aljanabi, S.M. & Martinez, I. (1997) Universal and rapid salt-extraction of high quality genomic DNA for PCR-based techniques. *Nucleic Acids Research*, **25**, 4692–4693.
- Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., *et al.* (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic acids research*, **25**, 3389–3402.
- Amrein, H., Hedley, M.L. & Maniatis, T. (1994) The role of specific protein-RNA and protein-protein interactions in positive and negative control of pre-mRNA splicing by *Transformer 2*. *Cell*, **76**, 735–746.
- Amrein, H., Maniatis, T. & Nöthiger, R. (1990) Alternatively spliced transcripts of the sex-determining gene *tra-2* of *Drosophila* encode functional proteins of different size. *The EMBO journal*, **9**, 3619–3629.
- Asplen, M.K., Whitfield, J.B., Boer, J.G. DE & Heimpel, G.E. (2009) Ancestral state reconstruction analysis of hymenopteran sex determination mechanisms. *Journal of Evolutionary Biology*, **22**, 1762–1769.
- Baker, B.S. & Wolfner, M.F. (1988) A molecular analysis of *doublesex*, a bifunctional gene that controls both male and female sexual differentiation in *Drosophila melanogaster*. *Genes & Development*, **2**, 477–489.
- Bertone, M.A., Courtney, G.W. & Wiegmann, B.M. (2008) Phylogenetics and temporal diversification of the earliest true flies (Insecta: Diptera) based on multiple nuclear genes. *Systematic Entomology*, **33**, 668–687.
- Beukeboom, L.W. (1995) Sex determination in Hymenoptera: A need for genetic and molecular studies. *BioEssays*, **17**, 813–817.
- Beukeboom, L.W. (2001) Single-locus complementary sex determination in the ichneumonid *Venturia canescens* (Gravenhorst) (Hymenoptera). *Netherlands Journal of Zoology*, **51**, 1–15.
- Beukeboom, L.W. (2012) Microbial manipulation of host sex determination. *BioEssays*, **34**, 484–488.
- Beukeboom, L.W., Ellers, J. & Alphen, J.J. Van. (2000) Absence of single-locus complementary sex determination in the braconid wasps *Asobara tabida* and *Alysia manducator*. *Heredity*, **84**, 29–36.
- Beukeboom, L.W. & Kamping, A. (2006) No patrigenes required for femaleness in the haplodiploid wasp *Nasonia vitripennis*. *Genetics*, **172**, 981–989.
- Beukeboom, L.W., Kamping, A., Louter, M., Pijnacker, L.P., Katju, V., Ferree, P.M., *et al.* (2007a) Haploid females in the parasitic wasp *Nasonia vitripennis*. *Science*, **315**, 206.

- Beukeboom, L.W., Kamping, A. & Zande, L. van de. (2007b) Sex determination in the haplodiploid wasp *Nasonia vitripennis* (Hymenoptera: Chalcidoidea): a critical consideration of models and evidence. *Seminars in Cell & Developmental Biology*, **18**, 371–378.
- Beukeboom, L.W. & Perrin, N. (2014) *The evolution of sex determination*. Oxford Univ Press.
- Beukeboom, L.W. & Pijnacker, L.P. (2000) Automictic parthenogenesis in the parasitoid *Venturia canescens* (Hymenoptera: Ichneumonidae) revisited. *Genome*, **43**, 939–944.
- Beukeboom, L.W. & Zande, L. van de. (2010) Genetics of sex determination in the haplodiploid wasp *Nasonia vitripennis* (Hymenoptera: Chalcidoidea). *Journal of Genetics*, **89**, 333–339.
- Beye, M., Hasselmann, M., Fondrk, M.K., Page, R.E. & Omholt, S.W. (2003) The gene *csd* is the primary signal for sexual development in the honeybee and encodes an SR-type protein. *Cell*, **114**, 419–429.
- Biémont, C. & Bouletreau, M. (1980) Hybridization and inbreeding effects on genome coadaptation in a haplo-diploid hymenoptera: *Cothonaspis bouleardi* (Eucoilidae). *Experientia*, **36**, 45–47.
- Boer, J.G. de, Kuijper, B., Heimpel, G.E. & Beukeboom, L.W. (2012) Sex determination meltdown upon biological control introduction of the parasitoid *Cotesia rubecula*? *Evolutionary Applications*, **5**, 444–454.
- Boer, J.G. de, Ode, P.J., Rendahl, A.K., Vet, L.E.M., Whitfield, J.B. & Heimpel, G.E. (2008) Experimental support for multiple-locus complementary sex determination in the parasitoid *Cotesia vestalis*. *Genetics*, **180**, 1525–1535.
- Boggs, R.T., Gregor, P., Idriss, S., Belote, J.M. & McKeown, M. (1987) Regulation of sexual differentiation in *D. melanogaster* via alternative splicing of RNA from the *transformer* gene. *Cell*, **50**, 739–747.
- Bonasio, R., Li, Q., Lian, J., Mutti, N.S., Jin, L., Zhao, H., *et al.* (2012) Genome-wide and caste-specific DNA methylomes of the ants *Camponotus floridanus* and *Harpegnathos saltator*. *Current Biology*, **22**, 1755–1764.
- Bonasio, R., Zhang, G., Ye, C., Mutti, N.S., Fang, X., Qin, N., *et al.* (2010) Genomic comparison of the ants *Camponotus floridanus* and *Harpegnathos saltator*. *Science*, **329**, 1068–1071.
- Bopp, D. (2010) About females and males: continuity and discontinuity in flies. *Journal of Genetics*, **89**, 315–323.
- Bopp, D., Saccone, G. & Beye, M. (2014) Sex Determination in Insects: Variations on a Common Theme. *Sexual Development*, **8**, 20–28.
- Bull, J.J. (1983) *Evolution of sex determining mechanisms*. Benjamin/Cummings Pub. Co., Menlo

Park, California.

- Burghardt, G., Hediger, M., Siegenthaler, C., Moser, M., Dübendorfer, A. & Bopp, D. (2005) The *transformer2* gene in *Musca domestica* is required for selecting and maintaining the female pathway of development. *Development Genes and Evolution*, **215**, 165–176.
- Burtis, K.C. & Baker, B.S. (1989) *Drosophila doublesex* gene controls somatic sexual differentiation by producing alternatively spliced mRNAs encoding related sex-specific polypeptides. *Cell*, **56**, 997–1010.
- Carabajal Paladino, L., Muntaabski, I., Lanzavecchia, S., Bagousse-Pinguet, Y. Le, Viscarret, M., Juri, M., *et al.* (2015) Complementary sex determination in the parasitic wasp *Diachasmimorpha longicaudata*. *PLoS ONE*, **10**, e0119619.
- Carmell, M.A., Xuan, Z., Zhang, M.Q. & Hannon, G.J. (2002) The Argonaute family: tentacles that reach into RNAi, developmental control, stem cell maintenance, and tumorigenesis. *Genes & Development*, **16**, 2733–2742.
- Carton, Y., Bouletreau, M., Alphen, J. Van & Lenteren, J. Van. (1986) The *Drosophila* parasitic wasps. In *The Genetics and Biology of Drosophila* (ed. by Ashburner, M., Carson, H. & Thompson, J.). Academic Press: London, pp. 347–394.
- Chen, P., Xu, S.L., Zhou, W., Guo, X.G., Wang, C.L., Wang, D.L., *et al.* (2014) Cloning and expression analysis of a *transformer* gene in *Daphnia pulex* during different reproduction stages. *Animal Reproduction Science*, **146**, 227–237.
- Chen, S.-L., Dai, S.-M., Lu, K.-H. & Chang, C. (2008) Female-specific *doublesex* dsRNA interrupts yolk protein gene expression and reproductive ability in oriental fruit fly, *Bactrocera dorsalis* (Hendel). *Insect Biochemistry and Molecular Biology*, **38**, 155–165.
- Cho, S., Huang, Z.Y. & Zhang, J. (2007) Sex-specific splicing of the honeybee *doublesex* gene reveals 300 million years of evolution at the bottom of the insect sex-determination pathway. *Genetics*, **177**, 1733–1741.
- Concha, C., Li, F. & Scott, M.J. (2010) Conservation and sex-specific splicing of the *doublesex* gene in the economically important pest species *Lucilia cuprina*. *Journal of Genetics*, **89**, 279–285.
- Concha, C. & Scott, M.J. (2009) Sexual development in *Lucilia cuprina* (Diptera, Calliphoridae) is controlled by the *transformer* gene. *Genetics*, **182**, 785–798.
- Cook, J.M. (1993) Sex determination in the Hymenoptera : a review of models and evidence. *Heredity*, **71**, 421–435.
- Cowan, D.P. & Stahlhut, J.K. (2004) Functionally reproductive diploid and haploid males in an inbreeding hymenopteran with complementary sex determination. *Proceedings of the*

- National Academy of Sciences of the United States of America*, **101**, 10374–10379.
- Crozier, R.H. (1971) Heterozygosity and sex determination in haplo-diploidy. *The American Naturalist*, **105**, 399–412.
- Cunha, A.B. Da & Kerr, W.E. (1957) A genetical theory to explain sex-determination by arrhenotokous parthenogenesis. *Forma et Functio*, **1**, 33–36.
- Davies, N.J. & Tauber, E. (2015) WaspAtlas: a *Nasonia vitripennis* gene database and analysis platform. *Database*, **2015**, bav103.
- Dedeine, F., Vavre, F., Fleury, F., Loppin, B., Hochberg, M.E. & Bouletreau, M. (2001) Removing symbiotic *Wolbachia* bacteria specifically inhibits oogenesis in a parasitic wasp. *Proceedings of the National Academy of Sciences of the United States of America*, **98**, 6247–6252.
- Dedeine, F., Vavre, F., Shoemaker, D.D. & Boulétreau, M. (2004) Intra-individual coexistence of a *Wolbachia* strain required for host oogenesis with two strains inducing cytoplasmic incompatibility in the wasp *Asobara tabida*. *Evolution*, **58**, 2167–2174.
- Devlin, R.H. & Nagahama, Y. (2002) Sex determination and sex differentiation in fish: an overview of genetic, physiological, and environmental influences. *Aquaculture*, **208**, 191–364.
- Dobson, S.L. & Tanouye, M.A. (1998) Evidence for a Genomic Imprinting Sex Determination mechanism in *Nasonia vitripennis* (Hymenoptera; Chalcidoidea). *Genetics*, **149**, 233–242.
- Dübendorfer, A., Hediger, M., Burghardt, G. & Bopp, D. (2002) *Musca domestica*, a window on the evolution of sex-determining mechanisms in insects. *The International Journal of Developmental Biology*, **46**, 75–79.
- Duron, O., Bouchon, D., Boutin, S., Bellamy, L., Zhou, L., Engelstadter, J., *et al.* (2008) The diversity of reproductive parasites among arthropods: *Wolbachia* do not walk alone. *BMC Biology*, **6**, 27.
- Erickson, J.W. & Quintero, J.J. (2007) Indirect effects of ploidy suggest X chromosome dose, not the X:A ratio, signals sex in *Drosophila*. *PLoS Biology*, **5**, e332.
- Foret, S., Kucharski, R., Pellegrini, M., Feng, S., Jacobsen, S.E., Robinson, G.E., *et al.* (2012) DNA methylation dynamics, metabolic fluxes, gene splicing, and alternative phenotypes in honey bees. *Proceedings of the National Academy of Sciences of the United States of America*, **109**, 4968–4973.
- Fukui, T., Kawamoto, M., Shoji, K., Kiuchi, T., Sugano, S., Shimada, T., *et al.* (2015) The endosymbiotic bacterium *Wolbachia* selectively kills male hosts by targeting the masculinizing gene. *PLoS Pathogens*, **11**, e1005048.

- Gailey, D.A., Billeter, J.-C., Liu, J.H., Bauzon, F., Allendorfer, J.B. & Goodwin, S.F. (2006) Functional conservation of the *fruitless* male sex-determination gene across 250 Myr of insect evolution. *Molecular Biology and Evolution*, **23**, 633–643.
- Gempe, T., Hasselmann, M., Schiøtt, M., Hause, G., Otte, M. & Beye, M. (2009) Sex determination in honeybees: two separate mechanisms induce and maintain the female pathway. *PLoS Biology*, **7**, e1000222.
- Geuverink, E. & Beukeboom, L.W. (2014) Phylogenetic distribution and evolutionary dynamics of the sex determination genes *doublesex* and *transformer* in insects. *Sexual Development*, **8**, 38–49.
- Giorgini, M., Monti, M.M., Caprio, E., Stouthamer, R. & Hunter, M.S. (2009) Feminization and the collapse of haplodiploidy in an asexual parasitoid wasp harboring the bacterial symbiont *Cardinium*. *Heredity*, **102**, 365–371.
- Glastad, K.M., Hunt, B.G. & Goodisman, M.A.D. (2014) Evolutionary insights into DNA methylation in insects. *Current Opinion in Insect Science*, **1**, 25–30.
- Glastad, K.M., Hunt, B.G., Yi, S. V & Goodisman, M.A.D. (2011) DNA methylation in insects: on the brink of the epigenomic era. *Insect Molecular Biology*, **20**, 553–565.
- Godfray, H.C.J. (1994) *Parasitoids: behavioral and evolutionary ecology*. Princeton University Press, Princeton, NJ.
- Goecks, J., Mortimer, N.T., Mobley, J.A., Bowersock, G.J., Taylor, J. & Schlenke, T.A. (2013) Integrative approach reveals composition of endoparasitoid wasp venoms. *PLoS one*, **8**, e64125.
- Goll, M.G. & Bestor, T.H. (2005) Eukaryotic Cytosine Methyltransferases. *Annual Review of Biochemistry*, **74**, 481–514.
- Goll, M.G., Kirpekar, F., Maggert, K.A., Yoder, J.A., Hsieh, C.-L., Zhang, X., *et al.* (2006) Methylation of tRNA^{Asp} by the DNA Methyltransferase Homolog Dnmt2. *Science*, **311**, 395–398.
- Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., *et al.* (2011) Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology*, **29**, 644–652.
- Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. (2013) QUAST: quality assessment tool for genome assemblies. *Bioinformatics*, **29**, 1072–1075.
- Haag, E. (2005) The evolution of nematode sex determination: *C. elegans* as a reference point for comparative biology. In *WormBook: The Online Review of C. elegans Biology*. Pasadena, CA.

- Hall, A.B., Basu, S., Jiang, X., Qi, Y., Timoshevskiy, V.A., Biedler, J.K., *et al.* (2015) A male-determining factor in the mosquito *Aedes aegypti*. *Science*, **348**, 1268–1270.
- Hasselmann, M. & Beye, M. (2004) Signatures of selection among sex-determining alleles of the honey bee. *Proceedings of the National Academy of Sciences of the United States of America*, **101**, 4888–4893.
- Hasselmann, M., Gempe, T., Schiøtt, M., Nunes-Silva, C.G., Otte, M. & Beye, M. (2008a) Evidence for the evolutionary nascence of a novel sex determination pathway in honeybees. *Nature*, **454**, 519–522.
- Hasselmann, M., Lechner, S., Schulte, C. & Beye, M. (2010) Origin of a function by tandem gene duplication limits the evolutionary capability of its sister copy. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 13378–13383.
- Hasselmann, M., Vekemans, X., Pflugfelder, J., Koeniger, N., Koeniger, G., Tingek, S., *et al.* (2008b) Evidence for convergent nucleotide evolution and high allelic turnover rates at the *complementary sex determiner* gene of Western and Asian honeybees. *Molecular Biology and Evolution*, **25**, 696–708.
- Hazelrigg, T. & Tu, C. (1994) Sex-specific processing of the *Drosophila* *exuperantia* transcript is regulated in male germ cells by the *tra-2* gene. *Proceedings of the National Academy of Sciences of the United States of America*, **91**, 10752–10756.
- Hediger, M., Burghardt, G., Siegenthaler, C., Buser, N., Hilfiker-Kleiner, D., Dübendorfer, A., *et al.* (2004) Sex determination in *Drosophila melanogaster* and *Musca domestica* converges at the level of the terminal regulator *doublesex*. *Development Genes and Evolution*, **214**, 29–42.
- Hediger, M., Henggeler, C., Meier, N., Perez, R., Saccone, G. & Bopp, D. (2010) Molecular characterization of the key switch *F* provides a basis for understanding the rapid divergence of the sex-determining pathway in the housefly. *Genetics*, **184**, 155–170.
- Hedley, M.L. & Maniatis, T. (1991) Sex-specific splicing and polyadenylation of *dsx* pre-mRNA requires a sequence that binds specifically to *tra-2* protein in vitro. *Cell*, **65**, 579–586.
- Heimpel, G.E. & Boer, J.G. de. (2008) Sex determination in the Hymenoptera. *Annual Review of Entomology*, **53**, 209–230.
- Heinrichs, V., Ryner, L.C. & Baker, B.S. (1998) Regulation of sex-specific selection of *fruitless* 5' splice sites by *transformer* and *transformer-2*. *Molecular and cellular biology*, **18**, 450–458.
- Herpin, A. & Schartl, M. (2015) Plasticity of gene-regulatory networks controlling sex determination: of masters, slaves, usual suspects, newcomers, and usurpators. *EMBO reports*, **16**, 1260–1274.

- Hertel, K.J., Lynch, K.W., Hsiao, E.C., Liu, E.H. & Maniatis, T. (1996) Structural and functional conservation of the *Drosophila doublesex* splicing enhancer repeat elements. *RNA*, **2**, 969–981.
- Hey, J. & Gargiulo, M.K. (1985) Sex-ratio changes in *Leptopilina heterotoma* in response to inbreeding. *Journal of Heredity*, **76**, 209–211.
- Hodgkin, J. (2002) The remarkable ubiquity of DM domain factors as regulators of sexual phenotype: ancestry or aptitude? *Genes & Development*, **16**, 2322–2326.
- Hoedjes, K.M., Smid, H.M., Schijlen, E.G.W.M., Vet, L.E.M. & Vugt, J.J.F.A. van. (2015) Learning-induced gene expression in the heads of two *Nasonia* species that differ in long-term memory formation. *BMC Genomics*, **16**, 162.
- Holt, R.A., Subramanian, G.M., Halpern, A., Sutton, G.G., Charlab, R., Nusskern, D.R., *et al.* (2002) The genome sequence of the malaria mosquito *Anopheles gambiae*. *Science*, **298**, 129–149.
- Hoshijima, K., Inoue, K., Higuchi, I., Sakamoto, H. & Shimura, Y. (1991) Control of *doublesex* alternative splicing by *transformer* and *transformer-2* in *Drosophila*. *Science*, **252**, 833–836.
- Hung, M.S., Karthikeyan, N., Huang, B., Koo, H.C., Kiger, J. & Shen, C.J. (1999) *Drosophila* proteins related to vertebrate DNA (5-cytosine) methyltransferases. *Proceedings of the National Academy of Sciences of the United States of America*, **96**, 11940–11945.
- Hussain, M., O'Neill, S.L. & Asgari, S. (2013) *Wolbachia* interferes with the intracellular distribution of Argonaute 1 in the dengue vector *Aedes aegypti* by manipulating the host microRNAs. *RNA Biology*, **10**, 1868–1875.
- Inoue, K., Hoshijima, K., Higuchi, I., Sakamoto, H. & Shimura, Y. (1992) Binding of the *Drosophila* transformer and transformer-2 proteins to the regulatory elements of doublesex primary transcript for sex-specific RNA processing. *Proceedings of the National Academy of Sciences of the United States of America*, **89**, 8092–8096.
- Jia, L.-Y., Xiao, J.-H., Xiong, T.-L., Niu, L.-M. & Huang, D.-W. (2016) The *transformer* genes in the fig wasp *Ceratosolen solmsi* provide new evidence for duplications independent of complementary sex determination. *Insect Molecular Biology*, **25**, 191–201.
- Jones, D.T., Taylor, W.R. & Thornton, J.M. (1992) The rapid generation of mutation data matrices from protein sequences. *Bioinformatics*, **8**, 275–282.
- Jones, P.A. (2012) Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nature Reviews Genetics*, **13**, 484–492.
- Jurkowski, T.P., Meusburger, M., Phalke, S., Helm, M., Nellen, W., Reuter, G., *et al.* (2008)

- Human DNMT2 methylates tRNA(Asp) molecules using a DNA methyltransferase-like catalytic mechanism. *RNA*, **14**, 1663–7160.
- Kageyama, D. & Traut, W. (2004) Opposite sex-specific effects of *Wolbachia* and interference with the sex determination of its host *Ostrinia scapularis*. *Proceedings of the Royal Society B*, **271**, 251–258.
- Kato, Y., Kobayashi, K., Oda, S., Tatarazako, N., Watanabe, H. & Iguchi, T. (2010) Sequence divergence and expression of a *transformer* gene in the branchiopod crustacean, *Daphnia magna*. *Genomics*, **95**, 160–165.
- Kato, Y., Kobayashi, K., Watanabe, H. & Iguchi, T. (2011) Environmental sex determination in the branchiopod crustacean *Daphnia magna*: Deep conservation of a *Doublesex* gene in the sex-determining pathway. *PLoS Genetics*, **7**, e1001345.
- Keeling, C.I., Yuen, M.M., Liao, N.Y., Roderick Docking, T., Chan, S.K., Taylor, G.A., *et al.* (2013) Draft genome of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins, a major forest pest. *Genome Biology*, **14**, R27.
- Keyes, L.N., Cline, T.W. & Schedl, P. (1992) The primary sex determination signal of *Drosophila* acts at the level of transcription. *Cell*, **68**, 933–943.
- Kijimoto, T., Moczek, A.P. & Andrews, J. (2012) Diversification of doublesex function underlies morph-, sex-, and species-specific development of beetle horns. *Proceedings of the National Academy of Sciences of the United States of America*, **109**, 20526–20531.
- Kirkness, E.F., Haas, B.J., Sun, W., Braig, H.R., Perotti, M.A., Clark, J.M., *et al.* (2010) Genome sequences of the human body louse and its primary endosymbiont provide insights into the permanent parasitic lifestyle. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 12168–12173.
- Kiuchi, T., Koga, H., Kawamoto, M., Shoji, K., Sakai, H., Arai, Y., *et al.* (2014) A single female-specific piRNA is the primary determiner of sex in the silkworm. *Nature*, **509**, 633–636.
- Koch, V., Nissen, I., Schmitt, B. & Beye, M. (2014) Independent evolutionary origin of *fem* paralogous genes and complementary sex determination in hymenopteran insects. *PLoS ONE*, **9**, e91883.
- Kong, L., Lv, W., Huang, Y., Liu, Z., Yang, Y. & Zhao, Y. (2015) Cloning, expression and localization of the *Daphnia carinata transformer* gene *DcarTra* during different reproductive stages. *Gene*, **566**, 248–256.
- Kopp, A. (2012) *Dmrt* genes in the development and evolution of sexual dimorphism. *Trends in Genetics*, **28**, 175–184.
- Kraaijeveld, K., Anvar, Y., Frank, J., Schmitz, A., Bast, J., Wilbrandt, J., *et al.* (2016) Decay of sexual

- trait genes in an asexual parasitoid wasp. *Genome Biology and Evolution*, eww273.
- Kraaijeveld, K., Franco, P., Knijff, P. De, Stouthamer, R. & Alphen, J.J.M. Van. (2011) Clonal genetic variation in a *Wolbachia*-infected asexual wasp: horizontal transmission or historical sex? *Molecular Ecology*, **20**, 3644–3652.
- Kremer, N., Charif, D., Henri, H., Bataille, M., Prévost, G., Kraaijeveld, K., *et al.* (2009) A new case of *Wolbachia* dependence in the genus *Asobara*: evidence for parthenogenesis induction in *Asobara japonica*. *Heredity*, **103**, 248–256.
- Kremer, N., Charif, D., Henri, H., Gavory, F., Wincker, P., Mavingui, P., *et al.* (2012) Influence of *Wolbachia* on host gene expression in an obligatory symbiosis. *BMC Microbiology*, **12**, S7.
- Krzywinska, E., Dennison, N.J., Lycett, G.J. & Krzywinski, J. (2016) A maleness gene in the malaria mosquito *Anopheles gambiae*. *Science*, **353**, 67–69.
- Kucharski, R., Maleszka, J., Foret, S. & Maleszka, R. (2008) Nutritional control of reproductive status in honeybees via DNA methylation. *Science*, **319**, 1827–1830.
- Kuhn, S., Sievert, V. & Traut, W. (2000) The sex-determining gene *doublesex* in the fly *Megaselia scalaris* : Conserved structure and sex-specific splicing. *Genome*, **43**, 1011–1020.
- Kulathinal, R.J., Skwarek, L., Morton, R.A. & Singh, R.S. (2003) Rapid evolution of the sex-determining gene *transformer*: structural diversity and rate heterogeneity among sibling species of *Drosophila*. *Molecular Biology and Evolution*, **20**, 441–452.
- Kumar, S., Stecher, G. & Tamura, K. (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, **33**, 1870–1874.
- Lagos, D., Koukidou, M., Savakis, C. & Komitopoulou, K. (2007) The *transformer* gene in *Bactrocera oleae*: the genetic switch that determines its sex fate. *Insect Molecular Biology*, **16**, 221–230.
- Lagos, D., Ruiz, M.F., Sánchez, L. & Komitopoulou, K. (2005) Isolation and characterization of the *Bactrocera oleae* genes orthologous to the sex determining *Sex-lethal* and *doublesex* genes of *Drosophila melanogaster*. *Gene*, **348**, 111–121.
- Lambkin, C.L., Sinclair, B.J., Pape, T., Courtney, G.W., Skevington, J.H., Meier, R., *et al.* (2013) The phylogenetic relationships among infraorders and superfamilies of Diptera based on morphological evidence. *Systematic Entomology*, **38**, 164–179.
- Laohakieat, K., Aketarawong, N., Iwasawin, S., Thitamadee, S. & Thanaphum, S. (2016) The study of the *transformer* gene from *Bactrocera dorsalis* and *B. correcta* with putative core promoter regions. *BMC Genetics*, **17**, 34.
- Lattorff, H.M.G., Moritz, R.F. a & Fuchs, S. (2005) A single locus determines thelytokous parthenogenesis of laying honeybee workers (*Apis mellifera capensis*). *Heredity*, **94**, 533–

537.

- Le, S.Q. & Gascuel, O. (2008) An improved general amino acid replacement matrix. *Molecular Biology and Evolution*, **25**, 1307–1320.
- Li, F., Vensko, S.P., Belikoff, E.J. & Scott, M.J. (2013) Conservation and sex-specific splicing of the *transformer* gene in the calliphorids *Cochliomyia hominivorax*, *Cochliomyia macellaria* and *Lucilia sericata*. *PLoS ONE*, **8**, e56303.
- Li-Byarlay, H., Li, Y., Stroud, H., Feng, S., Newman, T.C., Kaneda, M., *et al.* (2013) RNA interference knockdown of *DNA methyl-transferase 3* affects gene alternative splicing in the honey bee. *Proceedings of the National Academy of Sciences*, **110**, 12750–12755.
- Liu, G., Wu, Q., Li, J., Zhang, G. & Wan, F. (2015) RNAi-mediated knock-down of *transformer* and *transformer 2* to generate male-only progeny in the oriental fruit fly, *Bactrocera dorsalis* (Hendel). *PLoS ONE*, **10**, e0128892.
- Lyko, F., Foret, S., Kucharski, R., Wolf, S., Falckenhayn, C., Maleszka, R., *et al.* (2010) The honey bee epigenomes: differential methylation of brain DNA in queens and workers. *PLoS Biology*, **8**, e1000506.
- Lynch, J.A. & Desplan, C. (2006) A method for parental RNA interference in the wasp *Nasonia vitripennis*. *Nature Protocols*, **1**, 486–494.
- Ma, W.-J., Kuijper, B., Boer, J.G. de, Zande, L. van de, Beukeboom, L.W., Wertheim, B., *et al.* (2013) Absence of Complementary Sex Determination in the parasitoid wasp genus *Asobara* (Hymenoptera: Braconidae). *PLoS ONE*, **8**, e60459.
- Ma, W.-J., Pannebakker, B.A., Beukeboom, L.W., Schwander, T. & Zande, L. van de. (2014a) Genetics of decayed sexual traits in a parasitoid wasp with endosymbiont-induced asexuality. *Heredity*, **113**, 424–431.
- Ma, W.-J., Pannebakker, B.A., Zande, L. van de, Schwander, T., Wertheim, B. & Beukeboom, L.W. (2015) Diploid males support a two-step mechanism of endosymbiont-induced thelytoky in a parasitoid wasp. *BMC Evolutionary Biology*, **15**, 84.
- Ma, W.-J., Vavre, F. & Beukeboom, L.W. (2014b) Manipulation of arthropod sex determination by endosymbionts: diversity and molecular mechanisms. *Sexual Development*, **8**, 59–73.
- Madigan, S.J., Edeen, P., Esnayra, J., Mckeown, M., Madigan, S.J., Edeen, P., *et al.* (1996) *att*, a target for regulation by *tra2* in the testes of *Drosophila melanogaster*, encodes alternative RNAs and alternative proteins. *Molecular and Cellular Biology*, **16**, 4222–4230.
- Mannino, M.C., Rivarola, M., Scannapieco, A.C., González, S., Farber, M., Cladera, J.L., *et al.* (2016) Transcriptome profiling of *Diachasmimorpha longicaudata* towards useful

- molecular tools for population management. *BMC Genomics*, **17**, 793.
- Marshall Graves, J.A. (2008) Weird animal genomes and the evolution of vertebrate sex and sex chromosomes. *Annual Review of Genetics*, **42**, 565–586.
- Martín, I., Ruiz, M.F. & Sánchez, L. (2011) The gene *transformer-2* of *Sciara* (Diptera, Nematocera) and its effect on *Drosophila* sexual development. *BMC Developmental Biology*, **11**, 19.
- Mateo Leach, I., Pannebakker, B.A., Schneider, M.V., Driessen, G., Zande, L. Van De & Beukeboom, L.W. (2009) Thelytoky in Hymenoptera with *Venturia canescens* and *Leptopilina clavipes* as case studies. In *Lost Sex* (ed. by Schön, I., Martens, K. & Dijk, P.). Springer Netherlands, Dordrecht, pp. 347–375.
- Matson, C.K. & Zarkower, D. (2012) Sex and the singular DM domain: insights into sexual regulation, evolution and plasticity. *Nature Reviews Genetics*, **13**, 163–174.
- Mattox, W. & Baker, B.S. (1991) Autoregulation of the splicing of transcripts from the *transformer-2* gene of *Drosophila*. *Genes & Development*, **5**, 786–796.
- Mattox, W., McGuffin, M.E. & Baker, B.S. (1996) A negative feedback mechanism revealed by functional analysis of the alternative isoforms of the *Drosophila* splicing regulator *transformer-2*. *Genetics*, **143**, 303–314.
- Mattox, W., Palmer, M.J. & Baker, B.S. (1990) Alternative splicing of the sex determination gene *transformer-2* is sex-specific in the germ line but not in the soma. *Genes & Development*, **4**, 789–805.
- McAllister, B.F. & McVean, G. a. (2000) Neutral evolution of the sex-determining gene *transformer* in *Drosophila*. *Genetics*, **154**, 1711–1720.
- McGuffin, M.E., Chandler, D., Somaiya, D., Dauwalder, B. & Mattox, W. (1998) Autoregulation of *transformer-2* alternative splicing is necessary for normal male fertility in *Drosophila*. *Genetics*, **149**, 1477–1486.
- Misof, B., Liu, S., Meusemann, K., Peters, R.S., Donath, A., Mayer, C., *et al.* (2014) Phylogenomics resolves the timing and pattern of insect evolution. *Science*, **346**, 763–767.
- Mita, K., Kasahara, M., Sasaki, S., Nagayasu, Y., Yamada, T., Kanamori, H., *et al.* (2004) The genome sequence of silkworm, *Bombyx mori*. *DNA research*, **11**, 27–35.
- Mitsui, H., Achterberg, K. Van, Nordlander, G. & Kimura, M.T. (2007) Geographical distributions and host associations of larval parasitoids of frugivorous Drosophilidae in Japan. *Journal of Natural History*, **41**, 1731–1738.
- Morrow, J.L., Riegler, M., Frommer, M. & Shearman, D.C.A. (2014) Expression patterns of sex-determination genes in single male and female embryos of two *Bactrocera* fruit fly

- species during early development. *Insect Molecular Biology*, **23**, 754–767.
- Murata, Y.U., Ideo, S.H., Watada, M.A., Mitsui, H.I. & Kimura, M.A.T. (2009) Genetic and physiological variation among sexual and parthenogenetic populations of *Asobara japonica* (Hymenoptera: Braconidae), a larval parasitoid of drosophilid flies, **5759**, 171–178.
- Navajas-Perez, R., la Herrán, R. de, López González, G., Jamilena, M., Lozano, R., Ruiz Rejón, C., *et al.* (2005) The evolution of reproductive systems and sex-determining mechanisms within *Rumex* (Polygonaceae) Inferred from nuclear and chloroplastidial sequence data. *Molecular Biology and Evolution*, **22**, 1929–1939.
- Negri, I., Franchini, A., Gonella, E., Daffonchio, D., Mazzoglio, P.J., Mandrioli, M., *et al.* (2009) Unravelling the *Wolbachia* evolutionary role: the reprogramming of the host genomic imprinting. *Proceedings of the Royal Society B*, **276**, 2485–2491.
- Nene, V., Wortman, J.R., Lawson, D., Haas, B., Kodira, C., Tu, Z.J., *et al.* (2007) Genome sequence of *Aedes aegypti*, a major arbovirus vector. *Science*, **316**, 1718–1723.
- Niehuis, O., Hartig, G., Grath, S., Pohl, H., Lehmann, J., Tafer, H., *et al.* (2012) Genomic and morphological evidence converge to resolve the enigma of Strepsiptera. *Current Biology*, **22**, 1309–1313.
- Nissen, I., Müller, M. & Beye, M. (2012) The *Am-tra2* gene is an essential regulator of female splice regulation at two levels of the sex determination hierarchy of the honeybee. *Genetics*, **192**, 1015–1026.
- Niu, B.-L., Meng, Z.-Q., Tao, Y.-Z., Lu, S.-L., Weng, H.-B., He, L.-H., *et al.* (2005) Cloning and alternative splicing analysis of *Bombyx mori transformer-2* gene using silkworm EST database. *Acta Biochimica et Biophysica Sinica*, **37**, 728–736.
- Niyibigira, E.I., Overholt, W.A. & Stouthamer, R. (2004a) *Cotesia flavipes* Cameron (Hymenoptera: Braconidae) does not exhibit complementary sex determination (ii) Evidence from laboratory experiments. *Applied Entomology and Zoology*, **39**, 717–725.
- Niyibigira, E.I., Overholt, W.A. & Stouthamer, R. (2004b) *Cotesia flavipes* Cameron and *Cotesia sesamiae* (Cameron) (Hymenoptera : Braconidae) do not exhibit complementary sex determination: Evidence from field populations. *Applied Entomology and Zoology*, **39**, 705–715.
- Normark, B.B. (2003) The evolution of alternative genetic systems in insects. *Annual Review of Entomology*, **48**, 397–423.
- Nygaard, S., Zhang, G., Schiøtt, M., Li, C., Wurm, Y., Hu, H., *et al.* (2011) The genome of the leaf-cutting ant *Acromyrmex echinatior* suggests key adaptations to advanced social life and fungus farming. *Genome Research*, **21**, 1339–1348.

- O'Neil, M.T. & Belote, J.M. (1992) Interspecific comparison of the *transformer* gene of *Drosophila* reveals an unusually high degree of evolutionary divergence. *Genetics*, **131**, 113–128.
- O'Neill, S.L., Hoffmann, A.A. & Werren, J.H. (1997) *Influential passengers; inherited microorganisms and arthropod reproduction*. Oxford Univ Press, Oxford.
- Ohbayashi, F., Suzuki, M.G., Mita, K., Okano, K. & Shimada, T. (2001) A homologue of the *Drosophila doublesex* gene is transcribed into sex-specific mRNA isoforms in the silkworm, *Bombyx mori*. *Comparative Biochemistry and Physiology Part B*, **128**, 145–158.
- Oliveira, D.C.S.G., Werren, J.H., Verhulst, E.C., Giebel, J.D., Kamping, A., Beukeboom, L.W., *et al.* (2009) Identification and characterization of the *doublesex* gene of *Nasonia*. *Insect Molecular Biology*, **18**, 315–324.
- Pane, A., Salvemini, M., Delli Bovi, P., Polito, C. & Saccone, G. (2002) The *transformer* gene in *Ceratitis capitata* provides a genetic basis for selecting and remembering the sexual fate. *Development*, **129**, 3715–3725.
- Pannebakker, B.A., Beukeboom, L.W., Alphen, J.J.M. van, Brakefield, P.M. & Zwaan, B.J. (2004a) The genetic basis of male fertility in relation to haplodiploid reproduction in *Leptopilina clavipes* (Hymenoptera: Figitidae). *Genetics*, **168**, 341–349.
- Pannebakker, B.A., Garrido, N.R.T., Zwaan, B.J. & Alphen, J.J.M. van. (2008) Geographic variation in host-selection behaviour in the *Drosophila* parasitoid *Leptopilina clavipes*. *Entomologia Experimentalis et Applicata*, **127**, 48–54.
- Pannebakker, B.A., Pijnacker, L.P., Zwaan, B.J. & Beukeboom, L.W. (2004b) Cytology of *Wolbachia*-induced parthenogenesis in *Leptopilina clavipes* (Hymenoptera: Figitidae). *Genome*, **47**, 299–303.
- Pannebakker, B.A., Zwaan, B.J., Beukeboom, L.W. & Alphen, J.J.M. Van. (2004c) Genetic diversity and *Wolbachia* infection of the *Drosophila* parasitoid *Leptopilina clavipes* in western Europe. *Molecular Ecology*, **13**, 1119–1128.
- Patalano, S., Vlasova, A., Wyatt, C., Ewels, P., Camara, F., Ferreira, P.G., *et al.* (2015) Molecular signatures of plastic phenotypes in two eusocial insect species with simple societies. *Proceedings of the National Academy of Sciences of the United States of America*, **112**, 13970–13975.
- Peng, W., Zheng, W., Handler, A.M. & Zhang, H. (2015) The role of the *transformer* gene in sex determination and reproduction in the tephritid fruit fly, *Bactrocera dorsalis* (Hendel). *Genetica*, **143**, 717–727.
- Permpoon, R., Aketarawong, N. & Thanaphum, S. (2011) Isolation and characterization of *Doublesex* homologues in the *Bactrocera* species: *B. dorsalis* (Hendel) and *B. correcta*

- (Bezzi) and their putative promoter regulatory regions. *Genetica*, **139**, 113–127.
- Peters, R.S., Meusemann, K., Petersen, M., Mayer, C., Wilbrandt, J., Ziesmann, T., *et al.* (2014) The evolutionary history of holometabolous insects inferred from transcriptome-based phylogeny and comprehensive morphological data. *BMC Evolutionary Biology*, **14**, 52.
- Peters, R.S., Meyer, B., Krogmann, L., Borner, J., Meusemann, K., Schütte, K., *et al.* (2011) The taming of an impossible child: a standardized all-in approach to the phylogeny of Hymenoptera using public database sequences. *BMC Biology*, **9**, 55.
- Plantard, O., Rasplus, J.-Y., Mondor, G., Clainche, I. Le & Solignac, M. (1998) *Wolbachia*-induced thelytoky in the rose gallwasp *Diplolepis spinosissimae* (Giraud) (Hymenoptera: Cynipidae), and its consequences on the genetic structure of its host. *Proceedings of the Royal Society B*, **265**, 1075–1080.
- Poirie, M., Periquet, G. & Beukeboom, L.W. (1992) The Hymenopteran way of determining sex. *Seminars in Developmental Biology*, **3**, 357–361.
- Price, D.C., Egizi, A. & Fonseca, D.M. (2015) The ubiquity and ancestry of insect *doublesex*. *Scientific Reports*, **5**, 13068.
- Privman, E., Wurm, Y. & Keller, L. (2013) Duplication and concerted evolution in a master sex determiner under balancing selection. *Proceedings of the Royal Society B*, **280**, 20122968.
- Raddatz, G., Guzzardo, P.M., Olova, N., Fantappiè, M.R., Rampp, M., Schaefer, M., *et al.* (2013) Dnmt2-dependent methylomes lack defined DNA methylation patterns. *Proceedings of the National Academy of Sciences of the United States of America*, **110**, 8627–8631.
- Ramakers, C., Ruijter, J.M., Deprez, R.H.L. & Moorman, A.F.M. (2003) Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neuroscience Letters*, **339**, 62–66.
- Reumer, B.M., Alphen, J.J.M. van & Kraaijeveld, K. (2012) Occasional males in parthenogenetic populations of *Asobara japonica* (Hymenoptera: Braconidae): low *Wolbachia* titer or incomplete coadaptation? *Heredity*, **108**, 341–346.
- Richards, S., Gibbs, R.A., Weinstock, G.M., Brown, S.J., Denell, R., Beeman, R.W., *et al.* (2008) The genome of the model beetle and pest *Tribolium castaneum*. *Nature*, **452**, 949–955.
- Rideout, E.J., Narsaiya, M.S. & Grewal, S.S. (2015) The sex determination gene *transformer* regulates male-female differences in *Drosophila* body size. *PLoS Genetics*, **11**, e1005683.
- Ruiz, M.F., Alvarez, M., Eirín-López, J.M., Sarno, F., Kremer, L., Barbero, J.L., *et al.* (2015) An unusual role for *doublesex* in sex determination in the dipteran *Sciara*. *Genetics*, **200**, 1181–1199.

- Ruiz, M.F., Eirín-López, J.M., Stefani, R.N., Perondini, A.L.P., Selivon, D. & Sánchez, L. (2007a) The gene *doublesex* of *Anastrepha* fruit flies (Diptera, Tephritidae) and its evolution in insects. *Development Genes and Evolution*, **217**, 725–731.
- Ruiz, M.F., Milano, A., Salvemini, M., Eirín-López, J.M., Perondini, A.L.P., Selivon, D., *et al.* (2007b) The gene *transformer* of *Anastrepha* fruit flies (Diptera, Tephritidae) and its evolution in insects. *PLoS ONE*, **2**, e1239.
- Ruiz, M.F., Stefani, R.N., Mascarenhas, R.O., Perondini, A.L.P., Selivon, D. & Sánchez, L. (2005) The gene *doublesex* of the fruit fly *Anastrepha obliqua* (Diptera, Tephritidae). *Genetics*, **171**, 849–854.
- Ruther, J., Matschke, M., Garbe, L.-A. & Steiner, S. (2009) Quantity matters: male sex pheromone signals mate quality in the parasitic wasp *Nasonia vitripennis*. *Proceedings of the Royal Society B*, **276**, 3303–3310.
- Ryner, L.C., Goodwin, S.F., Castrillon, D.H., Anand, A., Vilella, A., Baker, B.S., *et al.* (1996) Control of male sexual behavior and sexual orientation in *Drosophila* by the *fruitless* gene. *Cell*, **87**, 1079–1089.
- Saccone, G., Peluso, I., Testa, G., DiPaola, F., Pane, A. & Polito, C. (1996) *Drosophila sex-lethal* and *doublesex* homologous genes in *Ceratitis capitata*: searching for sex-specific genes to develop a medfly transgenic sexing strain. In *Proceedings of the Enhancement of the Sterile Insect Technique through Genetic Transformation using Nuclear Techniques*. FAO/IAEA, Vienna, Austria, pp. 16–32.
- Salvemini, M., D’Amato, R., Petrella, V., Aceto, S., Nimmo, D., Neira, M., *et al.* (2013) The orthologue of the fruitfly sex behaviour gene *fruitless* in the mosquito *Aedes aegypti*: evolution of genomic organisation and alternative splicing. *PLoS ONE*, **8**, e48554.
- Salvemini, M., Mauro, U., Lombardo, F., Milano, A., Zazzaro, V., Arcà, B., *et al.* (2011) Genomic organization and splicing evolution of the *doublesex* gene, a *Drosophila* regulator of sexual differentiation, in the dengue and yellow fever mosquito *Aedes aegypti*. *BMC Evolutionary Biology*, **11**, 41.
- Salvemini, M., Robertson, M., Aronson, B., Atkinson, P., Polito, L.C. & Saccone, G. (2009) *Ceratitis capitata transformer-2* gene is required to establish and maintain the autoregulation of *Cctra*, the master gene for female sex determination. *The International Journal of Developmental Biology*, **53**, 109–120.
- Sánchez, L. (2008) Sex-determining mechanisms in insects. *The International Journal of Developmental Biology*, **52**, 837–856.
- Sandrock, C. & Vorburger, C. (2011) Single-locus recessive inheritance of asexual reproduction in a parasitoid wasp. *Current Biology*, **21**, 433–437.

- Sarno, F., Ruiz, M.F., Eirín-López, J.M., Perondini, A.L.P., Selivon, D. & Sánchez, L. (2010) The gene *transformer-2* of *Anastrepha* fruit flies (Diptera, Tephritidae) and its evolution in insects. *BMC Evolutionary Biology*, **10**, 140.
- Sarno, F., Ruiz, M.-F. & Sánchez, L. (2011) Effect of the *transformer-2* gene of *Anastrepha* on the somatic sexual development of *Drosophila*. *The International Journal of Developmental Biology*, **55**, 975–979.
- Scali, C., Catteruccia, F., Li, Q. & Crisanti, A. (2005) Identification of sex-specific transcripts of the *Anopheles gambiae doublesex* gene. *The Journal of Experimental Biology*, **208**, 3701–3709.
- Schetelig, M.F., Milano, A., Saccone, G. & Handler, A.M. (2012) Male only progeny in *Anastrepha suspensa* by RNAi-induced sex reversion of chromosomal females. *Insect Biochemistry and Molecular Biology*, **42**, 51–57.
- Schidlo, N.S., Pannebakker, B.A., Zwaan, B.J., Beukeboom, L.W. & Alphen, J.J.M. van. (2002) Curing thelytoky in the *Drosophila* parasitoid *Leptopilina clavipes* (Hymenoptera: Figitidae). *Proceedings Experimental and Applied Entomology*, **13**, 93–96.
- Schmieder, S., Colinet, D. & Poirié, M. (2012) Tracing back the nascence of a new sex-determination pathway to the ancestor of bees and ants. *Nature Communications*, **3**, 895.
- Schneider, M. V., Beukeboom, L.W., Driessen, G., Lapchin, L., Bernstein, C. & Alphen, J.J.M. Van. (2002) Geographical distribution and genetic relatedness of sympatrical thelytokous and arrhenotokous populations of the parasitoid *Venturia canescens* (Hymenoptera). *Journal of Evolutionary Biology*, **15**, 191–200.
- Sciabica, K.S. & Hertel, K.J. (2006) The splicing regulators Tra and Tra2 are unusually potent activators of pre-mRNA splicing. *Nucleic acids research*, **34**, 6612–6620.
- Sharma, A., Heinze, S.D., Wu, Y., Kohlbrenner, T., Morilla, I., Brunner, C., Wimmer, E.A., Zande, L. van de, Robinson, M.D., Beukeboom, L.W., Bopp, D. (submitted) Male sex in houseflies is determined by a paralog of the generic splice factor CWC22.
- Shearman, D.C.A. & Frommer, M. (1998) The *Bactrocera tryoni* homologue of the *Drosophila melanogaster* sex-determination gene *doublesex*. *Insect Molecular Biology*, **7**, 355–366.
- Shukla, J.N. & Nagaraju, J. (2010) *Doublesex*: a conserved downstream gene controlled by diverse upstream regulators. *Journal of Genetics*, **89**, 341–356.
- Shukla, J.N. & Palli, S.R. (2012a) Sex determination in beetles: production of all male progeny by parental RNAi knockdown of *transformer*. *Scientific Reports*, **2**, 602.
- Shukla, J.N. & Palli, S.R. (2012b) *Doublesex* target genes in the red flour beetle, *Tribolium*

- castaneum. *Scientific Reports*, **2**, 948.
- Shukla, J.N. & Palli, S.R. (2013) *Tribolium castaneum* Transformer-2 regulates sex determination and development in both males and females. *Insect Biochemistry and Molecular Biology*, **43**, 1125–1132.
- Siera, S.G. & Cline, T.W. (2008) Sexual back talk with evolutionary implications: stimulation of the *Drosophila* sex-determination gene *sex-lethal* by its target *transformer*. *Genetics*, **180**, 1963–1981.
- Sievert, V., Kuhn, S. & Traut, W. (1997) Expression of the sex determining cascade genes *Sex-lethal* and *doublesex* in the phorid fly *Megaselia scalaris*. *Genome*, **40**, 211–214.
- Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E. V & Zdobnov, E.M. (2015) BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, **31**, 3210–3212.
- Simpson, J.T., Wong, K., Jackman, S.D., Schein, J.E., Jones, S.J.M. & Birol, I. (2009) ABySS: A parallel assembler for short read sequence data. *Genome Research*, **19**, 1117–1123.
- Skinner, S.W. & Werren, J.H. (1980) The genetics of sex determination in *Nasonia vitripennis* (Hymenoptera, Pteromalidae). *Genetics*, **94**, s98.
- Smith, C.D., Zimin, A., Holt, C., Abouheif, E., Benton, R., Cash, E., *et al.* (2011a) Draft genome of the globally widespread and invasive Argentine ant (*Linepithema humile*). *Proceedings of the National Academy of Sciences of the United States of America*, **108**, 5673–5678.
- Smith, C.R., Smith, C.D., Robertson, H.M., Helmkampf, M., Zimin, A., Yandell, M., *et al.* (2011b) Draft genome of the red harvester ant *Pogonomyrmex barbatus*. *Proceedings of the National Academy of Sciences of the United States of America*, **108**, 5667–5672.
- Standage, D.S., Berens, A.J., Glastad, K.M., Severin, A.J., Brendel, V.P. & Toth, A.L. (2016) Genome, transcriptome and methylome sequencing of a primitively eusocial wasp reveal a greatly reduced DNA methylation system in a social insect. *Molecular Ecology*, **25**, 1769–1784.
- Stille, B. & Dävring, L. (1980) Meiosis and reproductive strategy in the parthenogenetic gall wasp *Diplolepis rosae* (L.) (Hymenoptera, Cynipidae). *Hereditas*, **92**, 353–362.
- Stouthamer, R. (1997) *Wolbachia*-induced parthenogenesis. In *Influential passengers: inherited microorganisms and arthropod reproduction* (ed. by O'Neill, S.L., Hoffmann, A.A. & Werren, J.H.). Oxford University Press, Oxford, UK, pp. 102–124.
- Stouthamer, R., Breeuwer, J.A.J. & Hurst, G.D.D. (1999) *Wolbachia pipientis*: microbial manipulator of arthropod reproduction. *Annual Review of Microbiology*, **53**, 71–102.
- Suen, G., Teiling, C., Li, L., Holt, C., Abouheif, E., Bornberg-Bauer, E., *et al.* (2011) The genome

- sequence of the leaf-cutter ant *Atta cephalotes* reveals insights into its obligate symbiotic lifestyle. *PLoS Genetics*, **7**, e1002007.
- Sugimoto, T.N., Fujii, T., Kayukawa, T., Sakamoto, H. & Ishikawa, Y. (2010) Expression of a *doublesex* homologue is altered in sexual mosaics of *Ostrinia scapularis* moths infected with *Wolbachia*. *Insect Biochemistry and Molecular Biology*, **40**, 847–854.
- Sugimoto, T.N. & Ishikawa, Y. (2012) A male-killing *Wolbachia* carries a feminizing factor and is associated with degradation of the sex-determining system of its host. *Biology Letters*, **8**, 412–415.
- Sugimoto, T.N., Kayukawa, T., Shinoda, T., Ishikawa, Y. & Tsuchida, T. (2015) Misdirection of dosage compensation underlies bidirectional sex-specific death in *Wolbachia*-infected *Ostrinia scapularis*. *Insect Biochemistry and Molecular Biology*, **66**, 72–76.
- Suomalainen, E., Saura, A. & Lokki, J. (1987) *Cytology and evolution in parthenogenesis*. CRC Press Boca Raton Inc, Florida.
- Suzuki, M.G. (2010) Sex determination: insights from the silkworm. *Journal of Genetics*, **89**, 357–363.
- Suzuki, M.G., Funaguma, S., Kanda, T., Tamura, T. & Shimada, T. (2003) Analysis of the biological functions of a *doublesex* homologue in *Bombyx mori*. *Development Genes and Evolution*, **213**, 345–354.
- Suzuki, M.G., Imanishi, S., Dohmae, N., Nishimura, T., Shimada, T. & Matsumoto, S. (2008) Establishment of a novel in vivo sex-specific splicing assay system to identify a trans-acting factor that negatively regulates splicing of *Bombyx mori dsx* female exons. *Molecular and Cellular Biology*, **28**, 333–343.
- Suzuki, M.G., Ohbayashi, F., Mita, K. & Shimada, T. (2001) The mechanism of sex-specific splicing at the *doublesex* gene is different between *Drosophila melanogaster* and *Bombyx mori*. *Insect Biochemistry and Molecular Biology*, **31**, 1201–1211.
- Suzuki, M.G., Suzuki, K., Aoki, F. & Ajimura, M. (2012) Effect of RNAi-mediated knockdown of the *Bombyx mori transformer-2* gene on the sex-specific splicing of *Bmdsx* pre-mRNA. *The International Journal of Developmental Biology*, **56**, 693–699.
- Suzuki, M.G., Tochigi, M., Sakaguchi, H., Aoki, F. & Miyamoto, N. (2015) Identification of a *transformer* homolog in the acorn worm, *Saccoglossus kowalevskii*, and analysis of its activity in insect cells. *Development Genes and Evolution*, **225**, 161–169.
- Suzuki, M.M. & Bird, A. (2008) DNA methylation landscapes: provocative insights from epigenomics. *Nature Reviews Genetics*, **9**, 465–476.
- Tamura, K., Dudley, J., Nei, M. & Kumar, S. (2007) MEGA4: Molecular Evolutionary Genetics

- Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution*, **24**, 1596–1599.
- The International Aphid Genomics Consortium. (2010) Genome sequence of the pea aphid *Acyrtosiphon pisum*. *PLoS Biology*, **8**, e1000313.
- Thomson, T. & Lin, H. (2009) The biogenesis and function of PIWI proteins and piRNAs: progress and prospect. *Annual review of cell and developmental biology*, **25**, 355–376.
- Tian, M. & Maniatis, T. (1992) Positive control of pre-mRNA splicing in vitro. *Science*, **256**, 237–240.
- Toyota, K., Kato, Y., Sato, M., Sugiura, N., Miyagawa, S., Miyakawa, H., *et al.* (2013) Molecular cloning of *doublesex* genes of four Cladocera (water flea) species. *BMC Genomics*, **14**, 239.
- Trautwein, M.D., Wiegmann, B.M., Beutel, R., Kjer, K.M. & Yeates, D.K. (2012) Advances in insect phylogeny at the dawn of the postgenomic era. *Annual Review of Entomology*, **57**, 449–468.
- Tweedie, S., Ng, H.H., Barlow, A.L., Turner, B.M., Hendrich, B. & Bird, A. (1999) Vestiges of a DNA methylation system in *Drosophila melanogaster*? *Nature Genetics*, **23**, 389–90.
- Verhulst, E.C., Beukeboom, L.W. & Zande, L. van de. (2010a) Maternal control of haplodiploid sex determination in the wasp *Nasonia*. *Science*, **328**, 620–623.
- Verhulst, E.C., Lynch, J. a, Bopp, D., Beukeboom, L.W. & Zande, L. van de. (2013) A new component of the *Nasonia* sex determining cascade is maternally silenced and regulates *transformer* expression. *PLoS ONE*, **8**, e63618.
- Verhulst, E.C. & Zande, L. van de. (2015) Double nexus-*Doublesex* is the connecting element in sex determination. *Briefings in Functional Genomics*, **14**, 396–406.
- Verhulst, E.C., Zande, L. van de & Beukeboom, L.W. (2010b) Insect sex determination: it all evolves around *transformer*. *Current Opinion in Genetics & Development*, **20**, 376–383.
- Wachi, N., Nomano, F.Y., Mitsui, H., Kasuya, N. & Kimura, M.T. (2015) Taxonomy and evolution of putative thelytokous species of *Leptopilina* (Hymenoptera: Figitidae) from Japan, with description of two new species. *Entomological Science*, **18**, 41–54.
- Walsh, T.K., Brisson, J.A., Robertson, H.M., Gordon, K., Jaubert-Possamai, S., Tagu, D., *et al.* (2010) A functional DNA methylation system in the pea aphid, *Acyrtosiphon pisum*. *Insect Molecular Biology*, **19**, 215–228.
- Wang, X.-Y., Zheng, Z.-Z., Song, H.-S. & Xu, Y.-Z. (2014) Conserved RNA cis-elements regulate alternative splicing of Lepidopteran *doublesex*. *Insect Biochemistry and Molecular Biology*, **44**, 1–11.

- Wang, Y., Jorda, M., Jones, P.L., Maleszka, R., Ling, X., Robertson, H.M., *et al.* (2006) Functional CpG methylation system in a social insect. *Science*, **314**, 645–647.
- Weeks, a R., Marec, F. & Breeuwer, J. a. (2001) A mite species that consists entirely of haploid females. *Science*, **292**, 247924–82.
- Werren, J.H., Baldo, L. & Clark, M.E. (2008) *Wolbachia*: master manipulators of invertebrate biology. *Nature Reviews Microbiology*, **6**, 741–751.
- Werren, J.H., Richards, S., Desjardins, C.A., Niehuis, O., Gadau, J., Colbourne, J.K., *et al.* (2010) Functional and evolutionary insights from the genomes of three parasitoid *Nasonia* species. *Science*, **327**, 343–348.
- Wexler, J.R., Plachetzki, D.C. & Kopp, A. (2014) Pan-metazoan phylogeny of the DMRT gene family: a framework for functional studies. *Development Genes and Evolution*, **224**, 175–181.
- Whiting, P.W. (1939) Sex determination and reproductive economy in *Habrobracon*. *Genetics*, **24**, 110–111.
- Whiting, P.W. (1943) Multiple alleles in complementary sex determination of *Habrobracon*. *Genetics*, **28**, 365–382.
- Whiting, P.W. (1960) Polyploidy in *Mormoniella*. *Genetics*, **45**, 949–970.
- Wiegmann, B.M., Trautwein, M.D., Winkler, I.S., Barr, N.B., Kim, J.-W., Lambkin, C., *et al.* (2011) Episodic radiations in the fly tree of life. *Proceedings of the National Academy of Sciences*, **108**, 5690–5695.
- Wilgenburg, E. van, Driessen, G. & Beukeboom, L.W. (2006) Single locus complementary sex determination in Hymenoptera: an “unintelligent” design? *Frontiers in Zoology*, **3**, 1.
- Wilkins, A.S. (1995) Moving up the hierarchy: a hypothesis on the evolution of a genetic sex determination pathway. *BioEssays*, **17**, 71–77.
- Woyke, J. (1963) What happens to diploid drone larvae in a honeybee colony. *Journal of Apicultural Research*, **2**, 73–75.
- Wurm, Y., Wang, J., Riba-Grognuz, O., Corona, M., Nygaard, S., Hunt, B.G., *et al.* (2011) The genome of the fire ant *Solenopsis invicta*. *Proceedings of the National Academy of Sciences of the United States of America*, **108**, 5679–5684.
- Xiao, J.-H., Yue, Z., Jia, L.-Y., Yang, X.-H., Niu, L.-H., Wang, Z., *et al.* (2013) Obligate mutualism within a host drives the extreme specialization of a fig wasp genome. *Genome Biology*, **14**, R141.
- Ye, Y.H., Woolfit, M., Huttley, G.A., Rancès, E., Caragata, E.P., Popovici, J., *et al.* (2013) Infection

- with a virulent strain of *Wolbachia* disrupts genome wide-patterns of cytosine methylation in the mosquito *Aedes aegypti*. *PLoS ONE*, **8**, e66482.
- Zande, L. van de & Verhulst, E.C. (2014) Genomic imprinting and maternal effect genes in haplodiploid sex determination. *Sexual Development*, **8**, 74–82.
- Zemach, A., McDaniel, I.E., Silva, P. & Zilberman, D. (2010) Genome-wide evolutionary analysis of eukaryotic DNA methylation. *Science*, **328**, 916–919.
- Zhang, G., Hussain, M., O’Neill, S.L. & Asgari, S. (2013) *Wolbachia* uses a host microRNA to regulate transcripts of a methyltransferase, contributing to dengue virus inhibition in *Aedes aegypti*. *Proceedings of the National Academy of Sciences of the United States of America*, **110**, 10276–10281.
- Zhou, Y., Gu, H. & Dorn, S. (2006) Single-locus sex determination in the parasitoid wasp *Cotesia glomerata* (Hymenoptera: Braconidae). *Heredity*, **96**, 487–492.
- Zug, R. & Hammerstein, P. (2012) Still a host of hosts for *Wolbachia*: analysis of recent data suggests that 40% of terrestrial arthropod species are infected. *PLoS ONE*, **7**, e38544.
- Zwier, M. V., Verhulst, E.C., Zwahlen, R.D., Beukeboom, L.W. & Zande, L. van de. (2012) DNA methylation plays a crucial role during early *Nasonia* development. *Insect Molecular Biology*, **21**, 129–138.

English summary

Sex determination and sex differentiation

The separation of and sexual differentiation into female and male sexes is a widespread phenomenon in the animal kingdom. Each of the two sexes produces haploid gametes of which the nuclei fuse in the fertilized egg to yield start the development of a male or female diploid embryo. This process must be of high fidelity, as intersexes usually have a lower fitness. As the individual matures, differentiation of sexual organs and secondary sexual traits occurs. This eventually yields a mature female with ovaries or a male with testes.

Both parents can control the action of genes by activating or silencing them on their provided chromosome sets. Only the mother provides the cytoplasm of the offspring and is able to deposit gene transcripts into her oocytes. This seemingly leaves the male with fewer options to control his offspring, but also he may be able to provide additional information, e.g. by imprinting genes or providing short RNA molecules through the sperm. After fertilization, the embryo starts its development in the presence of a maternal and paternal chromosome set and egg cytoplasm containing parentally derived products. These products in the egg degenerate over time while zygotic transcription starts.

Chromosomal and haplodiploid sex determination

Sex determination systems based on the presence of sex chromosomes are particularly common. In these systems a particular chromosome, containing a sex determining locus, determines the sex of the offspring. In male heterogametic systems all males have both an X and a Y chromosome, whereas females carry a homozygous combination of two X chromosomes. In supplying either an X or a Y chromosome to their spermatocytes, males determine at fertilization whether the embryo will become female or male. Conversely, in female heterogametic systems, the females have a Z and a W chromosome, whereas the males have two Z chromosomes. In these systems females determine the sex of the offspring through their oocytes with either a Z or a W chromosome. Variations to these heterogametic systems occur when either the male or the female lacks the minor sex chromosome, i.e. they have an XO or ZO genotype. In such systems the heterogametic sex determines the sex of the offspring alike the XY and ZW systems by providing either a sex chromosome or not.

Haplodiploid sex determination is a markedly different system, though it is also based on chromosomal configurations of the embryo. Haplodiploid species consist of haploid males and diploid females. Females produce haploid oocytes that do not need to be fertilized to develop into an adult individual. Unfertilized eggs develop into haploid males, whereas fertilized eggs develop into diploid females. The two sexes do not differ for a specific sex chromosome, but in the number of copies of the whole complement. Males only inherit a chromosome set from their mother and subsequently pass this on to their daughters. It means that there is no sex determining locus on a specific sex chromosome as any chromosome may occur in both sexes. Haplodiploidy allows behavioural control over fertilization as females can facultatively allow a sperm to her egg upon oviposition and they can thus control the sex of the offspring.

Yet, in sexually reproducing haplodiploids, the inclusion of a paternal chromosome set is, bar some exceptional cases, the only genetic option to generate a female.

Haplodiploid mechanisms of sex determination

One genetic mechanism of haplodiploid sex determination consists of a paternal genome and maternally provided elements activating the female-specific pathway upon zygotic transcription in the embryo. A hallmark of insect sex determination is the sex-specific splicing that occurs upon the start of female or male development. The molecular genetics of haplodiploid sex determination have thus far only been studied in two species of Hymenoptera: the honeybee (*Apis mellifera*) and the jewel wasp (*Nasonia vitripennis*). These mechanisms are very different in genetic versus epigenetic components and this variation gave the first indication of the fast evolution of sex determination mechanisms in the Hymenoptera.

The honeybee possesses a mechanism of Complementary Sex Determination (CSD), where heterozygosity at one or multiple loci determines the sex of the offspring. Heterozygous diploid individuals will develop into females, whereas hemizygous haploid individuals or homozygous diploid individuals will become males. This type of mechanism can easily be detected by inbreeding that will increase the level of homozygosity, and hence the ratio of male to female offspring. CSD has been described for many species in many different lineages of the Hymenoptera. The precise molecular mechanism has only been identified in the honeybee. The *csd* gene is a duplication of the key sex determination gene *feminizer* (*fem*), an ortholog of *transformer* (*tra*). Only a heterozygotic state of the *csd* gene can enforce female-specific splicing of the *fem* gene and start the path to female differentiation.

The molecular mechanism of sex determination in *N. vitripennis* is based on two very different principles. Maternal provision of female-specific *tra* transcripts is required to start female-specific splicing of this gene in the embryo. This alone is however not enough to start female development. An element which is silenced in the maternal genome, yet active in the paternal genome, is needed in combination with the female-specific *tra* provision to enable female-specific splicing of *tra*. This imprinting element has not yet been identified, but is not *tra* itself.

The insect sex determination gene cascade

A major question in evolutionary biology is how sex determination pathways evolve. Which genes are involved, how are they regulated, and do these genes have other functions as well? *Tra* is not only a key gene in the sex determination pathway of *A. mellifera* and *N. vitripennis*, but plays a pivotal role in development of all insect species in which its orthologs have been found. Only the female-specific splice variant of *tra* can be translated into a functional protein. This TRA protein forms a complex with TRA2 protein, encoded by the *tra2* gene that does not possess any splice forms restricted to a particular sex and is more conserved than other genes in the sex determination cascade. *Tra2* also has other developmental functions. The combination

of TRA2 and TRA ensures that mRNA transcribed from the downstream gene *doublesex* (*dsx*) is spliced in the female mode. DSX is a transcription factor and the female-specific protein will induce female differentiation, while the male-specific protein leads to male differentiation. In *A. mellifera* and *N. vitripennis* the zygotic state of *tra* is in the male mode. A feminizing primary signal is needed to initiate female-specific zygotic transcription of *tra* to start female development. The primary signals, which control this switch of the *tra-dsx* axis, are highly diverged, which hampers identification based on homology.

Dsx has thus far been detected in all investigated insect orders (Chapter 2). This conservation is confirmed in this thesis by its conserved sex-specific transcripts in *Asobara tabida* sexual development (Chapter 4). Its function as the ultimate switch leading to sexual differentiation may be very specific and not easily taken over by another gene. This specificity may not be the case for *tra*, as this gene appears to be absent (or evades detection) in multiple branches of insects (Chapter 2). *Tra* seems to have been lost multiple times from the sex determination cascade or, alternatively, may have been added multiple times in different groups. *Tra* has been duplicated in several families of the Hymenoptera, including non-CSD species, as paralogs were observed in *Leptopilina* wasps (Chapter 2 and 5). Yet it is unclear whether these duplications are involved in sex determination, be it as a CSD locus or as a maternal effect factor.

Maternal provision in sexually reproducing wasps

Maternal provision of both *tra2* (Chapter 3) and female-specific *tra* was found to be required to start female development in *N. vitripennis*. Inclusion of *tra2* mRNA into the egg is a conserved mode of gene regulation in sex determination. Decreased levels of *tra2* mRNA in embryos resulted in diploid male development, but also demonstrated a strong mortality in early life stages. These results indicate that early presence of *tra2* transcripts in the embryo also has other developmental functions in haploid embryos. Furthermore I observed that diploid embryos with a paternal chromosome set have a reduced offspring number if either *tra2* or *tra* is absent in early life stages.

Whereas female-specific *tra* maternal provisioning is crucial for *N. vitripennis*, it was not found in the wasp *A. tabida* (Chapter 4). *A. tabida* females do provide alternatively spliced variants of *tra* to their eggs that code for a shorter protein, which is, however, unlike the male form. The lack of female-specific *tra* in the early stages of diploid female development leaves the question how feminizing activity would be started. The alternative splice forms of *tra* could play a role in activating female-specific transcription of *tra* in the early embryo. An active paternal factor would be required to distinguish between haploid and diploid development. The crucial role of *tra2* maternal provision does appear conserved in *A. tabida* sex determination, as I could detect its presence in embryos prior to zygotic transcription and upon zygotic activation of the sex determination cascade.

Endosymbiont interference with sex determination

In insects whose sex determination mechanism is based on a zygotic state of *tra* in the male mode, a cue for female development is required that switches the sex determination cascade into the female mode. This cue does not necessarily come from the insect itself. Endosymbionts are widespread in many insect orders and can have drastic effects on the reproductive mode of their hosts. Endosymbionts are only transmitted via cytoplasm and, because of this one-sided transmission, benefit from turning genetic male embryos into females or excluding male embryos and switching the progeny sex ratio toward more females. In haplodiploid species they can induce asexual reproduction, meaning that infected diploid females produce infected haploid oocytes that are diploidized and feminized to develop into females.

The parasitoid wasps *Asobara japonica* and *Leptopilina clavipes* consist of both sexual and asexual populations. Asexual wasps of both species are infected with *Wolbachia* endosymbionts. At least for *A. japonica* it is documented that this feminization occurs as a separate step after the diploidization action. The *Wolbachia* infection can be removed with antibiotics treatments which results in the production of haploid male wasps.

Sexual females of *A. japonica* and *L. clavipes* do not provide female-specific *tra* transcripts to their oocytes (Chapter 5 and 6). This phenomenon was found to be shared with the sexual females of *A. tabida* (Chapter 4). Asexual females of both *A. japonica* and *L. clavipes*, however, do add female-specific *tra* transcripts to their oocytes in the presence of *Wolbachia*. When I removed *Wolbachia* from asexual *A. japonica* females, they did not provide the female-specific *tra* transcripts anymore and reverted to the pattern of *tra* provision observed in the sexual females (Chapter 6). Hence, *Wolbachia* appears to have taken over control of sex determination by manipulating the maternal provision of the feminizing *tra*.

Parental and endosymbiont effects on sex determination – who is in control?

Maternal provision and the sex-specific transcripts of sex determination genes during early stages of development yield valuable information for uncovering the molecular mechanisms of sex determination. Female-specific *tra* was found to not be required at the onset of female development under sexual reproduction of three wasp species (*A. tabida*, *A. japonica* and *L. clavipes*) (Chapter 4, 5 and 6). Yet the requirement of *tra2* maternal provision for female development is likely a widespread phenomenon. Additional maternal elements, other than *tra2*, may be required to induce the female-specific pathway, in example the alternative splice variants of *tra* or its paralogs. However, this maternal provision alone is not sufficient to start female development in sexually reproducing wasps. The paternal genome would require an epigenetic difference from the maternal genome to start female-specific zygotic transcription in diploid fertilized eggs. This paternal dependency may hypothetically also be manipulated by *Wolbachia* mimicking the activity of the paternal genome. Alternatively, rather than a distinction between two chromosome sets, the dosage effects resulting from the diploid state may initiate female development. The interference of endosymbionts with the splicing and maternal

provision of *tra* suggests the existence of an arms race between maternal and endosymbiotic elements over haplodiploid sex determination. This warrants the question: who is in control of sex determination, and who could take over control of sex determination?

Dutch summary

Geslachtsbepaling en geslachtsdifferentiatie

De scheiding in verschillende geslachten en de differentiatie tot mannelijke en vrouwelijke individuen is een wijdverspreid verschijnsel in het dierenrijk. Beide geslachten produceren haploïde gameten. De ontwikkeling van een mannelijk of vrouwelijk diploïd embryo start als de kernen van deze gameten fuseren in de bevruchte eicel. Deze tweedeling in ontwikkeling moet gewaarborgd zijn, aangezien interseksuele individuen vaak een lagere fitness hebben. Differentiatie van geslachtsorganen en secundaire geslachtskenmerken treedt op gedurende de verdere ontwikkeling. Dit leidt uiteindelijk tot een volgroeid vrouwelijk individu met ovaria of een volgroeid mannelijk individu met testes.

Beide ouders kunnen de activiteit van genen controleren door deze te aan- of uit te zetten op de set chromosomen die zij aan het embryo doorgeven. Alleen de moeder geeft haar cytoplasma mee aan haar nakomelingen en kan hier gen-transcripten in meegeven. Dit lijkt de vader minder mogelijkheden te geven om de ontwikkeling van zijn nageslacht te beïnvloeden. Maar ook hij kan mogelijkwijs extra elementen meegeven, bijvoorbeeld door genen in te prenten of door korte RNA moleculen met de spermatozoa mee te leveren. Na de bevruchting zal het embryo de eigen ontwikkeling beginnen in aanwezigheid van zowel een maternale als een paternale chromosoomset en een cytoplasma met daarin producten van beide ouders. Deze producten degenereren langzaam terwijl de transcriptie vanaf het genoom de zygote zelf begint.

Chromosomale en haplodiploïde geslachtsbepaling

Geslachtsbepalingssystemen gebaseerd op geslachtschromosomen zijn zeer algemeen. Een bepaald chromosoom in deze systemen bevat een geslachtsbepalingslocus en dit bepaalt, zoals de naam al zegt, het geslacht van de nakomelingen. In mannelijk heterogametische systemen hebben alle mannelijke individuen een X- en een Y-chromosoom, terwijl de vrouwelijke individuen twee X-chromosomen bezitten. Doordat de mannelijke individuen zowel een X- als een Y-chromosoom kunnen meegeven in hun spermatocyten wordt hierdoor bij de bevruchting bepaald of het embryo mannelijk of vrouwelijk wordt. Omgekeerd hebben vrouwelijke individuen in vrouwelijk heterogametische systemen een Z- als een W-chromosoom en de mannelijke individuen twee Z-chromosomen. De oöcyten van de vrouwelijke individuen bepalen in deze systemen het geslacht van de nakomelingen door een Z-chromosoom of een W-chromosoom mee te geven. Er zijn variaties op deze heterogametische systemen, waarbij of het mannelijke type of het vrouwelijke type een geslachtschromosoom mist. In dit geval is er sprake van een XO- of een ZO-genotype..

Haplodiploïde geslachtsbepaling werkt volgens een ander principe, maar is ook gebaseerd op de chromosomale samenstelling in het embryo. Haplodiploïde soorten bestaan uit haploïde mannelijke individuen en diploïde vrouwelijke individuen. Deze vrouwelijke individuen produceren haploïde oöcyten die niet bevrucht hoeven te worden om te ontwikkelen tot een volwassen nakomeling. Onbevruchte eieren ontwikkelen zich tot haploïde mannelijke individuen, terwijl bevruchte eieren zich ontwikkelen tot diploïde vrouwelijke individuen. De

twee geslachten zijn niet gescheiden door een specifiek geslachtschromosoom, maar verschillen in de hoeveelheid chromosoomsets. Mannelijke individuen krijgen alleen een chromosoomset van hun moeder en geven deze vervolgens alleen door aan dochters, want bij haplodiploïden produceren mannen geen zonen. Dit betekent, dat er geen geslachtsbepalingslocus op een specifiek geslachtschromosoom aanwezig is aangezien elk chromosoom in elk geslacht kan voorkomen. Haplodiploïde geslachtsbepaling wordt gecontroleerd door de moeder, die kan kiezen haar eieren te laten bevruchten door ontvangen spermatozoa, en hiermee het geslacht van haar nakomelingen kan bepalen. Eigenlijk is in seksueel voortplantende haplodiploïden, een enkele uitzondering daargelaten, de toevoeging van een paternaal genoom de enige optie om een vrouwelijk individu te verkrijgen.

Haplodiploïde mechanismen van geslachtsbepaling

Het genetische mechanisme van haplodiploïde geslachtsbepaling is gebaseerd op een paternaal genoom en maternaal toegevoegde elementen waardoor vrouwelijke ontwikkeling tijdens de start van zygotische transcriptie in het embryo in gang wordt gezet. Het belangrijkste kenmerk van geslachtsbepaling in insecten is de geslachtsspecifieke splicing die optreedt bij de start van vrouwelijke en mannelijke ontwikkeling. De moleculaire genetica van haplodiploïde geslachtsbepaling is tot dusver slechts bestudeerd in twee soorten Hymenoptera: de honingbij (*Apis mellifera*) en de juweelwesp (*Nasonia vitripennis*). Deze mechanismen verschillen tussen genetische en epigenetische componenten en deze variatie gaf de eerste aanwijzing voor de snelle evolutie van geslachtsbepalingsmechanismen in de Hymenoptera.

De honingbij bezit Complementaire Sex Determinatie (CSD), waarbij heterozygotie op een of meerdere loci het geslacht van de nakomelingen bepaalt. Heterozygote diploïde individuen ontwikkelen zich vrouwelijk, terwijl hemizygotie haploïde individuen en homozygote diploïde individuen uiteindelijk mannelijk worden. Dit type mechanisme kan gedetecteerd worden door inteeltstudies, die het niveau van homozygotie verhogen, en daarmee de ratio mannelijke ten opzicht van vrouwelijke ontwikkeling zullen verhogen. CSD-mechanismen zijn gevonden in veel soorten uit verschillende families van de Hymenoptera. Het precieze moleculaire mechanisme is echter alleen opgehelderd voor in de honingbij. Het *csd* gen is hier een duplicatie van het belangrijke geslachtsbepalingsgen *feminizer* (*fem*), een ortholoog van *transformer* (*tra*). Alleen een heterozygote staat van het *csd* gen kan ervoor zorgen dat *fem*-transcripten op een vrouwelijk specifieke manier geproduceerd worden en hierdoor start vrouwelijke differentiatie.

Het moleculaire mechanisme van geslachtsbepaling in *N. vitripennis* is gebaseerd op twee andere principes. Maternale provisie van vrouwelijk specifieke *tra*-transcripten is nodig om de vrouwelijke transcriptievorm van dit gen te starten in het embryo. Dit alleen is niet afdoende om vrouwelijke ontwikkeling te activeren. Een element, dat geïnactiveerd is op het maternale genoom, maar actief is op het paternale genoom, is nodig in combinatie met vrouwelijk

specifieke *tra*-provisie om de vrouwelijk specifieke zygotische transcriptie van *tra* te starten. Het geïnactiveerde element is nog niet geïdentificeerd, maar het is niet *tra* zelf.

De geslachtsbepaling gen-cascade van insecten

Een belangrijke vraag in de evolutionaire biologie is hoe geslachtsbepalingsroutes evolueren. Welke genen zijn hierbij betrokken; hoe worden zij gereguleerd; en hebben deze genen ook andere functies? *Tra* is niet alleen een belangrijk gen in de geslachtsbepalingsmechanismen van *A. mellifera* en *N. vitripennis*, maar speelt een centrale rol in de ontwikkeling van alle insectensoorten, waarin orthologen zijn gedetecteerd. Alleen de vrouwelijke splice variant van *tra* wordt vertaald in een functioneel eiwit. Dit TRA-eiwit vormt een complex met het TRA2-eiwit, dat gecodeerd wordt door het *tra2* gen. *Tra2* bezit geen geslachtsspecifieke splice varianten en is meer geconserveerd dan andere genen in de geslachtsbepalingscascade. Verder heeft het ook nog andere functies in vroege ontwikkeling. De combinatie van TRA en TRA2 zorgt ervoor, dat het mRNA van het volgende gen in de cascade, *doublesex (dsx)*, wordt gespliced in de vrouwelijke modus. DSX is een transcriptie factor, en het vrouwelijk specifieke eiwit induceert vrouwelijke differentiatie, terwijl het mannelijke eiwit leidt tot mannelijke differentiatie. De zygotische staat van *tra* is in de mannelijke modus in *A. mellifera* en *N. vitripennis*. Een feminiserend primair signaal is nodig om vrouwelijk specifieke zygotische transcriptie van *tra* te initiëren om vrouwelijke ontwikkeling te starten. De primaire signalen, die deze switch van de *tra-dsx*-as controleren, zijn sterk gedivergeerd, wat hun identificatie gebaseerd op homologie bemoeilijkt.

Dsx is tot dusver gedetecteerd in alle bestudeerde insecten-orde (Hoofdstuk 2). Deze conservering wordt bevestigd in deze thesis door de geconserveerde geslachtsspecifieke transcripten in *Asobara tabida* vrouwelijke en mannelijke ontwikkeling (Hoofdstuk 4). De functie van *dsx* als de laatste switch leidend tot seksuele differentiatie is mogelijk zeer specifiek en niet makkelijk over te nemen door een ander gen. Deze functionele conservering is mogelijk niet het geval voor het *tra* gen, aangezien dit gen afwezig lijkt te zijn (of niet te detecteren is) in verschillende groepen insecten (Hoofdstuk 2). *Tra* lijkt meermaals verloren te zijn gegaan uit de geslachtsbepalingscascade of, in een alternatief scenario, meermaals te zijn toegevoegd aan de geslachtsbepalingscascade. *Tra* is gedupliceerd in verschillende families van de Hymenoptera, ook in soorten zonder CSD, aangezien ik paralogen heb gevonden in *Leptopilina* wespen (Hoofdstuk 2 en 5). Het is echter nog onduidelijk of de wijdverbreide duplicaties betrokken zijn bij geslachtsbepaling, of dat nu is als een CSD locus of als een factor met maternale effecten.

Maternale provisie in seksueel voortplantende wespen

Maternale provisie van zowel *tra2* (Hoofdstuk 3) en vrouwelijk specifieke *tra* is nodig voor de start van vrouwelijke ontwikkeling in *N. vitripennis*. De toevoeging van *tra2* mRNA in de eieren is een geconserveerd element van gen-regulatie in geslachtsbepaling. Verminderde hoeveelheden *tra2* mRNA in embryo's resulteerde in de ontwikkeling van diploïde mannelijke individuen in *N.*

vitripennis (Hoofdstuk 3). Verder was er ook een sterke vermindering van hoeveelheid nakomelingen te zien. Dit geeft aan dat de vroege aanwezigheid van *tra2*-transcripten in het embryo ook andere functies in de ontwikkeling van haploïde embryo's heeft. Daarnaast observeerde ik een sterkere daling in het aantal diploïde nakomelingen met een paternale chromosoomset na vermindering van zowel *tra*- als *tra2*-transcripten in vroege stadia van ontwikkeling.

Ondanks dat vrouwelijk specifieke *tra*-transcripten cruciaal zijn tijdens de vroege ontwikkeling van *N. vitripennis* en maternaal meegeleverd worden, is dit niet het geval in de wesp *A. tabida* (Hoofdstuk 4). In plaats daarvan geven *A. tabida* moeders alternatieve splice varianten van *tra* mee aan hun eieren. Deze coderen voor een korter eiwit, dat echter een belangrijk domein bevat en niet zo kort is als de mannelijk specifieke eiwitten. Het gebrek aan vrouwelijk specifieke *tra* in de vroege stadia van diploïd vrouwelijke ontwikkeling opent echter de vraag hoe de vrouwelijk activiteit gestart wordt in dit systeem. De alternatieve splice vormen van *tra* kunnen mogelijk een rol spelen in de activatie van de vrouwelijk specifieke transcriptie van *tra* in het vroege embryo. Daarnaast zou echter een factor nodig zijn om het verschil tussen haploïde en diploïde ontwikkeling aan te geven, mogelijk een actieve paternale factor of een herkenning van dosis tussen een enkel of een dubbel genoom. De cruciale rol van *tra2* maternale provisie lijkt wel geconserveerd te zijn in *A. tabida* geslachtsbepaling. Deze transcripten kon ik detecteren in embryos voorafgaand aan zygotische transcriptie en tijdens de zygotische activatie van de geslachtsbepalingcascade (Hoofdstuk 4).

Verstoring van geslachtsbepaling door endosymbionten

Bij insecten waarvan het geslachtsbepalingsmechanisme is gebaseerd op een zygotische staat van *tra* in de mannelijke modus, is een aanzet voor vrouwelijke ontwikkeling nodig die de geslachtsbepalingcascade in de vrouwelijke modus plaatst. Deze aanzet hoeft niet perse vanuit het insect zelf te komen. Endosymbionten zijn wijdverspreid in veel insect-orde en kunnen drastische effecten hebben op de voortplanting van hun gastheren. Endosymbionten worden alleen doorgegeven via het cytoplasma. Vanwege deze eenzijdige transmissie is het in hun voordeel om genetisch mannelijke embryo's op een pad van vrouwelijke ontwikkeling te zetten of om deze mannelijke embryo's te verwijderen en het percentage vrouwelijke nakomelingen hoger te maken. In haplodiploïde soorten kunnen endosymbionten ervoor zorgen, dat de gastheren zich asexueel voortplanten. Geïnfecteerde diploïde moeders leggen geïnfecteerde haploïde eieren, die diploïd gemaakt en gefeminiseerd worden tot een volgende generatie van geïnfecteerde diploïde vrouwelijke individuen.

De parasitaire wespen *Asobara japonica* en *Leptopilina clavipes* hebben elk seksueel en asexueel voortplantende populaties. Asexuele wespen van beide soorten zijn geïnfecteerd door *Wolbachia* endosymbionten. Voor *A. japonica* is gedocumenteerd, dat de feminisering een aparte stap na de diploïdisatie is. De *Wolbachia* infectie kan verwijderd worden met antibiotica-behandelingen, die resulteren in de productie van haploïde mannelijke wespen.

Seksueel voortplantende *A. japonica* en *L. clavipes* wespen geven geen vrouwelijk specifieke *tra*-transcripten mee aan hun oöcyten (Hoofdstuk 5 en 6). Dit fenomeen komt overeen met het patroon van *tra*-transcripten, dat ik detecteerde in de seksuele voortplanting van *A. tabida* (Hoofdstuk 4). Aseksuele *A. japonica* en *L. clavipes* wespen geven echter wel vrouwelijk specifieke *tra*-transcripten mee aan hun oöcyten in de aanwezigheid van *Wolbachia*. Wanneer ik *Wolbachia* verwijderde uit de aseksuele *A. japonica* wespen, gaven zij geen vrouwelijk specifieke *tra*-transcripten meer mee, maar waren ze in plaats daarvan terug gegaan naar het patroon van *tra*-transcripten dat wordt meegegeven in seksuele wespen (Hoofdstuk 6). Hieruit concluderend lijkt *Wolbachia* de controle over geslachtsbepaling te hebben overgenomen door de maternale provisie van vrouwelijk specifieke *tra*-transcripten te manipuleren.

Ouderlijke en endosymbiont effecten op geslachtsbepaling – wie heeft de controle?

Maternale provisie en de geslachtsspecieke transcripten van geslachtsbepalingsgenen gedurende vroege stadia van ontwikkeling bieden kostbare informatie aangaande de nog te ontdekken moleculaire mechanismen van geslachtsbepaling. Aanwezigheid van vrouwelijk specifiek *tra* is niet noodzakelijk voor vrouwelijke ontwikkeling in drie seksueel voortplantende wesp soorten (*A. tabida*, *A. japonica* en *L. clavipes*) (Hoofdstuk 4, 5 en 6). Maternale provisie van *tra2* lijkt daarentegen wijdverbreid noodzakelijk te zijn voor vrouwelijke ontwikkeling. Andere maternale elementen, bijvoorbeeld alternatieve splice varianten van *tra* of paralogen daarvan, kunnen ook vereist zijn voor het starten van de vrouwelijke geslachtsbepalingcascade. Echter, alleen maternale provisie is niet afdoende om vrouwelijke ontwikkeling te starten in seksueel voortplantende wespen. Het paternale genome zou een epigenetisch verschil met het maternale genoom moeten bezitten om vrouwelijk specifieke transcriptie te starten in diploïde bevruchte eieren. Deze afhankelijkheid van de chromosoom set van de vader kan hypothetisch ook gemanipuleerd worden door *Wolbachia*, waarbij de endosymbiont de activiteit van het paternale genoom kan nabootsen. Alternatief kan het, in plaats van een verschil tussen het maternale en paternale genoom, echter ook het dosis effect van de diploïde staat van het embryo zijn die herkend wordt en vrouwelijke ontwikkeling start. De verstoring van *tra* maternale provisie suggereert, dat gastheer en endosymbiont elkaar proberen te overheersen in een conflict over de controle van haplodiploïde geslachtsbepaling. Dit roept de vraag op: wie heeft zeggenschap over geslachtsbepaling, en wie zou het beheer kunnen overnemen?

Curriculum vitae

Elzemies Geuverink was born on the 15th of February 1985 in Enschede, The Netherlands. She started her BSc Biology at the University of Groningen in 2003 and graduated in 2006 with a specialization in Ecology. In 2006 she continued with her MSc Ecology studies. Her MSc research included a project on gametic isolation in *Nasonia* wasps, a study on above- and belowground interactions in a salt marsh ecosystem and the implementation of in situ hybridization techniques in *Nasonia*. She graduated with honours (*cum laude*) in 2008 and started as an Early Stage Researcher at the University of Jyväskylä (Finland) within the Marie Curie Initial Training Network "Speciation". In Jyväskylä she studied the variation in developmental mode and reproduction in *Pygospio* worms (at the University of Jyväskylä) and *Littorina* snails (at the University of Sheffield). In 2010 she started the PhD project presented in this thesis at the University of Groningen. During 2014-2015 she worked as a teaching assistant, and currently she is working as a researcher in the Evolutionary Genetics, Development & Behaviour cluster at the University of Groningen studying the mechanisms with which *Wolbachia* interferes with sex determination in parasitoid wasps.

List of publications

Geuverink E & Beukeboom LW (2014) Phylogenetic distribution and evolutionary dynamics of the sex determination genes *doublesex* and *transformer* in insects. *Sexual Development*, **8**, 38-49

Kesäniemi JE, Geuverink E & Knott KE (2012) Polymorphism in developmental mode and its effect on population genetic structure of a spionid polychaete, *Pygospio elegans*. *Integrative and Comparative Biology*, **52**, 181-196

The Marie Curie Speciation Network (2012) What do we need to know about speciation? *Trends in Ecology and Evolution*, **27**, 27-39

Veen GF, Geuverink E & Olff H (2012) Large grazers modify effects of aboveground-belowground interactions on small-scale plant community composition. *Oecologia*, **168**, 511-518

Geuverink E, Gerritsma S, Pannebakker BA & Beukeboom LW (2009) A role for sexual conflict in the evolution of reproductive traits in *Nasonia* wasps? *Animal Biology*, **59**, 417-434

Submitted:

Geuverink E, Rensink AH, Rondeel I, Beukeboom LW, Van de Zande L & Verhulst EC (submitted) Maternal provision of *transformer-2* is required for female development and embryo viability in *Nasonia vitripennis*

Acknowledgements

At some point - a long long time ago- the completion of this thesis became a running joke. Somehow (and it felt as a strange fast-forward in the very last stages), the end is near. I want to thank all the people that helped make this thesis possible, but also all the people who worked on all the projects that did not end up in here.

Leo welcomed me back in Groningen, after a not so fruitful stint around Europe. I hope you haven't regretted this, even at times when you wished I could set my own deadlines. Thanks for the enthusiasm at even the tiniest discoveries and the support you showed throughout my PhD project, but also in earlier years. Louis, you could always point out exactly what was wrong, or more likely, misplaced or twisted around, in my manuscripts. Hopefully I've learned from this (better than these sentences appear to indicate), and thanks for all the feedback on whichever crazy idea I came up with. Eveline, even when you were not in our lab anymore I could count on your eagerness to discuss any project and to provide advice, at the strangest hours and locations. And then I dared to set the defence on the one day you were not available.

I want to thank the reading committee, Martin Beye, Jacintha Ellers and Jean-Christophe Billeter, for judging this manuscript and doing this in such a limited amount of time that we could actually make the ambitiously set defence date.

This restricted schedule to make the defence date caused me many anxious moments. Thanks, Anna and Syl, for calming me down and dealing with my neuroticism not only during this period but throughout my time in Groningen. The Evolutionary Genetics group has been a comfy home for many years. I want to thank all the PhD students who shared this place over the years and everyone else who wandered in for discussion, borrels and all the silliness.

Each chapter in this thesis consisted of a whole team of people that made the studies possible. Ken, Pieter, Bregje, Bart and the Van Gogh team, thank you for the advice and discussion on the experiments and the analyses of the genomic data. In particular I want to thank Wen-Juan, who I could count on to question anything at any time. Discussions with Daniel, the Nöthiger "crew" and many others were a necessity to me as a molecular biology newbie. Almost every chapter started with pilot studies or even entire first versions of experiments: thanks to Ammerins, Maria, Charlotte and Marin for conducting those tests. Many students joined us for a project for which I would like to thank Sean, Teun, Inge and Sandra. Marloes arrived at a critical moment and ensured that the work for this thesis could be completed. Rogier and Emma, thanks for the piles of bottles and paperwork that set the groundwork for this operation.

Gemma, sorry for timing my defence so poorly that you had to miss it. Thanks for all the work-outside-of-work discussion and even letting me crash in your office. This thesis and many other pieces of writing would have taken a lot longer without Jan van den Burg and the Herdershut. I hope we can stay there for many more highly effective writing camps.

Many thanks go to Pascal, my family and friends for listening to me going on and on about work in this final stage of my PhD. I apologize for nearly driving you insane by my unavailability for

anything that was not lab-located and my general absentmindedness. I even dragged you on fieldwork trips and into the lab on weekends to help with the wasp cultures, and I want to thank my parents for fuelling the madness so much so that they go on their own field trips.

The last thank you note is reserved for those I spend most of the time with ... the tiny workers ... all the countless wasps that were sacrificed.

And on that note I can now say: this thesis is finished, finally.

