

University of Groningen

A metal ion regulated artificial metalloenzyme

Bersellini, Manuela; Roelfes, Gerard

Published in:
 Dalton Transactions

DOI:
[10.1039/c7dt00533d](https://doi.org/10.1039/c7dt00533d)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
 Bersellini, M., & Roelfes, G. (2017). A metal ion regulated artificial metalloenzyme. *Dalton Transactions*, 46(13), 4325-4330. <https://doi.org/10.1039/c7dt00533d>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Cite this: *Dalton Trans.*, 2017, **46**, 4325

Received 13th February 2017,
Accepted 27th February 2017

DOI: 10.1039/c7dt00533d

rsc.li/dalton

A metal ion regulated artificial metalloenzyme†

Manuela Bersellini and Gerard Roelfes*

Regulation of enzyme activity is essential in living cells. The rapidly increasing number of designer enzymes with new-to-nature activities makes it necessary to develop novel strategies for controlling their catalytic activity. Here we present the development of a metal ion regulated artificial metalloenzyme created by combining two anchoring strategies, covalent and supramolecular, for introducing a regulatory and a catalytic site, respectively. This artificial metalloenzyme is activated in the presence of Fe^{2+} ions, but only marginally in the presence of Zn^{2+} .

Introduction

Control over enzymatic activity is vital for regulating the complex network of biosynthetic pathways in living cells. An important strategy employed by nature to regulate enzyme activity is allosteric regulation, that is binding of an effector molecule or metal ion at a site in the protein other than the active pocket.^{1–3} As a result of this interaction, the catalytic activity of an enzyme and, hence, its activity in a biosynthetic pathway can be regulated by the presence or absence of chemical signals in the environment. Such allosteric regulation would also be desirable for controlling the activities of designer enzymes with new-to-nature activities. Additionally, such artificial allosteric enzymes could also be of interest for other applications, for example as sensors. In this work we present an artificial metalloenzyme that is selectively activated by a specific metal ion, that is Fe^{2+} , but not by addition of Zn^{2+} .

Artificial metalloenzymes are hybrid catalysts that incorporate non-natural metal cofactors into biological scaffolds, thereby giving access to enzyme-like catalysts with new-to-nature catalytic activities.^{4–7} In recent years, this strategy has been utilized to create a wide variety of artificial metalloenzymes that display non-natural activities within a biological environment. However, examples of such hybrid enzymes that can be activated by mechanisms akin to allosteric control remain scarce. Indeed, the only example for this type of activation comes from Ward and colleagues, who has recently demonstrated that the activity of an artificial asymmetric trans-

fer hydrogenase could be upregulated by proteolysis when a natural protease is used as an external stimulus.⁸

We have introduced a novel class of artificial metalloenzymes based on the transcriptional regulator LmrR (Lactococcal multidrug resistance Regulator). For this protein a new active site can be created within the hydrophobic cavity that is present at the dimer interface.^{9,10} An attractive feature of LmrR as biomolecular scaffold for artificial metalloenzymes is that it is the only scaffold reported so far amenable to three different anchoring strategies for incorporation of non-natural cofactors: covalent and supramolecular anchoring and unnatural amino acid incorporation.^{11–14} The resulting hybrid catalysts have been applied in a variety of catalytic reactions including enantioselective Diels Alder, water addition to α,β -unsaturated ketones and Friedel–Crafts alkylation reactions.

For the creation of a metal regulated artificial metalloenzyme, we took advantage of the design versatility of LmrR and combined two anchoring strategies for binding transition metal complexes: incorporation of a catalytic site *via* supramolecular interactions and introduction of a regulatory site *via* covalent anchoring of a ligand. The incorporation of the catalytic site *via* the supramolecular approach is based on the binding of Cu^{2+} complexes of planar aromatic ligands (such as phenanthroline) between the two tryptophans located inside the hydrophobic pocket at the dimer interface of LmrR at positions 96 and 96' (Fig. 1a). This approach was applied to catalysis of the enantioselective Friedel–Craft alkylation of α,β -unsaturated imidazoles to indoles using Cu^{2+} phenanthroline complexes, such as $[\text{Cu}(\text{phen})(\text{NO}_3)_2]$, obtaining good conversions and excellent enantioselectivities of >90% in several cases.¹⁴

The introduction of the regulatory site *via* covalent anchoring of a metal binding moiety to LmrR is based on the site-selective modification of the protein using unique cysteines as bioconjugation handles for alkylation reactions.^{11,12} Since

Stratingh Institute for Chemistry, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands. E-mail: j.g.roelfes@rug.nl; http://roelfesgroup.nl

† Electronic supplementary information (ESI) available: Full experimental details and characterization, results of control and additional experiments. See DOI: 10.1039/c7dt00533d




Fig. 1 (a) Surface representation of the front entrance of LmrR (PDB 3F8F).¹⁰ Zoom in of part of the hydrophobic pocket at the dimer interface. Tryptophans in position 96 and 96' are shown in blue. (b) Schematic representation of the concept of the metal regulated artificial metalloenzyme.

LmrR is a homodimeric protein, the introduction of a ligand in a specific position *via* covalent anchoring will lead to functionalization of both monomers, resulting in a dimeric protein containing two metal binding moieties.

Results and discussion

Our design of a metal ion regulated artificial metalloenzyme is depicted in Fig. 1b. In absence of an effector metal ion, the metal binding moieties covalently bound to LmrR will bind the catalytically active complex [Cu(phen)(NO₃)₂]. As a result, the two free coordination sites on the Cu²⁺ ion required for substrate binding are blocked, rendering the artificial metalloenzyme inactive. Conversely, addition of transition metal ions that bind stronger than Cu²⁺, would trap the metal binding moieties into stable chelate complexes, allowing the active Cu²⁺ complex to bind and activate the substrates (Fig. 1b).

Previous work with LmrR based metalloenzymes showed that enantioselective reactions take place inside the hydrophobic pocket in close proximity to the front entrance. Therefore it was envisioned that the formation of chelate metal complexes in this area might also have an effect on the catalysis.^{11,12}

2-2'-Bipyridine was selected as metal binding moiety due to its ability to strongly bind late-first row transition metal ions with high affinity and in different stoichiometries.^{15–19} The bipyridine ligands were introduced on solvent exposed positions on the α 4 helices, which delimit the front entrance of the hydrophobic pocket. Introduction of bipyridine ligands on these helices was envisioned to allow the formation of chelate complexes involving one bipyridine moiety of each monomer.

Positions 93 and 104, located in the central part of the front entrance, and position 86, which is located towards the edge of the pocket, were chosen for the introduction of the bipyridine ligands (Fig. 2a).

Plasmids encoding C-terminally Strep-tagged cysteines mutants of LmrR (F93C, E104C, H86C) were prepared by standard site-directed mutagenesis technique (Quick-Change) from a vector available from previous work.¹¹

LmrR mutants were subsequently expressed in *E. coli* BL21 (DE3) C43 and purified by affinity chromatography (Strep-tactin Sepharose column) and cation exchange chromatography (Heparin column) to remove residual bound DNA. Target proteins were obtained in good yields (10–20 mg L⁻¹) with excellent purity as judged by Tricine-SDS-PAGE, UPLC-MS and analytical size exclusion chromatography (Fig. S1†).

Bipyridine units were introduced by selective alkylation of cysteines with a bromo acetamide derivative of 2-2'-bipyridine (Fig. 2b).^{11,12} UPLC-MS and Tricine-SDS-PAGE were used to confirm identity and purity of LmrR bipyridine conjugates and analytical size exclusion chromatography demonstrated that the dimeric structure of LmrR was not significantly altered by introduction of the metal binding moieties (Fig. S2–S4†).

With the LmrR bipyridine conjugates in hand, first the binding of these conjugates to Zn²⁺ and Fe²⁺ was studied by UV-visible titrations. These two ions are examples of divalent late-first row transition metal ions with different coordination properties. Upon addition of either metal ion, the appearance of a shoulder around 310 nm was observed, indicative of a change in the π - π^* transition of the bipyridine upon metal coordination (Fig. 3a and S5†). This red shift has previously been reported for bipyridine ligands in solution and for proteins containing a bipyridine moiety.^{18,22,23} In the titration with Fe²⁺ also the appearance of two additional absorption bands between 490 and 530 nm was observed, which are characteristic of Fe²⁺ bipyridine complexes (Fig. 3a and S5†).^{17,19,24,25} The resulting titration




Fig. 2 (a) Surface and helix representation of LmrR front entrance. Residues H86, F93, E104 are shown in blue, green and red, respectively. (b) Reaction scheme of the synthesis of protein–metal binding moieties conjugates. Ligand used for alkylation reaction on LmrR cysteine mutants: *N*-((2,2'-bipyridin-5-yl)methyl)-2-bromoacetamide (**1**).^{20,21}




Fig. 3 (a) UV-visible titrations of LmrR E104C_bpy with $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (top) and $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ (bottom). Titrations were performed in 50 mM NaH_2PO_4 pH 7.0, 500 mM NaCl with 22.5 μM LmrR_bipyridine conjugate solutions (dimer) with subsequent additions from 1 mM solutions of metal salts in milliQ water. Inset containing the fitting of the titration data obtained with non-linear regression. (b) Job's plot graphs of LmrR E104C_bpy with $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (top) and $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ (bottom). Plots of ΔA at 530 nm for Fe^{2+} and 311 nm for Zn^{2+} against the molar fraction (x) of the Fe^{2+} and Zn^{2+} , respectively. 100 μM stock solutions of LmrR conjugates in 50 mM NaH_2PO_4 pH 7.0, 500 mM NaCl and 200 μM of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ or $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ in milliQ water. Total concentration in the cuvette was kept constant at 60 μM .

curves could be fitted to a 1 : 1 binding model, corresponding to the formation of complexes containing one metal ion per LmrR dimer (Table S2†). The stoichiometry of the metal : protein complexes was further confirmed by Job's plot (Fig. 3b and S6†), in which the maximum complex formation was obtained with equimolar concentrations of metal ions and LmrR dimer. ICP-AES experiments performed after incubation of the LmrR_E104C_bipyridine conjugate with Fe^{2+} and Zn^{2+} , followed by dialysis to remove unbound metal ions, also suggested the presence of one equivalent of metal ion per LmrR dimer (Table S4†).

Combined these observations suggest the formation of a chelate metal complex between one metal ion and two bipyridines in the LmrR scaffold. Exogenous water molecules or ligands supplied by amino acid side chains might complete the coordination sphere around the metal. The formation of a chelate complex between the two bipyridines and the metal ions was expected for the conjugates containing bipyridine units located in the central part of the front entrance (LmrR F93C_bpy and LmrR E104C_bpy), given the proximity of the two amino acids in the protein dimer (10–15 Å from the reported crystal structure).¹⁰ However, for the conjugate in which the bipyridine units are located towards the edge of the hydrophobic pocket (LmrR H86C_bpy) the formation of a chelate metal complex appeared unlikely due to the distance between the two residues in position 86 (28–30 Å).¹⁰ Tentatively, chelation of the metal ions could be achieved due

to the known flexibility of the LmrR scaffold.¹⁰ The cost for formation of these rather unfavoured complexes is reflected in a lower binding affinity for both regulatory metals complexes (Table S2†) and an apparent lower stability of the metal-bound protein, which was prone to precipitation. However, also the possibility of formation of chelate metal complexes with bipyridine units from separate LmrR dimers should be considered, as suggested by the presence of higher order aggregates in the size exclusion chromatography of some Fe^{2+} conjugates (LmrR E104C_bpy and LmrR H86C_bpy) (Fig. S4†).

The catalytic activity of the LmrR_bipyridine conjugates and their metal complexes, was evaluated in the enantioselective, vinylogous Friedel–Crafts alkylation of 1-(1-methyl-1*H*-imidazol-2-yl)but-2-en-1-one (**1**) with 2-methyl-1*H*-indole (**2**).^{13,14,26–29}

The metal complexes of LmrR bipyridine conjugates with Fe^{2+} and Zn^{2+} (LmrR_bpy_M²⁺) were prepared fresh prior to catalysis by incubation of 1 equivalent of the corresponding metal salt with 1 equivalent of the LmrR_bpy conjugate (dimer) in 20 mM MOPS, pH 7.0, 500 mM NaCl (4 °C, 30 minutes). The mixtures were centrifuged to remove possible precipitate and the supernatant containing the LmrR_bipyridine_M²⁺ complex was used for the supramolecular assembly of the artificial metalloenzyme. The artificial metalloenzymes were prepared *in situ* by self-assembly from [Cu(phen)(NO₃)₂] (22 μM) with a slight excess (1.3 equivalents) of LmrR_bipyridine_M²⁺ complex (30 μM) in 20 mM MOPS, pH 7.0, 500 mM NaCl. The mixture was incubated at 4 °C for 30 minutes and the catalytic reaction was initiated by addition of substrates **1** and **2** at a final concentration of 1 mM. Reactions were incubated at 4 °C for 72 h, after which the products were isolated and analyzed by chiral HPLC.

As previously reported, the reaction of the α - β unsaturated acyl imidazole **1** with 2-methyl indole **2** catalyzed by the assembly of [Cu(phen)(NO₃)₂] with wt-LmrR is protein accelerated and results in excellent enantioselectivity up to 92% (Table 1, entries 1 and 2).¹⁴ As expected, using the LmrR_bipyridine conjugates in absence of any regulatory metal ion resulted in no significant catalytic activity of the artificial metalloenzymes with LmrR F93C_bpy and LmrR E104C_bpy, and low activity in case of LmrR H86C_bpy (Table 1, entries 3–5). This is due to the fact that the free coordination sites of the Cu²⁺ ion in the [Cu(phen)(NO₃)₂] complex are sequestered by the conjugated bipyridine ligands, thus preventing substrate binding (Fig. 1b). In case of LmrR H86C_bpy, this sequestration is apparently not perfect, for reasons not understood at present.

Formation of metal complexes with either Fe^{2+} or Zn^{2+} prior to catalysis, resulted in almost no product formation for the conjugate LmrR F93C_bpy (Table 1, entries 6 and 7). The observed lack of catalytic activity might be due to bulkiness of the chelate metal complexes located right in the middle of the front entrance of the hydrophobic pocket that does not allow the substrates to reach the active Cu²⁺ center located inside. This hypothesis is supported by the observation that no specific binding of the dye Hoechst 33342 (H33342), which is known to bind inside the hydrophobic pocket of LmrR *via* the

Table 1 Results of vinylogous Friedel–Crafts alkylation reactions catalyzed by LmrR_bpy conjugates and their respective Fe²⁺ and Zn²⁺ complexes

Entry	Catalyst	Yield ^a (%)	TON ^b	ee ^c (%)
1	Cu(phen)(NO ₃) ₂ (only)	39 ± 4	16 ± 2	<5
2	wt-LmrR	58 ± 3	24 ± 1	92 ± 1
3	LmrR F93C_bpy	≤1	—	n.d.
4	LmrR E104C_bpy	≤1	—	n.d.
5	LmrR H86C_bpy	7 ± 1	3 ± 1	81 ± 4
6	LmrR F93C_bpy + Fe ²⁺	2 ± 0	1 ± 0	n.d.
7	LmrR F93C_bpy + Zn ²⁺	2 ± 1	1 ± 1	n.d.
8	LmrR E104C_bpy + Fe ²⁺	34 ± 4	14 ± 2	75 ± 6
9	LmrR E104C_bpy + Zn ²⁺	4 ± 0	2 ± 0	n.d.
10	LmrR H86C_bpy + Fe ²⁺	26 ± 6	11 ± 2	88 ± 3
11	LmrR H86C_bpy + Zn ²⁺	14 ± 2	6 ± 1	80 ± 10

Typical conditions: 2.2 mol% [Cu(phen)(NO₃)₂] (22 μM) loading with 1.3 eq. LmrR_bpy conjugates (30 μM), 1 mM of substrate **1** and **2** in 20 mM MOPS buffer pH 7.0, 500 mM NaCl, for 72 h at 4 °C. All the results listed correspond to the average of two independent experiments, each carried out in duplicate. Errors are listed as standard deviations. ^a Yields were determined by HPLC and using 2-phenylquinoline as internal standard. ^b Turnover numbers (TON) were determined by dividing the concentration of product by the catalyst concentration. ^c For yields below 5% ee's were not determined.

front entrance, to LmrR F93C_bpy was observed using fluorescence spectroscopy (Fig. S7†).¹⁰

We were pleased to find that upon incubation with the regulatory metal ions (Fe²⁺ or Zn²⁺) prior to addition of the active [Cu(phen)(NO₃)₂] complex, two of conjugates, *i.e.* LmrR E104C_bpy and LmrR H86C_bpy, showed catalytic activity (Table 1, entries 8–10), albeit lower than with wt-LmrR. These results suggest that the activity of these artificial metalloenzymes can be regulated by the presence of metal ions.

It was observed consistently that activation with Fe²⁺ yielded more active artificial metalloenzymes when compared to the addition of Zn²⁺: higher turnover numbers (TON) were achieved for the artificial metalloenzymes containing Fe²⁺ complexes compared to the respective Zn²⁺ complexes (Table 1, entries 8–11). It was hypothesized that the difference in activity might reflect the different interactions of the active transition metal complex ([Cu(phen)(NO₃)₂]) with the two types of metal complexes (Fe²⁺ and Zn²⁺ complex). Nevertheless, as described above, the Fe²⁺ complexes of these two conjugates, in addition the dimeric form of LmrR, appeared to form higher order aggregates as well. At this moment it is not clear what effect these aggregates have on catalysis.

As this difference in activity is particularly pronounced for the LmrR E104C_bpyridine conjugate, this variant was selected for further investigation. According to the Irving–Williams series, six-coordinate octahedral Cu²⁺ complexes are thermodynamically more stable than other high spin divalent first-row transition metal complexes. Hence, in our design, the Cu²⁺ from the active [Cu(phen)(NO₃)₂] complex could replace Zn²⁺ from the high spin bipyridine complexes, forming a mixed complex with phenanthroline and the covalently bound bipyridine as ligands. As this additional interaction would prevent substrate binding, the resulting artificial metallo-

enzyme becomes inactive for the desired reaction. Conversely, Fe²⁺ is known to predominantly form low spin complexes with bipyridine ligands, which are kinetically more stable and, thus, less prone to ligand substitution. Therefore, exchange of Fe²⁺ by Cu²⁺ will not occur and the artificial metalloenzyme will remain active.^{30,31}

To test this hypothesis, reactions were performed using an excess of Fe²⁺ or Zn²⁺ ions. The LmrR_E104C_bpy conjugates were incubated with 1, 3 and 6 equivalents of metal ions and in case of Zn²⁺ also 10 equivalents (with respect to LmrR dimer) before the artificial metalloenzyme was assembled by addition of [Cu(phen)(NO₃)₂]. As expected, when additional Fe²⁺ equivalents were added, no significant change in catalytic activity was observed. In contrast, when going from 1 to 10 equivalents of Zn²⁺, an increase in activity was observed from 7 to 12 TON. With increasing amounts of Zn²⁺, also an erosion of the ee was observed, which most likely is due to structural changes of the chiral active site as a result of non-specific Zn²⁺ binding to the protein.

The possibility of metal exchange between the regulatory metal ion (Fe²⁺ or Zn²⁺) and the catalytic metal ion (Cu²⁺) was further investigated by ICP-AES measurements. LmrR_E104C_bpy_M²⁺ complexes were incubated with Cu²⁺ ions, followed by dialysis to remove non-bound metal ions (Fig. 4).

Quantification of the regulatory metal (Fe²⁺ or Zn²⁺) bound to the protein after incubation with [Cu(phen)(NO₃)₂] did not show a change in the total amount of Fe²⁺ or Zn²⁺ (Table S3†) because metal exchange will result in the formation a mixed Cu²⁺ phenanthroline bipyridine complex, while the regulatory metal might still be bound to the second bipyridine in the LmrR scaffold (Fig. 5a). In order to push the equilibrium towards a complete metal replacement, leading to a Cu²⁺ bisbipyridine complex and the release of the regulatory metal ion,




Fig. 4 Comparison of turnover numbers (black) and enantioselectivity (purple) in the vinylogous Friedel–Crafts alkylation reactions catalyzed by LmrR_bpy_E104C_bpy conjugate and respective Fe²⁺ and Zn²⁺ complexes in presence of increasing concentration of Fe²⁺ (left) and Zn²⁺ (right). Same reaction conditions as in Table 1.




Fig. 5 Schematic representation of metal ion exchange between regulatory metal ion (M²⁺) and the catalytic metal ion (Cu²⁺) with [Cu(phen)(NO₃)₂] and Cu(NO₃)₂.

incubation of the LmrR_bpy_M²⁺ complexes with Cu(NO₃)₂ was performed (Fig. 5b).

Quantification of the regulatory metal (Fe²⁺ or Zn²⁺) bound to the protein after incubation with Cu(NO₃)₂ showed a significant decrease in the amount of Zn²⁺ bound to the protein (Table S3 and Fig. S8[†]), confirming the hypothesis of partial replacement of Zn²⁺ with Cu²⁺ with subsequent inactivation of the catalyst. The observation that the metal displacement reaction does not seem to happen for the Fe²⁺ complex supports the formation of a stable low spin Fe²⁺ bisbipyridine complex within the LmrR scaffold.³¹

Finally the reversibility of the designed metal regulated artificial metalloenzyme was tested by inverting the order of

addition of the regulatory and catalytic metal ions. LmrR_bipyridine conjugates were first incubated with [Cu(phen)(NO₃)₂] at 4 °C for 30 minutes and then 1 equivalent of Fe²⁺ or Zn²⁺ salt was added. After incubation of the resulting mixture for 30 minutes substrates **1** and **2** were added at a final concentration of 1 mM. In parallel, exactly the same procedure was applied to the LmrR_E104C_bpy conjugates by incubating first the regulatory metal ions (Fe²⁺ or Zn²⁺) and then the active catalyst [Cu(phen)(NO₃)₂]. Reactions were incubated at 4 °C for 72 h, after which the products were isolated and analyzed by chiral HPLC. We were delighted to observe that catalytic activity (as well as enantioselectivity) could be restored to a large extent upon addition of Fe²⁺ salt to the inactive metalloenzyme (Table 2). In contrast, addition of Zn²⁺ salt did not restore the activity. This result suggests that selective regulation of activity of the designed artificial metalloenzyme by Fe²⁺ ions is reversible.

Conclusions

The results presented here show that we successfully designed, synthesized and characterized a metal ion regulated artificial metalloenzyme. By combining a regulatory site to bind an effector metal ion and an active site to recruit a catalytically active metal complex in the LmrR scaffold, the activity of the artificial metalloenzyme (LmrR_E104C_bpy) for a vinylogous Friedel–Crafts alkylation could be regulated by incubation with Fe²⁺ ions. Reminiscent of allosteric regulation in natural enzymes, we achieved selective activation by Fe²⁺ but not by Zn²⁺, taking advantage of the different coordination properties of these transition metals. This study represents the first example of a metal ion regulated artificial metalloenzyme and presents a significant advance toward controlling the activity of designer enzymes in hybrid bio-synthetic pathways.

Acknowledgements

This work was supported by European Research Council (ERC Starting Grant 280010). Financial support from the Ministry of Education, Culture, and Science (Gravitation Program

Table 2 Effect of the order of incubation on the results of vinylogous Friedel–Crafts alkylation reactions catalyzed by the [Cu(phen)(NO₃)₂]/LmrR_bpy_E104C_bpy conjugate

Entry	1 st incubation	2 nd incubation	Yield ^a (%)	TON ^b	ee ^c (%)
1	[Cu(phen)(NO ₃) ₂]	—	≤1	≤1	n.d.
2	Fe ²⁺	[Cu(phen)(NO ₃) ₂]	57 ± 14	23 ± 5	72 ± 14
3	[Cu(phen)(NO ₃) ₂]	Fe ²⁺	33 ± 5	14 ± 2	78 ± 2
4	Zn ²⁺	[Cu(phen)(NO ₃) ₂]	13 ± 2	6 ± 1	50 ± 1
5	[Cu(phen)(NO ₃) ₂]	Zn ²⁺	11 ± 3	5 ± 1	50 ± 13

Typical conditions: 2.2 mol% [Cu(phen)(NO₃)₂] (22 μM) loading with 1.3 eq. LmrR_bpy conjugates (30 μM), 1 mM of substrates **1** and **2** in 20 mM MOPS buffer pH 7.0, 500 mM NaCl, for 72 h at 4 °C. All the results listed correspond to the average of two independent experiments, each carried out in duplicate. Errors are listed as standard deviations. ^a Yields were determined by HPLC and using 2-phenylquinoline as internal standard. ^b Turnover numbers (TON) were determined by dividing the concentration of product by the catalyst concentration. ^c For yields below 5% ee's were not determined.

no. 024.001.035) is gratefully acknowledged. The authors thank I. Drienovská and Dr L. Villarino for useful discussion and suggestions.

Notes and references

- N. M. Goodey and S. J. Benkovic, *Nat. Chem. Biol.*, 2008, **4**, 474–482.
- Q. Cui and M. Karplus, *Protein Sci.*, 2008, **17**, 1295–1307.
- J. Fastrez, *ChemBioChem*, 2009, **10**, 2824–2835.
- F. Rosati and G. Roelfes, *ChemCatChem*, 2010, **2**, 916–927.
- J. Bos and G. Roelfes, *Curr. Opin. Chem. Biol.*, 2014, **19**, 135–143.
- J. C. Lewis, *Curr. Opin. Chem. Biol.*, 2015, **25**, 27–35.
- E. M. Brustad and F. H. Arnold, *Curr. Opin. Chem. Biol.*, 2011, **15**, 201–210.
- Z. Liu, V. Lebrun, T. Kitanosono, H. Mallin, V. Köhler, D. Häussinger, D. Hilvert, S. Kobayashi and T. R. Ward, *Angew. Chem., Int. Ed.*, 2016, **55**, 11587–11590.
- J. Lubelski, A. De Jong, R. Van Merkerk, H. Agustiandari, O. P. Kuipers, J. Kok and A. J. M. Driessen, *Mol. Microbiol.*, 2006, **61**, 771–781.
- P. K. Madoori, H. Agustiandari, A. J. M. Driessen and A.-M. W. H. Thunnissen, *EMBO J.*, 2009, **28**, 156–166.
- J. Bos, F. Fuseti, A. J. M. Driessen and G. Roelfes, *Angew. Chem., Int. Ed.*, 2012, **51**, 7472–7475.
- J. Bos, A. García-Herraz and G. Roelfes, *Chem. Sci.*, 2013, **4**, 3578–3582.
- I. Drienovská, A. Rioz-Martínez, A. Draksharapu and G. Roelfes, *Chem. Sci.*, 2014, **6**, 770–776.
- J. Bos, W. R. Browne, A. J. M. Driessen and G. Roelfes, *J. Am. Chem. Soc.*, 2015, **137**, 9796–9799.
- G. Atkinson and J. E. Bauman, *Inorg. Chem.*, 1962, **1**, 900–904.
- R. H. Holyer, C. D. Hubbard, S. F. A. Kettle and R. G. Wilkins, *Inorg. Chem.*, 1965, **4**, 929–935.
- P. Krumholz, *J. Am. Chem. Soc.*, 1949, **71**, 3654–3656.
- R. M. Franzini, R. M. Watson, G. K. Patra, R. M. Breece, D. L. Tierney, M. P. Hendrich and C. Achim, *Inorg. Chem.*, 2006, **45**, 9798–9811.
- M. Kurihara, K. Ozutsumi and T. Kawashima, *J. Solution Chem.*, 1995, **24**, 719–734.
- A. Basnet, P. Thapa, R. Karki, Y. Na, Y. Jahng, B.-S. Jeong, T. C. Jeong, C.-S. Lee and E.-S. Lee, *Bioorg. Med. Chem.*, 2007, **15**, 4351–4359.
- R. Ballardini, V. Balzani, M. Clemente-León, A. Credi, M. T. Gandolfi, E. Ishow, J. Perkins, J. F. Stoddart, H.-R. Tseng and S. Wenger, *J. Am. Chem. Soc.*, 2002, **124**, 12786–12795.
- J. Xie, W. Liu and P. G. Schultz, *Angew. Chem., Int. Ed.*, 2007, **46**, 9239–9242.
- J. H. Mills, S. D. Khare, J. M. Bolduc, F. Forouhar, V. K. Mulligan, S. Lew, J. Seetharaman, L. Tong, B. L. Stoddard and D. Baker, *J. Am. Chem. Soc.*, 2013, **135**, 13393–13399.
- K. D. Barker, A. L. Eckermann, M. H. Sazinsky, M. R. Hartings, C. Abajian, D. Georganopoulou, M. A. Ratner, A. C. Rosenzweig and T. J. Meade, *Bioconjugate Chem.*, 2009, **20**, 1930–1939.
- H. K. Munch, S. T. Heide, N. J. Christensen, T. Hoeg-Jensen, P. W. Thulstrup and K. J. Jensen, *Chem. – Eur. J.*, 2011, **17**, 7198–7204.
- A. J. Boersma, B. L. Feringa and G. Roelfes, *Angew. Chem., Int. Ed.*, 2009, **48**, 3346–3348.
- D. A. Evans, K. R. Fandrick, H.-J. Song, K. A. Scheidt and R. Xu, *J. Am. Chem. Soc.*, 2007, **129**, 10029–10041.
- D. A. Evans, K. R. Fandrick and H.-J. Song, *J. Am. Chem. Soc.*, 2005, **127**, 8942–8943.
- T. B. Poulsen and K. A. Jørgensen, *Chem. Rev.*, 2008, **108**, 2903–2915.
- M. A. Halcrow, *Crystals*, 2016, **6**, 58.
- F. H. Burstall and R. S. Nyholm, *J. Chem. Soc.*, 1952, 3570–3579.