

University of Groningen

## Biochemical characterization of $\beta$ -galactosidases and engineering of their product specificity

Yin, Huifang

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2017

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Yin, H. (2017). *Biochemical characterization of  $\beta$ -galactosidases and engineering of their product specificity*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# Chapter 1

## **$\beta$ -Galactosidase enzymes and their galactooligosaccharide products**

## **Introduction**


Lactose ( $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 4)-D-glucose) is a disaccharide composed of one galactose molecule linked to a glucose molecule, which can be found in the milk of mammals [1], [2]. The content of lactose in bovine milk ranges from 4.4% to 5.2%, while human milk contains 7% lactose [1]. Human newborns have the ability to produce  $\beta$ -galactosidase (E.C. 3.2.1.23) enzymes to digest lactose, however, adults usually have lost the ability to produce this enzyme [3]. The absence or deficiency of  $\beta$ -galactosidase enzymes can lead to lactose intolerance upon the consumption of dairy products [4], [5]. Lactose can be removed from milk to solve this problem and be used as an ingredient in the food industry. This also provides opportunities for the development of high value-added lactose derivatives such as galactooligosaccharides (GOS). The market price of GOS is 10-12 times higher than that of lactose while the GOS prebiotic effects are widely accepted [6], [7], [8].

The concept of prebiotics was first introduced in 1995 by Gibson and Roberfroid [9], and now is defined as “a non-digestible compound that, through its metabolism by microorganisms in the gut, modulates composition and/or activity of the gut microbiota, thus conferring a beneficial physiological effect on the host” [10]. GOS are important commercial prebiotics which are added into infant formula as alternatives for human milk oligosaccharides (hMOS) [11], [12]. GOS are composed of a number of galactose units linked to a terminal glucose or galactose residue via different glycosidic bonds, with degrees of polymerization (DP) from 2 to 10 units [13]. Nowadays GOS are produced from lactose using microbial  $\beta$ -galactosidase enzymes [14], [15], [16].

## **Structures of $\beta$ -galactosidases and reaction mechanism**

$\beta$ -Galactosidase enzymes belong to glycoside hydrolase (GH) families 1, 2, 35, and 42, which belong to the GH-A superfamily (Carbohydrate Active Enzymes

database, <http://www.cazy.org/>) [17].  $\beta$ -Galactosidases in GH1, GH2 and GH42 are found predominantly in bacteria and fungi, whereas the enzymes in GH35 have been found in bacteria, fungi, animals and plants [18]. As shown in Table 1, crystal structures are available for several  $\beta$ -galactosidase enzymes. These  $\beta$ -galactosidase enzymes catalyze the hydrolysis of terminal  $\beta$ -galactose residues from various substrates. The  $\beta$ -galactose residues subsequently serve as growth substrates and are used for carbon metabolism, energy generation, and maintaining a normal physiological activity of the organisms. As shown in Figure 1, the domain organization of  $\beta$ -galactosidases from different GH families is quite different. For  $\beta$ -galactosidases in the GH2 family, the catalytic domain is the third domain from the N-terminus, while for  $\beta$ -galactosidases in GH35 and GH42 the catalytic domains are the first domains from the N-terminus [19], [20], [21], [22], [23]. The catalytic domain is a  $(\beta/\alpha)_8$  TIM barrel that contains one glutamic acid residue as the nucleophile and another glutamic acid residue as the proton donor [17],[24]. The  $\beta$ -galactosidases in the GH1, 2, 35, and 42 families display a retaining mechanism in their reaction. The catalytic nucleophile first attacks the anomeric center of lactose with the assistance of the proton donor, forming a galactosyl-enzyme intermediate while releasing glucose. The second step depends on the identity of the acceptor substrate: if water serves as the acceptor, the




**Figure 1.** Domain distribution of  $\beta$ -galactosidases from different Glycoside Hydrolase (GH) families [17], [18], [19], [20], [21].

intermediate undergoes hydrolysis and releases galactose; if lactose serves as acceptor substrate, a DP3 GOS ( $\beta$ -D-Galp-(1→x)- $\beta$ -D-Galp-(1→4)-D-Glcp) is

**Table 1.**  $\beta$ -Galactosidase enzymes and their protein 3D structures.

Protein name	GH family	Organism	PDB code	UniProt code	Reference
BglA	1	<i>Thermotoga maritima</i>	1OD0	Q08638	[26]
LacZ	2	<i>Arthrobacter</i> sp. C2-2	1YQ2	Q8KRF6	[27]
BgaD	2	<i>Bacillus circulans</i> ATCC 31382	4YPJ	E5RWQ2	[28]
LacZ	2	<i>Escherichia coli</i> K12	4V40	P00722	[29]
BgaL	2	<i>Paracoccus</i> sp. 32d	5EUV	D1LZK0	[30]
BgaA	2	<i>Streptococcus pneumoniae</i> serotype 4	4CU6	A0A0H2UP19	[31]
Lac4	2	<i>Kluyveromyces lactis</i> CBS2359	3OB8	P00723	[32]
BgaC	35	<i>Bacillus circulans</i> ATCC 31382	4MAD	O31341	[33]
Bgl35A	35	<i>Cellovibrio japonicus</i> Ueda107	4D1I	B3PBE0	[34]
BgaC	35	<i>Streptococcus pneumoniae</i> TIGR4	4E8C	A0A0H2UN19	[17]
LacA	35	<i>Aspergillus oryzae</i> RIB40	4IUG	Q2UCU3	[35]
Glb1	35	<i>Homo sapiens</i>	3THC	P16278	[36]
LacA	35	<i>Penicillium</i> sp.	1TG7	Q700S9	[37]
Tbg4	35	<i>Solanum lycopersicum</i>	3W5F	O81100	[16]
Bga1	35	<i>Trichoderma reesei</i>	3OG2	Q70SY0	[38]
Bca	42	<i>Bacillus circulans</i> subsp. <i>alkalophilus</i>	3TTS		[39]
Gal42A	42	<i>Bifidobacterium animalis</i> subsp. <i>lactis</i> BI-04 ATCC SD5219	4UNI		[21]
LacZ2	42	<i>Bifidobacterium bifidum</i> S17	4UCF	E3EPA1	[40]
GanB	42	<i>Geobacillus stearothermophilus</i> T-6	4OIF	F8TRX0	[41]
R- $\beta$ -Gal	42	<i>Rahnella</i> sp. R3	5E9A	A0A0B4U8I5	[42]
A4- $\beta$ -Gal	42	<i>Thermus</i> sp. A4	1KWG	O69315	[43]

formed by transgalactosylation (Figure 2) [25], [26], [27]. This DP3 GOS may serve again as acceptor substrate and undergo another round of transgalactosylation. The transgalactosylation reaction thus results in GOS mixtures containing structures varying in size and in linkage types. Carbohydrates other than lactose can also serve as acceptors in the transgalactosylation reaction. The linkage type and degree of polymerization (DP) of GOS structures strongly depend on the  $\beta$ -galactosidase enzyme origin.




**Figure 2.** Reaction scheme of  $\beta$ -galactosidase enzymes. This figure has been adapted from Bultema *et al* [25]. The acid/base catalyst and the nucleophile are both Glutamic acid residues. The hydrolysis reaction uses water as acceptor substrate, while the transgalactosylation reaction uses lactose and other carbohydrates as acceptor substrate.

## Identification of GOS structures produced by $\beta$ -galactosidases

As mentioned above, the characterized  $\beta$ -galactosidase enzymes adopt the same reaction mechanism. However, they synthesize different GOS products reflecting variations in their crystal structures, especially in their active sites [46],[47]. One of the aims in our work is to understand the relation between the active site structures and the synthesized GOS compounds. This requires the availability and use of reliable techniques to characterize the composition and structures of the GOS compounds. Techniques like high-performance anion-exchange chromatography (HPAEC) coupled with pulsed amperometric detection (PAD), NMR spectroscopy, Mass Spectrometry (MS), and methylation analysis have been widely used in the identification of the GOS structures [48], [49], [50], [51], [52]. This has resulted in the identification of an increasing number of GOS structures. Rodriguez-Colinas *et al.* identified 5 structures in the GOS mixture produced by  $\beta$ -galactosidase from *Kluyveromyces lactis* [53]. Urrutia *et al.* found 9 structures in the GOS mixture produced by  $\beta$ -galactosidase from *Aspergillus oryzae* [54]. Yanahira *et al.* isolated 11 GOS structures from the products of  $\beta$ -galactosidase of *Bacillus circulans* ATCC 31382 [55]. Our laboratory has made big strides to identify GOS compounds, using a series of techniques. We identified 43 structures in the commercial Vivinal GOS produced with  $\beta$ -galactosidase of *B. circulans* ATCC 31382 [56],[57]. Recently van Leeuwen *et al.* compared 6 commercial GOS products with Vivinal GOS and found 13 new structures [58]. Taken together, a total of 60 structures have been characterized in the GOS produced by various  $\beta$ -galactosidase enzymes (Figure 3). GOS are generally used in the dairy industry, especially in infant formula, as alternatives for hMOS to give beneficial effects to babies [59]. Till now, more than 200 hMOS structures have been revealed [60], [61], [62], and the structural complexity of hMOS is considered as important for their multiple biological functions [63]. Although the GOS structures are less diverse than those of hMOS,

with the development of sensitive and accurate identification techniques and the discovery and engineering of new  $\beta$ -galactosidases, the potential of discovering new GOS structures is still growing.



**Figure 3.** The GOS structures synthesized from lactose in the transgalactosylation reaction of  $\beta$ -galactosidases. Structure 1 is galactose, structure 2 is glucose, structure 5 is lactose, all others are GOS structures. The numbers refer to GOS structures identified in previous studies by van Leeuwen *et al* [56], [57], [58].



## Efforts to improve the GOS yield and change the linkage specificity

$\beta$ -Galactosidase enzymes incubated with lactose catalyze two types of reactions: hydrolysis and transgalactosylation. Hydrolysis results in the production of galactose, and transgalactosylation produces GOS mixtures. Due to the hydrolysis reaction, the GOS yield cannot reach 100% in industrial processes (e.g. *B. circulans*  $\beta$ -galactosidase has a GOS yield of ~63.5% with 50% (w/w) lactose, incubated at 60 °C for 20 h [64]). Relatively high lactose concentrations are generally needed to improve the transgalactosylation/hydrolysis ratio, but also results in remaining lactose, in addition to glucose and galactose [65]. There is a clear wish to improve the GOS yield and lower the hydrolysis products and lactose content in the final product mixture. Although GOS are generally used as alternatives for hMOS [66], their structural diversity is much less than that of hMOS. It may be of interest to try and change the linkage specificity of the  $\beta$ -galactosidase enzymes to produce different type of GOS mixtures or to enrich the GOS structure complexity or avoid GOS components implicated in allergenic effects [67]. There are two approaches to improve the GOS yield and/or change the linkage specificity of the products. The first approach is to change the reaction conditions, such as the reaction solvents, temperatures, substrate concentrations and so on; i.e. process engineering. The second approach is to modify the enzymes themselves; i.e. enzyme engineering. It has been shown that bio-solvents derived from dimethylamide and glycerol changed the regioselectivity of Biolacta N5 (commercial available  $\beta$ -galactosidase from *B. circulans*, Daiwa Kasei), and improved the yield of GOS products with ( $\beta$ 1 $\rightarrow$ 6) linkages [68], [69]. A study of the  $\beta$ -galactosidases from *B. circulans*, *A. oryzae*, *K. lactis*, and *Kluyveromyces fragilis* found that higher temperatures resulted in higher GOS yields, and that high lactose substrate concentrations resulted in even higher GOS yields [70]. Other studies also found that the temperatures, pH values, lactose concentrations and enzyme origins not only contributed to the GOS yield,

but also influenced the composition of the GOS mixtures [71], [72], [73], [74], [75]. Besides, different enzyme immobilization techniques such as adsorption on celite, covalent coupling to chitosan, aggregation by cross-linking, and immobilized on magnetic polysiloxane-polyvinyl also contributed to the improvement of the GOS yield [75], [76], [77],[78], [79]. On the other hand, protein engineering is a powerful tool to optimize the enzyme catalytic efficiency and change the product specificity [46], [80], [81]. For example, deletion mutagenesis showed that removal of 580 amino acids from the C-terminus of the  $\beta$ -galactosidase from *Bifidobacterium bifidum* greatly improved its transgalactosylation ability [82]. The native enzyme only has transgalactosylation activity at 13.7% lactose concentration while the truncated enzyme has a relatively high yield of GOS (39%) at 10% initial lactose [82]. A single mutation (F426Y) in  $\beta$ -glucosidase from *Pyrococcus furiosus* increased the transglycosylation/hydrolysis ratio, increasing the GOS yield from 40% to 45%. A double mutant (F426Y/M424K) improved GOS synthesis at 10% lactose from 18% to 40% due to the increase in the ratio of transglycosylation to hydrolysis [83]. A mutagenesis approach was also applied to the  $\beta$ -galactosidase from *Geobacillus stearothermophilus*; mutation R109W increased the transglycosylation/hydrolysis ratio, and the yield of trisaccharide  $\beta$ -D-Galp-(1 $\rightarrow$ 3)- $\beta$ -D-Galp-(1 $\rightarrow$ 4)-D-Glcp enhanced from 2% to 23% at a lactose concentration of 18% [84]. Double mutants F571L/N574S and F571L/N574A of *Thermotoga maritima*  $\beta$ -galactosidase increased the transgalactosylating efficiency of the wild-type enzyme up to 2-fold [85]. In another mutagenesis study of  $\beta$ -galactosidase from *Sulfolobus solfataricus*, the GOS yield was enhanced by 11% by mutating phenylalanine to tyrosine (F441Y), which may be caused by the introduction of new H-bonds [86]. A recent study showed that incubation of the C-terminally truncated  $\beta$ -galactosidase from *B. circulans* with monobodies, synthetic binding peptides which can modulate the catalytic properties of enzymes, altered the enzyme specificity in such a way that it barely

produced any GOS higher than DP5 [87], whereas in the absence of monobodies this enzyme is capable of producing GOS up to at least DP8 [50].

## **The beneficial impact of GOS**

GOS are well-known prebiotics and have been used in infant formula for decades [12], [66], [88]. A study has shown that the administration of highly purified (>95%) short-chain GOS in human subjects improved the lactose digestion and tolerance [89]. Another study showed that the use of GOS in the first 6 months of infant nutrition reduced the total number of infections, and cumulative incidence of infections [90]. The dietary intervention with GOS in the first two years of babies feeding reduced both the manifestation of allergies, as well as infections [91]. GOS produced by whole cells of *Bifidobacterium bifidum* NCIMB 41171 significantly increased the bifidobacterial population in the stool of human adults [92]. Purified GOS greatly inhibited the adhesion of pathogens to the epithelial cell surface [93]. A study found that the consumption of short-chain GOS and long-chain fructooligosaccharides (FOS) in the early life of newborns modulated the microbiota in a similar way as that of hMOS, namely by increasing the number of *Bifidobacterium* [94].

The main end products of microbial fermentation of GOS are short-chain fatty acids, which can prevent host colon cancer and other intestinal disorders [95]. It is generally considered that GOS benefit the host in the following ways. Firstly, the structures of GOS are similar to some pathogen receptors, thus they act as decoys resulting in pathogen binding and excretion instead of adhesion in the gut [96], [97]. Secondly, GOS inhibit the growth of toxic bacteria such as *Clostridium difficile*, and stimulate the growth of beneficial bacteria such as *Bifidobacterium adolescentis* and *B. bifidum* [98]. Thirdly, GOS modulate the human gut microbiota and increase the percentage of probiotic bacteria in the gut [99], [98]. These probiotic bacteria improve human health in various ways:

inhibitory compounds such as bacteriocins produced by probiotic bacteria inhibit the growth of certain pathogens [100]; probiotic bacteria compete for the limiting nutrients with pathogens, and also produce short-chain fatty acids that result in a lowering of the pH and inhibition of the growth of certain pathogens [100], [96]; probiotic bacteria adhere to the intestinal mucosa and block the adherence of enteropathogens [100], [101]; probiotic bacteria modulate the development of the immune system [102].

### **Outline of the thesis**

GOS are generally considered as prebiotics and are widely used in the food industry and pharmacy. They are composed of galactose molecules and one glucose molecule through different glycosidic linkages. The linkage types and degree of polymerization (DP) of GOS structures strongly depend on the enzyme origin. Although there have been many studies of  $\beta$ -galactosidase enzymes and the produced GOS mixtures, there are still many questions that remain unanswered. It is already known that  $\beta$ -galactosidases produce different GOS mixtures, however, the enzyme active site structural details that determine the composition of GOS mixtures has remained unknown. What are the roles of specific amino acid residues in the  $\beta$ -galactosidase active site? What are the structural determinants for the GOS yield and linkage specificity? Can we engineer  $\beta$ -galactosidases to change their GOS linkage specificity and thus change or enrich the GOS structural complexity? Can we use  $\beta$ -galactosidases to synthesize products other than GOS?

In this thesis, we tried to answer these questions. We investigated the active site structural basis for the linkage specificity, to provide insights into the structure-function relationship and provide guidance for the engineering of  $\beta$ -galactosidase enzymes. In chapter 2, a more detailed analysis of GOS profiles of three commercial  $\beta$ -galactosidases from *B. circulans*, *K. lactis* and *A. oryzae* is

reported. The GOS yields, the linkage specificity, and the diversity of GOS produced by these enzymes are clearly different from each other. The presence of the monosaccharides glucose and galactose in the reaction mixture changed the GOS profile and yield, however, the influence of monosaccharides also depended on the enzyme origin.

In chapter 3, residue R484 near the +1 subsite of the C-terminally truncated  $\beta$ -galactosidase from *B. circulans* (BgaD-D) was subjected to site saturation mutagenesis. The mutant enzymes displayed significantly altered enzyme specificity, leading to a GOS mixture with mainly ( $\beta 1 \rightarrow 4$ ) and ( $\beta 1 \rightarrow 3$ ) linkages, while the wild-type enzyme gave a GOS mainly composed of ( $\beta 1 \rightarrow 4$ ) linkages. Besides, the mutant GOS mixtures also contained 14 structures that are not found in the GOS produced by the wild-type enzyme. Chapter 4 investigated the functional roles of selected amino acid residues in the BgaD-D active site using site-directed mutagenesis. A detailed biochemical characterization and product profile analysis was presented, showing that these amino acid residues in the active site were crucial to the enzyme activity, linkage specificity, transgalactosylation versus hydrolysis, acceptor substrate selection.

In chapter 5, lactulose was used as substrate for the BgaD-D wild-type and R484H mutant enzymes to expand the application of  $\beta$ -galactosidase enzymes. The oligosaccharides derived from lactulose were identified and several new-to-nature compounds characterized. They were tested as sole carbon source for growth by Bifidobacteria, reflecting their potential prebiotic properties. Chapter 6 summarizes the results reported in this thesis and gives perspectives for future research.

## References

- 1 Gänzle, M. G., Haase, G. and Jelen, P. (2008) Lactose: Crystallization, hydrolysis and value-added derivatives. *Int. Dairy J.* **18**, 685–694.

- 2 Schaafsma, G. (2008) Lactose and lactose derivatives as bioactive ingredients in human nutrition. *Int. Dairy J.* **18**, 458–465.
- 3 Jelen, P. and Tossavainen, O. (2003) Low lactose and lactose-free milk and dairy products - Prospects, technologies and applications. *Aust. J. Dairy Technol.* **58**, 161-165.
- 4 Swagerty, D. L., Walling, A. D. and Klein, R. M. (2002) Lactose intolerance. *Am. Fam. Physician* **65**, 1845–1850.
- 5 Heyman, M. B. and others. (2006) Lactose intolerance in infants, children, and adolescents. *Pediatrics* **118**, 1279–1286.
- 6 Lifran, E. V, Hourigan, J. a and Sleight, R. W. (2009) Lactose derivatives: turning waste into functional foods. *Aust. J. Dairy Technol.* **64**, 89–93.
- 7 Rycroft, C. E., Jones, M. R., Gibson, G. R. and Rastall, R. A. (2001) A comparative in vitro evaluation of the fermentation properties of prebiotic oligosaccharides. *J. Appl. Microbiol.* **91**, 878–887.
- 8 Rastall, R. A. (2010) Functional Oligosaccharides: Application and Manufacture. *Annu. Rev. Food Sci. Technol.* **1**, 305–339.
- 9 Gibson, G. R. and Roberfroid, M. B. (1995) Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *J. Nutr.* **125**, 1401-1412.
- 10 Valcheva, R. and Dieleman, L. A. (2016) Prebiotics: Definition and protective mechanisms. *Best Pract. Res. Clin. Gastroenterol.* **30**, 27–37.
- 11 Fanaro, S., Boehm, G., Garssen, J., Knol, J., Mosca, F., Stahl, B. and Vigi, V. (2005) Galacto-oligosaccharides and long-chain fructo-oligosaccharides as prebiotics in infant formulas: a review. *Acta Paediatr. Suppl.* **94**, 22–26.
- 12 Boehm, G., Stahl, B., Jelinek, J., Knol, J., Miniello, V. and Moro, G. E. (2005) Prebiotic carbohydrates in human milk and formulas. *Acta Paediatr. Suppl.* **94**, 18–21.
- 13 Sanz, M. L., Sanz, J. and Martinez-Castro, I. (2002) Characterization of O-trimethylsilyl oximes of disaccharides by gas chromatography-mass spectrometry. *Chromatographia* **56**, 617–622.
- 14 Gosling, A., Stevens, G. W., Barber, A. R., Kentish, S. E. and Gras, S. L. (2010) Recent advances refining galactooligosaccharide production from lactose. *Food Chem.* **121**, 307–318.
- 15 Husain, Q. (2010)  $\beta$  Galactosidases and their potential applications: a review. *Crit. Rev. Biotechnol.* **30**, 41–62.
- 16 Huerta, L. M., Vera, C., Guerrero, C., Wilson, L. and Illanes, A. (2011) Synthesis

- of galacto-oligosaccharides at very high lactose concentrations with immobilized  $\beta$ -galactosidases from *Aspergillus oryzae*. *Process Biochem.* **46**, 245–252.
- 17 Lombard, V., Golaconda Ramulu, H., Drula, E., Coutinho, P. M. and Henrissat, B. (2014) The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res.* **42**, D490-495.
- 18 Eda, M., Matsumoto, T., Ishimaru, M. and Tada, T. (2016) Structural and functional analysis of tomato  $\beta$ -galactosidase 4: insight into the substrate specificity of the fruit softening-related enzyme. *Plant J.* **86**, 300–307.
- 19 Cheng, W., Wang, L., Jiang, Y. L., Bai, X. H., Chu, J., Li, Q., Yu, G., Liang, Q. L., Zhou, C. Z. and Chen, Y. (2012) Structural insights into the substrate specificity of *Streptococcus pneumoniae*  $\beta(1,3)$ -galactosidase BgaC. *J. Biol. Chem.* **287**, 22910–22918.
- 20 Song, J., Abe, K., Imanaka, H., Imamura, K., Minoda, M., Yamaguchi, S. and Nakanishi, K. (2011) Causes of the production of multiple forms of  $\beta$ -galactosidase by *Bacillus circulans*. *Biosci. Biotechnol. Biochem.* **75**, 268–278.
- 21 Song, J., Imanaka, H., Imamura, K., Minoda, M., Yamaguchi, S. and Nakanishi, K. (2013) The discoidin domain of *Bacillus circulans*  $\beta$ -galactosidase plays an essential role in repressing galactooligosaccharide production. *Biosci. Biotechnol. Biochem.* **77**, 73–79.
- 22 Song, J., Imanaka, H., Imamura, K., Minoda, M., Katase, T., Hoshi, Y., Yamaguchi, S. and Nakanishi, K. (2011) Cloning and expression of a  $\beta$ -galactosidase gene of *Bacillus circulans*. *Biosci. Biotechnol. Biochem.* **75**, 1194–1197.
- 23 Viborg, A. H., Fredslund, F., Katayama, T., Nielsen, S. K., Svensson, B., Kitaoka, M., Lo Leggio, L. and Abou Hachem, M. (2014) A  $\beta 1$ -6/ $\beta 1$ -3 galactosidase from *Bifidobacterium animalis* subsp. *lactis* B1-04 gives insight into sub-specificities of  $\beta$ -galactoside catabolism within *Bifidobacterium*. *Mol. Microbiol.* **94**, 1024–1040.
- 24 Davies, G. and Henrissat, B. (1995) Structures and mechanisms of glycosyl hydrolases. *Structure* **3**, 853–859.
- 25 Juers, D. H., Matthews, B. W. and Huber, R. E. (2012) LacZ  $\beta$ -galactosidase: structure and function of an enzyme of historical and molecular biological importance. *Protein Sci.* **21**, 1792–1807.
- 26 Torres, D. P. M., Gonçalves, M. P. F., Teixeira, J. A. and Rodrigues, L. R. (2010) Galacto-Oligosaccharides: Production, properties, applications, and significance as prebiotics. *Compr. Rev. Food Sci. Food Saf.* **9**, 438–454.
- 27 Bultema, J. B., Kuipers, B. J. H. and Dijkhuizen, L. (2014) Biochemical characterization of mutants in the active site residues of the  $\beta$ -galactosidase

- enzyme of *Bacillus circulans* ATCC 31382. FEBS Open Bio **4**, 1015–1020.
- 28 Zechel, D. L., Boraston, A. B., Gloster, T., Boraston, C. M., Macdonald, J. M., Tilbrook, D. M. G., Stick, R. V. and Davies, G. J. (2003) Iminosugar glycosidase inhibitors: Structural and thermodynamic dissection of the binding of isofagomine and 1-deoxynojirimycin to  $\beta$ -glucosidases. *J. Am. Chem. Soc.* **125**, 14313–14323.
  - 29 Skálová, T., Dohnálek, J., Spiwok, V., Lipovová, P., Vondrácková, E., Petroková, H., Dusková, J., Strnad, H., Králová, B., and Hasek, J. (2005) Cold-active  $\beta$ -galactosidase from *Arthrobacter* sp. C2-2 forms compact 660 kDa hexamers crystal structure at 1.9 Å resolution. *J. Mol. Biol.* **353**, 282–294.
  - 30 Ishikawa, K., Kataoka, M., Yanamoto, T., Nakabayashi, M., Watanabe, M., Ishihara, S. and Yamaguchi, S. (2015) Crystal structure of  $\beta$ -galactosidase from *Bacillus circulans* ATCC 31382 (BgaD) and the construction of the thermophilic mutants. *FEBS J.* **282**, 2540–2552.
  - 31 Juers, D. H., Jacobson, R. H., Wigley, D., Zhang, X. J., Huber, R. E., Tronrud, D. E. and Matthews, B. W. (2000) High resolution refinement of  $\beta$ -galactosidase in a new crystal form reveals multiple metal-binding sites and provides a structural basis for alpha-complementation. *Protein Sci.* **9**, 1685–1699.
  - 32 Rutkiewicz-Krotewicz, M., Pietrzyk-Brzezinska, A. J., Sekula, B., Cieśliński, H., Wierzbicka-Woś, A., Kur, J. and Bujacz, A. (2016) Structural studies of a cold-adapted dimeric  $\beta$ -D-galactosidase from *Paracoccus* sp. 32d. *Acta Crystallogr. Sect. D, Struct. Biol.* **72**, 1049–61.
  - 33 Singh, A. K., Pluvinage, B., Higgins, M. A., Dalia, A. B., Woodiga, S. A., Flynn, M., Lloyd, A. R., Weiser, J. N., Stubbs, K. A., Boraston, A. B., et al. (2014) Unravelling the multiple functions of the architecturally intricate *Streptococcus pneumoniae*  $\beta$ -galactosidase, BgaA. *PLoS Pathog.* **10**, e1004364.
  - 34 Pereira-Rodríguez, A., Fernández-Leiro, R., González-Siso, M. I., Cerdán, M. E., Becerra, M. and Sanz-Aparicio, J. (2012) Structural basis of specificity in tetrameric *Kluyveromyces lactis*  $\beta$ -galactosidase. *J. Struct. Biol.* **177**, 392–401.
  - 35 Henze, M., You, D. J., Kamerke, C., Hoffmann, N., Angkawidjaja, C., Ernst, S., Pietruszka, J., Kanaya, S. and Elling, L. (2014) Rational design of a glycosynthase by the crystal structure of  $\beta$ -galactosidase from *Bacillus circulans* (BgaC) and its use for the synthesis of N-acetyllactosamine type 1 glycan structures. *J. Biotechnol.* **191**, 78–85.
  - 36 Larsbrink, J., Thompson, A. J., Lundqvist, M., Gardner, J. G., Davies, G. J. and Brumer, H. (2014) A complex gene locus enables xyloglucan utilization in the model saprophyte *Cellobivrio japonicus*. *Mol. Microbiol.* **94**, 418–433.
  - 37 Maksimainen, M. M., Lampio, A., Mertanen, M., Turunen, O. and Rouvinen, J. (2013) The crystal structure of acidic  $\beta$ -galactosidase from *Aspergillus oryzae*. *Int. J. Biol. Macromol.* **60**, 109–115.



- 38 Ohto, U., Usui, K., Ochi, T., Yuki, K., Satow, Y. and Shimizu, T. (2012) Crystal structure of human  $\beta$ -galactosidase: Structural basis of G<sub>M1</sub> gangliosidosis and morquio B diseases. *J. Biol. Chem.* **287**, 1801–1812.
- 39 Rojas, a L., Nagem, R. a P., Neustroev, K. N., Arand, M., Adamska, M., Eneyskaya, E. V, Kulminskaya, a a, Garratt, R. C., Golubev, a M. and Polikarpov, I. (2004) Crystal structures of beta-galactosidase from *Penicillium* sp. and its complex with galactose. *J. Mol. Biol.* **343**, 1281–1292.
- 40 Maksimainen, M., Hakulinen, N., Kallio, J. M., Timoharju, T., Turunen, O. and Rouvinen, J. (2011) Crystal structures of *Trichoderma reesei*  $\beta$ -galactosidase reveal conformational changes in the active site. *J. Struct. Biol.* **174**, 156–163.
- 41 Maksimainen, M., Paavilainen, S., Hakulinen, N. and Rouvinen, J. (2012) Structural analysis, enzymatic characterization, and catalytic mechanisms of  $\beta$ -galactosidase from *Bacillus circulans* sp. *alkalophilus*. *FEBS J.* **279**, 1788–1798.
- 42 Godoy, A. S., Camilo, C. M., Kadowaki, M. A., Muniz, H. dos S., Espirito Santo, M., Murakami, M. T., Nascimento, A. S. and Polikarpov, I. (2016) Crystal structure of  $\beta$ 1→6-galactosidase from *Bifidobacterium bifidum* S17: trimeric architecture, molecular determinants of the enzymatic activity and its inhibition by  $\alpha$ -galactose. *FEBS J.* **283**, 4097–4112.
- 43 Solomon, H. V., Tabachnikov, O., Feinberg, H., Govada, L., Chayen, N. E., Shoham, Y. and Shoham, G. (2013) Crystallization and preliminary crystallographic analysis of GanB, a GH42 intracellular  $\beta$ -galactosidase from *Geobacillus stearothermophilus*. *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.* **69**, 1114–1119.
- 44 Fan, Y., Hua, X., Zhang, Y., Feng, Y., Shen, Q., Dong, J., Zhao, W., Zhang, W., Jin, Z. and Yang, R. (2015) Cloning, expression and structural stability of a cold-adapted  $\beta$ -galactosidase from *Rahnella* sp. R3. *Protein Expr. Purif.* **115**, 158–164.
- 45 Hidaka, M., Fushinobu, S., Ohtsu, N., Motoshima, H., Matsuzawa, H., Shoun, H. and Wakagi, T. (2002) Trimeric crystal structure of the glycoside hydrolase family 42  $\beta$ -galactosidase from *Thermus thermophilus* A4 and the structure of its complex with galactose. *J. Mol. Biol.* **322**, 79–91.
- 46 Böttcher, D., and Bornscheuer, U. T. (2010) Protein engineering of microbial enzymes. *Curr. Opin. Microbiol.* **13**, 274–282.
- 47 Vera, C., Córdova, A., Aburto, C., Guerrero, C., Suárez, S. and Illanes, A. (2016) Synthesis and purification of galacto-oligosaccharides: state of the art. *World J. Microbiol. Biotechnol.* **32**, 197.
- 48 Cataldi, T. R., Campa, C. and De Benedetto, G. E. (2000) Carbohydrate analysis by high-performance anion-exchange chromatography with pulsed amperometric detection: the potential is still growing. *Fresenius. J. Anal. Chem.* **368**, 739–758.

- 49 Corradini, C., Cavazza, A. and Bignardi, C. (2012) High-performance anion-exchange chromatography coupled with pulsed electrochemical detection as a powerful tool to evaluate carbohydrates of food interest: principles and applications. *Int. J. Carbohydr. Chem.* **2012**, 1–13.
- 50 Coulier, L., Timmermans, J., Bas, R., Van Den Dool, R., Haaksman, I., Klarenbeek, B., Slaghek, T. and Van Dongen, W. (2009) In-depth characterization of prebiotic galacto-oligosaccharides by a combination of analytical techniques. *J. Agric. Food Chem.* **57**, 8488–8495.
- 51 Moreno, F. J., Montilla, A., Villamiel, M., Corzo, N. and Olano, A. (2014) Analysis, structural characterization, and bioactivity of oligosaccharides derived from lactose. *Electrophoresis* **35**, 1519–1534.
- 52 Van Leeuwen, S. S., Kuipers, B. J. H., Dijkhuizen, L. and Kamerling, J. P. (2014) Development of a  $^1\text{H}$  NMR structural-reporter-group concept for the analysis of prebiotic galacto-oligosaccharides of the  $[\beta\text{-D-Galp-(1}\rightarrow\text{x)}]_n\text{-D-Glcp}$  type. *Carbohydr. Res.* **400**, 54–58.
- 53 Rodriguez-Colinas, B., De Abreu, M. A., Fernandez-Arrojo, L., De Beer, R., Poveda, A., Jimenez-Barbero, J., Haltrich, D., Ballesteros Olmo, A. O., Fernandez-Lobato, M. and Plou, F. J. (2011) Production of galacto-oligosaccharides by the  $\beta$ -galactosidase from *Kluyveromyces lactis*: Comparative analysis of permeabilized cells versus soluble enzyme. *J. Agric. Food Chem.* **59**, 10477–10484.
- 54 Urrutia, P., Rodriguez-Colinas, B., Fernandez-Arrojo, L., Ballesteros, A. O., Wilson, L., Illanes, A. and Plou, F. J. (2013) Detailed analysis of galactooligosaccharides synthesis with  $\beta$ -galactosidase from *Aspergillus oryzae*. *J. Agric. Food Chem.* **61**, 1081–1087.
- 55 Yanahira, S., Kobayashi, T., Suguri, T., Nakakoshi, M., Miura, S., Ishikawa, H. and Nakajima, I. (1995) Formation of oligosaccharides from lactose by *Bacillus circulans*  $\beta$ -galactosidase. *Biosci. Biotechnol. Biochem.* **59**, 1021–1026.
- 56 van Leeuwen, S. S., Kuipers, B. J. H., Dijkhuizen, L. and Kamerling, J. P. (2014)  $^1\text{H}$  NMR analysis of the lactose/ $\beta$ -galactosidase-derived galacto-oligosaccharide components of Vivinal® GOS up to DP5. *Carbohydr. Res.* **400**, 59–73.
- 57 van Leeuwen, S. S., Kuipers, B. J. H., Dijkhuizen, L. and Kamerling, J. P. (2016) Corrigendum to “ $^1\text{H}$  NMR analysis of the lactose/ $\beta$ -galactosidase-derived galacto-oligosaccharide components of Vivinal® GOS up to DP5” [*Carbohydr. Res.* 400 (2014) 59–73]. *Carbohydr. Res.* **419**, 69–70.
- 58 van Leeuwen, S. S., Kuipers, B. J. H., Dijkhuizen, L. and Kamerling, J. P. (2016) Comparative structural characterization of 7 commercial galacto-oligosaccharide (GOS) products. *Carbohydr. Res.* **425**, 48–58.
- 59 Sabater, C., Prodanov, M., Olano, A., Corzo, N. and Montilla, A. (2016)

- Quantification of prebiotics in commercial infant formulas. *Food Chem.* **194**, 6–11.
- 60 Pettherick, A. (2010) Mother's milk : A rich opportunity. *Nature* **468**, 8–10.
- 61 Kunz, C., Rudloff, S., Baie, W., Klein, N. and St, S. (2000) Oligosaccharides in human milk: structural, functional, and metabolic aspects. *Annu. Rev. Nutr.* **20**, 699–722.
- 62 Boehm, G. and Stahl, B. (2007) Oligosaccharides from milk. *J. Nutr.* **137**, 847S–849S.
- 63 Bode, L. (2012) Human milk oligosaccharides: Every baby needs a sugar mama. *Glycobiology* **22**, 1147–1162.
- 64 Yin, H., Pijning, T., Meng, X., Dijkhuizen, L. and van Leeuwen, S. S. (2017) Engineering of the *Bacillus circulans*  $\beta$ -galactosidase product specificity. *Biochemistry* **56**, 704–711.
- 65 Kovács, Z., Benjamins, E., Grau, K., Rehman, A. U., Ebrahimi, M., and Zermak, P. (2014) Recent developments in manufacturing oligosaccharides with prebiotic functions. *Adv. Biochem. Eng. Biotechnol.* **143**, 257–295
- 66 Boehm, G., Fanaro, S., Jelinek, J., Stahl, B. and Marini, A. (2003) Prebiotic concept for infant nutrition. *Acta Paediatr. Suppl.* **91**, 64–67.
- 67 Moriya, T., Nagahata, N., Odaka, R., Nakamura, H., Yoshikawa, J., Kurashima, K. and Saito, T. (2017) Synthesis of an allergy inducing tetrasaccharide "4P-X". *Carbohydr. Res.* **439**, 44–49.
- 68 Pérez-Sánchez, M., Sandoval, M., Cortés-Cabrera, A., García-Marín, H., Sinisterra, J. V., García, J. I., and J. Hernaiz, M. J. (2011) Solvents derived from glycerol modify classical regioselectivity in the enzymatic synthesis of disaccharides with Biolacta  $\beta$ -galactosidase. *Green Chem.* **13**, 2810–2817.
- 69 Pérez-Sánchez, M., Sandoval, M. and Hernáiz, M. J. (2012) Bio-solvents change regioselectivity in the synthesis of disaccharides using Biolacta  $\beta$ -galactosidase. *Tetrahedron* **68**, 2141–2145.
- 70 Boon, M. A., Janssen, A. E. M. and Riet, K. Van. (2000) Effect of temperature and enzyme origin on the enzymatic synthesis of oligosaccharides. *Enzyme Microb. Technol.* **26**, 271–281.
- 71 Warmerdam, A., Boom, R. M. and Janssen, A. E. (2013)  $\beta$ -Galactosidase stability at high substrate concentrations. *Springerplus* **2**, 402.
- 72 Martínez-Villaluenga, C., Cardelle-Cobas, A., Corzo, N., Olano, A. and Villamiel, M. (2008) Optimization of conditions for galactooligosaccharide synthesis during lactose hydrolysis by  $\beta$ -galactosidase from *Kluyveromyces lactis* (Lactozym 3000

- L HP G). Food Chem. **107**, 258–264.
- 73 Reuter, S., Rusborg Nygaard, A. and Zimmermann, W. (1999)  $\beta$ -Galactooligosaccharide synthesis with  $\beta$ -galactosidases from *Sulfolobus solfataricus*, *Aspergillus oryzae*, and *Escherichia coli*. Enzyme Microb. Technol. **25**, 509–516.
- 74 Vera, C., Guerrero, C., Conejeros, R. and Illanes, A. (2012) Synthesis of galacto-oligosaccharides by  $\beta$ -galactosidase from *Aspergillus oryzae* using partially dissolved and supersaturated solution of lactose. Enzyme Microb. Technol. **50**, 188–194.
- 75 Zhou, Q. Z. K. and Chen, X. D. (2001) Effects of temperature and pH on the catalytic activity of the immobilized  $\beta$ -galactosidase from *Kluyveromyces lactis*. Biochem. Eng. J. **9**, 33–40.
- 76 Gaur, R., Pant, H., Jain, R. and Khare, S. K. (2006) Galacto-oligosaccharide synthesis by immobilized *Aspergillus oryzae*  $\beta$ -galactosidase. Food Chem. **97**, 426–430.
- 77 Neri, D. F. M., Balcão, V. M., Costa, R. S., Rocha, I. C. a P., Ferreira, E. M. F. C., Torres, D. P. M., Rodrigues, L. R. M., Carvalho, L. B. and Teixeira, J. a. (2009) Galacto-oligosaccharides production during lactose hydrolysis by free *Aspergillus oryzae*  $\beta$ -galactosidase and immobilized on magnetic polysiloxane-polyvinyl alcohol. Food Chem. **115**, 92–99.
- 78 Benjamins, E., Boxem, L., KleinJan-Noeverman, J. and Broekhuis, T. A. (2014) Assessment of repetitive batch-wise synthesis of galacto-oligosaccharides from lactose slurry using immobilised  $\beta$ -galactosidase from *Bacillus circulans*. Int. Dairy J. **38**, 160–168.
- 79 Warmerdam, A., Benjamins, E., de Leeuw, T. F., Broekhuis, T. A., Boom, R. M. and Janssen, A. E. M. (2013) Galacto-oligosaccharide production with immobilized  $\beta$ -galactosidase in a packed-bed reactor vs. free  $\beta$ -galactosidase in a batch reactor. Food Bioprod. Process. **92**, 383–392.
- 80 Bommarius, A. S., Blum, J. K., and Abrahamson, M. J. (2011) Status of protein engineering for biocatalysts: How to design an industrially useful biocatalyst. Curr. Opin. Chem. Biol. **15**, 194-200.
- 81 Quin, M. B. and Schmidt-Dannert, C. (2011) Engineering of biocatalyst from evolution to creation. ACS Catal. **1**, 1017–1021.
- 82 Jørgensen, F., Hansen, O. C. and Stougaard, P. (2001) High-efficiency synthesis of oligosaccharides with a truncated  $\beta$ -galactosidase from *Bifidobacterium bifidum*. Appl. Microbiol. Biotechnol. **57**, 647–652.
- 83 Hansson, T., Kaper, T., van Der Oost, J., de Vos, W. M. and Adlercreutz, P. (2001) Improved oligosaccharide synthesis by protein engineering of beta-

- glucosidase CelB from hyperthermophilic *Pyrococcus furiosus*. Biotechnol. Bioeng. **73**, 203–210.
- 84 Placier, G., Watzlawick, H., Rabiller, C. and Mattes, R. (2009) Evolved  $\beta$ -galactosidases from *Geobacillus stearothermophilus* with improved transgalactosylation yield for galacto-oligosaccharide production. Appl. Environ. Microbiol. **75**, 6312–6321.
- 85 Talens-Perales, D., Polaina, J. and Marín-Navarro, J. (2016) Structural dissection of the active site of *Thermotoga maritima*  $\beta$ -galactosidase identifies key residues for transglycosylating activity. J. Agric. Food Chem. **64**, 2917–2924.
- 86 Wu, Y., Yuan, S., Chen, S., Wu, D., Chen, J. and Wu, J. (2013) Enhancing the production of galacto-oligosaccharides by mutagenesis of *Sulfolobus solfataricus*  $\beta$ -galactosidase. Food Chem. **138**, 1588–1595.
- 87 Tanaka, S., Takahashi, T., Koide, A., Ishihara, S., Koikeda, S. and Koide, S. (2015) Monobody-mediated alteration of enzyme specificity. Nat. Chem. Biol. **11**, 762–764.
- 88 Rivero-Urgell, M. and Santamaria-Orleans, A. (2001) Oligosaccharides: Application in infant food. Early Hum. Dev. **65**, 43–52.
- 89 Azcarate-Peril, M. A., Ritter, A. J., Savaiano, D., Monteagudo-Mera, A., Anderson, C., Magness, S. T. and Klaenhammer, T. R. (2017) Impact of short-chain galactooligosaccharides on the gut microbiome of lactose-intolerant individuals. Proc. Natl. Acad. Sci. U.S.A. **114**, E367–E375.
- 90 Arslanoglu, S., Moro, G. E. and Boehm, G. (2007) Early supplementation of prebiotic oligosaccharides protects formula-fed infants against infections during the first 6 months of life J. Nutr. **137**, 2420–2424.
- 91 Arslanoglu, S., Moro, G. E., Schmitt, J., Tandoi, L., Rizzardi, S. and Boehm, G. (2008) Early dietary intervention with a mixture of prebiotic oligosaccharides reduces the incidence of allergic manifestations and infections during the first two years of life. J. Nutr. **138**, 1091–1095.
- 92 Depeint, F., Tzortzis, G., Vulevic, J., I'Anson, K., and Gibson, G. R. (2008) Prebiotic evaluation of a novel galactooligosaccharide mixture produced by the enzymatic activity of *Bifidobacterium bifidum* NCIMB 41171, in healthy humans: a randomized, double-blind, crossover, placebo-controlled intervention study. Am. J. Clin. Nutr. **87**, 785-791.
- 93 Shoaf, K., Mulvey, G. L., Armstrong, G. D. and Hutkins, R. W. (2006) Prebiotic galactooligosaccharides reduce adherence of enteropathogenic *Escherichia coli* to tissue culture cells. Infect. Immun. **74**, 6920–6928.
- 94 Oozeer, R., van Limpt, K., Ludwig, T., Ben Amor, K., Martin, R., Wind, R. D., Boehm, G. and Knol, J. (2013) Intestinal microbiology in early life: specific

- 
- prebiotics can have similar functionalities as human-milk oligosaccharides. *Am. J. Clin. Nutr.* **98**, 561S–571S.
- 95 Cardelle-Cobas, A., Fernández, M., Salazar, N., Martínez-Villaluenga, C., Villamiel, M., Ruas-Madiedo, P. and de los Reyes-Gavilán, C. G. (2009) Bifidogenic effect and stimulation of short chain fatty acid production in human faecal slurry cultures by oligosaccharides derived from lactose and lactulose. *J. Dairy Res.* **76**, 317–325.
- 96 Gibson, G. R., McCartney, A. L. and Rastall, R. A. (2005) Prebiotics and resistance to gastrointestinal infections, *Br. J. Nutr.* **93**, Suppl. 1, S31–S3
- 97 Gourbeyre, P., Denery, S. and Bodinier, M. (2011) Probiotics, prebiotics, and synbiotics: impact on the gut immune system and allergic reactions. *J. Leukoc. Biol.* **89**, 685–695.
- 98 Macfarlane, G. T., Steed, H. and Macfarlane, S. (2007) Bacterial metabolism and health-related effects of galacto-oligosaccharides and other prebiotics. *J. Appl. Microbiol.* **104**, 305–344.
- 99 Tuohy, K. M., Rouzaud, G. C. M., Brück, W. M. and Gibson, G. R. (2005) Modulation of the human gut microflora towards improved health using prebiotics – assessment of efficacy. *Curr. Pharm. Des.*, **11**, 75–90.
- 100 Fooks, L. J. and Gibson, G. R. (2002) Probiotics as modulators of the gut flora. *Br. J. Nutr.* **88**, S39–S49.
- 101 Macfarlane, S., Macfarlane, G. T. and Cummings, J. H. (2006) Review article: Prebiotics in the gastrointestinal tract. *Aliment. Pharmacol. Ther.* **24**, 701–714.
- 102 Tuohy, K. M., Probert, H. M., Smejkal, C. W. and Gibson, G. R. (2003) Using probiotics and prebiotics to improve gut health. *Drug Discov. Today* **8**, 692–700.

