

## University of Groningen

### In vitro characterization of representative clinical South African *Staphylococcus aureus* isolates from various clonal lineages

Oosthuysen, W F; Orth, H; Lombard, C. J.; Sinha, B; Wasserman, E

*Published in:*  
New Microbes and New Infections

*DOI:*  
[10.1002/nmi2.53](https://doi.org/10.1002/nmi2.53)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2014

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Oosthuysen, W. F., Orth, H., Lombard, C. J., Sinha, B., & Wasserman, E. (2014). In vitro characterization of representative clinical South African *Staphylococcus aureus* isolates from various clonal lineages. *New Microbes and New Infections*, 2(4), 115-122. <https://doi.org/10.1002/nmi2.53>

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# In vitro characterization of representative clinical South African *Staphylococcus aureus* isolates from various clonal lineages

W. F. Oosthuysen<sup>1</sup>, H. Orth<sup>1,2</sup>, C. Lombard<sup>3</sup>, B. Sinha<sup>4</sup> and E. Wasserman<sup>1,5</sup>

1) Division of Medical Microbiology, Stellenbosch University, Tygerberg Hospital, Cape Town, 2) National Health Laboratory Services, Microbiology Laboratory, Tygerberg Academic Complex, Cape Town, 3) Biostatistics Unit, Medical Research Council, Cape Town, South Africa, 4) University Medical Centre Groningen, Groningen, The Netherlands and 5) Pathcare Laboratories, Cape Town, South Africa

## Abstract

Data concerning the virulence and pathogenesis of South African strains of *Staphylococcus aureus* are limited. We investigated host–pathogen interactions of randomly selected clinical *S. aureus* isolates representing various clones. We characterized the ability of isolates to adhere to fibronectin, fibrinogen, collagens IV and VI, to invade host cells and to induce cell death *in vitro*. We analysed the possible association of these results with characteristics such as methicillin resistance, Pantone–Valentine leucocidin (PVL) positivity and clonality. The *S. aureus* isolates displayed diversity in their abilities to adhere to various human ligands. All isolates were highly invasive except for ST121. PVL-negative isolates were significantly more invasive than the PVL-positive isolates ( $p$  0.004). Isolates of CC5, CC30 and CC121 were non-cytotoxic, whereas isolates of CC22, CC8, CC15, CC45 and CC88 were very cytotoxic. No statistical association was identified between cell death and methicillin resistance, bacterial PVL status, clonality or patient HIV status. The vast majority of isolates were invasive and induced significant cell death. PVL-negative isolates were more invasive than PVL-positive isolates, while methicillin-resistant isolates were not found to be more invasive or cytotoxic than methicillin-susceptible isolates.

**Keywords:** Cellular invasiveness, cytotoxic, methicillin-resistance, Pantone–Valentine leucocidin, *Staphylococcus aureus*, South Africa

**Original Submission:** 22 March 2014; **Revised Submission:** 6 May 2014; **Accepted:** 14 May 2014

**Article published online:** 26 June 2014

*New Microbe and New Infect* 2014; **2**: 115–122

**Corresponding author:** W. F. Oosthuysen, University of Wuerzburg, Institute of Hygiene and Microbiology, Josef-Schnieder-Straße 2/E1, Wuerzburg 97080, Germany  
**E-mail:** wilhelm.oosthuysen@hygiene.uni-wuerzburg.de

## Introduction

*Staphylococcus aureus* is a facultative intracellular bacterium and a significant human pathogen. It possesses many surface factors that aid with host colonization and cellular invasion as well as secreted virulence factors involved in host cell death induction [1].

Fibronectin, fibrinogen and collagen are three of many extracellular matrix molecules found in macromolecular structures. Both fibronectin and fibrinogen play significant roles in the adherence of *S. aureus* during infections associated with skin diseases such as atopic dermatitis [2]. Fibronectin is also a component of human plasma and connective tissue [3]. Fibrin-

ogen-binding is commonly associated with infective endocarditis [4], whereas collagen binding is commonly required for the colonization of cartilage [5]. Numerous bacterial surface proteins can be used during the process of adherence to host ligands and are called ‘microbial surface components recognizing adhesive matrix molecules’ or MSCRAMM, such as fibronectin-binding proteins A and B [6], staphylococcal protein A and clumping factors. Another group of bacterial proteins which is involved in this process are the SERAM molecules, or the ‘secreted expanded repertoire adhesive molecules’ [7], such as the extracellular adherence protein (Eap). Adherence to fibronectin by *S. aureus* can be mediated by fibronectin-binding proteins A and B (FnB A/B), which also aid in the binding of the organism to plasma clots [8]. Both genes are fundamental for the invasion of eukaryotic cells [7]. *Staphylococcus aureus* possess two distinct fibrinogen-binding proteins, namely clumping-factor A and B, of which clumping factor A is mainly used to adhere to substances containing fibrinogen [8].

Many groups have clearly demonstrated the role of FnB proteins as the main invasin of *S. aureus* and identified a

fibronectin-dependent bridging mechanism to the host cellular integrin  $\alpha_5\beta_1$  [9]. Fibronectin-binding proteins do not require any other *S. aureus*-specific co-receptors to confer invasiveness and this function can be accomplished by either FnbA or FnbB [10], which must be anchored into the bacterial cell wall, as truncation of these proteins results in deficient adherence and cellular invasiveness [11]. It has been shown that the extracellular adherence protein, with its broad binding capacity, can play a role in the cellular invasion of host cells [12].

In any given *S. aureus* strain, host cell death induction is difficult to predict and depends on many factors [13]. Various bacterial virulence factors are involved, of which  $\alpha$ -toxin [14] is described as the most prominent. Intracellular *S. aureus*, if viable, can exist free in the cytoplasm and kill endothelial cells, partly by apoptosis [15]. Metabolically active intracellular staphylococci are required for the induction of apoptosis in endothelial cells, which is dependent on *agr* and *sigB* [16]. Strains with invasive and haemolytic phenotypes are normally associated with caspase-dependent induction of apoptosis, while non-invasive haemolytic or non-haemolytic invasive isolates are not [16].

Another well-characterized virulence factor of *S. aureus* that can be responsible for host cell death induction, especially of human neutrophils [17], is the two-component leucotoxin, Pantón–Valentine leucocidin (PVL). This toxin has been associated with necrotizing pneumonia [18], skin-and-soft tissue infections (SSTI) [19] and necrotizing lesions of the skin and subcutaneous tissues [20] and is very common among diverse genetic backgrounds associated with community-acquired methicillin-resistant *S. aureus* (MRSA), especially the USA300 clone [21].

The aim of this research was to investigate the abilities of *S. aureus* isolates representative of clones causing infection in our patient population to adhere to immobilized ligands, to investigate their cellular invasiveness and host cell death induction abilities, and to identify any associations between adherence, invasiveness or cell death induction and bacterial characteristics, such as methicillin resistance, PVL positivity and clonality.

## Materials and Methods

### Selection of representative isolates ( $n = 25$ )

From a collection of 367 well-characterized clinical *S. aureus* isolates originating from patients in the Western Cape [22], South Africa, a representative isolate was randomly selected from each major and intermediate pulsed-field gel electrophoresis clone. Two isolates from minor clones statistically associated with HIV infection and two isolates selected from the

HIV-positive patients from the dominant MRSA and methicillin-susceptible *S. aureus* (MSSA) clones were also included to investigate any specific associations with HIV infection. An MRSA isolate with a non-typeable SCCmec element was also included as representative of a unique local clone.

### Bacterial strains

All bacterial isolates were stored at  $-80^{\circ}\text{C}$  until further testing. The following isolates were used as controls: NCTC8325-4 (adherence), Cowan I (invasive control isolate), *Staphylococcus carnosus* TM300 (non-invasive control and non-cytotoxic control isolate) and 6850 (cytotoxic control isolate).

### Adherence assay

Adherence was tested first in uncoated plates to establish a baseline. Then, 96-well plates (Sarstedt, Nümbrecht, Germany) were coated with a specific ligand using a modified method of Peacock *et al.* [23]. Standardized bacterial cultures were used at an  $\text{OD}_{600} = 1$  in triplicate with three independent experiments. The plate was inoculated with bacterial culture, incubated at  $37^{\circ}\text{C}/5\% \text{CO}_2$  overnight, washed with PBS and stained using 0.1% crystal violet solution. After this, the plate was washed with PBS, eluted with 1% SDS at room temperature overnight, and subsequently measured with an ELISA reader (TECAN Infinite Pro 200, Männedorf, Switzerland) at 620 nm. Adherence to a specific ligand was expressed as a percentage relative to the positive control after subtraction of the PBS negative control. The mean of the means and standard error of mean (SEM) were determined.

### Mammalian cell culture

293 cells (www.atcc.org) were used to investigate the cellular invasiveness. 293 cells were maintained in Dulbecco's modified Eagle's medium (DMEM)/F-12 (Invitrogen, Carlsbad, CA, USA) supplemented with 10% fetal calf serum (PAA Laboratories, Pasching, Austria) and  $1 \times$  Pen/strep mix (100 U/mL penicillin and 100  $\mu\text{g}/\text{mL}$  streptomycin) (Cambrex Bio Science, Verviers, Belgium) and maintained in humidified air ( $37^{\circ}\text{C}/5\% \text{CO}_2$ ). Ea.hy926 cells (www.atcc.org) were used to investigate cell death induction. Ea.hy926 cells were maintained in DMEM/F-12 supplemented with 10% fetal calf serum and  $1 \times$  penicillin/streptomycin mix (100 U/mL penicillin and 100  $\mu\text{g}/\text{mL}$  streptomycin) and maintained in humidified air ( $37^{\circ}\text{C}/5\% \text{CO}_2$ ).

### FACS invasion assay

The cellular invasiveness was determined by adapting a previously published FACS-based invasion assay [10]. Briefly, 293 cells were plated in 24-well plates at  $3 \times 10^5$  cells/well the day before the assay. The day of the assay, the cells were washed once with 500  $\mu\text{L}$  invasion medium (1% human serum

albumin and 10 mM HEPES in DMEM), 500  $\mu$ L invasion medium was added to each well and the plate was pre-cooled at 4°C for 20 min. Following this, 50  $\mu$ L bacterial suspension (adjusted to an  $OD_{540} = 1$ ) was added to each well (MOI ~30) and incubated for 1 h at 4°C. Thereafter, the cells were shifted to 37°C with 5%  $CO_2$  for 3 h to allow for bacterial invasion. The cells were harvested and analysed by FACS analyses as previously described [9].

All bacterial suspensions were adjusted to an  $OD_{540} = 1$ . Results obtained were expressed as the average of the mean values with the SEM. An arbitrary cut-off value of 50% was used to discriminate between invasive ( $\geq 50\%$ ) and non-invasive ( $< 50\%$ ) isolates.

#### Cell death/viability assays

Bacteria were grown as described previously and standardized to an  $OD_{540} = 1$ . Ea.hy926 cells were plated in 24-well plates at  $3 \times 10^5$  cells/well for the Nicoletti assay or 96-well plates at  $5 \times 10^4$  cells/well for the lactate dehydrogenase (LDH) and WST-I assays and infected at an MOI ~30. Results obtained were expressed as the average of the mean values with the SEM. An arbitrary cut-off value of 50% was used to discriminate between cytotoxic ( $\geq 50\%$ ) and non-/moderately cytotoxic ( $< 50\%$ ) isolates.

#### Nicoletti assay

An adapted version of this previously published protocol was used [16]. The percentage of intact cells, relative to the cells-only control (set at 100%), was determined and the percentage of dead cells was calculated.

#### LDH assay

This assay was performed using the LDH cytotoxicity kit (Roche Diagnostics, Mannheim, Germany) as per the manufacturers' recommendation. The absorbance was measured using a Tecan Reader Infinite F200 Pro (Tecan GmbH, Crailsheim, Germany) at 490 nm, with a reference reading at  $> 600$  nm. Cell death induction was expressed as a percentage relative to the 1% Triton-X control (set at 100%).

#### WST-I cell viability assay

The WST-I ready-to-use reagent (Roche Diagnostics) was used as per the manufacturers' recommendations. The absorbance was measured using a Tecan Reader Infinite 200 Pro (Tecan Group, Männedorf, Switzerland) at 490 nm, with a reference reading at  $> 600$  nm. Cell viability was expressed as a percentage relative to the cells-only control (set at 100%). Results obtained were expressed as % cell death induction for direct comparison with the Nicoletti and LDH assays.

#### Statistical investigations

Associations between adherence, invasion or cell death induction and methicillin-resistance, clonality, patient HIV status or bacterial PVL status. For associations between adherence, invasion or cell death induction and methicillin-resistance, two tests were performed: the two-sample t-test and the two-sample Wilcoxon rank-sum test (Mann–Whitney test). The Mann–Whitney test was also used to investigate associations between adherence, invasion or cell death induction and bacterial PVL status. For associations between adherence, invasion or cell death induction and clonality/patient HIV status, the Kruskal–Wallis test was performed. The singletons were grouped together. For all statistical investigations, a confidence interval of 95% was used and an association was regarded as statistically significant if a  $p < 0.05$  was obtained. All statistical analyses were performed with the software package STAT v.12.

*Correlation of in vitro data.* We investigated correlations between the *in vitro* data generated by calculating the correlation coefficient ( $r$ ) and the statistical significance thereof ( $p$  value). The software package STATISTICA v.10 was used.


## Results

#### Adherence assays


The selected isolates displayed variability in their adherence to the different immobilized ligands. Some isolates displayed a preference for adherence to fibronectin and fibrinogen (CC8 and CC5) (Fig. 1) whereas others (CC30, CC15 and ST239) displayed preferences for adherence to collagen IV and VI (Fig. 2). CC22 adhered strongly to immobilized fibrinogen, fibronectin, collagen IV and VI. Some isolates such as ST88, ST97 and ST45 isolates were poor binders (Figs. 1 and 2). MRSA isolates displayed stronger adherence to plasma ( $p$  0.025) (Table 1). PVL-negative isolates adhered more strongly in the absence of a ligand ( $p$  0.01) and fibronectin ( $p$  0.034) compared to PVL-positive isolates (Table 1).

#### Cellular invasiveness

The cellular invasiveness of the isolates was diverse (42.4–211.6%; mean 142.6%; median 142.2%) (Fig. 3). All isolates, irrespective of the multilocus sequence typing clonal complex (MLST CC), were invasive except for THW-366 (ST121, MSSA PVL+). Two isolates were classified as highly invasive ( $> 200\%$ ): THW-273 (ST612:CC8; SCCmec IV; PVL– MRSA) and THW-356 (ST45:CC45; MSSA PVL–). We were unable to determine the invasiveness of isolate THW-264 (ST5:CC5; SCCmec I; PVL– MRSA). No significant difference in invasive-



**FIG. 1.** *In vitro* adherence of the representative isolates to fibronectin (black bars) and fibrinogen (white bars). The number next to each isolate's name is the MLST ST:MLST CC. The data displayed are representative of three independent experiments performed in triplicate and relative to NCTC8325-4 as the control. MLST, Multi locus sequence typing.



**FIG. 2.** *In vitro* adherence of the representative isolates to collagen IV (black bars) and collagen VI (white bars). The number next to each isolate's name is the MLST ST:MLST CC. The data displayed are representative of three independent experiments performed in triplicate and relative to NCTC8325-4 as the control. MLST, Multi locus sequence typing.

ness was observed between isolates from HIV-positive or HIV-negative patients.

#### Cell death assays

**Nicoletti assay.** Host cell death induction was diverse (23.7–79.6%; mean = 65.4%; median = 69.6%), although 21 of the isolates killed >60% of cells (Fig. 4). The following isolates were classified as non-cytotoxic: THW-382 (ST36:CC30; SCCmec II; PVL-negative MRSA); THW-264 (ST5:CC5; SCCmec I; PVL-negative MRSA) and THW-366 (ST121:CC121; PVL-positive MSSA; SSTI). Thirteen isolates could be classified as very cytotoxic (>70% cell death) (CC22, CC8, CC15, CC5, CC97, CC88).

**LDH assay.** Host cell death induction was diverse (34.1–94.3%; mean = 61.4%; median = 59.6%) (see Supporting information,

Fig. S1). Twenty-two isolates were cytotoxic and three were non-cytotoxic, of which two were also classified as non-cytotoxic using the Nicoletti assay (THW264; THW-366). THW-99 (ST239:CC8; SCCmec NT) was also non-cytotoxic using this assay. Nine isolates could be classified as very cytotoxic (>70% cell death) (CC22, CC30, CC8, CC15, CC45, CC88).

**WST-1 assay.** Host cell death induction was diverse (68–27.1%; mean = 38.1%; median = 36%) (data not shown). Twenty-two isolates were cytotoxic and three were non-cytotoxic, of which the same two isolates were previously classified as non-cytotoxic using the Nicoletti and LDH assays. The following isolate was also non-cytotoxic using this assay: (1) THW-81 (ST239:CC8; PVL-negative MRSA).

**TABLE 1.** Statistical associations between adherence, invasion and cell death and methicillin-resistance, bacterial Panton–Valentine leucocidin status, clonality or patient HIV status

Ligand	p value— MRSA/ MSSA <sup>a</sup>	p value— MRSA/ MSSA <sup>b</sup>	p value— PVL <sup>a</sup>	p value— clonality <sup>c</sup>	p value— HIV <sup>c</sup>
Uncoated	0.767	<b>0.016</b> (MRSA)	<b>0.010</b> (PVL–)	0.194	0.322
Serum	0.613	0.365	0.364	0.152	0.506
Plasma	0.767	<b>0.025</b> (MRSA)	0.250	0.530	0.849
Fibronectin	0.567	0.335	<b>0.034</b> (PVL–)	0.201	0.581
Fibrinogen	0.680	0.080	0.468	0.160	0.210
Collagen IV	0.440	0.505	0.364	0.248	0.104
Collagen VI	0.553	0.758	0.904	0.292	0.055
Invasion	0.681	0.112	<b>0.004</b> (PVL–)	<b>0.004</b> (CC8)	0.628
Nicoletti	0.680	0.901	0.758	0.758	0.089
LDH	0.507	0.910	0.586	0.586	0.829
WST-I	0.577	0.319	0.525	0.525	0.101

Statistically significant values are displayed in bold ( $p < 0.05$ ). LDH, lactate dehydrogenase; MRSA, methicillin-resistant *Staphylococcus aureus*; MSSA, methicillin-susceptible *Staphylococcus aureus*; PVL, Panton-Valentine leucocidin.

aMann–Whitney U-test.

b $t$ -test.

cKruskal–Wallis test.

Using this assay, seven isolates could be characterized as very cytotoxic (>70% viable cells) (CC22, CC8, CC45, CC88).

Two isolates were classified as non-cytotoxic using all three assays: (1) THW-366 (ST121:CC121; PVL-positive MSSA; SSTI); (2) THW-264 (ST5:CC5; SCC<sub>mec</sub> I; PVL-negative MRSA).

### Statistical associations


MRSA isolates adhered more strongly to plasma proteins than MSSA isolates, whereas PVL-negative isolates adhered more strongly to fibronectin compared with PVL-positive isolates

(Table 1). CC8 isolates were the only isolates statistically associated with being invasive. PVL-negative isolates were significantly more invasive than the PVL-positive isolates (Table 1). Regarding correlation analyses, the Pearson correlation coefficient ( $r$ ) and respective  $p$ -values for correlations between the *in vitro* assays can be found in the Supporting information, Table S2.


## Discussion

Our collection of isolates displayed a diverse range of adherence to the different ligands. ST612 was the dominant MRSA clone identified and has previously been reported in Australia [24] and South Africa [25]. Fibronectin and fibrinogen may play a role in SSTI. Two of the ST612 isolates were isolated from SSTI. Elgalai *et al.* also described a collection of wound isolates that displayed a diverse range of adherence and similarly to our set of isolates, displayed stronger adherence to fibrinogen than to fibronectin [26]. Seidl *et al.* described no significant adherence potentials to immobilized fibronectin between isolates derived from persistent and resolving MRSA infections [27]. Unlike previously reported by Arciola *et al.* [28], adherence to both human collagen IV and VI varied greatly among the isolates. The reason for the stronger adherence seen by MRSA isolates to human serum proteins, as well as for PVL-negative isolates to fibronectin, remains unclear, as all isolates tested positive for *fnbA* and *fnbB* (data not shown). We might speculate that the clonal lineages associated with methicillin-resistance and PVL-negativity prefer a preference for adherence to these ligands.

We were able to establish that 23/25 isolates were invasive. No association could be identified between invasion and



**FIG. 3.** Invasiveness of representative South African *Staphylococcus aureus* isolates determined by FACS-based invasion assay on 293 cells. The number next to each isolate's name is the MLST ST:MLST CC. Invasiveness is expressed as a percentage relative to Cowan I for three independent experiments performed in duplicate. MLST, Multi locus sequence typing.



**FIG. 4.** Induction of host cell death of representative South African *Staphylococcus aureus* isolates determined by the Nicoletti assay on EA.hy926 cells. The number next to each isolate's name is the MLST ST:MLST CC. Cytotoxicity is expressed as a percentage relative to the cells only control of three independent experiments performed in duplicate. MLST, Multi locus sequence typing.

methicillin-resistance. An association between clonality and invasiveness was only identified for CC8 isolates, the only CC with a large enough number of isolates to allow for this analysis. All CC8 isolates were classified as being invasive and displayed a high degree of cellular invasiveness (range 133.3–200.5%). Fowler *et al.* also previously reported that most *S. aureus* genotypes are capable of causing invasive diseases, but that CC5 and CC30 isolates displayed greater levels of haematogenous complication [29]. Zautner *et al.* identified a variety of CCs associated with recurrent tonsillitis-causing isolates as a result of intracellular persisting *S. aureus*. They characterized these isolates and identified many of them to be invasive isolates from CC30, CC45, CC8, CC5, CC15 and CC22 [30]. Park *et al.* investigated the association between the invasiveness of clinical *S. aureus* isolates and their abilities to cause metastatic infections [31]. They found that invasiveness of endothelial cells is not a major determinant of metastatic infections. When investigating the expression patterns of human umbilical endothelial cells upon internalization of *S. aureus*, Stark *et al.* found that innate immune responses dominated, irrespective of the invasiveness of the bacterial isolates [32]. THW-366 (ST121; PVL+ MSSA; associated with SSTI; 42.4% invasiveness) was the only isolate classified as non-invasive and also the only *agr* IV isolate. This ST has previously been associated with community acquired MRSA SSTI in children in Portugal [33]. No data on the invasiveness were available from these studies. Our data contradict data published by Zautner *et al.*, who identified three ST121 isolates associated with recurrent tonsillitis. The isolates displayed invasiveness of 33.7%, 56.5% and 65.3% relative to Cowan I, the invasive control [30]. THW-264 (ST5:CC5; SCCmec I; PVL– MRSA) is another isolate of interest as we were unable to determine the invasiveness of this isolate due

to technical difficulties. We speculate that this isolate might be *pIs*<sup>+</sup>, since this gene is commonly associated with SCCmec I. It has previously been demonstrated that MRSA isolates expressing *pIs* displayed a significantly reduced ability to invade host cells [34]. We hypothesize that this isolate might be non-invasive. This isolate was collected from a paediatric patient suffering from conjunctivitis.

Twenty-three isolates were cytotoxic (>50% cell death using all three assays) and the remaining two isolates were non-cytotoxic (<50% cell death using all three assays). No association was identified between cell death and methicillin-resistance or clonality. Four isolates stood out as they were the only isolates classified as very cytotoxic according to all three assays, resulting in >70% cell death for every assay (THW-262 (ST612-MRSA-IV), THW-70 (ST612-MRSA-IV); THW-356 (ST45-MSSA) and THW-255 (ST88-MRSA-IV)). These data show that PVL– isolates (THW262; THW70; THW271; THW273; THW412; THW356) can be as cytotoxic as PVL+ isolates (THW146; THW393; THW255) and a very high degree of cytotoxicity was even more common in PVL– isolates. A possible cause of this phenotypic display of high levels of cytotoxicity might be due to the action of phenol soluble modulins, as it has been shown that phenol soluble modulins is required for phagosomal escape following phagocytosis by non-professional phagocytes, allowing for intracellular bacterial replication and dissemination [22].

The same applied for MSSA and MRSA isolates. Highly cytotoxic MRSA and MSSA isolates were identified. Three highly cytotoxic MRSA isolates were identified (THW262; THW70; THW255) and one highly cytotoxic MSSA isolate was identified (THW356). Also, one non-cytotoxic MSSA and MRSA isolate was identified. It is interesting to note that the most cytotoxic MRSA isolates all carry SCCmec IV. This is of

interest because ST88 is associated with community-acquired MRSA. ST612 has previously been described from hospital isolates in South Africa. Four isolates were classified as very cytotoxic according to the Nicoletti and LDH assays only. We can speculate that these isolates are capable of inducing host cell death using both apoptotic and necrotic mechanisms. These isolates were identified from a pus swab of the tibia of an HIV-positive man, fluid from a sinus fungal infection and pus from a hand infection from another HIV-positive female. Park *et al.* investigated if any association was present between cytotoxicity of clinical *S. aureus* isolates and their abilities to cause metastatic infections [31]. They found that the ability to induce host cell death and be cytotoxic to endothelial cells was not a major determinant of metastatic infections. During the investigation of ten MRSA (CC45 and CC5) isolates collected during a national clinical trial, Seidl *et al.* determined that all ten isolates were invasive in human endothelial cells, but only some of the isolates were cytotoxic [13]. Our data contradict those of Krut *et al.*, who published cytotoxicity data for a collection of *S. aureus* isolates, which displayed cytotoxicity to be a strain-specific characteristic, and more than half of the collection of isolates were classified as non-cytotoxic [35]. It is worth noting that the study conducted by Krut *et al.* focused on murine cell lines, whereas our study used a human cell line to investigate host cell death induction.

## Conclusion

Clinical *S. aureus* isolates collected at Tygerberg hospital displayed diversity in their abilities to adhere to various immobilized human ligands. MRSA isolates displayed stronger adherence to human plasma proteins than MSSA isolates, while PVL<sup>-</sup> isolates adhered more strongly to fibronectin and were as cytotoxic as PVL<sup>+</sup> isolates. The vast majority of clinical *S. aureus* isolates included in this study were invasive, although differences in the degree of invasiveness were common between different CCs. ST121 was the only ST in this collection classified as non-invasive. The vast majority of these isolates are also able to induce the death of host cells. No differences in cellular invasiveness or cytotoxicity of isolates from HIV-positive and/or HIV-negative persons were detectable. Isolates containing virulence factors such as PVL or methicillin-resistant isolates were not found to be more invasive or cytotoxic than PVL<sup>-</sup> or MSSA isolates.

### Study limitation

Isolates used were a convenience sample and bias could have been introduced through the selection. The statistical analyses

performed assume that a representative random sample of isolates were constituted.

## Acknowledgements

The work was supported by the Medical Research Council (MRC) of South Africa and the Harry Crossley Foundation (HCF) of Stellenbosch University. WFO was supported by the DFG/NRF International Research and Training Group (IRTG) 1522 'HIV and associated infectious diseases in Southern Africa' and by a travel grant from Stellenbosch University to attend ISSSI2012. This work was presented, in part, at the 15th International Symposium on Staphylococci and Staphylococcal Infections (ISSSI), Lyon, France, 26–30 August, 2012. We would like to thank Barbara Conrad and Heidi Linß for expert technical assistance with the invasion and Nicoletti assays.

## Conflict of Interest

None declared.

## Supporting Information

Additional Supporting Information may be found in the online version of this article:

**Table S1.** Strains selected as representative isolates

**Table S2.** Pearson correlation (*r*) for the correlation between different assays and the corresponding *p*-value. Statistically significant (*p* < 0.05) correlations are displayed in bold

**Fig. S1.** Induction of host cell death of representative South African *Staphylococcus aureus* isolates determined by lactate dehydrogenase (LDH) assay on EA.hy926 cells. The number next to each isolates name is the MLST ST:MLST CC. Cytotoxicity is expressed as a percentage relative to the Triton-X control of three independent experiments performed in triplicate.

## References

1. Que Y-A, Moreillon P. *Staphylococcus aureus* (including staphylococcal toxic shock). In: Mandell GL, Bennett JE, Dolin R, eds, *Principles and practice of infectious diseases*, 7th edn. Philadelphia: Elsevier, 2010; 2543–2578.
2. Cho S, Strickland I, Boguniewicz M, Leung DYM. Fibronectin and fibrinogen contribute to the enhanced binding of *Staphylococcus aureus* to atopic skin. *J Allergy Clin Immunol* 2001; 108: 269–274.



3. Switalski LM, Ryden C, Rubin K, Ljungh A, Hook M, Wadstrom T. Binding of fibronectin to *Staphylococcus aureus*. *Infect Immun* 1983; 42: 628–633.
4. Moreillon P, Entenza JM, Francioli P et al. Role of *Staphylococcus aureus* coagulase and clumping factor in pathogenesis of experimental endocarditis. *Infect Immun* 1995; 63: 4738–4743.
5. Speziale P, Raucci G, Visai L, Switalski LM, Timpl R, Hook M. Binding of collagen to *Staphylococcus aureus* Cowan I. *J Bacteriol* 1986; 167: 77–81.
6. Gordon RJ, Lowy FD. Pathogenesis of methicillin-resistant *Staphylococcus aureus* infection. *Clin Infect Dis* 2008; 46(suppl 5): S350–S359.
7. Hauck CR, Ohlsen K. Sticky connections: extracellular matrix protein recognition and integrin-mediated cellular invasion by *Staphylococcus aureus*. *Curr Opin Microbiol* 2006; 9: 5–11.
8. Foster TJ, Hook M. Surface protein adhesins of *Staphylococcus aureus*. *Trends Microbiol* 1998; 6: 484–488.
9. Sinha B, Francoise PP, Busse O et al. Fibronectin-binding proteins act as *Staphylococcus aureus* invasin via fibronectin bridging to the integrin  $\alpha_5\beta_1$ . *Cell Microbiol* 1999; 1: 101–117.
10. Sinha B, Francois P, Que Y-A et al. Heterologously expressed *Staphylococcus aureus* fibronectin-binding proteins are sufficient for invasion of host cells. *Infect Immun* 2000; 68: 6871–6878.
11. Grundmeier M, Hussain M, Becker P, Heilmann C, Peters G, Sinha B. Truncation of fibronectin-binding proteins in *Staphylococcus aureus* strain Newman leads to deficient adherence and host cell invasion due to loss of the cell wall anchor function. *Infect Immun* 2004; 72: 7155–7163.
12. Haggar A, Hussain M, Lonnie H, Herrmann M, Norrby-Teglund A, Flock JI. Extracellular adherence protein from *Staphylococcus aureus* enhances internalization into eukaryotic cells. *Infect Immun* 2003; 71: 2310–2317.
13. Seidl K, Bayer AS, McKinnell JA, Ellison S, Filler SG, Xiong YQ. *In vitro* endothelial cell damage is positively correlated with enhanced virulence and poor vancomycin responsiveness in experimental endocarditis due to methicillin-resistant *Staphylococcus aureus*. *Cell Microbiol* 2011; 13: 1530–1541.
14. Mestre MB, Colombo MI. Camp and epac are key players in the regulation of the signal transduction pathway involved in the  $\alpha$ -hemolysin autophagic response. *PLoS Pathog* 2012; 8: e1002664.
15. Menzies BE, Kourteva I. Internalization of *Staphylococcus aureus* by endothelial cells induces apoptosis. *Infect Immun* 1998; 66: 5994–5998.
16. Haslinger-Loffler B, Kahl BC, Grundmeier M et al. Multiple virulence factors are required for *Staphylococcus aureus*-induced apoptosis in endothelial cells. *Cell Microbiol* 2005; 7: 1087–1097.
17. Loffler B, Hussain M, Grundmeier M et al. *Staphylococcus aureus* Pantón–Valentine leukocidin is a very potent cytotoxic factor for human neutrophils. *PLoS Pathog* 2010; 6: e1000715.
18. Diep BA, Chan L, Tattevin P et al. Polymorphonuclear leukocytes mediate *Staphylococcus aureus* Pantón–Valentine leukocidin-induced lung inflammation and injury. *Proc Natl Acad Sci USA* 2010; 107: 5587–5592.
19. Kaltsas A, Guh A, Mediavilla JR et al. Frequency of Pantón–Valentine leukocidin-producing methicillin-sensitive *Staphylococcus* strains in patients with complicated skin and soft tissue infection in Bronx, New York. *J Clin Microbiol* 2011; 49: 2992–2995.
20. Lina G, Piemont Y, Godail-Gamot F et al. Involvement of Pantón–Valentine leukocidin-causing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clin Infect Dis* 1999; 29: 1128–1132.
21. McDougal LK, Steward CD, Killgore GE, Chaitram JM, McAllister SK, Tenover FC. Pulsed-field gel electrophoresis typing of oxacillin-resistant *Staphylococcus aureus* isolated from the United States: establishing a national database. *J Clin Microbiol* 2003; 41: 5113–5120.
22. Grosz M, Kolter J, Paprotka K et al. Cytoplasmic replication of *Staphylococcus aureus* upon phagosomal escape triggered by phenol-soluble modulins. *Cell Microbiol* 2014; 16: 451–465.
23. Peacock SJ, Day NPJ, Thomas MG, Berendt AR, Foster TJ. Clinical isolates of *Staphylococcus aureus* exhibit diversity in *fnb* genes and adhesion to human fibronectin. *J Infect* 2000; 41: 23–31.
24. Bell JM, Turnidge JD, Participants SA. High prevalence of oxacillin-resistant *Staphylococcus aureus* isolates from hospitalized patients in Asia-Pacific and South Africa: results from sentry antimicrobial surveillance program, 1998–1999. *Antimicrob Agents Chemother* 2002; 46: 879–881.
25. Jansen van Rensburg MJ, Eliya Madikane V, Whitelaw A, Chachage M, Haffeejee S, Gay Elisha B. The dominant methicillin-resistant *Staphylococcus aureus* clone from hospitals in Cape Town has an unusual genotype: ST612. *Clin Microbiol Infect* 2011; 17: 785–792.
26. Elgalal I, Foster HA. Comparison of adhesion of wound isolates of *Staphylococcus aureus* to immobilized proteins. *J Appl Microbiol* 2003; 94: 413–420.
27. Seidl K, Bayer AS, Fowler VG et al. Combinatorial phenotypic signatures distinguish persistent from resolving methicillin-resistant *Staphylococcus aureus* bacteremia isolates. *Antimicrob Agents Chemother* 2011; 55: 575–582.
28. Arciola CR, Campoccia D, Gamberini S, Baldassarri L, Montanaro L. Prevalence of *cna*, *fnbA* and *fnbB* adhesin genes among *Staphylococcus aureus* isolates from orthopedic infections associated to different types of implant. *FEMS Microbiol Lett* 2005; 246: 81–86.
29. Fowler VG Jr, Nelson CL, McIntyre LM et al. Potential associations between hematogenous complications and bacterial genotype in *Staphylococcus aureus* infection. *J Infect Dis* 2007; 196: 738–747.
30. Zautner AE, Krause M, Stropahl G et al. Intracellular persisting *Staphylococcus aureus* is the major pathogen in recurrent tonsillitis. *PLoS One* 2010; 5: e9452.
31. Park WB, Kim SH, Kang Ci et al. *In vitro* ability of *Staphylococcus aureus* isolates from bacteraemic patients with and without metastatic complications to invade vascular endothelial cells. *J Med Microbiol* 2007; 56: 1290–1295.
32. Stark L, Matussek A, Strindhall JAN et al. *Staphylococcus aureus* isolates from blood and anterior nares induce similar innate immune responses in endothelial cells. *APMIS* 2009; 117: 814–824.
33. Conceição T, Aires-de-Sousa M, Pona N et al. High prevalence of ST121 in community-associated methicillin-susceptible *Staphylococcus aureus* lineages responsible for skin and soft tissue infections in Portuguese children. *Eur J Clin Microbiol Infect Dis* 2010; 30: 293–297.
34. Juuti KM, Sinha B, Werbick C, Peters G, Kuusela PI. Reduced adherence and host cell invasion by methicillin-resistant *Staphylococcus aureus* expressing the surface protein *pls*. *J Infect Dis* 2004; 189: 1574–1584.
35. Krut O, Utermohlen O, Schlossherr X, Kronke M. Strain-specific association of cytotoxic activity and virulence of clinical *Staphylococcus aureus* isolates. *Infect Immun* 2003; 71: 2716–2723.