

University of Groningen

## Lipids in hepatic glycogen storage diseases

Derks, Terry G. J.; van Rijn, Margreet

*Published in:*  
Journal of Inherited Metabolic Disease

*DOI:*  
[10.1007/s10545-015-9811-2](https://doi.org/10.1007/s10545-015-9811-2)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2015

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Derks, T. G. J., & van Rijn, M. (2015). Lipids in hepatic glycogen storage diseases: Pathophysiology, monitoring of dietary management and future directions. *Journal of Inherited Metabolic Disease*, 38(3), 537-543. <https://doi.org/10.1007/s10545-015-9811-2>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Lipids in hepatic glycogen storage diseases: pathophysiology, monitoring of dietary management and future directions

Terry G. J. Derks · Margreet van Rijn

Received: 17 November 2014 / Revised: 30 December 2014 / Accepted: 6 January 2015 / Published online: 30 January 2015  
© The Author(s) 2015. This article is published with open access at Springerlink.com

**Abstract** Hepatic glycogen storage diseases (GSD) underscore the intimate relationship between carbohydrate and lipid metabolism. The hyperlipidemias in hepatic GSD reflect perturbed intracellular metabolism, providing biomarkers in blood to monitor dietary management. In different types of GSD, hyperlipidemias are of a different origin. Hypertriglyceridemia is most prominent in GSD type Ia and associated with long-term outcome morbidity, like pancreatitis and hepatic adenomas. In the ketotic subtypes of GSD, hypertriglyceridemia reflects the age-dependent fasting intolerance, secondary lipolysis and increased mitochondrial fatty acid oxidation. The role of high protein diets is established for ketotic types of GSD, but non-traditional dietary interventions (like medium-chain triglycerides and the ketogenic diet) in hepatic GSD are still controversial and necessitate further studies. Patients with these rare inherited disorders of carbohydrate metabolism meet several criteria of the metabolic syndrome, therefore close monitoring for cardiovascular diseases in ageing GSD patients may be justified.

## Abbreviations

BMI	Body mass index
CNGDF	Continuous nocturnal gastric drip-feeding
CPT I	Carnitine palmitoyltransferase I
EGP	Endogenous glucose production
ESGSD I	European Study on GSD I
G6Pase	Glucose-6-phosphatase

The manuscript was presented at the last SSIEM symposium in Innsbruck, Sept 2-5, 2014.

Communicated by: John H Walter

T. G. J. Derks (✉) · M. van Rijn  
Section of Metabolic Diseases, Beatrix Children's Hospital,  
University of Groningen, University Medical Center Groningen, PO  
Box 30 001, 9700 RB Groningen, The Netherlands  
e-mail: t.g.j.derks@umcg.nl

GNG	Gluconeogenesis
GSD	Glycogen storage disease
MCT	Medium-chain triglycerides
mFAO	Mitochondrial fatty acid oxidation
UCCS	Uncooked cornstarch

## Introduction

To date there are 14 types of glycogen storage diseases (GSD), which are classified according to enzyme or transporter deficiency and organ distribution (Laforêt et al 2011). The hepatic GSDs are inborn errors of glycogen synthesis or breakdown clinically characterized by hepatomegaly, failure to thrive and fasting intolerance, biochemically associated with hypoglycaemia. Both endogenous glucose production (EGP) and metabolic clearance rate of glucose determine blood glucose concentration during fasting. Impairment of EGP may be caused by decreased gluconeogenesis (GNG), decreased glycogenolysis or a combination. Instead of discussing all hepatic GSD types, this review focuses on lipid metabolism and monitoring of dietary management in GSD type I and GSD type III. These GSD types display important differences in both pathophysiology and effects on lipid metabolism. During follow-up, monitoring and prevention from significant long-term morbidity is the rule in GSD I and GSD III, whereas to date, the remaining subtypes (except GSD IV) are generally considered disorders without morbidity in adulthood.

GSD I is caused by a deficiency of the glucose-6-phosphatase (G6Pase) complex (Bali et al 1993). G6Pase is expressed in liver, kidney and intestine and plays a central role in both glycogenolysis and GNG. This explains the severe short-term fasting intolerance and hypoglycemia compared

to patients with other hepatic types of GSD. Besides severe nonketotic hypoglycemia, metabolic decompensation is characterized by secondary biochemical abnormalities (hyperlactacidaemia, hyperuricaemia and hyperlipidaemia). Untreated patients classically displayed protruding abdomen (hepatomegaly due to storage of glycogen and fat), short stature, truncal obesity, a rounded doll face, wasted muscles and bleeding tendency. Two major subtypes of GSD I are recognized. GSD type Ia (OMIM #232200) is caused by deficient activity of the catalytic unit of G6Pase, whereas GSD Ib (OMIM #232220) is caused by defect of the putative transporter. In addition to the classical phenotype, GSD Ib patients display recurrent bacterial infections, inflammatory bowel disease and thyroid auto-immunity, associated with neutropenia and neutrophil dysfunction.

The remaining hepatic GSD types (0, III, VI, IX and XI) are associated with fasting ketotic hypoglycemia and considered relatively mild compared to GSD I because GNG is intact (Derks and Smit 2014). In these types of GSD, ketone body (KB) concentrations reflect increased mitochondrial fatty acid oxidation (mFAO), which is proceeded by activation of GNG and secondary endogenous proteolysis from muscle tissue. GSD III (OMIM # 232400) is caused by a deficiency of the debranching enzyme and phenotypically classified as either type IIIa or type IIIb (Dagli et al 1993). GSD IIIa patients ( $\pm 85\%$ ) display symptoms and signs due to the enzyme deficiency in liver, skeletal muscle and heart, whereas the remaining patients with GSD IIIb ( $\pm 15\%$ ) have only liver-related phenotypes.

Dietary management in hepatic GSD is traditionally based on the provision of exogenous carbohydrates to compensate defective EGP, to achieve normoglycemia and to correct the secondary metabolic effects as much as possible. In infant GSD I patients, normoglycemia is maintained by frequent (every 1.5–3 h) lactose free formula feeds enriched with maltodextrin. In 1974 continuous nocturnal gastric drip-feeding (CNGDF) via a nasogastric tube was introduced. It was demonstrated that normoglycemia could be maintained (Burr et al 1974; Greene et al 1976), secondary metabolic alterations could be corrected and patients and parents were able to sleep throughout the entire night. In 1984 uncooked cornstarch (UCCS) was described for the first time in GSD patients and developed as either a good alternative or additional strategy to maintain metabolic control (Chen et al 1984; Smit et al 1984). Although UCCS may be introduced at 6 months of age, the tolerance may be reduced as a consequence of lower pancreatic amylase activity until 1 year of age (Hayde and Widhalm 1990). Both CNGDF and UCCS have advantages and disadvantages and differ in many ways, like the optimal age of introduction, duration of normoglycemia, (no) need to wake up during the night, caloric content, glycaemic index, risk of technical failure with subsequent metabolic risks, difficulty for parents and/or patients, invasiveness and

costs (Derks et al 2013). Dietary management in GSD III patients follows similar principles of exogenous carbohydrates, but the additional role of increasing dietary protein is recognized.

After dietary treatment became available, the phenotype of GSD patients changed from mortality to morbidity and the focus of attention moved towards (prevention of) long-term complications (Moses 2002). For GSD I patients, these complications mainly involve the liver (adenomas and their risks of bleeding and secondary conformation to hepatocellular carcinoma), kidney (glomerular and tubular dysfunction, developing into renal insufficiency), intestine (GSD related enteropathy), growth (end length, body mass index (BMI)), anemia, joints (gout) and bones (osteopenia and osteoporosis) (Bali et al 1993). For GSD III patients, the liver (fibrosis, cirrhosis and hepatocellular carcinoma), muscle (myopathy), heart (cardiomyopathy), growth and bones deserve special attention (Dagli et al 1993).

### Monitoring of plasma lipids in hepatic GSD

In GSD I patients, triglyceride concentration are considered the most useful parameter for chronic metabolic control. Like blood glucose concentrations, lactate concentrations fluctuate rapidly, liver enzymes are only slightly increased before initiation of dietary management and rapidly normalize, and uric acid concentrations can be influenced by allopurinol treatment. In our personal clinical experience with GSD Ia patients, increasing triglyceride concentrations can reflect decreased metabolic control, for instance when young patients outgrow their diet, by decreased compliance, during increased growth in puberty or co-morbidity like hypothyroidism. Based on an expert meeting as part of the European Study on GSD I (ESGSD I), follow-up guidelines for the frequency of routine laboratory investigations (including lipid profiles) are age dependent: age 0–3 years every 2 months; 3–20 years every 3 months; adults every 6 months (Rake et al 2002b).

Cross-sectional cohort data on serum lipid concentrations in GSD1 patients originate from the retrospective multicenter ESGSD I, which included 231 GSD Ia and 57 GSD Ib patients (Rake et al 2002a). Hypercholesterolemia and triglyceridemia were more common and more severe in GSD Ia patients compared to GSD Ib patients. In adult GSD Ia ( $n=40$ ) and GSD Ib ( $n=4$ ) patients hypercholesterolemia was noted in 43 % versus 25 %, whereas 98 % versus 75 % displayed hypertriglyceridemia. In this cohort, complications associated with hyperlipidaemia were rare; pancreatitis was reported in three patients and cholelithiasis in two patients. In the management guideline paper originating from 2002, serum triglyceride concentrations  $<6.0$  mmol/L was included as a biomedical target (Rake et al 2002b).

In the ESGSD I the overall prevalence of liver adenomas was 16 % and increased with age (Rake et al 2002a). More recently, Wang and co-workers retrospectively characterized the natural history and factors related to liver adenoma development in GSD1a patients (Wang et al 2011). Patients were stratified in two groups, i.e. patients with mean 5-year triglyceride concentrations  $\leq 500$  mg/dL (corresponding with 5.6 mmol/L) or  $> 500$  mg/dL, based on the consensus panel discussion at the 2010 Association for Glycogen Storage Disease Conference. The authors reported significantly increased adenoma progression in the latter group. By demonstrating the association between a biomarker (i.e. increased triglyceride concentrations) and outcome (i.e. increased prevalence of liver adenomas), the authors added a new dimension.

In general, hyperlipidemia disappears with age and is milder in the ketotic types of GSD, like GSD III. Bernier et al studied the lipid profiles in 44 GSD III patients ranging from 6 months to 30 years of age (Bernier et al 2008). They demonstrated that the overall prevalences of hypercholesterolemia (31 %) and hypertriglyceridemia (67 %) are lower than in GSD I patients. Hypertriglyceridemia significantly decreased with increasing age in GSD III patients, the median slope was  $-0.11$  mmol/L per year.

### Pathophysiology of lipid metabolism in hepatic GSD

From the point of pathophysiology, several important factors contribute to the difference between GSD I and GSD III.

The fasted state in GSD I is characterized by intracellular accumulation of glucose-6-phosphate with secondary shunting to glycolysis and pentosephosphate route. Glycolysis increases acetyl-CoA production and thereby lipogenesis and malonyl-CoA, the latter inhibiting carnitine palmitoyltransferase I (CPT I), the rate-controlling step of mFAO. Pharmacological inhibition of the glucose 6-phosphate transporter by S4048 induces hypoglycemia in wild type mice, increased hepatic glycogen and triglyceride content, and a markedly enhanced hepatic lipogenic gene expression (van Dijk et al 2001). Human stable isotope studies revealed that GSD Ia is associated with increased rates of de novo lipogenesis and cholesterologenesis (Bandsma et al 2008). After administration of S4048 to several knock-out mice models, the carbohydrate-response-element-binding protein (but not the sterol-regulatory-element-binding protein 1c or liver X receptor  $\alpha$ ) was identified as an important molecular link mediating the induction of hepatic lipogenic gene expression in GSD I, hence a potential target for pharmacological intervention in human patients (Grefhorst et al 2010). Several metabolites of glycolysis (like glucose-6-phosphate and fructose-2,6-bisphosphate) and the pentosephosphate route (xylulose-5-phosphate) act as activators of the carbohydrate-response-element-binding protein, illustrating

the complex interactions between intermediate metabolites and transcription factors (for a recent review, see Oosterveer and Schoonjans 2014).

Experimental data from stable isotope studies demonstrate that EGP is age dependent and decreases relatively with body weight (Bier et al 1977) and age (Huidekoper et al 2014). Because in GSD I patients both glycogenolysis and GNG are impaired, defective EGP and fasting intolerance are more pronounced and lifelong, compared to the ketotic GSD types like GSD III. In the latter group, hyperlipidemia is caused by increased fatty acid flux from adipose tissue to the liver, as an alternative source of energy by mFAO. The decreasing triglyceride concentrations with increasing age goes in parallel with decreasing EGP, reflecting improved fasting tolerance in ageing GSD III patients.

### The role of dietary carbohydrate, fat and protein

First reports about dietary energy percentages of macronutrients in GSD already originate from over four decades ago. Kelsch and Oliver demonstrated that frequent high-carbohydrate feeds improved most metabolic abnormalities in a GSD I patient (Kelsch and Oliver 1969). Fernandes and Pikaar recommended 50–70 and 40–50 % carbohydrates, 15–35 and 30–40 % fat and 15 and 20 % protein, in patients with GSD I and ketotic subtypes, respectively (Fernandes and Pikaar 1969). Fernandes also studied the effects of different disaccharides on blood lactate concentrations and these studies are the experimental base of dietary restrictions of fructose, lactose and saccharose (Fernandes 1974).

Since then, between treatment centers worldwide, major differences have arisen in the practical effectuation of dietary management in GSD (type I). Examples include the type of nocturnal management (UCCS versus CNGDF), the timing of the introduction of UCCS (at which age?), the methods by which exogenous carbohydrates are given and the extent of dietary restrictions of fructose, lactose and saccharose. Recent data from a meta-analysis of three studies suggest that UCCS was associated with significantly higher plasma cholesterol concentrations compared to CNGDF (Shah and O'Dell 2013). There was no significant difference in plasma triglyceride concentrations between these treatments. The difference in lipid profiles between UCCS and heat-treated modified starch has not been systematically studied. In general, these examples emphasize the need of international detailed (dietary) data collection and practical management guidelines appreciating that dietary management in hepatic GSD needs to be individually tailored.

Next to the recommendations about dietary energy percentages of macronutrients, data from stable isotope studies on EGP in humans are helpful to estimate the absolute carbohydrate requirements. Taking into account, that the


theoretical EGP decreases with age, the exogenous glucose requirements for GSD I patients can be estimated with regression formulas derived from these tracer studies. The historical studies by Bier and co-workers have generated a regression formula using body weight as a variable; recently Huidekoper and co-workers generated a regression formula based on age (Fig. 1). Compared to the latter method, the historical method may overestimate carbohydrate requirements because it obviously does not correct for overweight/obesity. At some time points in childhood, the calculation based on age is 25–30 % lower compared to the other method. In addition, both methods share another risk of overtreatment because the tracer studies have been performed in healthy overnight fasted control subjects, a situation different from the perturbed glucose homeostasis in GSD I patients. Experimental data in GSD Ia patients indicate that *in vivo* EGP is not completely absent and may reach ~60 % of normal (Huidekoper et al 2010). The mechanism is not understood, theoretically, residual *in vivo* glucose-6-phosphatase- $\alpha$  activity next to alternative glycogenolysis by (muscle) glucose-6-phosphatase- $\beta$  and/or the  $\alpha$ -glucosidase pathway may contribute to whole body EGP in GSD Ia patients.

Figure 2 presents total energy expenditure and EGP in childhood, both expressed in kcal/day. The graph illustrates that the caloric intake by replacement of EGP declines from 60 to 30 % in the first 3 years of life. This may indicate that especially in (early) childhood, dietary management is characterized by a delicate balance between *undertreatment* (with

hypoglycemia, poor metabolic control and risk of long-term complications) and *overtreatment* (with the risk of rebound hypoglycemia, peripheral body fat storage and obesity). In the European Study on GSD I, about 25 % of the GSD Ia and 33 % of the GSD Ib patients under 20 years of age display BMI above the 90th percentile, interestingly highest prevalence was observed in young patients between 2 and 5 years of age.

From a practical point of view, increasing one macronutrient (carbohydrates) without affecting the second (protein) inevitably lowers fat intake. Given the carbohydrate deficiency and hyperlipidemia in GSD I and GSD III, increasing dietary fat is somewhat counterintuitive. However, there are two positive reports about high-fat diets in GSD III patients. The combination of synthetic ketone bodies, 2:1 ketogenic diet and high-protein was associated with reversal of hypertrophic cardiomyopathy in a 2-months-old GSD IIIa patient (Valayannopoulos et al 2011). In two 7- and 5-year-old siblings, a high-carbohydrate diet was isocalorically replaced by a high fat (60 %) and high protein (25 %, which remained fairly constant), low-carbohydrate (15 %) diet, associated with reversal of hypertrophic cardiomyopathy (Brambilla et al 2014).


Several case reports describe the effects of medium chain triglycerides (MCT) in GSD I and III (Fernandes and Pikaar 1969; Cuttino et al 1970; Nagasaka et al 2007; Hanou et al 2008; Burns et al 2009; Das et al 2010; Bernstein et al 2010; El-Gharbawy et al 2014). In summary, laboratory parameters



**Fig. 1** Two regression models to calculate endogenous glucose production, based on body weight (*grey*, by Bier et al 1977) and age (*black*, by Huidekoper et al 2014). The *grey line* represents the EGP regression line according to Bier. Reference values for body weight in boys were retrieved from the World Health Organization and the Centers for Disease Control and Prevention, below 2 years and after 2 years of

age, respectively. EGP (expressed in mg/min) is calculated as:  $0.0014x^3 - 0.214x^2 + 10.411x - 9.084$ , with  $[x]$  representing body weight in kg. The *black line* represents EGP regression line according to Huidekoper. EGP (expressed in mg/kg/min) is calculated as:  $6.50 \times 2.72^{-0.145 \times Z} + 1.93$ , with  $[z]$  representing age in years

**Fig. 2** Daily total energy expenditure (*black*) and endogenous glucose production calculated by age (*grey*), both expressed in kcal/day, according to age. Reference values for total energy expenditure (expressed as kcal/kg/day) in boys were retrieved from the Joint FAO/WHO/UNU Expert Consultation on Human Energy Requirements. Reference values for body weight in boys were retrieved from the World Health Organization and the Centers for Disease Control and Prevention, below 2 years and after 2 years of age, respectively. EGP was calculated according to Huidekoper



of metabolic control (triglyceride concentrations in GSD I; transaminases and CK in GSD III) decrease after MCT-supplementation, demonstrating that hyperlipidemia is carbohydrate-induced in GSD I. However, an overall interpretation with conclusions and management advice based on these observations is complicated by differences in age, GSD type and severity (type of mutations) of the disorder, relative and absolute quantities of macronutrients, MCT-sources, indications for MCT-treatment, outcome-parameters, the absence of long-term clinically relevant follow-up data and the possibility of selection bias, unfavourable cases not being reported.

There are no experimental data to substantiate whether long-term dietary MCT may cause liver adenoma and hepatocellular carcinoma. The development of hepatic tumors is enhanced by a high fat enriched diet in liver-specific GSD I deficient mice (Rajas et al 2014), but it is not mentioned whether this depends on the type of fat, i.e. MCT versus long-chain triglycerides. In GSD I *cytoplasmatic* Acetyl-CoA accumulation and subsequent malonyl-CoA formation functionally inhibit mFAO at the level of CPT I. MCT are absorbed via the portal vein and the released free fatty acids enter mFAO without the carnitine shuttle. The beta-oxidation generates *intramitochondrial* Acetyl-CoA for direct ketogenesis. It can be hypothesized that increased ketogenesis by MCT decreases the metabolic clearance rate of glucose in GSD I patients, although the exact molecular mechanism is still unknown.

MCT can be supplied by naturally occurring dietary products, medical dietary supplements, or pharmaceutically (triheptanoin). For all methods, the potential risk of glycerol overload towards impaired GNG in GSD I patients is unknown. Triheptanoin is a medium-chain triglyceride of 7-carbon fatty acids observationally studied in patients with long-chain mitochondrial fatty acid oxidation disorders.

Besides the ease of use, the anapleurotic effect of triheptanoin GSD Ia could be beneficial.

The role of dietary protein in GSD I has not systematically been studied and has recently been reviewed for GSD III (Derks and Smit 2014).

### Discussion and future directions

In summary, hepatic GSD underscores the intimate relations between carbohydrate and lipid metabolism. Hyperlipidemia is most pronounced and often permanent in GSD I, in whom the intrahepatic degree of metabolic control is reflected by blood lipid concentrations. Hypertriglyceridemia is associated with pancreatitis and liver adenomas in GSD I patients. In ketotic GSD patients, hyperlipidemia reflects lipolysis from extrahepatic sources, associated with the age-dependent fasting intolerance.

Like often in rare diseases, medical emergencies in individual patients initiated (dietary) interventions and generated new hypotheses. Future experimental animal studies and well-designed collaborative dietary intervention studies are necessary to improve our understanding about manipulations in macronutrients, like the restrictions of simple sugars, traditional versus heat-treated modified starch, the role of protein, MCT-treatment, triheptanoin and the ketogenic diet in GSD type III. International patient registries will be necessary to study outcome and the effects of (elements of) dietary management.

In the next decades, dietary management of GSD patients will be individualized more than ever, integrating personal metabolic and genetic data to optimize metabolic control. Continuous glucose monitoring systems are used in GSD patients in the real life situation to improve dietary management and thereby parameters of metabolic control, including the lipid profile

(Hershkovitz et al 2001; White and Jones 2011; Kasapkara et al 2014). Nowadays, the most modern (and expensive) devices alarm at low glucose levels, significantly improving safety. Near infra red spectroscopy is a promising technique for non-invasive measurement of blood glucose (Xue et al 2014) and the integration of this technique in the glucose monitoring systems would add new dimensions to the safety and acceptability of dietary management of GSD patients.

The definition of the metabolic syndrome and their criteria has been revised several times, leading to multiple definitions in current use. However, the general concept is that patients have a combination of conditions, among which are insulin resistance or hyperglycemia, hyperlipidemia, hypertension and obesity. Regardless of which definition, GSD I patients meet some of the criteria applied. Currently there are conflicting experimental data (Ubels et al 2002; Bernier et al 2009) and no convincing epidemiological data substantiating strong associations between metabolic syndrome, type 2 diabetes and GSD. Close clinical monitoring for cardiovascular diseases, chronic kidney disease and diabetes in ageing GSD patients (regardless what subtype) seems to be justified, because these disorders share many metabolic pathways (Rajas et al 2013).

**Acknowledgments** Taking care of GSD-patients is teamwork and the discussions with the colleagues in our department have been very stimulating while preparing the presentation and this subsequent manuscript. Therefore the authors are thankful to Rixt van der Ende, Cynthia van Amerongen, Irene Hooegeveen, Joost Holslag, Chris Peter Sentner, Esther van Dam, Foekje de Boer, Peter Smit, Tom de Koning and Francjan van Spronsen.

#### Compliance with Ethics Guidelines

**Conflict of interest** In the last 5 years, Terry GJ Derks had received speaker's fees from Recordati Rare Diseases, Danone Nutricia and Vitaflo, research fees from Sigma Tau and Vitaflo, and training support from Sigma Tau and Genzyme.

Margreet van Rijn is a member of the European Nutrition Expert Panel (Merck Serono), Danone Research and Development consultant, Member of the Advisory Board Element (Danone-Nutricia), Financial research support of Nutricia Netherlands, Honoraria for lecturing and guideline development (Orphan Europe, Excemed, Nutricia).

**Human and animal rights and informed consent** This article does not contain any studies with human or animal subjects performed by the any of the authors.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

#### References

Bali DS, Chen YT, Goldstein JL (1993) Glycogen storage disease type I. In: Pagon RA, Adam MP, Ardinger HH et al (eds) GeneReviews(R). University of Washington, Seattle, Seattle

- Bandsma RH, Prinsen BH, van Der Velden MS et al (2008) Increased de novo lipogenesis and delayed conversion of large VLDL into intermediate density lipoprotein particles contribute to hyperlipidemia in glycogen storage disease type 1a. *Pediatr Res* 63:702–707
- Bernier AV, Sentner CP, Correia CE, Theriaque DW, Shuster JJ, Smit GP, Weinstein DA (2008) Hyperlipidemia in glycogen storage disease type III: effect of age and metabolic control. *J Inherit Metab Dis* 31: 729–732
- Bernier AV, Correia CE, Haller MJ, Theriaque DW, Shuster JJ, Weinstein DA (2009) Vascular dysfunction in glycogen storage disease type I. *J Pediatr* 154:588–591
- Bernstein LE, Burns CE, Wilkinson LJ, Boney A, Balliet J, Van Hove J (2010) Treatment of elevated triglycerides in glycogen storage disease type 1a and hypertriglyceridemia with medium chain triglycerides sources. *J Inherit Metab Dis* 33:S173–S173
- Bier DM, Leake RD, Haymond DW, Arnold KJ, Gruenke LD, Sperling MA, Kipnis DM (1977) Measurement of “true” glucose production rates in infancy and childhood with 6,6-dideuteroglucose. *Diabetes* 26:1016–1023
- Brambilla A, Mannarino S, Pretese R, Gasperini S, Galimberti C, Parini R (2014) Improvement of cardiomyopathy after high-fat diet in two siblings with glycogen storage disease type III. *JIMD Rep* 17:91–95
- Burns CE, Bernstein LE, Wilkinson LJ, Boney A, Myers S, Van Hove J (2009) Treatment of elevated triglycerides in glycogen storage disease type and hypertriglyceridemia with new medium chain triglycerides sources. *Mol Genet Metab* 98:56–57
- Burr IM, O'Neill JA, Karzon DT, Howard LJ, Greene HL (1974) Comparison of the effects of total parenteral nutrition, continuous intragastric feeding, and portacaval shunt on a patient with type I glycogen storage disease. *J Pediatr* 85:792–795
- Chen YT, Cornblath M, Sidbury JB (1984) Cornstarch therapy in type I glycogen-storage disease. *N Engl J Med* 310:171–175
- Cuttino JT Jr, Summer GK, Hill HD Jr, Mitchell BJ (1970) Response to medium chain triglycerides in von Gierke's disease. *Pediatrics* 46: 925–929
- Dagli A, Sentner CP, Weinstein DA (1993) Glycogen storage disease type III. In: Pagon RA, Bird TD, Dolan CR, Stephens K, Adam MP (eds) GeneReviews. University of Washington, Seattle
- Das AM, Lucke T, Meyer U, Hartmann H, Illsinger S (2010) Glycogen storage disease type I: impact of medium-chain triglycerides on metabolic control and growth. *Ann Nutr Metab* 56:225–232
- Derks TG, Smit GP (2014) Dietary management in glycogen storage disease type III: what is the evidence? *J Inherit Metab Dis*. doi:10.1007/s10545-014-9756-x
- Derks TG, Martens DH, Sentner CP, van Rijn M, de Boer F, Smit GP, van Spronsen FJ (2013) Dietary treatment of glycogen storage disease type Ia: uncooked cornstarch and/or continuous nocturnal gastric drip-feeding? *Mol Genet Metab* 109:1–2
- El-Gharbawy AH, Arnold GL, Perrott-Taylor N, Hughley T, Long K, Vockley J, Kishnani PS (2014) Optimizing metabolic control of glycogen storage disease type 3 (Gsd3): potential role for medium chain triglycerides (Mct). *Mol Genet Metab* 111:284–285
- Fernandes J (1974) The effect of disaccharides on the hyperlactacidemia of glucose-6-phosphatase deficient children. *Acta Paediatr Scand* 63:695–698
- Fernandes J, Pikaar NA (1969) Hyperlipemia in children with liver glycogen disease. *Am J Clin Nutr* 22:617–627
- Greene HL, Slonim AE, O'Neill JA Jr, Burr IM (1976) Continuous nocturnal intragastric feeding for management of type 1 glycogen-storage disease. *N Engl J Med* 294:423–425
- Grefhorst A, Schreurs M, Oosterveer MH et al (2010) Carbohydrate-response-element-binding protein (ChREBP) and not the liver X receptor alpha (LXRalpha) mediates elevated hepatic lipogenic gene expression in a mouse model of glycogen storage disease type 1. *Biochem J* 432:249–254

- Hanou CE, Wilkinson LJ, Bernstein LE (2008) Treatment of fatty acid oxidation disorders, glycogen storage disease type 1A and hypertriglyceridemia with new medium chain triglycerides sources. *J Inherit Metab Dis* 31:146–146
- Hayde M, Widhalm K (1990) Effects of cornstarch treatment in very young children with type I glycogen storage disease. *Eur J Pediatr* 149:630–633
- Hershkovitz E, Rachmel A, Ben-Zaken H, Phillip M (2001) Continuous glucose monitoring in children with glycogen storage disease type I. *J Inherit Metab Dis* 24:863–869
- Huidekoper HH, Visser G, Ackermans MT, Sauerwein HP, Wijburg FA (2010) A potential role for muscle in glucose homeostasis: in vivo kinetic studies in glycogen storage disease type 1a and fructose-1,6-bisphosphatase deficiency. *J Inherit Metab Dis* 33:25–31
- Huidekoper HH, Ackermans MT, Ruitter AF, Sauerwein HP, Wijburg FA (2014) Endogenous glucose production from infancy to adulthood: a non-linear regression model. *Arch Dis Child* 99:1098–1102
- Kasapkara ÇS, Cinasal Demir G, Hasanoğlu A, Tümer L (2014) Continuous glucose monitoring in children with glycogen storage disease type I. *Eur J Clin Nutr* 68:101–105
- Kelsch RC, Oliver WJ (1969) Studies on dietary correction of metabolic abnormalities in hepatorenal glycogenosis. *Pediatr Res* 3:160–170
- Laforêt P, Weinstein DA, Smit GPA (2011) The glycogen storage diseases and related disorders, chap 6. In Saudubray JM, van den Berghe G, Walter J (eds). *Inborn metabolic diseases*. Springer, Berlin
- Moses SW (2002) Historical highlights and unresolved problems in glycogen storage disease type I. *Eur J Pediatr* 161:S2–S9
- Nagasaka H, Hirano K, Ohtake A et al (2007) Improvements of hypertriglyceridemia and hyperlactemia in Japanese children with glycogen storage disease type 1a by medium-chain triglyceride milk. *Eur J Pediatr* 166:1009–1016
- Oosterveer MH, Schoonjans K (2014) Hepatic glucose sensing and integrative pathways in the liver. *Cell Mol Life Sci* 71:1453–1467
- Rajas F, Labrune P, Mithieux G (2013) Glycogen storage disease type 1 and diabetes: learning by comparing and contrasting the two disorders. *Diabetes Metab* 39:377–387
- Rajas F, Clar J, Gautier-Stein A, Mithieux G (2014) Lessons from new mouse models of glycogen storage disease type 1a in relation to the time course and organ specificity of the disease. *J Inherit Metab Dis*. doi:10.1007/s10545-014-9761-0
- Rake JP, Visser G, Labrune P, Leonard JV, Ullrich K, Smit GP (2002a) Glycogen storage disease type I: diagnosis, management, clinical course and outcome. Results of the European study on glycogen storage disease type I (ESGSD I). *Eur J Pediatr* 161(Suppl 1):S20–S34
- Rake JP, Visser G, Labrune P, Leonard JV, Ullrich K, Smit GP, European Study on Glycogen Storage Disease Type I (ESGSD I) (2002b) Guidelines for management of glycogen storage disease type I - European study on glycogen storage disease type I (ESGSD I). *Eur J Pediatr* 161(Suppl 1):S112–S119
- Shah KK, O'Dell SD (2013) Effect of dietary interventions in the maintenance of normoglycaemia in glycogen storage disease type 1a: a systematic review and meta-analysis. *J Hum Nutr Diet* 26:329–339
- Smit GP, Berger R, Potasnick R, Moses SW, Fernandes J (1984) The dietary treatment of children with type I glycogen storage disease with slow release carbohydrate. *Pediatr Res* 18:879–881
- Ubels FL, Rake JP, Slaets JP, Smit GP, Smit AJ (2002) Is glycogen storage disease 1a associated with atherosclerosis? *Eur J Pediatr* 161(Suppl 1):S62–S64
- Valayannopoulos V, Bajolle F, Arnoux J et al (2011) Successful treatment of severe cardiomyopathy in glycogen storage disease type III With D, L-3-hydroxybutyrate, ketogenic and high-protein diet. *Pediatr Res* 70:638–641
- van Dijk TH, van der Sluijs FH, Wiegman CH et al (2001) Acute inhibition of hepatic glucose-6-phosphatase does not affect gluconeogenesis but directs gluconeogenic flux toward glycogen in fasted rats. A pharmacological study with the chlorogenic acid derivative S4048. *J Biol Chem* 276:25727–25735
- Wang DQ, Fiske LM, Carreras CT, Weinstein DA (2011) Natural history of hepatocellular adenoma formation in glycogen storage disease type I. *J Pediatr* 159:442–446
- White FJ, Jones SA (2011) The use of continuous glucose monitoring in the practical management of glycogen storage disorders. *J Inherit Metab Dis* 34:631–642
- Xue J, Chen H, Xiong D, Huang G, Ai H, Liang Y, Yan X, Gan Y, Chen C, Chao R, Ye L (2014) Noninvasive measurement of glucose in artificial plasma with near-infrared and Raman spectroscopy. *Appl Spectrosc* 68:428–433