

University of Groningen

## Modeling two-dimensional infrared spectroscopy of hydrogen bonded systems

De Carvalho Vicente Da Cunha, Ana

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2017

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

De Carvalho Vicente Da Cunha, A. (2017). *Modeling two-dimensional infrared spectroscopy of hydrogen bonded systems*. [Thesis fully internal (DIV), University of Groningen]. Rijksuniversiteit Groningen.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## BIBLIOGRAPHY

- [1] O. Lafont, "Nicolas Lémery and acidity," *Rev. Hist. Pharm. (Paris)*, vol. 50, pp. 53–62, 2002.
- [2] N. Lémery, *Cours de chymie contenant la maniere de faire les operations qui sont en usage dans la medecine, par une methode facile. Avec des raisonnemens sur chaque operation, pour l'instruction de ceux qui veulent s'appliquer à cette science.* 6th ed., Michallet, 1687.
- [3] P.-J. Macquer, *Elemens de Chymie Theorique.* Herissantt, 1749.
- [4] J. Dalton, *A new system of chemical philosophy.* London, 1808.
- [5] Y. Marchal, ed., *The Hydrogen Bond and the Water Molecule.* Amsterdam: Elsevier, 1st ed., 2007.
- [6] G. Lewis, "The atom and the molecule," *J. Am. Chem. Soc.*, vol. 38, pp. 762–785, 1916.
- [7] L. Pauling, "The nature of the chemical bond. IV. The energy of single bonds and the relative electronegativity of atoms.," *J. Chem. Am. Soc.*, vol. 54, pp. 3570–3582, 1932.
- [8] I. Dzyaloshinskii, E. Lifshitz, and L. Pitaevskii, "The general theory of van der waals forces," *Advances in Physics*, vol. 10, pp. 165–209, 1961.
- [9] F. London, "The general theory of molecular forces," *Trans. Faraday Soc.*, vol. 33, pp. 8b–26, 1937.
- [10] R. Eisenschitz and F. London, "Über das verhältnis der van der waalsschen kräfte zu den homöopolaren bindungskräften," *Z. Physik*, vol. 60, pp. 491–527, 1930.
- [11] P. W. Atkins and L. Jones, *Chemical principles.* 4th ed., W.H. Freeman and Company, New York, 2008.
- [12] S. Hammes-Schiffer and S. J. Benkovic, "Relating protein motion to catalysis," *Annu. Rev. Biochem.*, vol. 75, p. 519, 2006.

- [13] K. A. Henzler-Wildman, M. Lei, V. Thai, S. J. Kerns, M. Karplus, and D. A. Kern, "Hierarchy of timescales in protein dynamics is linked to enzyme catalysis," *Nature*, vol. 450, p. 913, 2007.
- [14] J. P. Erzberger and J. M. Berger, "Evolutionary relationships and structural mechanisms of AAA+ proteins," *Annu. Rev. Biophys. Biomol. Struct.*, vol. 35, p. 93, 2006.
- [15] H. Frauenfelder, S. G. Sligar, and P. G. Wolynes, "The energy landscapes and motions of proteins," *Science*, vol. 254, p. 1598, 1991.
- [16] R. Jimenez, G. Salazar, J. Yin, T. Joo, and F. E. Romesberg, "Protein dynamics and the immunological evolution of molecular recognition," *P. Natl. Acad. Sci. USA*, vol. 101, p. 3803, 2004.
- [17] M. K. Hong, D. Braunstein, B. R. Cowen, H. Frauenfelder, I. E. T. Iben, J. R. Mourant, P. Ormos, R. Scholl, A. Schulte, P. J. Steinbach, A. H. Xie, and R. D. Young, "Conformational substates and motions in myoglobin - external influences on structure and dynamics," *Biophys. J.*, vol. 58, p. 429, 1990.
- [18] H. Frauenfelder, B. H. McMahon, R. H. Austin, K. Chu, and J. T. Groves, "The role of structure, energy landscape, dynamics, and allostery in the enzymatic function of myoglobin," *P. Natl. Acad. Sci. USA*, vol. 98, p. 2370, 2001.
- [19] M. C. Thielges, J. Y. Axup, D. Wong, H. S. Lee, J. K. Chung, P. G. Schultz, and M. D. Fayer, "Two-dimensional ir spectroscopy of protein dynamics using two vibrational labels: A site-specific genetically encoded unnatural amino acid and an active site ligand," *J. Phys. Chem. B*, vol. 115, p. 11294, 2011.
- [20] H. Ishikawa, K. Kwak, J. K. Chung, S. Kim, and M. D. Fayer, "Direct observation of fast protein conformational switching," *P. Natl. Acad. Sci. USA*, vol. 105, p. 8619, 2008.
- [21] E. L. Hahn, "Spin echoes," *Phys. Rev.*, vol. 80, pp. 580–594, 1950.
- [22] J. Jeener, B. H. Meier, P. Bachmann, and R. R. Ernst, "Investigation of exchange processes by two-dimensional NMR spectroscopy," *J. Chem. Phys.*, vol. 71, pp. 4546–4553, 1979.

- [23] P. Hamm, M. Lim, and R. M. Hochstrasser, "Structure of the amide I band of peptides measured by femtosecond nonlinear-infrared spectroscopy," *J. Phys. Chem. B*, vol. 102, pp. 6123–6138, 1998.
- [24] T. L. C. Jansen and J. Knoester, "Two-dimensional infrared population transfer spectroscopy for enhancing structural markers of proteins," *Biophys. J.*, vol. 94, pp. 1818–1825, 2008.
- [25] H. Torri, "Comparison of different optical processes of the two-dimensional infrared spectroscopy in the effect of intermolecular vibrational coupling calculated in the time domain: The case of liquid N,N-dimethylformamide," *Vibr. Spectr.*, vol. 42, pp. 140 – 146, 2006.
- [26] W. Zhuang, D. Abramavicius, T. Hayashi, and S. Mukamel, "Simulation protocols for coherent femtosecond vibrational spectra of peptides," *J. Phys. Chem. B*, vol. 110, pp. 3362–3374, 2006.
- [27] S. Ham, J. H. Kim, H. Lee, and M. H. Cho, "Correlation Between Electronic and Molecular Structure Distortions and Vibrational Properties. II. Amide I Modes of NMA-nD<sub>2</sub>O Complexes," *J. Chem. Phys.*, vol. 118, pp. 3491–3498, 2003.
- [28] H. Maekawa and N. H. Ge, "Comparative study of electrostatic models for the amide-I and -II modes: Linear and Two-dimensional infrared spectra," *J. Phys. Chem. B*, vol. 114, p. 1434, 2010.
- [29] L. Wang, C. T. Middleton, M. T. Zanni, and J. L. Skinner, "Development and Validation of Transferable Amide I Vibrational Frequency Maps for Peptides," *J. Phys. Chem. B*, vol. 115, pp. 3713–3724, 2011.
- [30] M. Reppert and A. Tokmakoff, "Electrostatic frequency shifts in amide I vibrational spectra: Direct parameterization against experiment," *J. Chem. Phys.*, vol. 138, p. 134116, 2013.
- [31] J. Wang, W. Zhuang, S. Mukamel, and R. M. Hochstrasser, "Two-Dimensional Infrared Spectroscopy as a Probe of the Solvent Electrostatic Field for a Twelve Residue Peptide," *J. Phys. Chem. B*, vol. 112, pp. 5930–5937, 2008.
- [32] E.-L. Karjalainen, T. Ersmark, and A. Barth, "Optimization of model parameters for describing the amide I spectrum of a large set of proteins," *J. Phys. Chem. B*, vol. 116, pp. 4831–4842, 2012.

- [33] M. W. D. Hanson-Heine, F. S. Husseini, J. D. Hirst, and N. A. Besley, "Simulation of the two-dimensional infrared spectroscopy of peptides using localized normal modes," *J. Chem. Theory Comput.*, vol. 12, pp. 1905–1918, 2016.
- [34] P. Bour and T. A. Keiderling, "Empirical modeling of the peptide amide I band IR intensity in water solution," *J. Chem. Phys.*, vol. 119, p. 11253, 2003.
- [35] S. Ham and M. Cho, "Amide I modes in the N-methylacetamide dimer and glycine dipeptide analog: Diagonal force constants," *J. Chem. Phys.*, vol. 118, pp. 6915–6922, 2003.
- [36] T. L. C. Jansen and J. Knoester, "A transferable electrostatic map for solvation effects on amide I vibrations and its application to linear and two-dimensional spectroscopy," *J. Chem. Phys.*, vol. 124, p. 044502, 2006.
- [37] J. Lessing, S. Roy, M. Reppert, M. Baer, D. Marx, T. Jansen, J. Knoester, and A. Tokmakoff, "Identifying residual structure in intrinsically disordered systems: A 2D IR spectroscopic study of the GVGXPGVG peptide," *J. Am. Chem. Soc.*, vol. 134, pp. 5032–5035, 2012.
- [38] R. D. Gorbunov, D. S. Kosov, and G. Stock, "Ab initio-based exciton model of amide I vibrations in peptides: Definition, conformational dependence, and transferability," *J. Chem. Phys.*, vol. 122, p. 224904, 2005.
- [39] T. M. Watson and J. D. Hirst, "Theoretical studies of the amide I vibrational frequencies of [leu]-enkephalin," *Molec. Phys.*, vol. 103, pp. 1531–1546, 2005.
- [40] H. Torii, "Amide I vibrational properties affected by hydrogen bonding out-of-plane of the peptide group," *J. Phys. Chem. Lett.*, vol. 6, pp. 727–733, 2015.
- [41] T. M. Watson and J. D. Hirst, "Influence of Electrostatic Environment of the Vibrational Frequencies of Proteins," *J. Phys. Chem. A*, vol. 107, pp. 6843–6849, 2003.
- [42] F. Ingrosso, G. Monard, M. H. Farag, A. Bastida, and M. F. Ruiz-Lopez, "Importance of polarization and charge transfer effects to model the infrared spectra of peptides in solution," *J. Chem. Theory Comput.*, vol. 7, pp. 1840–1849, 2011.

- [43] M. Reppert and A. Tokmakoff, "Quantitative multi-site frequency maps for amide I vibrational spectroscopy," *J. Chem. Phys.*, vol. 143, p. 061102, 2015.
- [44] N. H. Andersen, *Protein structure, stability and folding, methods in molecular biology*. Humana Press, 2001.
- [45] M. L. Huggins, "term. paper. some speculations regarding molecular structure. (unpublished paper)," *Chem.* 120, 1919.
- [46] W. M. Latimer and W. H. Rodebush, "Polarity and ionization from the standpoint of Lewis theory of valence.," *J. Am. Chem. Soc.*, vol. 42, pp. 1419–1433, 1920.
- [47] G. C. Pimentel and A. L. McClellan, *The hydrogen bond*. W. H. Freeman (Reinhol Publishing Corporation), 1960.
- [48] M. H. Chadwell, "The molecular structure of water," *Chem. Rev.*, vol. 4, pp. 375–398, 1927.
- [49] S. Tsuzuki and H. P. Löthi, "Interaction energies of van der Waals and hydrogen bonded systems calculated using density functional theory: Assessing the PW91 model," *J. Chem. Phys.*, vol. 114, pp. 3949–3957, 2001.
- [50] S. Tsuzuki, T. Uchimaru, K. Matsumura, M. Mikami, and K. Tanabe, "Effects of basis set and electron correlation on the calculated interaction energies of hydrogen bonding complexes: MP2/cc-pV5Z calculations of H<sub>2</sub>O-MeOH, H<sub>2</sub>O-Me<sub>2</sub>O, H<sub>2</sub>O-H<sub>2</sub>CO, MeOH-MeOH, and HCOOH-HCOOH complexes," *J. Chem. Phys.*, vol. 110, pp. 11906–11910, 1999.
- [51] J. W. Larson and T. B. McMahon, "Gas-phase bihalide and pseudobihalide ions. An ion cyclotron resonance determination of hydrogen bond energies in XHY- species (X, Y = F, Cl, Br, CN)," *Inorg. Chem.*, vol. 23, pp. 2029–2033, 1984.
- [52] O. Markovitch and N. Agmon, "Structure and energetics of the hydronium hydration shells," *J. Phys. Chem. A*, vol. 111, pp. 2253–2256, 2007.
- [53] J. Emsley, "Very strong hydrogen bonding," *Chem. Soc. Rev.*, vol. 9, pp. 91–124, 1980.

- [54] J. Zielkiewicz, "Structural properties of water: Comparison of the SPC, SPCE, TIP4P, and TIP5P models of water," *J. Chem. Phys.*, vol. 123, p. 104501, 2005.
- [55] A. Tokmakoff, "Shining light on the rapidly evolving structure of water," *Science*, vol. 317, p. 54, 2007.
- [56] M. Cowan, B. Bruner, N. Huse, J. R. Dwyer, B. Chugh, E. T. J. Nibbering, T. Elsaesser, and R. J. D. Miller, "Ultrafast memory loss and energy redistribution in the hydrogen bond network of liquid H<sub>2</sub>O," *Nature*, vol. 434, p. 199, 2005.
- [57] J. B. Asbury, T. Steinel, C. Stromberg, J. K. Gaffney, I. Piletic, and M. Fayer, "Hydrogen bond breaking probed with multidimensional stimulated vibrational echo correlation spectroscopy," *J. Chem. Phys.*, vol. 119, p. 12981, 2003.
- [58] J. B. Asbury, T. Steinel, K. Kwak, S. Corcelli, C. P. Lawrence, J. L. Skinner, and M. Fayer, "Dynamics of water probed with vibrational echo correlation spectroscopy," *J. Chem. Phys.*, vol. 121, p. 12431, 2004.
- [59] C. J. Fecko, J. D. Eaves, A. T. J. J. Loparoand, and P. L. Geissler, "Ultrafast hydrogen-bond dynamics in the infrared spectroscopy of water," *Science*, vol. 301, p. 1698, 2003.
- [60] R. Laenen, G. M. Gale, and N. Lascoux, "IR spectroscopy of hydrogen-bonded methanol: Vibrational and structural relaxation on the femtosecond time scale," *J. Phys. Chem. A*, vol. 103, p. 10708, 1999.
- [61] S. Woutersen, U. Emmerichs, and H. Bakker, "A femtosecond midinfrared pump-probe study of hydrogen-bonding in ethanol," *J. Chem. Phys.*, vol. 107, p. 1483, 1997.
- [62] K. Mazur, M. Bonn, and J. Hunger, "Hydrogen bond dynamics in primary alcohols: A femtosecond infrared study," *J. Phys. Chem. B*, vol. 119, p. 1558, 2015.
- [63] S. Roy, M. S. Pshenichnikov, and T. L. C. Jansen, "Analysis of 2D CS spectra for systems with non-gaussian dynamics," *J. Phys. Chem. B*, vol. 115, p. 5431, 2011.

- [64] C. P. van der Vegte, S. Knop, P. Vöhringer, J. Knoester, and T. L. C. Jansen, "OH-stretching in synthetic hydrogen-bonded chains," *J. Phys. Chem. B*, vol. 118, p. 6256, 2014.
- [65] S. Knop, T. L. C. Jansen, J. Lindner, and P. Vöhringer, "On the nature of OH-stretching vibrations in hydrogen-bonded chains: Pump frequency dependent vibrational lifetime," *Phys. Chem. Chem. Phys.*, vol. 13, p. 4641, 2011.
- [66] J. M. Gulbis, Z. Kelman, J. Hurwitz, M. O'Donnell, and J. Kuriyan, "Structure of the c-terminal region of p21<sup>WAF1/CIP1</sup> complexed with human PCNA," *Cell*, vol. 87, pp. 297–306.
- [67] Z. Ganim, S. A. W. Chung, H. S., L. P. DeFlores, K. C. Jones, and A. Tokmakoff, "Amide I two-dimensional infrared spectroscopy of proteins," *Acc. Chem. Research*, vol. 41, pp. 432–441, 2008.
- [68] A. Tokmakoff, M. J. Lang, D. S. Larsen, G. R. Fleming, V. Chernyak, and S. Mukamel, "Two-dimensional raman spectroscopy of vibrational interaction in liquids," *Phys. Rev. Lett.*, vol. 79, p. 2702, 1997.
- [69] S. Woutersen and P. Hamm, "Nonlinear two-dimensional vibrational spectroscopy of peptides," *J. Phys.: Condens. Matter*, vol. 14, p. 1035, 2002.
- [70] H. S. Chung, M. Khalil, and A. Tokmakoff, "Nonlinear infrared spectroscopy of protein conformational change during thermal unfolding," *J. Phys. Chem. B*, vol. 108, p. 15332, 2004.
- [71] H. S. Chung, M. Khalil, A. W. Smith, Z. Ganim, and A. Tokmakoff, "Conformational changes during the nanosecond to millisecond unfolding of ubiquitin," *Proc. Natl. Acad. Sci. U.S.A.*, vol. 102, p. 612, 2005.
- [72] H. S. Chung, Z. Ganim, K. C. Jones, and A. Tokmakoff, "Transient 2D IR spectroscopy of ubiquitin unfolding dynamics," *Proc. Natl. Acad. Sci. U.S.A.*, vol. 104, p. 14237, 2007.
- [73] A. T. Krummel, P. Mukherjee, and M. T. Zanni, "Inter and intrastrand vibrational coupling in dna studied with heterodyned 2D IR spectroscopy," *J. Phys. Chem. B*, vol. 107, p. 9165, 2003.



- [74] Y. S. Kim and R. M. Hochstrasser, "Chemical exchange 2D IR of hydrogen-bond making and breaking," *P. Natl. Acad. Sci. USA*, vol. 102, p. 11185, 2005.
- [75] C. R. Baiz, C. S. Peng, M. Reppert, K. C. Jones, and A. Tokmakoff, "Coherent two-dimensional infrared spectroscopy: Quantitative analysis of protein secondary structure in solution," *Analyst*, vol. 137, pp. 1739–1799, 2012.
- [76] M. Fayer, *Ultrafast Infrared Vibrational Spectroscopy*. CRC Press, 1 ed., 2013.
- [77] A. Barth and C. Zscherp, "What vibrations tell about proteins," *Q. Rev. Biophys.*, vol. 35, pp. 369–430, 2002.
- [78] S. Woutersen and P. Hamm, "Time-resolved two-dimensional vibrational spectroscopy of a short  $\alpha$ -helix in water," *J. Chem. Phys.*, vol. 115, pp. 7737–7743, 2001.
- [79] N. Demirdoven, C. M. Cheatum, H. S. Chung, M. Khalil, J. Knoester, and A. Tokmakoff, "Two-dimensional infrared spectroscopy of antiparallel  $\beta$ -sheet secondary structure," *J. Am. Chem. Soc.*, vol. 126, pp. 7981–7990, 2004.
- [80] S. D. Moran, A. M. Woys, L. E. Buchanan, E. Bixby, S. M. Decatur, and M. T. Zanni, "Two-dimensional ir spectroscopy and segmental  $^{13}\text{C}$  labeling reveals the domain structure of human  $\gamma$ d-crystallin amyloid fibrils," *P. Natl. Acad. Sci. USA*, vol. 109, pp. 3329–3334, 2012.
- [81] A. M. Woys, A. M. Almeida, L. Wang, C.-C. Chiu, M. McGovern, J. J. de Pablo, J. L. Skinner, S. H. Gellman, and M. T. Zanni, "Parallel  $\beta$ -sheet vibrational couplings revealed by 2D IR spectroscopy of an isotopically labeled macrocycle: Quantitative benchmark for the interpretation of amyloid and protein infrared spectra," *J. Am. Chem. Soc.*, vol. 134, pp. 19118–19128, 2012.
- [82] T. L. C. Jansen and J. Knoester, "Waiting time dynamics in two-dimensional infrared spectroscopy," *Acc. Chem. Research*, vol. 42, pp. 1405–1411, 2009.

- [83] C. Liang and T. L. C. Jansen, “An efficient  $N^3$ -scaling propagation scheme for simulating two-dimensional infrared and visible spectra,” *J. Chem. Theory Comput.*, vol. 8, pp. 1706–1713, 2012.
- [84] A. B. J. and T. E. Wainwright, “Studies in molecular dynamics. I. General method,” *J. Chem. Phys.*, vol. 31, pp. 459–466, 1959.
- [85] A. Rahman, “Correlations in the motion of atoms in liquid argon,” *Phys. Rev.*, vol. 136, pp. A405–A411, 1964.
- [86] J. B. Gibson, A. N. Goland, M. Milgram, and G. H. Vineyard, “Dynamics of radiation damage,” *Phys. Rev.*, vol. 120, pp. 1229–1253, 1960.
- [87] K. J. Kollath and M. C. Standage, *Stark Effect*. 1979.
- [88] B. M. Auer and J. L. Skinner, “IR and Raman spectra of liquid water: Theory and interpretation,” *J. Chem. Phys.*, vol. 128, p. 224511, 2008.
- [89] S. Mukamel, *Principles of non-linear optical spectroscopy*. New York: Oxford University Press, 1995.
- [90] K. A. Beauchamp, Y.-S. Lin, R. Das, and V. S. Pande, “Are Protein Force Fields Getting Better? A Systematic Benchmark on 524 Diverse NMR Measurements,” *J. Chem. Theory Comput.*, vol. 8, pp. 1409–1414, 2012.
- [91] P. Hamm and M. Zanni, *Concepts and Methods of 2D Infrared Spectroscopy*. Cambridge University Press, 1 ed., 2011.
- [92] J. Bredenbeck, J. Helbing, J. R. Kumita, G. A. Woolley, and P. Hamm, “ $\alpha$ -helix formation in a photoswitchable peptide tracked from picoseconds to microseconds by time-resolved IR spectroscopy,” *P. Natl. Acad. Sci. USA*, vol. 102, pp. 2379–2384, 2005.
- [93] J. Bredenbeck and P. Hamm, “Peptide structure determination by two-dimensional infrared spectroscopy in the presence of homogeneous and inhomogeneous broadening,” *J. Chem. Phys.*, vol. 119, pp. 1569–1578, 2003.
- [94] C. Fang, J. Wang, A. K. Charnley, W. Barber-Armstrong, A. B. Smith III, S. M. Decatur, and R. M. Hochstrasser, “Two-dimensional infrared measurements of the coupling between amide modes of an  $\alpha$ -helix,” *Chem. Phys. Lett.*, vol. 382, pp. 586 – 592, 2003.

- [95] P. Mukherjee, A. T. Krummel, E. C. Fulmer, I. Kass, I. T. Arkin, and M. T. Zanni, "Site-specific vibrational dynamics of the CD3zeta membrane peptide using heterodyned two-dimensional infrared photon echo spectroscopy," *J. Chem. Phys.*, vol. 120, pp. 10215–10224, 2004.
- [96] D. B. Strasfeld, Y. L. Ling, S. Shim, and M. T. Zanni, "Tracking fiber formation in human islet amyloid polypeptide with automated 2D IR spectroscopy," *J. Am. Chem. Soc.*, vol. 130, pp. 6698–6699, 2008.
- [97] A. Woys, Y. Lin, A. S. Reddy, W. Xiong, J. de Pablo, J. Skinner, and M. Zanni, "2D IR line shapes probe ovispirin peptide conformation and depth in lipid bilayers," *J. Am. Chem. Soc.*, vol. 132, pp. 2832–2838, 2010.
- [98] Z. Ganim, K. C. Jones, and A. Tokmakoff, "Insulin dimer dissociation and unfolding revealed by amide I two-dimensional infrared spectroscopy," *Phys. Chem. Chem. Phys.*, vol. 12, pp. 3579–3588, 2010.
- [99] N. Sengupta, H. Maekawa, W. Zhuang, C. Toniolo, S. Mukamel, D. J. Tobias, and N. H. Ge, "Sensitivity of 2D IR Spectra to Peptide Helicity: A Concerted Experimental and Simulation Study of an Octapeptide," *J. Phys. Chem. B*, vol. 113, pp. 12037–12049, 2009.
- [100] C. Liang, M. Louhivuori, S. J. Marrink, T. L. C. Jansen, and J. Knoester, "Vibrational Spectra of a Mechanosensitive Membrane Channel," *J. Phys. Chem. Lett.*, vol. 4, pp. 448–452, 2013.
- [101] C. Liang, J. Knoester, and T. L. C. Jansen, "Proton transport in a membrane protein channel: Two-dimensional infrared spectrum modeling," *J. Phys. Chem. B*, vol. 116, pp. 6336–6345, 2012.
- [102] J. Wang, J. Chen, and R. M. Hochstrasser, "Local Structure of  $\beta$ -Hairpin Isotopomers by FTIR, 2D IR, and Ab Initio Theory," *J. Phys. Chem. B*, vol. 110, pp. 7545–7555, 2006.
- [103] S. Bagchi, C. Falvo, S. Mukamel, and R. M. Hochstrasser, "2D IR experiments and simulations of the coupling between amide-i and ionizable side chains in proteins: Application to the villin headpiece," *J. Phys. Chem. B*, vol. 113, pp. 11260–11273, 2009.
- [104] A. Ghosh, J. Qiu, W. F. DeGrado, and R. M. Hochstrasser, "Tidal surge in the M2 proton channel sensed by 2D IR spectroscopy," *P. Nat. Acad. Sci.*, vol. 108, pp. 6115–6120, 2011.

- [105] A. G. Dijkstra and J. Knoester, "Collective oscillations and the linear and two-dimensional infrared spectra of inhomogeneous  $\beta$ -sheets," *J. Phys. Chem. B*, vol. 109, p. 9787, 2005.
- [106] A. G. Dijkstra, T. L. C. Jansen, and J. Knoester, "Two-dimensional spectroscopy of extended molecular systems: Applications to energy transport and relaxation in an  $\alpha$ -Helix," *J. Phys. Chem. A*, vol. 114, pp. 7315–7320, 2010.
- [107] M. Reppert and A. Tokmakoff, "Computational amide I 2D IR spectroscopy as a probe of protein structure and dynamics," *Annu. Rev. Phys. Chem.*, vol. 67, pp. 16.1–16.28, 2016.
- [108] T. L. C. Jansen and J. Knoester, "Nonadiabatic effects in the two-dimensional infrared spectra of peptides application to alanine dipeptide," *J. Phys. Chem. B*, vol. 110, pp. 22910–22916, 2006.
- [109] T. Hayashi, W. Zhuang, and S. Mukamel, "Electrostatic DFT map for the complete vibrational amide band of NMA," *J. Phys. Chem. A*, vol. 109, pp. 9747–9759, 2005.
- [110] K. Lazonder, M. S. Pshenichnikov, and D. A. Wiersma, "Easy interpretation of optical two-dimensional correlation spectra," *Opt. Lett.*, vol. 31, p. 3354, 2006.
- [111] T. L. C. Jansen and J. Knoester, "Calculation of two-dimensional infrared spectra of ultrafast chemical exchange with numerical langevin simulations," *J. Phys. Chem.*, p. 234502, 2007.
- [112] J. F. Cahoon, K. R. Sawyer, J. P. Schlegel, and C. B. Harris, "Determining transition-state geometries in liquids using 2D IR," *Science.*, pp. 1820–1823, 2008.
- [113] S. Woutersen, Y. Mu, G. Stock, and P. Hamm, "Hydrogen-bond lifetime measured by time-resolved 2D IR spectroscopy: N-methylacetamide in methanol," *J. Chem. Phys.*, vol. 266, pp. 137–147, 2001.
- [114] J. Zheng, K. Kwak, J. B. Asbury, X. Chen, I. R. Piletic, and M. D. Fayer, "Ultrafast dynamics of solute-solvent complexation observed at thermal equilibrium in real time," *Science.*, vol. 309, pp. 1338–1343, 2005.

- [115] A. Barth, "Infrared spectroscopy of proteins," *Bioch. Et Biophys. Acta*, vol. 1767, p. 1073, 2007.
- [116] Z. Ganim and A. Tokmakoff, "Spectral Signatures of Heterogeneous Protein Ensembles Revealed by MD Simulations of 2D IR Spectra," *Biophys. J.*, vol. 91, p. 2636, 2006.
- [117] E. G. Buchanan, H. William, C. S. Hyuk, G. Guo, S. H. Gellman, C. Muller, and T. Zwier, "Single-conformation infrared spectra of model peptides in the amide I and amide II regions: Experiment-based determination of local mode frequencies and inter-mode coupling," *J. Chem. Phys.*, vol. 137, 2012.
- [118] J. K. Carr, A. V. Zabuga, S. Roy, T. R. Rizzo, and J. L. Skinner, "Assessment of amide I spectroscopic maps for a gas-phase peptide using IR-UV double-resonance spectroscopy and density functional theory calculations," *J. Chem. Phys.*, vol. 140, p. 224111, 2014.
- [119] W. Humphrey, A. Dalke, and K. Schulten, "VMD - Visual Molecular Dynamics," *J. Mol. Graphics*, vol. 14, pp. 33–38, 1996.
- [120] T. L. C. Jansen, A. G. Dijkstra, T. M. Watson, J. D. Hirst, and J. Knoester, "Modeling the amide I bands of small peptides," *J. Chem. Phys.*, vol. 125, p. 044312, 2006.
- [121] H. Torii and M. Tasumi, "Ab initio molecular orbital study of the amide I vibrational interactions between the peptide groups in di- and tripeptides and considerations on the conformation of the extended helix," *J. Raman Spectroscopy*, vol. 29, pp. 81–86, 1998.
- [122] K. S. Witold and H. H. Mantsch, "New insight into protein secondary structure from resolution-enhanced infrared spectra," *Biochem. et Biophys. Acta*, vol. 952, pp. 115 – 130, 1988.
- [123] T. L. C. Jansen, A. G. Dijkstra, T. M. Watson, J. D. Hirst, and J. Knoester, "Erratum: "modeling the amide I bands of small peptides" [j. chem. phys.125, 044312 (2006)]," *J. Chem. Phys.*, vol. 136, p. 209901, 2012.
- [124] E. Małolepsza and J. E. Straub, "Empirical maps for the calculation of amide I vibrational spectra of proteins from classical molecular dynamics simulations," *J. Phys. Chem. B*, vol. 118, pp. 7848–7855, 2014.

- [125] A. S. Bondarenko and T. L. C. Jansen, "Application of two-dimensional infrared spectroscopy to benchmark models for the amide I band of proteins," *J. Chem. Phys.*, vol. 142, p. 212437, 2015.
- [126] P. J. Artymiuk, C. Blake, D. Rice, and K. Wilson, "The structures of the monoclinic and orthorhombic forms of hen egg-white lysozyme at 6 Å resolution," *Acta Crys. Sec. B*, vol. 38, pp. 778–783, 1982.
- [127] E. Chatani, R. Hayashi, H. Moriyama, and T. Ueki, "Conformational strictness required for maximum activity and stability of bovine pancreatic ribonuclease a as revealed by crystallographic study of three phe120 mutants at 1.4 Å resolution," *Prot. Science*, vol. 11, pp. 72–81, 2002.
- [128] A. Deacon, T. Gleichmann, A. J. Kalb Gilboa, H. Price, J. Raftery, G. Bradbrook, J. Yariv, and J. R. Helliwell, "The structure of concanavalin a and its bound solvent determined with small-molecule accuracy at 0.94 [aring] resolution," *J. Chem. Soc., Faraday Trans.*, vol. 93, pp. 4305–4312, 1997.
- [129] A. D. MacKerell, N. Banavali, and N. Frollo, "Development and current status of the CHARMM force field for nucleic acids," *Biopol.*, vol. 56, pp. 257–265, 2000.
- [130] W. L. Jorgensen and J. Tirado-Rives, "Potential energy functions for atomic-level simulations of water and organic and biomolecular systems," *P. Natl. Acad. Sci. USA*, vol. 102, pp. 6665–6670, 2005.
- [131] D. van der Spoel, E. Lindahl, B. Hess, G. Groenhof, A. E. Mark, and H. J. C. Berendsen, "Gromacs: fast, flexible, and free," *J. Comput. Chem.*, vol. 26, pp. 1701–1718, 2005.
- [132] H. J. C. Berendsen, J. R. Grigera, and T. P. Straatsma, "The missing term in effective pair potentials," *J. Phys. Chem.*, vol. 91, pp. 6269–6271, 1987.
- [133] G. Bussi, D. Donadio, and M. Parrinello, "Canonical sampling through velocity rescaling," *J. Chem. Phys.*, vol. 126, p. 014101, 2007.
- [134] M. Parrinello and A. Rahman, "Polymorphic transitions in single crystals: A new molecular dynamics method," *J. of App. Phys.*, vol. 52, pp. 7182–7190, 1981.

- [135] C. L. Brooks III, "Computer simulation of liquids," *J. Sol. Chem.*, vol. 18, pp. 99–99, 1989.
- [136] T. Darden, D. York, and L. Pedersen, "Particle mesh Ewald: An N·log(N) method for Ewald sums in large systems," *J. Chem. Phys.*, vol. 98, pp. 10089–10092, 1993.
- [137] B. Hess, H. Bekker, H. J. C. Berendsen, and J. G. E. M. Fraaije, "LINCS: A Linear Constraint Solver for Molecular Simulations," *J. Comput. Chem.*, vol. 18, pp. 1463–1472, 1997.
- [138] C. Oostenbrink, A. Villa, A. E. Mark, and W. F. Van Gunsteren, "A biomolecular force field based on the free enthalpy of hydration and solvation: The gromos force-field parameter sets 53a5 and 53a6," *J. Comput. Chem.*, vol. 25, pp. 1656–1676, 2004.
- [139] S. Roy, J. Lessing, G. Meisl, Z. Ganim, A. Tokmakoff, J. Knoester, and T. L. C. Jansen, "Solvent and conformation dependence of amide I vibrations in peptides and proteins containing proline," *J. Chem. Phys.*, vol. 135, 2011.
- [140] P. Hamm and S. Woutersen, "Coupling of the Amide I modes of the glycine dipeptide," *Bull. Chem. Soc. Jpn.*, vol. 75, p. 985, 2002.
- [141] J. F. Kruijer, C. P. van der Vegte, and T. L. C. Jansen, "Suppressing sampling noise in linear and two-dimensional spectral simulations," *J. Chem. Phys.*, vol. 142, p. 054201, 2015.
- [142] "MATLAB and Statistics Toolbox Release 2013b, The MathWorks, Inc., Natick, Massachusetts, United States."
- [143] S. Roy, T. L. C. Jansen, and J. Knoester, "Structural Classification of the Amide I Sites of a  $\beta$ -Hairpin with Isotope Label 2D IR Spectroscopy," *Phys. Chem. Chem. Phys.*, vol. 12, pp. 9347–9357, 2010.
- [144] W. L. Jorgensen, J. Chandrasekhar, J. D. Madura, R. W. Impey, and M. L. Klein, "Comparison of Simple Potential Functions for Simulating Liquid Water," *J. Chem. Phys.*, vol. 79, pp. 926–935, 1983.
- [145] M. W. Mahoney and W. L. Jorgensen, "Diffusion constant of the TIP5P model of liquid water," *J. Chem. Phys.*, vol. 114, pp. 363–366, 2001.

- [146] J. R. Schmidt, S. T. Roberts, J. J. Loparo, A. Tokmakoff, M. D. Fayer, and J. L. Skinner, "Are Water Simulation Models Consistent with Steady-State and Ultrafast Vibrational Spectroscopy Experiments?," *Chem. Phys.*, vol. 341, pp. 143–157, 2007.
- [147] C. J. Burnham and S. S. Xantheas, "Development of transferable interaction models for water. IV. A flexible all-atom polarizable potential (TTM2-F) based on geometry dependent charges derived from an *ab initio* monomer dipole moment surface," *J. Chem. Phys.*, vol. 116, p. 5115, 2002.
- [148] C. Tainter, P. A. Pieniazek, Y.-S. Lin, and J. L. Skinner, "Robust three-body water simulation model," *J. Chem. Phys.*, vol. 134, p. 184501, 2011.
- [149] G. Corongiu and E. Clementi, "Molecular dynamics simulations with a flexible and polarizable potential: Density of states for liquid water at different temperatures," *J. Chem. Phys.*, vol. 98, p. 4984, 1993.
- [150] J. A. C. Rullmann and P. T. v. Duijnen, "A Polarizable water model for calculation of hydration energies.," *Mol. Phys.*, vol. 63, p. 451, 1988.
- [151] P. J. van Maaren and D. van der Spoel, "Molecular dynamics simulations of water with novel shell-model potentials," *J. Phys. Chem. B*, vol. 105, pp. 2618–2626, 2001.
- [152] S. W. Rick, S. J. Stuart, and B. J. Berne, "Dynamical fluctuating charge force field: Application to liquid water," *J. Chem. Phys.*, vol. 101, p. 6141, 1994.
- [153] G. Lamoureux, A. D. MacKerell, and B. Roux, "A simple polarizable model of water based on classical Drude oscillators," *J. Chem. Phys.*, vol. 119, pp. 5185–5197, 2003.
- [154] T. L. C. Jansen, "Linear absorption and two-dimensional infrared spectra of N-methylacetamide in chloroform revisited: Polarizability and multipole effects," *J. Phys. Chem. B*, vol. 118, pp. 8162–8169, 2014.
- [155] S. Patel, A. D. MacKerell, and C. L. Brooks, "CHARMM fluctuating charge force field for proteins: II protein/solvent properties from molecular dynamics simulations using a nonadditive electrostatic model.," *J. Comput. Chem.*, vol. 25, pp. 1504–1514, 2004.



- [156] H. A. Stern, G. A. Kaminski, J. L. Banks, R. Zhou, B. J. Berne, and R. Friesner, "Fluctuating Charge, Polarizable dipole, and Combined Models: Parametrization from ab initio quantum chemistry," *J. Phys. Chem. B*, vol. 103, p. 4730, 1999.
- [157] W. Xie, J. Pu, A. D. MacKerell, and J. Gao, "Development of a polarizable intermolecular potential function (PIPF) for liquid amides and alkanes," *J. Chem. Theory Comput.*, vol. 3, pp. 1878–1889, 2007.
- [158] P. E. M. Lopes, J. Huang, J. Shim, Y. Luo, H. Li, B. Roux, and A. D. MacKerell, "Polarizable force field for peptides and proteins based on the classical drude oscillator," *J. Chem. Theory Comput.*, vol. 9, pp. 5430–5449, 2013.
- [159] R. Chelli and P. Procacci, "A transferable polarizable electrostatic force field for molecular mechanics based on the chemical potential equalization principle," *J. Chem. Phys.*, vol. 117, p. 9175, 2002.
- [160] E. Harder, V. M. Anisimov, T. Whitfield, A. D. MacKerell, and B. Roux, "Understanding the dielectric properties of liquid amides from a polarizable force field," *J. Phys. Chem. B*, vol. 112, pp. 3509–3521, 2008.
- [161] W. L. Jorgensen and P. Schyman, "Treatment of halogen bonding in the OPLS-AA force field: Application to potent Anti-HIV agents," *J. Chem. Theory Comput.*, vol. 8, pp. 3895–3901, 2012.
- [162] M. Drescher, M. Huber, and V. Subramaniam, "Hunting the chameleon: Structural conformations of the intrinsically disordered protein alpha-synuclein," *ChemBioChem*, vol. 13, pp. 761–768, 2012.
- [163] V. N. Uversky, "A protein-chameleon: Conformational plasticity of  $\alpha$ -synuclein, a disordered protein involved in neurodegenerative disorders," *J. Biomol. Struct. Dyn.*, vol. 21, pp. 211–234, 2003.
- [164] M. Wells, H. Tidow, T. J. Rutherford, P. Markwick, M. Ringkjøbing Jensen, E. Mylonas, D. I. Svergun, M. Blackledge, and A. R. Fersht, "Structure of tumor suppressor P-53 and its intrinsically disordered N-terminal transactivation domain," *Proc. Nat. Acad. Sci. USA*, vol. 105, pp. 5762–5767, 2008.

- [165] K. Kwac and M. Cho, "Molecular dynamics simulation study of N-methylacetamide in water. i. amide I mode frequency fluctuation," *J. Chem. Phys.*, vol. 119, pp. 2247–2255, 2003.
- [166] S. Krimm and Y. Abe, "Intermolecular interaction effects in the amide I vibrations of  $\beta$  polypeptides," *Proc. Nat. Acad. Sci. USA*, vol. 69, pp. 2788–2792, 1972.
- [167] M. Lima, R. Chelli, V. V. Volkov, and R. Righini, "Two-dimensional infrared spectroscopy of a structured liquid: neat formamide," *J. Chem. Phys.*, vol. 130, p. 204518, 2009.
- [168] P. Bour and T. Keiderling, "Empirical modeling of the peptide amide I band IR intensity in water solution," *J. Chem. Phys.*, vol. 119, pp. 11253–11262, 2003.
- [169] A. V. Cunha, A. S. Bondarenko, and T. L. C. Jansen, "Assessing spectral simulation protocols for the amide I band of proteins," *J. Chem. Theory Comput.*, vol. 12, pp. 3982–3992, 2016.
- [170] T. Hayashi, W. Zhuang, and S. Mukamel, "Electrostatic DFT map for the complete vibrational amide band of NMA," *J. Phys. Chem. A*, vol. 109, pp. 9747–9759, 2005.
- [171] J. Jeon and M. Cho, "Direct quantum mechanical/molecular mechanical simulations of two-dimensional vibrational responses: N-methylacetamide in water," *New. J. Phys.*, vol. 12, p. 065001, 2010.
- [172] C. Liang and T. L. C. Jansen, "An efficient  $N^3$ -scaling propagation scheme for simulating two-dimensional infrared and visible spectra," *J. Chem. Theory Comput.*, vol. 8, p. 1706, 2012.
- [173] Y. S. Lin, J. M. Shorb, P. Mukherjee, M. T. Zanni, and J. L. Skinner, "Empirical amide I vibrational frequency map: Application to 2D IR line shapes for isotope-edited membrane peptide bundles," *J. Phys. Chem. B*, vol. 113, pp. 592–602, 2009.
- [174] H. Maekawa, C. Toniolo, A. Moretto, Q. B. Broxterman, and N.-H. Ge, "Different spectral signatures of octapeptide 310- and  $\alpha$ -helices revealed by two-dimensional infrared spectroscopy," *J. Phys. Chem. B*, vol. 110, pp. 5834–5837, 2006.

- [175] E. Malolepsza and J. E. Straub, “Empirical maps for the calculation of amide I vibrational spectra of proteins from classical molecular dynamics simulations,” *J. Phys. Chem. B.*, vol. 118, pp. 7848–7855, 2014.
- [176] M. Reppert, A. R. Roy, and A. Tokmakoff, “Isotope-enriched protein standards for computational amide I spectroscopy,” *J. Chem. Phys.*, vol. 142, p. 125104, 2015.
- [177] K. Shinokita, A. V. Cunha, T. L. C. Jansen, and M. S. Pshenichnikov, “Hydrogen bond dynamics in bulk alcohols,” *J. Chem. Phys.*, vol. 142, p. 212450, 2015.
- [178] M. F. DeCamp, L. DeFlores, J. M. McCracken, A. Tokmakoff, K. Kwac, and M. Cho, “Amide I vibrational dynamics of N-methylacetamide in polar solvents: The role of electrostatic interactions,” *J. Phys. Chem. B.*, vol. 109, pp. 11016–11026, 2005.
- [179] H. Torri, “Vibrational interactions in the amide I subspace of the oligomers and hydration clusters of n-methylacetamide,” *J. Phys. Chem. A.*, vol. 108, pp. 7272–7280, 2004.
- [180] H. Torri, “Effects of intermolecular vibrational coupling and liquid dynamics on the polarized raman and two-dimensional infrared spectral profiles of liquid N,N-Dimethylformamide analyzed with a time-domain computational method,” *J. Phys. Chem. A.*, vol. 110, pp. 4822–4832, 2006.
- [181] W. L. Jorgensen, A. D. S. Maxwell, and J. Tirado-Rives, “Development and testing of the OPLS All-atom force field on conformational energetics and properties of organic liquids,” *J. Am. Chem. Soc.*, vol. 118, p. 11225, 1996.
- [182] H. J. C. Berendsen, J. P. M. Postma, W. F. van Gunsteren, A. DiNola, and J. R. Haak, “Molecular dynamics with coupling to an external bath,” *J. Chem. Phys.*, vol. 81, p. 3684, 1984.
- [183] C. Caleman, P. J. van Maaren, M. Hong, J. S. Hub, L. T. Costa, and D. van der Spoel, “Force field benchmark of organic liquids: Density, enthalpy of vaporization, heat capacities, surface tension, isothermal compressibility, volumetric expansion coefficient, and dielectric constant,” *J. Chem. Theory Comput.*, vol. 8, pp. 61–74, 2012.

- [184] D. J. Tobias, C. L. Brooks, and S. H. Fleischman, "Conformational flexibility in free energy simulations," *Chem. Phys. Lett.*, vol. 156, pp. 256–260, 1989.
- [185] B. Hess, H. Bekker, H. J. C. Berendsen, and J. G. E. M. Fraaije, "LINCS: A linear constraint solver for molecular simulations," *J. Comput. Chem.*, vol. 18, p. 1463, 1998.
- [186] H. Torii and M. Tasumi, "Model calculations on the amide-I infrared bands of globular proteins," *J. Chem. Phys.*, vol. 96, pp. 3379–3387, 1992.
- [187] T. L. C. Jansen, A. G. Dijkstra, T. M. Watson, J. D. Hirst, and J. Knoester, "Modeling the amide I bands of small peptides," *J. Chem. Phys.*, vol. 125, p. 044312, 2006.
- [188] W. M. Haynes, *Handbook of chemistry and physics: a ready-reference book of chemical and physical data*. CRC Press, 18th ed., 2009.
- [189] J. Helbing and P. Hamm, "Compact implementation of fourier transform two-dimensional IR spectroscopy without phase ambiguity," *J. Opt. Soc. Am. B*, vol. 28, p. 171, 2011.
- [190] R. Bloem, S. Garrett-Roe, H. Strzalka, P. Hamm, and P. Donaldson, "Enhancing signal detection and completely eliminating scattering using quasi-phase-cycling in 2D IR experiments," *Opt. Express*, vol. 18, pp. 27067–27078, 2010.
- [191] E. Backus, S. Garret-Roe, and P. Hamm, "Phasing problem of heterodyne-detected two-dimensional infrared spectroscopy," *Opt. Lett.*, vol. 33, pp. 2665–2667, 2008.
- [192] A. Luzar and D. Chandler, "Hydrogen-bond kinetics in liquid water," *Nature*, vol. 379, pp. 55–57, 1996.
- [193] T. L. C. Jansen, B. M. Auer, M. Yang, and J. L. Skinner, "Two-dimensional infrared spectroscopy and ultrafast anisotropy decay of water," *J. Chem. Phys.*, vol. 132, p. 224503, 2010.
- [194] S. Woutersen and H. J. Bakker, "Resonant intermolecular transfer of vibrational energy in liquid water," *Nature*, vol. 402, pp. 507–509, 1999.

- [195] M. R. Panman, D. J. Shaw, B. Ensing, and S. Woutersen, “Local orientational order in liquids revealed by resonant vibrational energy transfer,” *Phys. Rev. Lett.*, vol. 113, p. 207801, 2014.
- [196] D. J. Thouless, “Electrons in disordered systems and the theory of localization,” *Phys. Rep.*, vol. 13, pp. 93–142, 1974.
- [197] J.-H. Choi, S. Ham, and M. Cho, “Inter-peptide interaction and delocalization of amide I vibrational excitons in myoglobin and flavodoxin,” *J. Chem. Phys.*, vol. 117, pp. 6821–6832, 2002.
- [198] B. C. Starcher, “Elastin and the lung,,” *Thorax*, vol. 41, pp. 577–585, 1986.
- [199] J. A. Foster and S. W. Curtiss, “The regulation of lung elastin synthesis,” *Am. J. Physiol.*, vol. 259, pp. L13–L23, 1990.
- [200] B. C. Starcher, “Lung elastin and matrix,” *Chest*, vol. 117, pp. 229S–234S, 2000.
- [201] L. Robert, M. P. Jacob, C. Frances, G. Godeau, and W. Hornebeck, “Interaction between elastin and elastases and its role in the aging of the arterial wall, skin and other connective tissues. a review,” *Mech. Ageing Dev.*, vol. 28, pp. 155–166, 1984.
- [202] H. Oxlund, J. Manschot, and A. Viidik, “The role of elastin in the mechanical properties of skin,” *J. Biomech.*, vol. 21, pp. 213–218, 1988.
- [203] C. Frances and L. Robert, “Elastin and elastic fibers in normal and pathologic skin,” *Int. J. Dermatol.*, vol. 23, pp. 166–179, 1984.
- [204] A. Tsamis, J. Krawiec, and D. Vorp, “Elastin and collagen fibre microstructure of the human aorta in aging and disease: A review,” *J. Roy. Soc. Interface*, vol. 10, p. 20121004, 2013.
- [205] A. Patel, B. Fine, M. Sandig, and K. Mequanint, “Elastin biosynthesis: the missing link in tissue-engineered blood vessels,” *Cardiovasc. Res.*, vol. 71, pp. 40–49, 2006.
- [206] L. Debelle and A. M. Tamburro, “Elastin: molecular description and function,” *Int. J. Biochem. Cell Biol.*, vol. 31, pp. 261–272, 1999.

- [207] D. W. Urry, M. M. Long, B. A. Cox, T. Ohnishi, L. W. Mitchell, and M. Jacobs, "The synthetic polypentapeptide of elastin coacervates and forms filamentous aggregates," *Biochim. Biophys. Acta*, vol. 371, pp. 597–602, 1974.
- [208] R. Herald, R. C. Anthony, and R. R. Anthony, "Short elastin-like peptides exhibit the same temperature-induced structural transitions as elastin polymers: implications for protein engineering," *J. Mol. Biol.*, vol. 283, pp. 255–264, 1998.
- [209] C. Bellingham, M. Lillie, J. Gosline, G. Wright, B. Starcher, A. Bailey, K. Woodhouse, and F. W. Keeley, "Recombinant human elastin polypeptides self-assemble into biomaterials with elastin-like properties," *Biopolymers*, vol. 70, pp. 445–455, 2003.
- [210] L. N. Dana, C. Ashutosh, and A. S. Lori, "Applications of elastin-like polypeptides in tissue engineering," *Adv. Drug Deliv. Rev.*, vol. 62, pp. 1479–1485, 2010.
- [211] W. F. Daamen, J. H. Veerkamp, J. C. M. van Hest, and T. H. van Kuppevelt, "Elastin as a biomaterial for tissue engineering," *Biomaterials*, vol. 28, pp. 4378–4398, 2007.
- [212] I. Massodi, G. L. Bidwell, and D. Raucher, "Evaluation of cell penetrating peptides fused to elastin-like polypeptide for drug delivery," *J. Control. Release*, vol. 108, pp. 396–408, 2005.
- [213] S. Dirk, "Thermo- and pH-responsive polymers in drug delivery," *Adv. Drug Deliv. Rev.*, vol. 58, pp. 1655–1670, 2006.
- [214] M. Khaled and D. Urry, "Nuclear Overhauser enhancement demonstration of the type II  $\beta$ -turn in repeat peptides of tropoelastin," *Biochem. Biophys. Res. Commun.*, vol. 70, pp. 485–491, 1976.
- [215] C. M. Venkatachalam and D. W. Urry, "Development of a linear helical conformation from its cyclic correlate.  $\beta$ -spiral model of the elastin poly (pentapeptide)(VPGVG)<sub>n</sub>," *Macromolecules*, vol. 14, pp. 1225–1229, 1981.
- [216] D. W. Urry, "Entropic elastic processes in protein mechanisms. I. Elastic structure due to an inverse temperature transition and elasticity due to internal chain dynamics," *J. Protein Chem.*, vol. 7, pp. 1–34, 1988.

- [217] L. Bin, O. V. A. Darwin, and D. Valerie, "The molecular basis for the inverse temperature transition of elastin," *J. Mol. Biol.*, vol. 305, pp. 581–592, 2001.
- [218] A. Krukau, I. Brovchenko, and A. Geiger, "Temperature-induced conformational transition of a model elastin-like peptide GVG (VPGVG)(3) in water," *Biomacromolecules*, vol. 8, pp. 2196–2202, 2007.
- [219] C. Nicolini, R. Ravindra, B. Ludolph, and R. Winter, "Characterization of the temperature- and pressure-induced inverse and reentrant transition of the minimum elastin-like polypeptide GVG(VPGVG) by DSC, PPC, CD, and FT-IR spectroscopy," *Biophys. J.*, vol. 86, pp. 1385–1392, Mar. 2004.
- [220] V. Serrano, W. Liu, and S. Franzen, "An infrared spectroscopic study of the conformational transition of elastin-like polypeptides," *Biophys. J.*, vol. 93, pp. 2429–2435, 2007.
- [221] Y. Cho, L. B. Sagle, S. Imura, Y. Zhang, J. Kherb, J. M. Chilkoti, A. and Scholtz, and P. S. Cremer, "Hydrogen bonding of  $\beta$ -turn structure is stabilized in D<sub>2</sub>O," *J. Am. Chem. Soc.*, vol. 131, pp. 15188–15193, 2009.
- [222] Z. Ahmed, J. P. Scaffidi, and S. A. Asher, "Circular dichroism and UV-resonance raman investigation of the temperature dependence of the conformations of linear and cyclic elastin," *Biopolymers*, vol. 91, pp. 52–60, 2009.
- [223] G. Thomas, B. Prescott, and D. Urry, "Raman amide bands of type-II  $\beta$ -turns in cyclo-(VPGVG) 3 and poly-(VPGVG), and implications for protein secondary-structure analysis," *Biopolymers*, vol. 26, pp. 921–934, 1987.
- [224] D. Kurková, J. Kríz, P. Schmidt, J. Dybal, J. C. Rodríguez-Cabello, and M. Alonso, "Structure and dynamics of two elastin-like polypentapeptides studied by NMR spectroscopy," *Biomacromolecules*, vol. 4, pp. 589–601, 2003.
- [225] R. Glaves, M. Baer, E. Schreiner, R. Stoll, and D. Marx, "Conformational dynamics of minimal elastin-like polypeptides: The role of proline revealed by molecular dynamics and nuclear magnetic resonance," *Chem. Phys. Chem.*, vol. 9, 2008.

- [226] N. Harald and H. A. Klok, "Secondary structure formation and LCst behavior of short elastin-like peptides," *Biomacromol.*, vol. 9, pp. 2755–2763, 2008.
- [227] R. Rousseau, E. Schreiner, A. Kohlmeyer, and D. Marx, "Temperature-dependent conformational transitions and hydrogen-bond dynamics of the elastin-like octapeptide GVG (VPGVG): a molecular-dynamics study," *Biophys. J.*, vol. 86, pp. 1393–1407, 2004.
- [228] M. Reppert, S. Roy, J. O. B. Tempkin, A. R. Dinner, and A. Tokmakoff, "Refining disordered peptide ensembles with computational amide I spectroscopy: Application to elastin-like peptides," *J. Phys. Chem. B*, 2016.
- [229] M. Buck, "Trifluoroethanol and colleagues: cosolvents come of age. Recent studies with peptides and proteins," *Q. Rev. Biophys.*, vol. 31, pp. 297–355, 1998.
- [230] D. Roccatano, G. Colombo, M. Fioroni, and A. E. Mark, "Mechanism by which 2,2,2-trifluoroethanol/water mixtures stabilize secondary-structure formation in peptides: A molecular dynamics study," *P. Natl. Acad. Sci. USA*, vol. 99, pp. 12179–12184, 2002.
- [231] R. L. M. Teeuwen, H. Zuilhof, F. A. de Wolf, and J. C. M. van Hest, "Temperature-controlled positioning of fusion proteins in microreactors," *Soft Matter*, vol. 5, pp. 2261–2268, 2009.
- [232] D. Meyer and A. Chilkoti, "Purification of recombinant proteins by fusion with thermally-responsive polypeptides," *Nat. Biotechnol.*, vol. 17, pp. 1112–1115, 1999.
- [233] V. V. Andrushchenko, H. J. Vogel, and E. J. Prenner, "Optimization of the hydrochloric acid concentration used for trifluoroacetate removal from synthetic peptides," *J. Pept. Sci.*, vol. 13, pp. 37–43, 2007.
- [234] M. D. Hanwell, D. E. Curtis, D. C. Lonie, T. Vandermeersch, E. Zurek, and G. R. Hutchison, "Avogadro: an advanced semantic chemical editor, visualization, and analysis platform," *J. Cheminf.*, vol. 4, p. 1, 2012.
- [235] D. Van der Spoel, E. Lindahl, B. Hess, G. Groenhof, A. Mark, and H. Berendsen, "Gromacs: fast, flexible, and free," *J. Comput. Chem.*, vol. 26, pp. 1701–1718, 2005.



- [236] M. Parrinello and A. Rahman, "Polymorphic transitions in single crystals: A new molecular dynamics method," *J. Appl. Phys.*, vol. 52, pp. 7182–7190, 1981.
- [237] G. Bussi, D. Donadio, and M. Parrinello, "Canonical sampling through velocity rescaling," *J. Chem. Phys.*, vol. 126, p. 014101, 2007.
- [238] C. L. Brooks, "Computer simulation of liquids," *J. Solution Chem.*, vol. 18, pp. 99–99, 1989.
- [239] T. Darden, D. York, and L. Pedersen, "Particle mesh Ewald: An  $N \cdot \log(N)$  method for Ewald sums in large systems," *J. Chem. Phys.*, vol. 98, pp. 10089–10092, 1993.
- [240] A. V. Cunha, A. S. Bondarenko, and T. L. C. Jansen, "Assessing spectral simulation protocols for the amide I band of proteins," *J. Chem. Theory Comput.*, vol. 12, pp. 3982–3992, 2016.
- [241] Q. Guo, P. Pagano, Y. Li, A. Kohen, and C. Cheatum, "Line shape analysis of two-dimensional infrared spectra," *J. Chem. Phys.*, vol. 142, 2015.
- [242] S. Roy, T. L. C. Jansen, and J. Knoester, "Structural classification of the amide I sites of a [small beta]-hairpin with isotope label 2D IR spectroscopy," *Phys. Chem. Chem. Phys.*, vol. 12, pp. 9347–9357, 2010.
- [243] Y. S. Kim, L. Liu, P. H. Axelsen, and R. M. Hochstrasser, "2D IR provides evidence for mobile water molecules in  $\beta$ -amyloid fibrils," *P. Natl. Acad. Sci. USA*, vol. 106, pp. 17751–17756, 2009.
- [244] E. Vass, M. Kurz, R. K. Konat, and M. Hollósi, "FTIR and CD spectroscopic studies on cyclic penta- and hexa-peptides. detailed examination of hydrogen bonding in  $\beta$ - and  $\gamma$ -turns determined by NMR," *Spectrochim. Acta Mol. Biomol.*, vol. 54, pp. 773–786, 1998.
- [245] O. Satoshi, K. Akiko, and K. Shohei, "Solution conformation and amyloid-like fibril formation of a polar peptide derived from a  $\beta$ -hairpin in the OspA single-layer  $\beta$ -sheet1," *J. Mol. Biol.*, vol. 301, pp. 477–489, 2000.
- [246] M. Ramirez-Alvarado, L. Serrano, and F. Blanco, "Conformational analysis of peptides corresponding to all the secondary structure elements of protein L B1 domain: Secondary structure propensities are not conserved in proteins with the same fold," *Protein Sci.*, vol. 6, pp. 162–174, 1997.

- [247] M. Jackson and H. H. Mantsch, "The use and misuse of FTIR spectroscopy in the determination of protein structure," *Crit. Rev. Biochem. Mol. Biol.*, vol. 30, pp. 95–120, 1995.
- [248] F. Demmel, W. Doster, W. Petry, and A. Schulte, "Vibrational frequency shifts as a probe of hydrogen bonds: thermal expansion and glass transition of myoglobin in mixed solvents," *Eur. Biophys. J.*, vol. 26, pp. 327–335, 1997.
- [249] A. Barth, "Infrared spectroscopy of proteins," *Biochim. Biophys. Acta*, vol. 1767, pp. 1073–1101, 2007.
- [250] H. J. Bakker, P. C. M. Planken, and A. Lagendijk, "Role of solvent on vibrational energy transfer in solution," *Nature*, vol. 347, pp. 745–747, 1990.
- [251] V. M. Kenkre, A. Tokmakoff, and M. D. Fayer, "Theory of vibrational relaxation of polyatomic molecules in liquids," *J. Chem. Phys.*, vol. 101, pp. 10618–10629, 1994.
- [252] S. Woutersen, Y. Mu, G. Stock, and P. Hamm, "Subpicosecond conformational dynamics of small peptides probed by two-dimensional vibrational spectroscopy," *P. Natl. Acad. Sci. USA*, vol. 98, pp. 11254–11258, 2001.
- [253] M. Fioroni, M. D. Diaz, K. Burger, and S. Berger, "Solvation phenomena of a tetrapeptide in water/trifluoroethanol and water/ethanol mixtures: a diffusion NMR, intermolecular NOE, and molecular dynamics study," *J. Am. Chem. Soc.*, vol. 124, pp. 7737–7744, 2002.
- [254] A. Yang, B. Hitz, and B. Honig, "Free energy determinants of secondary structure formation: III.  $\beta$ -turns and their role in protein folding," *J. Mol. Biol.*, vol. 259, pp. 873–882, 1996.
- [255] E. Hutchinson and J. Thornton, "A revised set of potentials for  $\beta$ -turn formation in proteins," *Protein Sci.*, vol. 3, pp. 2207–2216, 1994.
- [256] B. Li, D. O. V. Alonso, B. J. Bennion, and V. Daggett, "Hydrophobic hydration is an important source of elasticity in elastin-based biopolymers," *J. Am. Chem. Soc.*, vol. 123, pp. 11991–11998, 2001.

- [257] W. J. Cook, H. Einspahr, T. L. Trapane, D. W. Urry, and C. E. Bugg, "Crystal structure and conformation of the cyclic trimer of a repeat pentapeptide of elastin, cyclo-(L-valyl-L-prolylglycyl-L-valylglycyl)<sub>3</sub>," *J. Am. Chem. Soc.*, vol. 102, pp. 5502–5505, 1980.
- [258] T. Yamaoka, T. Tamura, T. Seto, Y. and Tada, S. Kunugi, and D. Tirrell, "Mechanism for the phase transition of a genetically engineered elastin model peptide (VPGIG)<sub>40</sub> in aqueous solution," *Biomacromolecules*, vol. 4, pp. 1680–1685, 2003.
- [259] D. W. Urry, R. G. Shaw, and K. U. Prasad, "Polypentapeptide of elastin: temperature dependence of ellipticity and correlation with elastomeric force," *Biochem. Biophys. Res. Commun.*, vol. 130, pp. 50–57, 1985.
- [260] M. Cho, *Two-Dimensional Optical Spectroscopy*. CRC Press, 1 ed., 2009.
- [261] D. Laage and J. T. Hynes, "A molecular jump mechanism of water reorientation," *Science*, vol. 311, p. 832, 2006.
- [262] S. T. Roberts, K. Ramasesha, and A. Tokmakoff, "Structural rearrangements in water viewed through two-dimensional infrared spectroscopy," *Acc. Chem. Res.*, vol. 42, p. 1239, 2009.
- [263] Y. Umena, K. Kawakami, J.-R. Shen, and N. Kamiya, "Crystal structure of oxygen-evolving photosystem II at a resolution of 1.9 Å," *Nature*, vol. 473, p. 55, 2012.
- [264] T. Sun, "An antifreeze protein folds with an interior network of more than 400 semi-clathrate waters," *Science*, vol. 343, p. 795, 2014.
- [265] S. K. Pal, J. Peon, and A. H. Zewail, "Biological water at the protein surface: Dynamical solvation probed directly with femtosecond resolution," *P. Natl. Acad. Sci. USA*, vol. 99, p. 1763, 2002.
- [266] M. Chaplin, "Do we underestimate the importance of water in cell biology?," *Nat. Rev. Mol. Cell Biol.*, vol. 7, p. 861, 2006.
- [267] J. Israelachvili and H. Wennerström, "Role of hydration and water structure in biological and colloidal interactions," *Nature*, vol. 379, p. 219, 1996.

- [268] A. A. Bakulin, M. S. Pshenichnikov, H. J. Bakker, and C. Petersen, "Hydrophobic molecules slow down the hydrogen-bond dynamics of water," *J. Phys. Chem. A*, vol. 115, p. 1821, 2011.
- [269] D. Laage, G. Stirnemann, and J. T. Hynes, "Why water reorientation slows without iceberg formation around hydrophobic solutes," *J. Phys. Chem. B*, vol. 113, p. 2428, 2009.
- [270] Y. Rezus and H. Bakker, "Observation of immobilized water molecules around hydrophobic groups," *Phys. Rev. Lett.*, vol. 99, p. 148301, 2007.
- [271] K. J. Gaffney, P. H. Davis, I. R. Piletic, N. E. Levinger, and M. D. Fayer, "Hydrogen bond dissociation and reformation in methanol oligomers following hydroxyl stretch relaxation," *J. Phys. Chem. A*, vol. 106, p. 12012, 2002.
- [272] R. Laenen, C. Rauscher, and A. Laubereau, "Transient hole burning in the infrared in an ethanol solution," *J. Phys. Chem. A*, vol. 101, p. 3201, 1997.
- [273] R. Laenen and K. Simeonidis, "Energy relaxation and reorientation of the OH mode of simple alcohol molecules in different solvents monitored by transient IR spectroscopy," *Chem. Phys. Lett.*, vol. 299, p. 589, 1999.
- [274] R. Laenen and C. Rauscher, "Time-resolved infrared spectroscopy of ethanol monomers in liquid solution: molecular reorientation and energy relaxation times," *Chem. Phys. Lett.*, vol. 274, p. 63, 1997.
- [275] L. K. Iwaki and D. D. Dlott, "Three-dimensional spectroscopy of vibrational energy relaxation in liquid methanol," *J. Phys. Chem. A*, vol. 104, p. 9101, 2000.
- [276] K. Kwac and E. Geva, "A mixed quantum-classical molecular dynamics study of the hydroxyl stretch in methanol/carbon tetrachloride mixtures III: Nonequilibrium hydrogen-bond dynamics and infrared pump-probe spectra," *J. Phys. Chem. B*, vol. 117, p. 7737, 2013.
- [277] S. Woutersen, U. Emmerichs, and H. J. Bakker, "Femtosecond mid-IR pump-probe spectroscopy of liquid water: Evidence for a two-component structure," *Science*, vol. 278, p. 658, 1997.

- [278] U. Essmann, L. Perera, M. L. Berkowitz, T. Darden, H. Lee, and L. G. Pedersen, "A smooth particle mesh Ewald method," *J. Chem. Phys.*, vol. 103, p. 8577, 1995.
- [279] B. Auer, R. Kumar, J. Schmidt, and J. L. Skinner, "Hydrogen bonding and Raman, IR, and 2D IR spectroscopy of dilute HOD in liquid D<sub>2</sub>O.," *Proc. Natl. Acad. Sci. U.S.A.*, vol. 104, pp. 14215–14220, 2007.
- [280] C. P. van der Vegte, A. G. Dijkstra, J. Knoester, and T. L. C. Jansen, "Calculating two-dimensional spectra with the mixed quantum-classical Ehrenfest method," *J. Phys. Chem. A*, vol. 117, p. 5970, 2013.
- [281] F. Li and J. L. Skinner, "Infrared and Raman line shapes for ice Ih. I. Dilute HOD in H<sub>2</sub>O and D<sub>2</sub>O," *J. Chem. Phys.*, vol. 132, p. 204505, 2010.
- [282] T. L. C. Jansen, T. Hayashi, W. Zhuang, and S. Mukamel, "Stochastic Liouville equations for hydrogen-bonding fluctuations and their signatures in two-dimensional vibrational spectroscopy of water," *J. Chem. Phys.*, vol. 123, p. 114504, 2005.
- [283] W. L. Jorgensen and J. Tirado-Rives, "The OPLS Potential Functions for Proteins. Energy Minimizations for Crystals of Cyclic Peptides and Crambin," *J. Am. Chem. Soc.*, vol. 110, pp. 1657–1666, 1988.
- [284] O. Kristiansson, "Investigation of the OH stretching vibration of CD<sub>3</sub>OH in CCl<sub>4</sub>," *J. Mol. Struct.*, vol. 477, p. 105, 1999.
- [285] S. S. Farwanah, J. Yarwood, I. Cabaço, and M. Besnard, "Infrared studies of hydrogen-bonding of methanol in binary mixtures with acetonitrile," *J. Mol. Liq.*, vol. 56, p. 317, 1993.
- [286] R. Kumar, J. R. Schmidt, and J. L. Skinner, "Hydrogen bonding definitions and dynamics in liquid water," *J. Chem. Phys.*, vol. 126, p. 204107, 2007.
- [287] J. R. Schmidt, S. A. Corcelli, and J. L. Skinner, "Pronounced non-condon effects in the ultrafast infrared spectroscopy of water," *J. Chem. Phys.*, vol. 123, p. 044513, 2005.
- [288] C. P. Lawrence and J. L. Skinner, "Vibrational spectroscopy of HOD in liquid D<sub>2</sub>O. II. Infrared line shapes and vibrational stokes shift," *J. Chem. Phys.*, vol. 117, p. 8847, 2002.

- [289] T. Steinel, J. B. Asbury, S. A. Corcelli, C. P. Lawrence, J. Skinner, and M. Fayer, "Water dynamics: dependence on local structure probed with vibrational echo correlation spectroscopy," *Chem. Phys. Lett.*, vol. 386, p. 295, 2004.
- [290] K. B. Møller, R. Rey, and J. T. Hynes, "Hydrogen bond dynamics in water and ultrafast infrared spectroscopy: A theoretical study," *J. Phys. Chem. A*, vol. 108, p. 1275, 2004.
- [291] S. Yeremenko, M. S. Pshenichnikov, and D. Wiersma, "Interference effects in IR photon echo spectroscopy of liquid water," *Phys. Rev. A*, vol. 73, p. 021804, 2006.
- [292] J. J. Loparo, C. J. Fecko, J. D. Eaves, S. T. Roberts, and A. Tokmakoff, "Reorientational and configurational fluctuations in water observed on molecular length scales," *Phys. Rev. B*, vol. 70, p. 180201(R), 2004.
- [293] J. Lindner, P. Vöhringer, M. S. Pshenichnikov, D. Cringus, D. Wiersma, and M. Mostovoy, "Vibrational relaxation of pure liquid water," *Chem. Phys. Lett.*, vol. 421, p. 329, 2006.
- [294] J. Lindner, D. Cringus, M. S. Pshenichnikov, and P. Vöhringer, "Anharmonic bend stretch coupling in neat liquid water," *Chem. Phys.*, vol. 341, p. 326, 2007.
- [295] T. L. C. Jansen, D. Cringus, and M. S. Pshenichnikov, "Dissimilar dynamics of coupled water vibrations," *J. Phys. Chem. A*, vol. 113, p. 6260, 2009.
- [296] K. J. Gaffney, I. R. Piletic, and M. D. Fayer, "Orientational relaxation and vibrational excitation transfer in methanol-carbon tetrachloride solutions," *J. Chem. Phys.*, vol. 118, p. 2270, 2003.
- [297] D. Laage, G. Stirnemann, F. Sterpone, and J. T. Hynes, "Water jump reorientation: From theoretical prediction to experimental observation," *Acc. Chem. Res.*, vol. 45, p. 53, 2012.
- [298] D. Laage, G. Stirnemann, and J. T. Hynes, "Water jump reorientation and ultrafast vibrational spectroscopy," *J. Photochem. Photobiol., A*, vol. 234, p. 75, 2012.

- [299] Y. L. A. Rezus and H. J. Bakker, "On the orientational relaxation of HDO in liquid water," *J. Chem. Phys.*, vol. 123, p. 114502, 2005.
- [300] R. Ludwig, M. D. Zeidler, and T. C. Farrar, "Molecular dynamics in lower alcohols," *J. Phys. Chem.*, vol. 189, p. 19, 1995.
- [301] R. Ludwig and M. D. Zeidler, "NMR relaxation in ethanol and propanol and in their binary mixtures with carbon tetrachloride," *Mol. Phys.*, vol. 82, p. 313, 1994.
- [302] T. D. Ferris and T. C. Farrar, "The temperature dependence of the hydroxyl deuterium quadrupole coupling parameter and the rotational correlation time of the OD internuclear vector in neat ethanol-d<sub>1</sub>," *Mol. Phys.*, vol. 100, p. 303, 2002.
- [303] R. Buchner and J. Barthel, "Dynamics of methanol-tetrachloromethane mixtures - a dielectric relaxation study," *J. Mol. Liq.*, vol. 52, p. 131, 1992.
- [304] J. Barthel, K. Bachhuber, R. Buchner, J. B. Gill, and M. Kleebauer, "Dielectric spectra of some common solvents in the microwave region. Dipolar aprotic solvents and amides," *Chem. Phys. Lett.*, vol. 167, p. 62, 1990.
- [305] J. Hunger, A. Stoppa, A. Thoman, M. Walther, and R. Buchner, "Broad-band dielectric response of dichloromethane," *Chem. Phys. Lett.*, vol. 471, p. 85, 2009.
- [306] B. M. Ladanyi and M. S. Skaf, "Wave vector-dependent dielectric relaxation of methanol/water mixtures," *J. Phys. Chem.*, vol. 100, p. 1368, 1996.
- [307] A. A. Bakulin, D. Cringus, P. A. Pieniazek, J. L. Skinner, T. L. C. Jansen, and M. S. Pshenichnikov, "Dynamics of water confined in reversed micelles: Multidimensional vibrational spectroscopy study," *J. Phys. Chem. B*, vol. 117, p. 15545, 2013.
- [308] L. A. Woolf, "Insights into solute-solute-solvent interactions from transport property measurements with particular reference to methanol-water mixtures and their constituents," *Pure Appl. Chem.*, vol. 57, p. 1083, 1985.

- [309] I. Chang and H. Sillescu, "Heterogeneity at the glass transition: Translational and rotational self-diffusion," *J. Phys. Chem. B*, vol. 101, p. 8794, 1997.
- [310] C. Lederle, W. Hiller, C. Gainaru, and R. Böhmer, "Diluting the hydrogen bonds in viscous solutions of n-butanol with n-bromobutane: II. A comparison of rotational and translational motions," *J. Chem. Phys.*, vol. 134, p. 064512, 2011.
- [311] "Definition of the hydrogen bond (IUPAC Recommendations 2011)," *Pure Appl. Chem.*, vol. 83, pp. 1637–1641, 2011.
- [312] P. Schuster, G. Zundel, and C. Sandorfy, *The Hydrogen Bond - Recent Developments in Theory and Experiment*. Amsterdam: North-Holland, 1976.
- [313] M. Wahl, "C-H...O hydrogen bonding in biology," *Trend. Biochem. Sci.*, vol. 22, pp. 97–102, 1997.
- [314] M. Cho, "Coherent Two-Dimensional Optical Spectroscopy," *Chem. Rev.*, vol. 108, pp. 1331–1418, 2008.
- [315] S. Ashihara, N. Huse, A. Espagne, E. T. J. Nibbering, and T. Elsaesser, "Ultrafast Structural Dynamics of Water Induced by Dissipation of Vibrational Energy," *J. Phys. Chem.*, vol. 111, pp. 743–746, 2007.
- [316] A. J. Lock and H. J. Bakker, "Temperature dependence of vibrational relaxation in liquid Temperature dependence of vibrational relaxation in liquid H<sub>2</sub>O," *J. Chem. Phys.*, vol. 117, pp. 1708–1713, 2002.
- [317] D. Cringus, A. Bakulin, J. Lindner, P. Vöhringer, M. Pshenichnikov, and D. Wiersma, "Ultrafast Energy Transfer in Water - AOT Reverse Micelles," *J. Phys. Chem. B*, vol. 111, pp. 14193–14207, 2007.
- [318] D. Cringus, J. Lindner, M. Milder, M. Pshenichnikov, P. Vöhringer, and D. Wiersma, "Femtosecond water dynamics in reverse-micellar nanodroplets," *Chem. Phys. Lett.*, vol. 408, pp. 162–168, 2005.
- [319] L. J. Bellamy and R. J. Pace, "The nature of the hydrogen bond in alcohol: dimers and polymers," *Spectrochimica Acta*, vol. 22, pp. 525–533, 1966.



- [320] J. Asbury, T. Steinel, and M. Fayer, "Hydrogen bond networks: Structure and evolution after hydrogen bond breaking," *J. Phys. Chem. B*, vol. 108, pp. 6544–6554, 2004.
- [321] T. S. Gulmen and E. L. Sibert III, "Vibrational energy relaxation of the OH(D) stretch fundamental of methanol in carbon tetrachloride," *J. Chem. Phys.*, vol. 123, p. 204508, 2005.
- [322] K. Kwac and E. Geva, "A mixed quantum-classical molecular dynamics study of the hydroxyl stretch in methanol/carbon tetrachloride mixtures III: Nonequilibrium hydrogen-bond dynamics and infrared pump-probe spectra," *J. Phys. Chem. B.*, vol. 117, pp. 7737–7749, 2013.
- [323] D. S. Kuen and K. Feierabend, "Cavity-enhanced overtone spectroscopy of methanol in aprotic solvents: Probing solute-solvent interactions and self-associative behavior," *J. Phys. Chem. A*, vol. 118, pp. 2942–2951, 2014.
- [324] I. Piletic, K. Gaffney, and M. Fayer, "Structural dynamics of hydrogen bonded methanol oligomers: Vibrational transient hole burning studies of spectral diffusion," *J. Chem. Phys.*, vol. 119, pp. 423–434, 2003.
- [325] A. Yokozeki, D. J. Kasprzak, and M. B. Shiflett, "IR-spectroscopic studies of hydrogen-bonding solutions: Lineshape analysis of ethanol + hexane system," *Applied Energy*, vol. 84, pp. 863–873, 2007.
- [326] N. Levinger, P. Davis, and M. Fayer, "Vibrational relaxation of the free terminal hydroxyl stretch in methanol oligomers: Indirect pathway to hydrogen bond breaking," *J. Chem. Phys.*, vol. 115, pp. 9352–9360, 2001.
- [327] M. Matsumoto and K. Gubbins, "Hydrogen bonding in liquid methanol," *J. Chem. Phys.*, vol. 93, pp. 1981–1994, 1990.
- [328] K. Mizuno, Y. Miyashita, Y. Shindo, and H. Ogawa, "NMR and FT-IR Studies of Hydrogen Bonds in Ethanol-Water Mixtures," *J. Phys. Chem.*, vol. 99, pp. 3225–3228, 1995.
- [329] N. Nishi, S. Takahashi, M. Matsumoto, A. Tanaka, K. Muraya, T. Takamuku, and T. Yamaguchi, "Hydrogen Bonding Cluster Formation and Hydrophobic Solute Association in Aqueous Solution of Ethanol," *J. Phys. Chem.*, vol. 99, pp. 462–468, 1995.

- [330] M. Pagliai, G. Cardini, R. Righini, and V. Schettino, "Hydrogen bond dynamics in liquid methanol," *J. Chem. Phys.*, vol. 119, pp. 6655–6662, 2003.
- [331] D. J. Shaw, M. R. Panman, and S. Woutersen, "Evidence for Cooperative Vibrational Relaxation of the NH-, OH-, and OD-Stretching Modes in Hydrogen-Bonded Liquids Using Infrared Pump-Probe Spectroscopy," *Phys. Rev. Lett.*, vol. 103, p. 227401, 2009.
- [332] A. A. Vartia, K. R. Mitchell-Koch, G. Stirnemann, D. Laage, and W. H. Thompson, "On the reorientation and hydrogen-bond dynamics of Alcohols," *J. Phys. Chem.*, vol. 115, pp. 12173–12178, 2011.
- [333] D. S. Venables and C. A. Schmuttenmaer, "Spectroscopy and dynamics of mixtures of water with acetone, acetonitrile, and methanol," *J. Chem. Phys.*, vol. 113, pp. 11222–11236, 2000.
- [334] K. Mazur, M. Bonn, and J. Hunger, "Hydrogen bond dynamics in primary alcohols: A femtosecond infrared study," *J. Phys. Chem. B.*, vol. 119, pp. 1558–1566, 2015.
- [335] O. Mesele, A. Vartia, D. Laage, and W. Thompson, "Reorientation of Isomeric Butanols: The Multiple Effects of Steric Bulk Arrangement on Hydrogen-Bond Dynamics," *J. Phys. Chem.*, vol. 120, pp. 1546–1559, 2016.
- [336] R. Ludwig, M. D. Zeidler, and T. C. Farrar, "Molecular Dynamics in Lower Alcohols," *Z. Phys. Chem.*, vol. 189, pp. 19–27, 1995.
- [337] E. Kryachko and M. Nguyen, "Hydrogen bonding between phenol and acetonitrile," *J. Phys. Chem. A*, vol. 106, pp. 4267–4271, 2002.
- [338] S. Yeremenko, A. Baltuška, F. de Haan, M. S. Pshenichnikov, and D. A. Wiersma, "Frequency-resolved pump-probe characterization of femtosecond infrared pulses," *Opt. Lett.*, vol. 27, pp. 1171–1173, 2002.
- [339] J. Helbing and P. Hamm, "Compact implementation of Fourier transform two-dimensional IR spectroscopy without phase ambiguity," *J. Opt. Soc. Am. B*, vol. 28, pp. 171–178, 2011.
- [340] P. Hamm and M. Zanni, *Concepts and Methods of 2D Infrared Spectroscopy*. Cambridge: Cambridge University Press, 2011.

- [341] W. Humphrey, A. Dalke, and K. Schulten, "VMD: Visual molecular dynamics," *J. Mol. Graph.*, vol. 14, pp. 33–38, 1996.
- [342] W. P. de Boeij, M. S. Pshenichnikov, and D. A. Wiersma, "Ultrafast Solvation Dynamics Explored By Femtosecond Photon ECHO Spectroscopies," *Annu. Rev. Phys. Chem.*, vol. 49, pp. 99–123, 1998.
- [343] D. Cringus, S. Yeremenko, M. S. Pshenichnikov, and D. A. Wiersma, "Hydrogen Bonding and Vibrational Energy Relaxation in Water-Acetonitrile Mixtures," *J. Phys. Chem. B*, vol. 108, pp. 10376–10387, 2004.
- [344] A. Luzar and D. Chandler, "Hydrogen-bond kinetics in liquid water," *Nature*, vol. 379, pp. 55–57, 1996.