

## University of Groningen

### Bacterial transmission

Gusnaniar

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2017

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Gusnaniar (2017). *Bacterial transmission*. [Thesis fully internal (DIV), University of Groningen]. Rijksuniversiteit Groningen.

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# chapter ONE



**General introduction**


Advances in Colloid and Interface Science, 2017

(submitted)



## Bacterial Transmission

The biofilm mode of growth is greatly preferred by most bacterial strains and species [1,2]. The sequence of events leading to biofilm formation is generally considered to commence with bacterial transport by convective-diffusion from a liquid suspension of planktonic bacteria to a substratum surface or impingement from aerosols (see **Figure 1**) [3].



**Figure 1.** Transport in bacterial adhesion to a substratum surface and biofilm growth.

- (a) Bacterial transport from a flowing suspension by convective-diffusion.
- (b) Bacterial transport by impingement from aerosols.
- (c) A multilayered biofilm resulting from growth of adhering bacteria.

Initially, bacterial adhesion is reversible, but production of extracellular polymeric substances (EPS) can rapidly lead to an irreversible state and subsequent growth of a bacterial (sub)monolayer into a multilayered biofilm. Transmission of sessile bacteria from one substrate to another is a less commonly highlighted means of bacterial transport, but equally if not more prevalent in many environments. Bacterial transmission frequently occurs in hospital environments and nursing homes among hands of healthcare workers [4] and patients [5,6]. Bacterial transmission


also occurs during insertion of indwelling urinary [7] or vascular catheters, either from the peri-urethral area or subcutaneous layers of the skin [8,9], respectively and similarly may occur towards endoscopes [10,11]. Transmission of microorganisms from contaminated lens cases to contact lenses followed by transmission to human cornea is a well-known cause of microbial keratitis, posing a general healthcare threat due to the large number of people wearing contact lenses [12,13]. Toothbrushes are mentioned more and more as a source for microbial transmission [14,15]. Also in the domestic environment transmission is inevitable, but usually involves less pathogenic microorganisms than present in biomedical environments [16,17]. Finally, bacterial transmission occurs in industrial environments, including slaughterhouses [18,19], agriculture [20,21], forest [22,23] and sea-water environments [24,25].

## **Mechanism of Bacterial Transmission**

Mechanistically different from bacterial adhesion, transmission involves adhesion of donor bacteria to a receiver surface and subsequent detachment from the donor surface [26,27], a complicated process that will be influenced by intrinsic factors such as the bacterial species involved [28,29], and environmental factors, like the properties of both the donating and the receiving surfaces [30,31], contact time, moisture level [26] and the application of friction and pressure [32]. Although inherently more complex than adhesion, bacterial transmission can be described according to similar surface thermodynamic principles as microbial adhesion to surfaces [33]. As a result of these considerations, a hydrophobic surface reduces the ability of hydrophilic bacteria to adhere closely to the surface, leading to a higher bacterial transmission from a

donor [34]. However, hydrophobic bacteria adhere more strongly to hydrophobic surfaces than hydrophilic ones, leading to less transmission [35]. In addition to hydrophobicity, also surface roughness and structure is generally considered as a key factor in bacterial adhesion and therewith in bacterial transmission. A smooth surface supports more contact area between receiving and donating surfaces, and therewith creates a higher transmission [27]. Although many factors are influential upon bacterial transmission, it is important to distinguish between transmission from a donor surface contaminated with a (sub)monolayer of adhering bacteria (**Figure 2a**) or from a donor surface fully covered with a multilayered bacterial biofilm (**Figure 2b**).

In the latter case, transmission from a donor surface can occur either through cohesive failure in the biofilm or interfacial failure at the donor-biofilm interface. As EPS molecules can act as bridging molecules between donor and receiver surfaces they thus affect transmission depending on the balance between the cohesive strength and the adhesion force on either the donor or the receiver.



**Figure 2.** Distinction between bacterial transmission from a (sub) monolayer of contaminating bacteria *versus* transmission from a multilayered biofilm.

(a) Transmission from a bacterial (sub)monolayer, involving interfacial failure at the donor-bacterium interface.

(b) Transmission of bacteria from a multilayered biofilm, involving either cohesive failure in the biofilm (b1) or interfacial failure at the donor-biofilm interface (b2).

## **Aim of The Thesis**

The aim of this thesis is to study the effect of various environmental and intrinsic factors on bacterial transmission from a donor surface covered with a multilayered bacterial biofilm. The main environmental factors studied are the surface (nano-)structure of receiving surfaces and the pressure and shear forces applied during transmission. The main intrinsic factors are the bacterial species and in particular the impact of the visco-elastic properties of the EPS matrix on biofilm transmission.



## References

- [1] Hall-Stoodley L, Costerton J, Stoodley P. Bacterial biofilms: from the natural environment to infectious diseases. *Nat Rev Microbiol* 2004;2:95–108. doi:10.1038/nrmicro821.
- [2] Bjarnsholt T, Alhede M, Alhede M, Eickhardt-Sørensen SR, Moser C, Kühl M, et al. The in vivo biofilm. *Trends Microbiol* 2013;21:466–74. doi:10.1016/j.tim.2013.06.002.
- [3] Tiller JC, Lee SB, Lewis K, Klibanov AM. Polymer surfaces derivatized with poly(vinyl-N-hexylpyridinium) kill airborne and waterborne bacteria. *Biotechnol Bioeng* 2002; 79:465–71. doi:10.1002/bit.10299.
- [4] Shuman EK, Chenoweth CE. Recognition and prevention of healthcare-associated urinary tract infections in the intensive care unit. *Crit Care Med* 2010;38:S373-9. doi:10.1097/CCM.0b013e3181e6ce8f.
- [5] Tacconelli E, Cataldo MA, Dancer SJ, De Angelis G, Falcone M, Frank U, et al. ESCMID guidelines for the management of the infection control measures to reduce transmission of multidrug-resistant Gram-negative bacteria in hospitalized patients. *Clin Microbiol Infect* 2014;20:1–55. doi:10.1111/1469-0691.12427.
- [6] Albrich WC, Harbarth S. Health-care workers: source, vector, or victim of MRSA? *Lancet Infect Dis* 2008;8:289–301. doi:10.1016/S1473-3099(08)70097-5.
- [7] Nicolle LE. Catheter associated urinary tract infections. *Antimicrob Resist Infect Control* 2014;3:1–8. doi:10.1186/2047-2994-3-23.
- [8] Safdar N, Maki DG. The pathogenesis of catheter-related bloodstream infection with noncuffed short-term central venous catheters. *Intensive Care Med* 2004;30:62–7. doi:10.1007/s00134-003-2045-z.
- [9] Maki DG. In vitro studies of a novel antimicrobial luer-activated needleless connector for prevention of catheter-related bloodstream infection. *Clin Infect Dis* 2010;50:1580–7. doi:10.1086/652764.
- [10] Kovaleva J. Infectious complications in gastrointestinal endoscopy and their prevention. *Best Pract Res Clin Gastroenterol* 2016;30:689–704. doi:10.1016/j.bpg.2016.09.008.
- [11] Muscarella LF. Evaluation of the risk of transmission of bacterial biofilms and *Clostridium difficile* during gastrointestinal endoscopy. *Gastroenterol Nurs* 2010; 33:28–35. doi:10.1097/SGA.0b013e3181cd199f.
- [12] Qu W, Busscher HJ, Hooymans JMM, Van der Mei HC. Surface thermodynamics and adhesion forces governing bacterial transmission in

contact lens related microbial keratitis. *J Colloid Interface Sci* 2011;358:430–6. doi:10.1016/j.jcis.2011.03.062.

[13] Hall BJ, Jones L. Contact lens cases: the missing link in contact lens safety? *Eye Contact Lens* 2010; 36:101–5. doi:10.1097/ICL.0b013e3181d05555.

[14] Richards D. How clean is your toothbrush? *Evid Based Dent* 2012;13:111–111. doi:10.1038/sj.ebd.6400895.

[15] Frazelle MR, Munro CL. Toothbrush contamination: a review of the literature. *Nurs Res Pract* 2012;2012:1–6. doi:10.1155/2012/420630.

[16] Davis MF, Iverson SA, Baron P, Vasse A, Silbergeld EK, Lautenbach E, et al. Household transmission of methicillin-resistant *Staphylococcus aureus* and other staphylococci. *Lancet Infect Dis* 2012;12:703–16. doi:10.1016/S1473-3099(12)70156-1.

[17] Donofrio RS, Bechanko R, Hitt N, O'Malley K, Charnauski T, Bestervelt LL, et al. Are we aware of microbial hotspots in our household? *J Environ Health* 2012;75:12–9.

[18] Newell DG, Elvers KT, Dopfer D, Hansson I, Jones P, James S, et al. Biosecurity-based interventions and strategies to reduce *Campylobacter* spp. on poultry farms. *Appl Environ Microbiol* 2011;77:8605–14. doi:10.1128/AEM.01090-10.

[19] Herman L, Heyndrickx M, Grijspeerdt K, Vandekerchove D, Rollier I, De Zutter L. Routes for *Campylobacter* contamination of poultry meat: epidemiological study from hatchery to slaughterhouse. *Epidemiol Infect* 2003;131:1169–80. doi:10.1017/S0950268803001183.

[20] Darrasse A, Darsonval A, Boureau T, Brisset MN, Durand K, Jacques MA. Transmission of plant-pathogenic bacteria by nonhost seeds without induction of an associated defense reaction at emergence. *Appl Environ Microbiol* 2010;76:6787–96. doi:10.1128/AEM.01098-10.

[21] Barret M, Guimbaud JF, Darrasse A, Jacques MA. Plant microbiota affects seed transmission of phytopathogenic microorganisms. *Mol Plant Pathol* 2016; 17:791–5. doi:10.1111/mpp.12382.

[22] Goldberg TL, Gillespie TR, Rwego IB, Estoff EL, Chapman CA. Forest fragmentation as cause of bacterial transmission among nonhuman primates, humans, and livestock, Uganda. *Emerg Infect Dis* 2008;14:1375–82. doi:10.3201/eid14.9.071196.

[23] Wilcox BA, Colwell RR. Emerging and reemerging infectious diseases: biocomplexity as an interdisciplinary paradigm. *Ecohealth* 2005;2:244–57. doi:10.1007/s10393-005-8961-3.[24] Kunttu HMT, Valtonen ET, Jokinen EI, Suomalainen LR. Saprophytism of a fish pathogen as a transmission strategy. *Epidemics* 2009;1:96–100. doi:10.1016/j.epidem.2009.04.003.

- [25] Starliper CE. Bacterial coldwater disease of fishes caused by *Flavobacterium psychrophilum*. J Adv Res 2011; 2:97–108. doi:10.1016/j.jare.2010.04.001.
- [26] Pérez-Rodríguez F, Valero A, Carrasco E, García RM, Zurera G. Understanding and modelling bacterial transfer to foods: a review. Trends Food Sci Technol 2008; 19:131–44. doi:10.1016/j.tifs.2007.08.003.
- [27] Hizal F, Choi C-H, Busscher HJ, Van der Mei HC. Staphylococcal adhesion, detachment and transmission on nanopillared Si surfaces. ACS Appl Mater Interfaces 2016;8: 30430–9. doi:10.1021/acsami.6b09437.
- [28] Gomez-Suarez C, Busscher HJ, Van der Mei HC. Analysis of bacterial detachment from substratum surfaces by the passage of air-liquid interfaces. Appl Environ Microbiol 2001;67:2531–7. doi:10.1128/AEM.67.6.2531-2537.2001.
- [29] Rusin P, Maxwell S, Gerba C. Comparative surface-to-hand and fingertip-to-mouth transfer efficiency of gram-positive bacteria, gram-negative bacteria, and phage. J Appl Microbiol 2002;93:585–92. doi:10.1046/j.1365-2672.2002.01734.x.
- [30] Knobben BAS, Van der Mei HC, Van Horn JR, Busscher HJ. Transfer of bacteria between biomaterials surfaces in the operating room—an experimental study. J Biomed Mater Res Part A 2007;80A:790–9. doi:10.1002/jbm.a.30978.
- [31] Sattar SA, Springthorpe S, Mani S, Gallant M, Nair RC, Scott E, et al. Transfer of bacteria from fabrics to hands and other fabrics: development and application of a quantitative method using *Staphylococcus aureus* as a model. J Appl Microbiol 2001;90:962–70. doi:10.1046/j.1365-2672.2001.01347.x.
- [32] Montville R, Schaffner DW. Inoculum size influences bacterial cross contamination between surfaces. Appl Environ Microbiol 2003;69:7188–93. doi:10.1128/AEM.69.12.7188-7193.2003.
- [33] Sharma PK, Hanumantha Rao K. Adhesion of *Paenibacillus polymyxa* on chalcopyrite and pyrite: surface thermodynamics and extended DLVO theory. Colloids Surfaces B Biointerfaces 2003;29:21–38. doi:10.1016/S0927-7765(02)00180-7.
- [34] Boks NP, Norde W, Van der Mei HC, Busscher HJ. Forces involved in bacterial adhesion to hydrophilic and hydrophobic surfaces. Microbiology 2008;154:3122–33. doi:10.1099/mic.0.2008/018622-0.
- [35] Foong SCC, Dickson JS. Attachment of *Listeria monocytogenes* on ready-to-eat meats. J Food Prot 2004;67:456–62. doi:10.4315/0362-028X-67.3.456.