

University of Groningen

## Towards strengthening memory immunity in the ageing population

van der Heiden, Marieke

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2018

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

van der Heiden, M. (2018). *Towards strengthening memory immunity in the ageing population: Investigating the immunological fitness of middle-aged adults*. [Thesis fully internal (DIV), University of Groningen]. Rijksuniversiteit Groningen.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# TETANUS TOXOID CARRIER PROTEIN INDUCED T-HELPER CELL RESPONSES UPON VACCINATION OF MIDDLE-AGED ADULTS

Marieke van der Heiden<sup>1,2</sup>

Aafke Duizendstra<sup>1</sup>

Guy A. M. Berbers<sup>1</sup>

Annemieke M.H. Boots<sup>2</sup>

Anne-Marie Buisman<sup>1</sup>

<sup>1</sup> Centre for Infectious Disease Control (Cib), National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands

<sup>2</sup> Department of Rheumatology and Clinical Immunology, University of Groningen, University Medical Centre Groningen, Groningen, The Netherlands

*Vaccine*; 2017.08.056

---



## Abstract

Vaccines frequently induce suboptimal immune responses in the elderly, due to immunological ageing. Timely vaccination may be a strategy to overcome this problem, which classifies middle-aged adults as an interesting target group for future vaccine interventions. However, the immunological fitness of the middle-aged population is ill-defined. It is currently unknown whether effective T-cell help towards B-cells is initiated by conjugate-carrier vaccines at middle-age.

We characterized systemic Tetanus Toxoid (TT) specific T-helper cell responses in the circulation of middle-aged adults (50-65 years of age, n= 31) having received the MenACWY-TT vaccination. Blood samples were taken pre- as well as 7 days, 28 days, and 1 year post-vaccination. TT-specific T-cell responses were determined by IFN $\gamma$  Elispot and by the secretion of IFN $\gamma$ , IL13, IL10, IL17, and IL21 in cell culture supernatants. Circulating CD4+CXCR5+ICOS+IL21+ cells were analyzed by flow cytometry, and meningococcal and TT-specific IgG responses by bead-based immunoassays. The correlation between the T-cell help and humoral responses was evaluated.

Vaccination with a TT-carrier protein induced a mixed TT-specific Th1 (IFN $\gamma$ ), Th2 (IL13, IL10), and Th17 (IL17) response in most participants. Additionally, circulating CD4+CXCR5+ICOS+IL21+ cells were significantly increased 7 days post-vaccination. Pre-vaccination TT-specific cytokine production and post-vaccination Th2 responses correlated positively with the increase of CD4+CXCR5+ICOS+IL21+ cells. No correlation between T-cell help and antibody responses was found.

The characteristics of the T-cell response upon a TT-carrier vaccination suggests effective T-cell help towards B-cells in response to meningococcal polysaccharides, although the absence of a correlation with the antibody responses warrants further clarification. However, the robust T-helper cell response in middle-aged adults, decades after previous TT vaccinations, strengthens the classification of this age group for future vaccine interventions in the context of population ageing.

## Introduction

Immunological ageing is characterized by clear compositional changes in the T-cell compartment [1-5], which become already apparent in the 6<sup>th</sup> decade of life [2]. This immunological ageing increases the susceptibility towards infectious diseases and reduces the immune response to vaccines [6]. Therefore, timely vaccination is essential to circumvent the impact of this so called 'immunosenescence' and thereby enhance memory immunity in the ageing population. Consequently, future vaccination programs need to be developed towards a life-course scheme [7-9]. From this perspective middle-aged adults are an interesting group for studying appropriate future vaccine programs[6]. However, information about the immunological fitness and consequently vaccine responses in middle-aged individuals is lacking.

It is currently unknown whether conjugate-carrier vaccines are able to induce effective T-helper cell responses in middle-aged adults. In infants, carrier proteins are used to induce T-cell help to the B-cell epitopes in order to generate long-term protection and memory B-cell formation in response to bacterial polysaccharides [10]. Not only the strength but also the type of the T-helper cell response influences the quality of the T-cell help. In general Th2 responses provide more efficient B-cell help than do Th1 responses, which may affect the functionality of the humoral response [11-14]. In addition, T-cell help is mainly provided by follicular helper T-cells ( $T_{FH}$ -cells) within the germinal centers (GCs), where the B- and T-cells interact [15, 16]. With advancing age, altered numbers of these  $T_{FH}$ -cells, lower  $T_{FH}$ -cell functionality, and diminished formation of GCs were observed [17, 18], possibly negatively affecting the T-cell help after conjugate-carrier vaccination in older age groups.

Interestingly, it was recently described that an ongoing B:T-cell interaction in the GCs is mirrored by higher frequencies of  $T_{FH}$ -cells in the circulation at 6 to 8 days post-vaccination [15, 18-21]. Until now, no consensus on the exact phenotype of circulating  $T_{FH}$ -cells is established. Similar to GC  $T_{FH}$ -cells, circulating CXCR5+ cells were previously described to promote B-cell differentiation and antibody secretion [13, 22-24]. A key characteristic of  $T_{FH}$ -cells is the production of interleukin 21 (IL21), a cytokine involved in GC development, Ig isotype switching, hyper-mutation of the IgG heavy chain, IgG affinity maturation, and memory B-cell formation [13, 25-28]. Moreover, the inducible costimulatory molecule (ICOS) was found essential in the interaction of  $T_{FH}$ -cells with the follicular B-cells [17, 29, 30]. Finally, multiple studies described roles for programmed cell death 1 (PD1) in phenotype characterization of the peripheral  $T_{FH}$ -cells [13, 15, 18, 20, 22, 31]. Although some studies suggest a direct link between the  $T_{FH}$ -cell and the humoral responses [18, 20, 32], this is currently unclear.

In this study, we determined the induced T-cell responses initiated by a TT-carrier protein in the circulation of middle-aged adults (50-65 years of age) having received the TT-conjugated MenACWY vaccine (Nimenrix), in order to evaluate the T-cell help provided to B-cells in

response to the bacterial polysaccharides in this age group. This information is essential for the understanding of T-helper cell responses at older ages and adds to knowledge on the immunological fitness of middle-aged adults.

## Methods

### Study subjects and blood sampling

Thirty-one relatively healthy middle-aged (50-65 years of age) participants were randomly selected from a larger cohort. All participants were intramuscularly vaccinated with one dose of the TT-conjugated MenACWY vaccine (Nimenrix, GlaxoSmithKline (GSK)). Potential participants were excluded based on the following criteria: antibiotic use or fever ( $>38^{\circ}\text{C}$ ) within the last 14 days, serious diseases demanding immune suppressive medical treatment within the last 3 months, a known or suspected immune deficiency, a blood coagulation disorder, a neurologic disorder, administration of blood products in the past 6 months, serious surgery within the last 3 months, the use of hormone supplementation, pregnancy, a suspected allergy towards the vaccine components, history of serious adverse events after previous vaccinations, a previous meningococcal vaccination, a previous meningococcal episode, a tetanus vaccination within the last 5 years, and any vaccination in the month before enrolment. Additional health characteristics of these participants are presented in

#### **S. Table 1.**

The Medical Ethical Committee: Medical Research Ethics Committees United (Mec-U) (NTR4636) approved the study and written informed consent was obtained from all participants prior to the study. All procedures were in accordance with the Declaration of Helsinki.

Blood samples were taken pre-, as well as 7 days, 28 days, and 1 year post- vaccination. Peripheral blood mononuclear cells (PBMCs) were isolated using Vacutainer cell preparation tubes (CPT) containing sodium citrate (Becton Dickinson (BD)), according to the manufacturer's prescriptions. Cells were washed with RPMI-1640 medium (Gibco) supplemented with 1% heat inactivated fetal calf serum (FCS, Gibco), 1% Penicillin and Streptomycin (Lonza). PBMCs were counted and  $7,5 \cdot 10^6$  cells/mL were frozen in a 90% FCS 10% DMSO solution at  $-135^{\circ}\text{C}$  until further use. Serum was collected using serum clotting tubes (BD) and was stored at  $-20^{\circ}\text{C}$  until use.

From the initial cohort, PBMC samples obtained in sufficient quantities at all 4 time points were available from 31 participants for analysis. TT-specific T-cell responses were determined in 22 participants, whereas  $T_{\text{FH}}$ -cell markers were investigated in 20 participants. Samples from 11 participants provided enough cells for analysis in all assays (**Figure 1**).




Figure 1. Participants flow chart.

### Tetanus specific T-cell stimulation and IFN $\gamma$ Elispot

PBMC stimulation, supernatant collection and the IFN $\gamma$  Elispot method were consecutively performed as previously described [33]. Based on prior titration and kinetics experiments,  $3 \times 10^5$  PBMCs/well were stimulated for 5 days with medium containing 5% AB serum (Sigma Aldrich) only (negative control), or with addition of 6.67 Lf/mL purified tetanus toxoid (NVI Bilthoven, The Netherlands) or 5  $\mu$ g/mL Pokeweed mitogen (Sigma) as a positive control, in triplicate. Thereafter, stimulated cells were transferred to anti-IFN $\gamma$ -coated Elispot plates (Mabtech) for 18h. Elispot plates were scanned with the Epon ELISPOT Scanner and spots enumerated with the AELVIS software, using a standardized protocol. Antigen specific spots were calculated by subtraction of the spots in the negative controls. In order to avoid influences of assay variability, all 4 samples (pre-, 7d, 28d, and 1 year) of the study participants were measured in the same experiment employing the same batch of IFN $\gamma$  coated Elispot plates.

### Cytokine detection in supernatants

After 5 days of PBMC stimulation, cell culture supernatants were collected and stored at -80 °C until further use. Subsequently, IFN $\gamma$ , IL13, IL10, IL17, and IL21 concentrations in the supernatants were determined by commercially available multiplex bead-based immunoassay kits according to the manufacturer's instructions (Bio-Rad Laboratories). Samples with concentrations below the lower limit of quantification were assigned half the concentration of the lowest measurement. Cytokine concentrations in the negative controls were subtracted from those in the antigen stimulated samples.

### **Detection of CD4+CXCR5+ICOS+IL21+ cells by flow cytometry**

To assess the potency of circulating T-cells to produce IL21,  $3 \times 10^5$  PBMCs/well of samples pre- as well as 7 and 28 days post-vaccination were incubated with medium only (negative control) or with a cell stimulation cocktail containing phorbol 12-myristate 13-acetate (PMA) and ionomycin (IONO) in a suboptimal 2000x dilution (eBiosciences) in triplicate. Cells were incubated for 6 hours at 37 °C with 5% CO<sub>2</sub>. After one hour, GolgiPlug protein transport inhibitor containing Brefeldin A (1000x dilution, BD) was added to each well. After thorough washing, cells were incubated for 30 minutes with a mixture of Life-Death Zombie Aqua fluorescent dye (Biolegend) and surface antibodies in FACS buffer, containing PBS with 0.5%BSA and 2mM EDTA. The following antibodies were used for the surface staining: CD3 (UCHT1)-V450, CD4 (RPA-T4)-APC, CXCR5 (RF8B2)-PerCP-Cy5.5, PD1 (MIH4)- FITC (all BD), and ICOS (C398.4A)-APC-Cy7 (Biolegend). Thereafter, cells were washed with PBS, and permeabilized for 20 minutes with Cytotfix/Cytoperm (BD). A Perm/Wash solution (BD) was used to wash the cells. In addition, the cells were stained for 30 minutes with IL21 (3A3-N2.1)-PE (BD) and with CD4 (RPA-T4)-APC detecting intracellular IL21 and CD4, respectively. After washing, the cells were resuspended in cold FACS-fixation buffer (Biolegend) until measurement. Flowcytometry analysis was performed on a 4-lazer LSR Fortessa (BD) and data was analyzed using FlowJo V10.

### **Serological analysis**

Tetanus and meningococcal specific IgG concentrations were determined using the fluorescent-bead-based multiplex immunoassay (MIA) as previously described [34, 35].

### **Statistics**


The number of TT-specific IFN $\gamma$  spots, cytokine concentrations, CD4+CXCR5+ICOS+IL21+ cell responses, and the IgG responses at the different time points were compared with the Wilcoxon signed rank test, preceded by the Friedman test. Correlations were determined by the Spearman's rho correlation test. Graphpad V7 and SPSS V22.0 were used.

## **Results**

### **TT-specific IFN $\gamma$ -producing cells are boosted after vaccination**

Numbers of TT-specific IFN $\gamma$ -producing T-cells were enumerated pre- and post- vaccination (**Figure 2**). Pre-vaccination, already high numbers ( $>10$  spots/  $10^5$  PBMCs) of TT-specific IFN $\gamma$ -producing cells were found in the majority of the participants (16 out of 22). These numbers significantly increased 28 days post-vaccination; a strong trend was seen after 7 days. One year post-vaccination, the numbers of TT-specific IFN $\gamma$ -producing cells returned

to pre-vaccination values (**Figure 2a**). Although the response was highly variable (**Figure 2b**), most participants showed an enhanced IFN $\gamma$  T-cell response either at day 7 or at day 28 post-vaccination, indicating that vaccination with a TT-carrier protein boosted a short-term Th1 type response in middle-aged adults.



**Figure 2. Numbers of TT-specific IFN $\gamma$  producing cells.**


**a)** The numbers of IFN $\gamma$  producing cells/  $10^5$  PBMCs pre- and post-vaccination in the negative control (BL) and after stimulation with Tetanus Toxoid (TT). The horizontal line indicates the geometric mean concentration **b)**. Paired numbers of IFN $\gamma$  producing cells/  $10^5$  PBMCs pre- and post- vaccination. The different time points were compared with the Wilcoxon signed rank test preceded by the Friedman test. \*\*  $p < 0.01$ ,  $n = 22$ .

### Vaccination with a TT-carrier protein boosted mixed T-helper cytokine responses

Next, we analyzed TT-specific Th1 (IFN $\gamma$ ), Th2 (IL10, IL13), Th17 (IL17) and T<sub>FH</sub> (IL21) cytokine secretion profiles. Similar to the numbers of IFN $\gamma$ -producing cells, TT vaccination led to increased TT-specific IFN $\gamma$  production at days 7 and 28 post-vaccination (**Figure 3a**). Also, a marked increase of the IL13 production was seen post-vaccination (**Figure 3b**), which resulted in a relative decrease of the TT-specific IFN $\gamma$ /IL13 ratio at 28 days post-vaccination (**Figure 3c**).

Additionally, TT-specific IL10 production was largely enhanced 7 days post-vaccination in all participants, after which this response gradually declined 28 days post-vaccination (**Figure 3d**). Moreover, a significant but limited enhancement of TT-specific IL17 secretion was observed both at day 7 and 28 (**Figure 3e**). Of note, TT-specific IL21 production was below detection limit (data not shown). Overall, 7 days post-vaccination, a mixed cytokine profile was induced in most donors, although some participants clearly showed a Th1 (IFN $\gamma$ ) or Th2 (IL10, IL13) dominated response (**Figure 3f**).





**Figure 3. TT-specific cytokine responses**


Concentrations (pg/mL) of IFN $\gamma$  (**a**), IL13 (**b**), IL10 (**d**), and IL17 (**e**) in cell culture supernatants after 5 days of Tetanus Toxoid stimulation. The red line indicates the geometric mean concentration. **c**. The IFN $\gamma$ /IL13 ratio pre- and post- vaccination. The different time points were compared with the Wilcoxon signed rank test preceded by the Friedman test. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , \*\*\*\* $p < 0.0001$   $n = 20$ . **f**. The delta TT-specific cytokine responses at day 7 (day 7 – day 0) per donor.

### **The TT-carrier protein induced circulating CD4+CXCR5+ICOS+IL21+ cells**

Although we did not detect TT-specific IL21 production, we next investigated the induction of circulating CD4+CXCR5+ICOS+IL21+ cells following vaccination by TT-specific or polyclonal stimulation with PMA/IONO (**Figure 4a**). Since PMA/IONO stimulation caused the internalization of CD4 by T-cells, both intra- and extra-cellular staining of CD4 was performed to detect the total CD4 T-cell pool. Firstly, no intracellular IL21 production was observed after TT-specific stimulation (data not shown). Stimulation with PMA/IONO induced IL21 production in both the CXCR5+ and CXCR5- subsets, although this production was significantly higher in the CXCR5+ cells (**Figure 4c**). Also, ICOS expression was higher in the CXCR5+ cells (**Figure 4b**). Similarly, higher frequencies of IL21+ICOS+ cells were detected in the CXCR5+ subset (**Figure 4d**). Seven days post vaccination a significant increase in the frequency of IL21+ICOS+ cells within the CXCR5+ compartment was observed (**Figure 4d**). This response was variable between the participants, with 17 responders and 3 non-responders. Additionally, some samples (n=12) were analyzed for PD1 expression (**S. Figure 1b**). A strong significant correlation between the percentages of CXCR5+IL21+ICOS+ cells and CXCR5+IL21+ICOS+PD1+ cells 7 days post-vaccination was found (**S. Figure 1c**). These results indicate a significant increase in activated T-cells that have high potential for B-cell help after vaccination with a TT-carrier protein in middle-aged adults.

### **TT-specific T-helper cell cytokines correlate with the CD4+CXCR5+ICOS+IL21+ cell response 7 days post-vaccination**

We further assessed whether the increase in the CD4+CXCR5+ICOS+IL21+ cells at 7 days post-vaccination correlated with TT-specific T-helper cytokine production, both pre- and post-vaccination. Seven days post-vaccination, a strong correlation was observed between the increase in CD4+CXCR5+ICOS+IL21+ cells and the increase in TT-specific IL10 and IL13 production (**Table 1a. sub 1**). The increase in CD4+CXCR5+ICOS+IL21+ cells did not correlate with the TT-specific Th17 or Th1 response, although a trend was seen for the latter. Next we analyzed the correlation between pre-vaccination TT-specific cytokine production and the increase in CD4+CXCR5+ICOS+IL21+ cells at day 7. A significant correlation was found between the production of pre-vaccination TT-specific IFN $\gamma$  and the increase in CD4+CXCR5+ICOS+IL21+ cells 7 days post-vaccination (**Table 1a. sub 2**). Moreover, a positive trend was found for the other T-helper subsets, suggesting that the presence of pre-vaccination TT-specific T-helper cells enhanced the CD4+CXCR5+ICOS+IL21+ cells responses.



**Figure 4. Induction of circulating CD4+CXCR5+ICOS+IL21+ cells post-vaccination.**

**a).** Gating strategies used for characterization of the CD4+CXCR5+ICOS+IL21+ cells. A representative sample is shown. **b).** Percentages of ICOS+ cells within the CXCR5+ and CXCR5- populations pre- and post-vaccination (7 and 28 days). **c).** Percentages of IL21+ cells within the CXCR5+ and CXCR5- populations pre- and post-vaccination (7 and 28 days). **d).** Percentages of ICOS+IL21+ cells within the CXCR5+ and CXCR5- populations pre- and post-vaccination (7 and 28 days). The horizontal lines in the graph represent the geometric mean percentages. The different time points were compared with the Wilcoxon signed rank test preceded by the Friedman test. CXCR5+ and CXCR5- cells at the different time points were compared with the Mann Whitney U test. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , \*\*\*\* $p < 0.0001$ , n=20.


**Table 1a. Correlations between the TT-specific cytokine responses and the induction of circulating CD4+CXCR5+ICOS+IL21+ cells.**

Parameter	variable	Spearman's rho	p-value
1. <u>delta TT-specific cytokine response at day 7</u> (day 7 – day 0, pg/ml)	IFN $\gamma$	0.650	0.058
	IL13	0.700	<b>0.036</b>
	IL10	0.850	<b>0.004</b>
	IL17	0.367	0.332
2. <u>pre-vaccination TT-specific cytokine response</u> (pg/ml)	IFN $\gamma$	0.795	<b>0.010</b>
	IL13	0.503	0.138
	IL10	0.661	0.053
	IL17	0.571	0.084

The delta % CD4+CXCR5+ICOS+IL21+ cells at days 7 (7 days – pre-vaccination) was used in the correlations. n=10. The Spearman's rho correlation test was used.

## TT-specific T-helper cell responses do not correlate with antigen specific IgG responses

The TT-conjugated meningococcal vaccine induced clear antibody responses to TT and the meningococcal C polysaccharides (MenC-PS) (**Figure 5**). Yet, no correlations between the TT-specific T-helper cell responses or the CD4+CXCR5+ICOS+IL21+ cells response with the MenC PS-specific and TT-specific IgG responses were detected, using delta (**Table 1b**), absolute, and fold change T-cell responses (data not shown). On a side note, positive correlations were found between the pre- and post-vaccination IgG concentrations for TT (7d: rho: 0.591 p=0.006; 28d: rho: 0.711 p=<0.0001, 1y: rho: 0.713 p<0.0001), but not for MenC. Adjusting for pre-vaccination immunity did not change our results. Similar results were obtained for the other meningococcal serotypes, and for the IgG subclass responses (data not shown).

**Figure 5. Anti TT and MenC-PS specific IgG responses**

TT (a) and MenC-PS (b) specific IgG responses pre- and post- vaccination. The different time points were compared with the Wilcoxon signed rank test preceded by the Friedman test. \*\* p<0.01 \*\*\*\*p<0.0001 n=20.

**Table 1b. Correlations between the induced circulating CD4+CXCR5+ICOS+IL21+ cell response, TT-specific cytokine responses and the TT and MenC-PS specific IgG responses.**

T1-T0 response	parameter	Time point	Spearman's rho	p-value
<u>CXCR5+ICOS+IL21+</u>	TT IgG	28 days	-0.135	0.569
	TT IgG	1 year	-0.271	0.248
	MenC IgG	28 days	0.056	0.816
	MenC IgG	1 year	0.039	0.870
<u>IL10</u>	TT IgG	28 days	0.168	0.478
	TT IgG	1 year	0.026	0.915
	MenC IgG	28 days	0.194	0.413
	MenC IgG	1 year	0.206	0.384
<u>IL13</u>	TT IgG	28 days	0.265	0.259
	TT IgG	1 year	0.277	0.238
	MenC IgG	28 days	0.152	0.523
	MenC IgG	1 year	0.217	0.359
<u>IFN<math>\gamma</math></u>	TT IgG	28 days	0.245	0.298
	TT IgG	1 year	0.084	0.724
	MenC IgG	28 days	0.314	0.177
	MenC IgG	1 year	0.307	0.188
<u>IL17</u>	TT IgG	28 days	0.198	0.402
	TT IgG	1 year	0.176	0.458
	MenC IgG	28 days	-0.060	0.801
	MenC IgG	1 year	0.048	0.840

The delta % of CD4+CXCR5+ICOS+IL21+ cells at days 7 (7 days – pre-vaccination) was used in the correlations, as well as the delta cytokine production (7 days- pre-vaccination). n= 20. The Spearman's rho correlation test was used.

## Discussion

Within this study, we show that vaccination of middle-aged adults with a TT-carrier protein, used to initiate T-cell help in response to meningococcal polysaccharides, induced a mixed Th1, Th2, and Th17 T-cell response, although some participants showed a clear Th1 (IFN $\gamma$ ) or Th2 (IL10, IL13) dominated response. In addition, the enhanced frequencies of circulating CD4+CXCR5+ICOS+IL21+ cells 7 days post-vaccination suggests the induction of activated T-cells with T<sub>FH</sub> cells properties by a TT-carrier protein in older adults. These T-helper cell responses are suggestive of effective T-cell help towards the B-cells in response to the meningococcal polysaccharides, although we did not find a direct correlation between the T-cell response induced by the carrier protein and the meningococcal specific antibody responses.

Although a mixed cytokine response was observed, our data suggests a slightly skewed Th2 response post-vaccination, mainly represented by a strong IL10 response 7 days post-

vaccination and a decreased post-vaccination IFN $\gamma$ /IL13 ratio. The strong IL10 response may both regulate the Th1 response [36], and stimulate the antibody responses towards the bacterial polysaccharides [12, 13]. This positive effect of IL10 was strengthened by the strong positive correlation between the IL10 production and the circulating CD4+CXCR5+ICOS+IL21+ cell response at day 7 post-vaccination.

Antigen specific detection of IL21 production by T-cells is notoriously difficult, since B-cells directly consume secreted IL21 and the T<sub>FH</sub>-cells need vigorous *in vitro* stimulation to produce IL21 [31, 37, 38]. Therefore, we assessed the potency of IL21 production by T-cells following PMA/IONO stimulation and intercellular detection of IL21. Indeed, IL21+ cells were found in both the CXCR5+ and CXCR5- subset, as was also shown by others [22, 23, 31]. Yet, higher frequencies of IL21+ cells were observed in the CXCR5+ subset as compared to the CXCR5-counterpart, which is in agreement with others [23]. Although it was previously shown that IL21 production of both subsets provided help towards B-cells, this was significantly increased in CXCR5+ cells due to the elevated expression of ICOS, a B-cell co-stimulator [22]. Therefore, IL21 production alone is perceived insufficient to determine efficient T<sub>FH</sub>-cell responses [22, 31]. Post-vaccination, we observed a tendency towards increased expression of ICOS in the CXCR5+ subset (data not shown), which is in line with findings by Herati *et al.* after influenza vaccination in the elderly [18]. Taken together, our findings of increased frequencies of CXCR5+ICOS+IL21+ cells 7 days post-vaccination suggest the induction of T<sub>FH</sub>-cells. This was strengthened by the strong correlation between the frequencies of CXCR5+ICOS+IL21+ cells and the CXCR5+ICOS+IL21+PD1+ cells 7 days post-vaccination, since circulating PD1+ T<sub>FH</sub> cells share functional properties with GC T<sub>FH</sub> cells [20, 23, 32].

In addition, we observed a correlation between the increased frequencies of CXCR5+ICOS+IL21+ cells at day 7 and the presence of TT-specific pre-vaccination cytokine production, indicating that pre-vaccination T-cell immunity enhances T-cell responses upon conjugate-carrier vaccination. Although only 3 of the participants received a TT vaccination after the year 2000, pre-vaccination TT-specific T-cell cytokine production was observed in most participants, which indicates that TT-specific T-helper cells persist for years or even decades after historical vaccinations in most participants. However, due to the exclusion of individuals with recent (<5 years) TT vaccinations in this study, the effects of recent tetanus vaccinations on the T-cell response towards the TT-carrier protein remains unknown. Of note, pre-vaccination TT-specific IgG concentrations did not correlate with TT-specific cytokine production and thus cannot be used as surrogate marker for the presence of TT-specific T-cell immunity.

In spite of the induction of T-cell responses that favour B-cell help by the TT-carrier protein, we did not observe an association between the strength of the T-cell response and the TT or meningococcal polysaccharide specific antibody concentrations. The absence of this correlation might be explained by several factors. First of all, the multivalent nature of the

meningococcal vaccination might complicate the detection of a correlation between the T-cell and the humoral response. Moreover, the different polysaccharide antigens induced diverse antibody responses within the same person (data not shown). Therefore, we suggest a large effect of the existing B-cell repertoire in response to the polysaccharide antigens, a notion corroborated by the findings of other studies [28, 39]. Additionally, the lack of a correlation between the T-helper cell and antibody responses at the time points analysed may be explained by potential differences in peripheral B- and T-cell kinetics. Furthermore, we propose that simply the presence but not the strength of the T-cell response is sufficient to induce an effective vaccine antibody response, also that is described for a conjugated meningococcal vaccination in infants [40]. Finally, although our results might suggest the induction of circulating activated T-cells with  $T_{FH}$  cell properties after vaccination, the exact functionality of these CD4+CXCR5+ICOS+IL21+ cells is currently unknown. In particular the relation of these CD4+CXCR5+ICOS+IL21+ cells to a possible  $T_{FH}$  cell response in the GCs, clearly awaits further confirmation, especially in relation to immunological ageing, since GC and  $T_{FH}$ -cell functionality might be altered with advancing age [17, 41]. It has been described that circulating  $T_{FH}$  cells did correlate with influenza specific antibody responses post-vaccination in young adults but not in the elderly [18], this might indicate that GC responses are diminished in older adults although circulating  $T_{FH}$  cells are present. In this study, we did not detect clear age related differences in the T-cell responses, which might be influenced by the limited age range, small sample size in our study, and possible confounding effects of pre-vaccination T-cell immunity.

In conclusion, vaccination with a TT-carrier protein in middle-aged adults induced a mixed TT-specific T-helper cell response. Moreover, the induction of circulating CD4+CXCR5+ICOS+IL21+ cells 7 days post-vaccination suggests a circulating T-cell response with high potential for B-cell help. Subsequently, the characteristics of these T-cells proposes effective T-cell help towards B-cells in response to the meningococcal polysaccharides, although a direct correlation with the antibody responses was not found, a finding that warrants further clarification especially in the context of immune ageing. Notwithstanding, the robust T-helper cell response, decades after previous TT vaccination, adds to the knowledge on the immunological fitness of middle-aged adults. Moreover, these results strengthen the classification of this age group as an interesting target group for future vaccine interventions, to stimulate memory immunity before reaching old age.

## Acknowledgements

We thank all the participants of the study and the nurses who performed the vaccinations and blood drawings. Furthermore, we thank the technicians who helped with the PBMC isolation as well as Lia de Rond for her laboratory assistance and Teun Guichelaar for critical reviewing of the manuscript.

## References

1. den Braber, I., et al., *Maintenance of peripheral naive T cells is sustained by thymus output in mice but not humans*. *Immunity*, 2012. **36**(2): p. 288-297.
2. Herndler-Brandstetter, D., H. Ishigame, and R.A. Flavell, *How to define biomarkers of human T cell aging and immunocompetence?* *Front Immunol*, 2013. **4**: p. 136.
3. Arnold, C.R., et al., *Gain and loss of T cell subsets in old age--age-related reshaping of the T cell repertoire*. *J Clin Immunol*, 2011. **31**(2): p. 137-46.
4. Massoud, A. and A.H. Massoud, *Biological and Phenotypic Alterations of T Cells in Aging*, in *Immunology of Aging*. 2014, Springer. p. 177-182.
5. Weng, N.P., A.N. Akbar, and J. Goronzy, *CD28(-) T cells: their role in the age-associated decline of immune function*. *Trends Immunol*, 2009. **30**(7): p. 306-12.
6. Boraschi, D., et al., *The gracefully aging immune system*. *Science translational medicine*, 2013. **5**(185): p. 185ps8-185ps8.
7. Michel, J.-P. and P.O. Lang, *Promoting life course vaccination*. *Rejuvenation Research*, 2011. **14**(1): p. 75-81.
8. Lang, P.O. and R. Aspinall, *Immunosenescence and herd immunity: with an ever-increasing aging population do we need to rethink vaccine schedules?* *Expert Review of Vaccines*, 2012. **11**(2): p. 167-176.
9. Lang, P.O., et al., *Immunosenescence: implications for vaccination programmes in adults*. *Maturitas*, 2011. **68**(4): p. 322-330.
10. Pollard, A.J., K.P. Perrett, and P.C. Beverley, *Maintaining protection against invasive bacteria with protein-polysaccharide conjugate vaccines*. *Nature Reviews Immunology*, 2009. **9**(3): p. 213-220.
11. Abbas, A.K., K.M. Murphy, and A. Sher, *Functional diversity of helper T lymphocytes*. *Nature*, 1996. **383**(6603): p. 787-793.
12. Kidd, P., *Th1/Th2 balance: the hypothesis, its limitations, and implications for health and disease*. *Alternative Medicine Review*, 2003. **8**(3): p. 223-246.
13. Abbas, A.K., A.H. Lichtman, and S. Pillai, *Basic immunology: functions and disorders of the immune system*. 2012: Elsevier Health Sciences.
14. Kawano, Y., T. Noma, and J. Yata, *Regulation of human IgG subclass production by cytokines. IFN-gamma and IL-6 act antagonistically in the induction of human IgG1 but additively in the induction of IgG2*. *The Journal of Immunology*, 1994. **153**(11): p. 4948-4958.
15. Vinuesa, C.G., et al., *Follicular Helper T Cells*. *Annual Review of Immunology*, 2016. **34**(1).
16. Nutt, S.L. and D.M. Tarlinton, *Germinal center B and follicular helper T cells: siblings, cousins or just good friends [quest]*. *Nature immunology*, 2011. **12**(6): p. 472-477.
17. Linterman, M.A., *How T follicular helper cells and the germinal centre response change with age*. *Immunology and cell biology*, 2014. **92**(1): p. 72-79.
18. Herati, R.S., et al., *Circulating CXCR5+ PD-1+ response predicts influenza vaccine antibody responses in young adults but not elderly adults*. *The Journal of Immunology*, 2014. **193**(7): p. 3528-3537.
19. Pallikkuth, S., et al., *Impaired peripheral blood T-follicular helper cell function in HIV-infected nonresponders to the 2009 H1N1/09 vaccine*. *Blood*, 2012. **120**(5): p. 985-993.
20. He, J., et al., *Circulating precursor CCR7 lo PD-1 hi CXCR5+ CD4+ T cells indicate Tfh cell activity and promote antibody responses upon antigen reexposure*. *Immunity*, 2013. **39**(4): p. 770-781.
21. Linterman, M.A. and D.L. Hill, *Can follicular helper T cells be targeted to improve vaccine efficacy?* *F1000Research*, 2016. **5**.
22. Chevalier, N., et al., *CXCR5 expressing human central memory CD4 T cells and their relevance for humoral immune responses*. *The Journal of Immunology*, 2011. **186**(10): p. 5556-5568.
23. Morita, R., et al., *Human blood CXCR5+ CD4+ T cells are counterparts of T follicular cells and contain specific subsets that differentially support antibody secretion*. *Immunity*, 2011. **34**(1): p. 108-121.




24. MacLeod, M.K., et al., *Memory CD4 T cells that express CXCR5 provide accelerated help to B cells*. The Journal of Immunology, 2011. **186**(5): p. 2889-2896.
25. Vogelzang, A., et al., *A fundamental role for interleukin-21 in the generation of T follicular helper cells*. Immunity, 2008. **29**(1): p. 127-137.
26. Bryant, V.L., et al., *Cytokine-mediated regulation of human B cell differentiation into Ig-secreting cells: predominant role of IL-21 produced by CXCR5+ T follicular helper cells*. The Journal of Immunology, 2007. **179**(12): p. 8180-8190.
27. Avery, D.T., et al., *IL-21-induced isotype switching to IgG and IgA by human naive B cells is differentially regulated by IL-4*. The Journal of Immunology, 2008. **181**(3): p. 1767-1779.
28. de Wit, J., et al., *Human B cells promote T-cell plasticity to optimize antibody response by inducing coexpression of TH1/TFH signatures*. Journal of Allergy and Clinical Immunology, 2015. **135**(4): p. 1053-1060.
29. Bauquet, A.T., et al., *The costimulatory molecule ICOS regulates the expression of c-Maf and IL-21 in the development of follicular T helper cells and TH-17 cells*. Nature immunology, 2009. **10**(2): p. 167-175.
30. Choi, Y.S., et al., *ICOS receptor instructs T follicular helper cell versus effector cell differentiation via induction of the transcriptional repressor Bcl6*. Immunity, 2011. **34**(6): p. 932-946.
31. Crotty, S., *Follicular helper CD4 T cells (T<sub>fh</sub>)*. Annual review of immunology, 2011. **29**: p. 621-663.
32. Locci, M., et al., *Human circulating PD-1+ CXCR3- CXCR5+ memory T<sub>fh</sub> cells are highly functional and correlate with broadly neutralizing HIV antibody responses*. Immunity, 2013. **39**(4): p. 758-769.
33. Schure, R.-M., et al., *T-cell responses before and after the fifth consecutive acellular pertussis vaccination in 4-year-old Dutch children*. Clinical and Vaccine Immunology, 2012. **19**(11): p. 1879-1886.
34. van Gageldonk, P.G., et al., *Development and validation of a multiplex immunoassay for the simultaneous determination of serum antibodies to Bordetella pertussis, diphtheria and tetanus*. Journal of immunological methods, 2008. **335**(1): p. 79-89.
35. de Voer, R.M., et al., *Simultaneous detection of Haemophilus influenzae type b polysaccharide-specific antibodies and Neisseria meningitidis serogroup A, C, Y, and W-135 polysaccharide-specific antibodies in a fluorescent-bead-based multiplex immunoassay*. Clinical and Vaccine Immunology, 2009. **16**(3): p. 433-436.
36. Ouyang, W., et al., *Regulation and functions of the IL-10 family of cytokines in inflammation and disease*. Annual review of immunology, 2011. **29**: p. 71-109.
37. Huang, J., et al., *ELISpot and ELISA analyses of human IL-21-secreting cells: Impact of blocking IL-21 interaction with cellular receptors*. Journal of immunological methods, 2015. **417**: p. 60-66.
38. Grasse, M., et al., *Booster vaccination against tetanus and diphtheria: insufficient protection against diphtheria in young and elderly adults*. Immunity & Ageing, 2016. **13**(1): p. 26.
39. Kamboj, K.K., et al., *Significant variation in serotype-specific immunogenicity of the seven-valent Streptococcus pneumoniae capsular polysaccharide-CRM197 conjugate vaccine occurs despite vigorous T cell help induced by the carrier protein*. Journal of Infectious Diseases, 2003. **187**(10): p. 1629-1638.
40. Fuery, A., P.C. Richmond, and A.J. Currie, *Human Infant Memory B Cell and CD4+ T Cell Responses to HibMenCY-TT Glyco-Conjugate Vaccine*. PloS one, 2015. **10**(7): p. e0133126.
41. Zhou, M., et al., *The effect of aging on the frequency, phenotype and cytokine production of human blood CD4+ CXCR5+ T follicular helper cells: comparison of aged and young subjects*. Immunity & Ageing, 2014. **11**(1): p. 12.

## Supplementary information

**Supplementary Table 1. Participant characteristics**

Parameter	
BMI (range)	26.1 (18.4 – 35.2)
CMV seropositive (number, %)	16 (51.6%)
<b>Diseases in last year (number, %)</b>	
Diabetes type II	2 (6.5%)
High blood pressure	8 (25.8%)
Vascular diseases	1 (3.2%)
Lung diseases	4 (12.9%)
Rheumatic diseases	2 (6.5%)
Gastro-intestinal diseases	1 (3.2%)
Other diseases	3 (9.7%)
No serious diseases	16 (51.6%)
<b>Medication last 6 months (number, %)</b>	
Medication for infections #	2 (6.5%)
Cholesterol lowering medication	7 (22.6%)
Diabetic medication	2 (6.5%)
Blood pressure lowering medication	8 (25.8%)
Immunosuppressive medication #	0 (0%)
No medication	17 (54.8%)
<b>Infections (number, %)</b>	
Influenza < 4 weeks	1 (3.2%)
Cold <4 weeks	5 (16.1%)
No infection < 4 weeks	25 (80.6%)
<b>Smoking (number, %)</b>	
Cigarette smoking	4 (12.9%)
Cigars, pipe smoking	4 (12.9%)
No smoking	22 (71%)
<b>Physical activity (numbers, %)</b>	
Weekly or more	19 (61.3%)
Less than weekly	5 (16.1%)
No activity	7 (22.6%)

# Medication (mainly corticosteroids and antibiotics) was used more than 3 months ago.



### Supplementary Figure 1.

**a).** Percentages of CXCR5+ cells at the different time point pre- and post- vaccination. The horizontal lines in the graphs represent the geometric mean percentages. The different time points were compared with the Wilcoxon signed rank test preceded by the Friedman test. CXCR5+ and CXCR5- cells at the different time points were compared with the Mann Whitney U test. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ ,  $n = 20$ .

**b).** Gating strategy for  $T_{FH}$ -cells with addition of the PD1 marker. A representative sample is shown. **c).** The correlation between the frequency of CXCR5+ICOS+IL21+PD1+ and CXCR5+ICOS+IL21+ cells 7 days post-vaccination. The correlation was determined by the Spearman's rho correlation.  $n = 12$ .