

University of Groningen

## Challenges of diagnosing glaucoma in myopic eyes

Qiu, Kunliang

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2018

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Qiu, K. (2018). *Challenges of diagnosing glaucoma in myopic eyes: Characteristics and determinants of the anatomical structures relevant to glaucoma*. University of Groningen.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## *Chapter 8*

### **Effect of optic disc-fovea distance on measurements of individual macular intraretinal layers in normal subjects**

**Kunliang Qiu**, Binyao Chen, Haoyu Chen, Enting Gao, Jianling Yang, Xinjian Chen, Mingzhi Zhang

*Retina*. 2018 Feb 27.

## ABSTRACT

**Purposes:** To investigate the effect of optic disc-fovea distance (DFD) on measurements of macular intraretinal layers with spectral-domain optical coherence tomography (SD-OCT) in normal subjects.

**Methods:** 182 eyes from 182 normal subjects were imaged with SD-OCT. The average thicknesses of 8 macular intraretinal layers were measured with an automatic segmentation algorithm. Partial correlation test and multiple regression analysis were used to determine the effect of DFD on thicknesses of intraretinal layers.

**Results:** DFD correlated negatively with the overall average thickness in all of the intraretinal layers ( $r \leq -0.17$ , all  $p \leq 0.025$ ) except the ganglion cell layer and photoreceptor. In multiple regression analysis, greater DFD was associated with thinner nerve fiber layer (6.78  $\mu\text{m}$  decrease per each mm increase in DFD,  $p < 0.001$ ), thinner ganglion cell-inner plexiform layer (2.16  $\mu\text{m}$  decrease per each mm increase in DFD,  $p = 0.039$ ), thinner ganglion cell complex (8.94  $\mu\text{m}$  decrease per each mm increase in DFD,  $p < 0.001$ ), thinner central macular thickness (18.16  $\mu\text{m}$  decrease per each mm increase in DFD,  $p < 0.001$ ) and thinner total macular thickness (15.94  $\mu\text{m}$  decrease per each mm increase in DFD,  $p < 0.001$ ).

**Conclusions:** Thinner measurements of macular intraretinal layers were significantly associated with greater DFD. A clinical assessment of macular intraretinal layers in the evaluation of various macular diseases should always be interpreted in the context of DFD.

## Introduction

Evaluation of macular structure is important and useful in diagnosing and evaluating the efficacy of treatment of various ocular diseases involving macular changes, such as macular edema (Barham et al., 2017). The introduction of optical coherence tomography (OCT) has facilitated the detection and follow up of subtle changes in macular structure quantitatively and reliably (Virgili et al., 2011). Recently, the spectral-domain OCT with faster scan speed and higher resolution has been widely used as an important technology for in vivo measurement of macular structure (Ruia et al., 2016; Lee et al., 2015). Besides, advances in segmentation algorithms have further allowed detailed separation and demarcation of individual intraretinal layers (Garvin et al., 2008; Loduca et al., 2010). Previous studies have demonstrated that thickness measurement of intraretinal layers is becoming a powerful and reliable surrogate marker for assessing and monitoring macular changes resulting from retinal diseases, glaucoma, and optic neuropathy (Huang et al., 2016; Cheng et al., 2016; Nolan et al., 2015; Hood et al., 2017). Thus, understanding of normal macular structure and its variability is essential for assessing optic neuropathy and maculopathy.

Several factors including axial length, gender, age, and disc area have been reported to be associated with thicknesses of individual retinal layers in normal subjects (Demirkaya et al., 2013; Higashide et al., 2016; Ooto et al., 2011; Mwanza et al., 2011). These findings have been valuable in the clinical assessment of glaucoma and various macular diseases. However, our knowledge about the variation of macular intraretinal structure is far from complete. The distance between the optic disc center and the fovea (DFD) is another biometric variable that may influence the macular thickness. Eyes with large DFD may be associated with a stretching of the posterior fundus which may cause a change of the retinal thickness. Moreover, DFD has been reported to be associated with axial length (Jonas et al., 2015). Thus, it is important to determine the effect of DFD on measurements of individual intraretinal layers. However, to our best

knowledge, the relationship between DFD and measurements of individual macular intraretinal layers has not been reported.

The purpose of this study was to determine the effect of DFD on measurements of macular intraretinal layers with an automatic retinal layer segmentation algorithm (Gao et al., 2015) in normal subjects.

## **Methods**

### *Subjects*

In this prospective, cross-sectional observational study, the normal subjects were consecutively recruited from the general clinic of Joint Shantou International Eye Center. All the included subjects received complete ophthalmic examinations including the measurement of best corrected visual acuity, axial length (IOL master; Carl-Zeiss Meditec, Dublin, CA), refraction, intraocular pressure (IOP), fundus examination and slit-lamp biomicroscopy. Each of the included eyes had no other concurrent ocular disease except a refractive error and mild cataract. One eye was randomly selected if both eyes were eligible. Subjects were excluded if refractive errors over +3.0 or under -6.0 diopters (D), the best corrected visual acuity was less than 20/40, the IOP over 21 mmHg, if they had a family history of glaucoma, or if they had a history of intraocular surgery, refractive surgery, macular degeneration, neurological disease, glaucoma, or diabetes. The present study followed the tenets of the declaration of Helsinki and was approved by the local ethical committee with written informed consent obtained from each subject before enrolment.

### *Visual field testing*

Visual field testing was performed with standard automated white-on-white threshold perimetry, using the 24-2 SITA standard strategy (Humphrey Field Analyzer II; Carl Zeiss Meditec, Inc.). Only reliable visual field tests with fixation

loss less than 20% and false positive and negative less than 15% were included in the study. All the included visual field tests were within normal limits in the glaucoma hemifield test and had a pattern standard deviation (PSD) p value > 5%.

### *Optical Coherence Tomography*

All of the included eyes underwent macular and optic disc imaging with the Topcon 3D OCT-2000 (software version 8.11; Topcon). The axial resolution for this spectral-domain OCT is 6  $\mu\text{m}$  and the scan speed is 50,000 A-scans per second (TOPCON CORPORATION 2009). Both the macular 3D Scan 512 $\times$ 128 protocol and Optic Disc 3D Scan 512 $\times$ 128 protocol were performed.


Measurements with eye movements during image acquisition were excluded and retaken. Each of the included images had a minimum image quality score of 45 which is recommended by the manual of 3D OCT-2000.<sup>17</sup> The disc area was recorded for subsequent analysis from the analysis printout of the optic disc scan protocol with the built-in OCT software.

The raw data from the macular scan protocol, which consists of a scan area of 6 $\times$ 6 mm<sup>2</sup> and 128 B-scans (512 A-scans per B-scan), was exported for subsequent thickness measurements of individual macular retinal layers. Each 3D-OCT volumetric macula-centered scan was automatically segmented by a graph search algorithm, which is a fast, three-dimension, automatic graph-theoretical segmentation approach (Gao et al., 2015; Garvin et al., 2009; Shi et al., 2015). With this automatic algorithm, the lowest location of the first surface (internal limiting membrane) in each image was determined and used as a center point of the Early Treatment Diabetic Retinopathy (ETDRS) grid. Subsequently, 11 intraretinal surfaces defining 10 retinal layers were segmented and thicknesses of individual layers were then measured on 9 regions according to the ETDRS grid (Figure 1). For analysis, the average thickness of each layer within 3 concentric rings (Figure 1B) was calculated for the following layers: nerve fiber layer (NFL), ganglion cell layer (GCL), inner plexiform layer (IPL), inner nuclear layer (INL), outer plexiform layer (OPL), outer nuclear layer (ONL), photoreceptor (IS/OS),

and retinal pigment epithelium (RPE). The central ring was a circle with a diameter of 1 mm, centered on the fovea. The pericentral ring is a concentric ring around the central ring with an inner diameter of 1 mm and an outer diameter of 3 mm. The peripheral ring is another concentric ring extending from the edge of the pericentral ring with an outer diameter of 6 mm centered on the fovea. As inner retinal layers are almost absent in the fovea, only measurements of outer retinal layers (OPL, ONL, IS/OS, and RPE) were analyzed in the central ring. Thicknesses of ganglion cell-inner plexiform layer (GCIPL, combined measurement of GCL and IPL), ganglion cell complex (GCC, combined measurement of NFL and GCIPL), and total macular thickness (NFL+GCL+IPL+INL+OPL+ONL+IS/OS) were also calculated for analysis.

#### *Measurement of DFD*

DFD was manually measured on fundus photographs by using ImageJ software (available in the public domain at <http://rsbweb.nih.gov/ij/>; [www.nih.gov](http://www.nih.gov), National Institutes of Health, Bethesda, MD, USA). Firstly, a rectangle was fitted to the height and width of the ONH manually. Two diagonal lines were drawn, and their crossing was considered as the ONH center. Subsequently, DFD was determined, based on the coordinates of the fovea and the center of the optic disc. To determine the measurement repeatability, 30 fundus images from 30 subjects were randomly selected. The DFD in each image was measured by the same observer for two times in two separate occasions.



**Figure 1.** Segmentation of individual intraretinal layers and thickness measurement in three concentric rings on the ETDRS chart.


### *Statistical Analysis*

The statistical analyses were performed by using the SPSS software (ver. 22.0; SPSS Inc, Chicago, IL). The coefficient of variation and intraclass correlation coefficient (ICC) were computed to evaluate the measurement repeatability. Partial correlation test was used to determine the effect of DFD on thicknesses of intraretinal layers after adjusting for other confounders (age, axial length, gender, disc area, and image quality). Stepwise multiple linear regression analysis was performed to evaluate factors associated with the overall measurements of NFL, GCIPL, GCC, and total macular thickness. A p value less than 0.05 was considered statistically significant.

### **Results**

Eleven subjects were excluded because of poor quality of fundus photographs (4 subjects), unreliable visual field tests (5 subjects), and poor OCT scan quality (2

subjects). Finally, 182 eyes from 182 subjects (106 females and 95 right eyes) were included in the analysis. The mean age and DFD were  $43.8 \pm 15.6$  years (range, 20 to 78 year) and  $4.90 \pm 0.29$  mm (range, 3.98 to 5.66 mm), respectively. The coefficient of variation and ICC of DFD measurement were 0.8% (95% confidence interval: 0.5%-0.9%) and 0.98 (95% confidence interval: 0.94–0.99), respectively. Figure 2 displays the distribution of DFD across all subjects. The mean refraction and axial length were  $-0.80 \pm 1.92$  (range, -6.0 to 2.69D) and  $23.63 \pm 1.11$  mm (range, 20.74 to 26.70 mm), respectively. No significant association was detected between axial length/refraction and DFD ( $r = -0.12$ ,  $p = 0.113$  and  $r = 0.11$ ,  $p = 0.160$ , respectively). The mean disc area was  $2.24 \pm 0.39$  mm<sup>2</sup> (range, 1.28 to 3.28 mm<sup>2</sup>). Table 1 summarizes the thickness measurements of the macular intraretinal layers by the automatic segmentation algorithm in the study population.



**Figure 2.** Histogram of DFD of all included eyes.

**Table 1.** Thickness measurements of the individual macular intraretinal layers/combined retinal layers (n=182).

	Mean±SD	Range
<b>NFL</b>		
Overall	37.7±4.3	28.6 to 52.3
Pericentral ring	22.8±1.9	16.8 to 27.8
Peripheral ring	36.7±3.7	29.3 to 48.7
<b>GCL</b>		
Overall	34.6±2.7	27.3 to 41.3
Pericentral ring	55.0±5.7	26.8 to 68.3
Peripheral ring	33.8±3.0	25.9 to 41.9
<b>IPL</b>		
Overall	37.0±2.8	28.3 to 48.4
Pericentral ring	39.1±3.4	30.5 to 47.5
Peripheral ring	40.1±3.2	29.5 to 53.2
<b>INL</b>		
Overall	35.0±2.2	28.1 to 40.6
Pericentral ring	43.0±3.2	34.9 to 53.1
Peripheral ring	36.1±2.4	28.1 to 42.6
<b>OPL</b>		
Overall	26.0±2.1	22.1 to 35.4
center	21.4±7.0	11.4 to 48.1
Pericentral ring	28.8±4.7	21.8 to 48.0
Peripheral ring	26.4±1.9	22.8 to 33.2
<b>ONL</b>		
Overall	77.9±7.0	46.8 to 96.5
center	114.4±11.8	82.9 to 143.8
Pericentral ring	92.8±9.4	67.3 to 116.5
Peripheral ring	76.1±7.3	39.7 to 95.7
<b>IS/OS</b>		
Overall	29.5±2.9	20.5 to 40.6
center	32.8±3.2	16.4 to 41.1
Pericentral ring	27.9±2.9	17.2 to 41.7
Peripheral ring	29.1±3.4	20.5 to 41.6
<b>RPE</b>		
Overall	36.7±2.5	31.2 to 43.4
center	41.1±3.9	28.1 to 41.1
Pericentral ring	40.5±3.4	32.0 to 47.7
Peripheral ring	36.6±3.1	30.0 to 36.6
<b>GCIPL</b>		
Overall	71.6±4.5	59.0 to 83.5
Pericentral ring	94.1±6.9	65.2 to 108.0
Peripheral ring	73.9±5.2	59.3 to 86.9

**Continued Table 1.** Thickness measurements of the individual macular intraretinal layers/combined retinal layers (n=182).

	Mean±SD	Range
<b>GCC</b>		
Overall	109.3±7.1	92.3 to 128.7
Pericentral ring	116.9±8.0	84.0 to 132.9
Peripheral ring	110.5±7.1	93.2 to 126.9
<b>Total macular thickness</b>		
Overall	262.2±13.5	216.1 to 290.7
Center	216.8±17.1	166.4 to 272.7
Pericentral ring	293.8±15.5	245.6 to 330.7
Peripheral ring	262.7±14.0	212.9 to 291.4

Table 2 demonstrates the associations between DFD and measurements of individual intraretinal layers/combined retinal layers in three different rings, adjusted for age, axial length, image quality, gender and disc area. For individual intraretinal layers, DFD correlated significantly and negatively with the overall average thickness in all of the retinal layers (all  $p \leq 0.025$ ) except the GCL and IS/OS. In the central ring, there was a significant and negative correlation between DFD and ONL ( $r = -0.15$ ,  $p = 0.041$ ). In the pericentral ring, DFD correlated negatively with NFL, IPL, INL, and ONL (all  $p \leq 0.065$ ). In the peripheral ring, DFD correlated significantly and negatively with NFL, IPL, OPL, ONL and RPE (all  $p \leq 0.023$ ). No significant relationship between GCL, IS/OS and DFD was detected in all three rings. Figure 3 shows the correlation between DFD and the overall average thickness measurements of the individual intraretinal layers. For the combined retinal layers, there were significant and negative correlations between GCIPL, GCC, total macular thickness and DFD in both regional and overall measurements (all  $p \leq 0.045$ ).

**Table 2.** Associations between DFD and individual intraretinal layers/combined retinal layers, adjusting for axial length, gender, age, disc area, and image quality (Partial correlation test, n=182).


	r	p
<b>NFL</b>		
Overall	-0.51	<0.001
Pericentral ring	-0.31	<0.001
Peripheral ring	-0.47	<0.001
<b>GCL</b>		
Overall	-0.02	0.774
Pericentral ring	-0.11	0.131
Peripheral ring	0.05	0.493
<b>IPL</b>		
Overall	-0.22	0.003
Pericentral ring	-0.27	<0.001
Peripheral ring	-0.27	<0.001
<b>INL</b>		
Overall	-0.26	<0.001
Pericentral ring	-0.26	<0.001
Peripheral ring	-0.23	0.002
<b>OPL</b>		
Overall	-0.24	0.002
center	-0.08	0.306
Pericentral ring	-0.11	0.140
Peripheral ring	-0.28	<0.001
<b>ONL</b>		
Overall	-0.17	0.025
center	-0.15	0.041
Pericentral ring	-0.14	0.065
Peripheral ring	-0.17	0.023
<b>IS/OS</b>		
Overall	-0.06	0.426
center	-0.09	0.241
Pericentral ring	-0.11	0.138
Peripheral ring	0.10	0.169

**Continued Table 2.** Associations between DFD and individual intraretinal layers/combined retinal layers, adjusting for axial length, gender, age, disc area, and image quality (Partial correlation test, n=182).

	r	p
<b>RPE</b>		
Overall	-0.24	0.001
Center	-0.08	0.307
Pericentral ring	-0.06	0.431
Peripheral ring	-0.24	0.001
<b>GCIPL</b>		
Overall	-0.16	0.039
Pericentral ring	-0.23	0.002
Peripheral ring	-0.15	0.045
<b>GCC</b>		
Overall	-0.38	<0.001
Pericentral ring	-0.27	<0.001
Peripheral ring	-0.34	<0.001
<b>Total macular thickness</b>		
Overall	-0.38	<0.001
Center	-0.31	<0.001
Pericentral ring	-0.34	<0.001
Peripheral ring	-0.32	<0.001

Table 3 presents the multiple linear regression analysis regarding the associations between various factors and the overall average thickness of NFL, GCIPL, GCC, central macular thickness, and total macular thickness. Greater DFD was independently and significantly associated with thinner NFL (6.78  $\mu\text{m}$  decrease per each mm increase in DFD,  $p<0.001$ ), thinner GCIPL (2.16  $\mu\text{m}$  decrease per each mm increase in DFD,  $p=0.039$ ), thinner GCC (8.94  $\mu\text{m}$  decrease per each mm increase in DFD,  $p<0.001$ ), thinner central macular thickness (18.16  $\mu\text{m}$

decrease per each mm increase in DFD,  $p < 0.001$ ) and thinner total macular thickness (15.94  $\mu\text{m}$  decrease per each mm increase in DFD,  $p < 0.001$ ).



**Figure 3.** Scatter plots of DFD versus overall average thickness of individual intraretinal layers/combined retinal layers. DFD and NFL overall average thickness (A); DFD and GCL overall average thickness (B); DFD and IPL overall average thickness (C); DFD and INL overall average thickness (D); DFD and OPL overall average thickness (E); DFD and ONL overall average thickness (F); DFD and IS/OS overall average thickness (G); DFD and RPE overall average thickness (H); DFD and total macular thickness (I).

**Table 3. Factors associated with overall average thickness of NFL, GCIPL, GCC, central macular, and total macular (Stepwise multiple linear regression analysis, n=182)**

	Overall NFL		Overall GCIPL		Overall GCC		Central macular thickness		Overall total macular thickness	
	$\beta$	p	$\beta$	p	$\beta$	p	$\beta$	p	$\beta$	p
DFD (mm)	-6.78	<0.001	-2.16	0.041	-8.97	<0.001	-18.16	<0.001	-15.94	<0.001
Axial length (mm)	1.40	<0.001	-1.56	<0.001	-	-	2.88	0.007	-3.65	<0.001
Age (per year)	-	-	-0.13	<0.001	-0.16	<0.001	-	-	-0.42	<0.001
Gender	2.45	<0.001	-	-	-	-	-6.57	0.006	-3.66	0.034
Disc area (mm <sup>2</sup> )	-	-	2.17	0.010	3.06	0.015	-	-	-	-
Image quality	0.18	0.005	-	-	-	-	-	-	-	-

## Discussion

The present study was performed to determine the effect of DFD on measurements of macular intraretinal layers in normal subjects. We demonstrated that thickness measurements of several individual retinal layers varied significantly with DFD. Thinner measurements of NFL, IPL, INL, OPL, ONL, RPE, GCIPL, GCC, and total macular thickness were significantly associated with greater DFD, independent of other covariates. Such findings are of potential significance in clinical evaluation of macular structural measurements obtained with OCT.

Variations about the thicknesses of individual intraretinal layers have been described previously (Demirkaya et al., 2013; Higashide et al., 2016; Ooto et al., 2011; Mwanza et al., 2011; Szigeti et al., 2015; Liu et al., 2015). Factors including age, axial length, gender, and disc area have been reported to be associated with thickness measurements of individual macular retinal layers (Demirkaya et al., 2013; Higashide et al., 2016; Ooto et al., 2011; Mwanza et al., 2011; Szigeti et al., 2015; Liu et al., 2015). To our best knowledge, however, the effect of DFD on thicknesses of macular intraretinal layers has not been studied. In the current study, negative relationships between DFD and the overall average thickness measurements of NFL, IPL, INL, OPL, ONL, and RPE were detected after adjusting for age, axial length, gender, and disc area. Our results suggest that DFD is one of the important factors determining the thickness measurements of macular intraretinal layers.

Why is DFD associated with measurements of macular intraretinal layers? In regard to macular NFL thickness, one possible explanation is the different scan area for the NFL measurement. As the fovea is farther away from the optic disc in eyes with a greater DFD, the OCT scan area (centered at the fovea) is farther away from the optic disc. Previous studies have demonstrated that the RNFL is thinner farther from the optic disc than it is closer to the optic disc margin (Gabriele et al., 2007). Therefore, one would expect to find thinner NFL

measurement in eyes with greater DFD. With respect to thicknesses of IPL, INL, OPL, ONL, and RPE, a possible explanation is the stretching of the posterior fundus in eyes with a greater DFD. The DFD has been found to be associated with the parapapillary zones (Jonas et al., 2015) and the peripapillary retinal nerve fiber distribution in healthy eyes (Hong et al., 2010). In the current study, we found that measurements of several intraretinal layers significantly decreased in eyes with greater DFD. On the basis of these findings, we speculate that the posterior fundus in eyes with greater DFD are stretched which may cause the decrease of thicknesses of the macular intraretinal layers. The relationship between DFD and the intraretinal layers may be established during the development of the retina.

The current results have potential clinical significance in the evaluation of glaucoma, optic neuropathy and retina disease involving macular changes. For example, evaluation of NFL, GCIPL, GCC, central macular thickness (central subfield thickness), and total retinal thickness in macular region has been reported to be useful in clinical evaluation of glaucoma, multiple sclerosis, Alzheimer's disease, and various maculopathy (Nolan et al., 2015; Cunha et al., 2016; Balk et al., 2016; Hood et al., 2017). However, individual variability of thicknesses of intraretinal layers may limit their use in clinical practice. In a systematic review, it is reported that central retinal thickness (with cutoffs ranging between 230  $\mu\text{m}$  and 300  $\mu\text{m}$ ) measured with OCT is not sensitive enough (0.81) nor specific enough (0.85) to detect the central type of clinically significant macular oedema (Virgili et al., 2015). Therefore, it is important to understand the variability of normal macular structure. Consistent with previous studies (Hong et al., 2010; Jonas et al., 2015), significant interindividual variation of DFD was observed in the current study population. More importantly, we found that DFD was significantly associated with thickness measurements of several macular intraretinal layers. Extrapolation from the regression analysis indicated that average NFL thickness decreases by 18.0% (6.78  $\mu\text{m}$ ) per mm of greater DFD; average GCC thickness decreases by 8.2% (8.94  $\mu\text{m}$ ) per mm of greater DFD; central macular thickness decreases by 8.4% (18.16  $\mu\text{m}$ ) per mm of greater DFD;

and average macular thickness decreases by 6.1% (15.94  $\mu\text{m}$ ) per mm of greater DFD. The data and results were adjusted for other confounders including axial length, age, gender, image quality, and disc area. Our current findings indicate that a clinical assessment of thickness measurements of macular retinal layers should take into consideration of DFD. Moreover, by using the OCT built-in manual measurement tool, it would not be difficult for a clinician to obtain DFD measurements on the OCT fundus image (Lujan et al., 2008).

In the present study, segmentation and measurements of macular intraretinal layers were performed by using an automatic graph search algorithm which has been validated in previous studies (Gao et al., 2015; Garvin et al., 2009). Associations between GCIPL thickness, GCC thickness, total macular thickness and axial length have been reported previously (Mwanza et al., 2011; Koh et al., 2012; Kim et al., 2011; Takeyama et al., 2014; Zhao et al., 2013; Hirasawa et al., 2013; Lam et al., 2007). Consistent with most of the previous studies, we found that thinner GCIPL was associated with longer axial length (Mwanza et al., 2011; Kim et al., 2011; ). In line with previous studies (Lam et al., 2007), regional variations of association between macular thickness and axial length were observed in the current study. Conflicting data regarding the relationship between axial length and GCC thickness has been reported (Kim et al., 2011; Takeyama et al., 2014; Zhao et al., 2013; Hirasawa et al., 2013).<sup>29-32</sup> Kim et al. (2011) reported that GCC thickness did not correlate with axial length. However, others reported that GCC thickness varied significantly with axial length (Takeyama et al., 2014; Zhao et al., 2013; Hirasawa et al., 2013). In the present study, no significant relationship between axial length and GCC thickness was detected. Several differences in study design could have contributed to these conflicting results, such as adjustment for different covariates and different study populations (inclusion of high myopic eyes or not).

The effect of ocular magnification has been reported previously (Leung et al., 2007; Higashide et al., 2016). According to previous reports, OCT measurements, OCT scan area, and measurements on fundus photos could be different due to

ocular magnification (Leung et al., 2007). As only the uncorrected measurements are available in a clinical setting with commercial devices and software, we decided to perform the analysis with and without correction for ocular magnification. By using Bennett's formula (Bennett et al., 1994), the actual measurements of DFD and disc area were calculated for analysis. We found significant but minor difference between corrected and uncorrected DFD measurements (4.90 vs 4.73 mm,  $p < 0.001$ ). We then repeated the analyses using the magnification corrected measurements. In the partial correlation analysis, we found similar pattern of associations regarding the relationship between DFD and measurements of individual macular intraretinal layers (Table 4). For the macular scan area, unfortunately, the actual scan area is not possible to obtain in the present study. However, to reduce this effect, analyses were repeated in a subgroup of eyes with a narrow range of axial length (25 to 75 percentiles, 22.88 to 24.33mm). Similar and stronger correlations were observed between DFD and thickness measurements of macular intraretinal layers (Table 5). On the basis of these findings, we believe that greater DFD is indeed associated with thinner thickness measurements of macular intraretinal layers.

The current study had some limitations. First, only ethnic Chinese were evaluated in the present study, and the findings may vary in other ethnic groups. Second, the measurement of DFD which was based on the two-dimensional images may be underestimated as some of the eyes may have a steeper posterior retinal curvature than emmetropic eyes. To minimize this effect, we excluded high myopic (refraction  $\leq -6D$ ) eyes in the present study (mean refraction:  $0.80 \pm 1.93D$ ). All the included eyes were reviewed carefully and eyes with myopic macular degeneration including a posterior staphyloma were excluded from the study. Moreover, a similar and stronger pattern of correlations was detected regarding the relationship between DFD and macular intraretinal layers in a subgroup of eyes with a narrow range of axial length (Table 5, axial length range: 22.88 to 24.33mm).

In conclusion, thinner measurements of NFL, IPL, INL, OPL, ONL, RPE, GCIPL, GCC, and total macular thickness were significantly associated with greater DFD, independent of other covariates. A clinical assessment of thickness measurements of macular intraretinal layers in the evaluation of glaucoma, optic neuropathy and retina disease involving macular changes should always be interpreted in the context of DFD.

**Table 4.** Associations between magnification corrected DFD and individual intraretinal layers/combined retinal layers, adjusting for axial length, gender, age, magnification corrected disc area, and image quality (Partial correlation test, n=182).

	r	p
<b>NFL</b>		
Overall	-0.50	<0.001
Pericentral ring	-0.31	<0.001
Peripheral ring	-0.47	<0.001
<b>GCL</b>		
Overall	-0.02	0.806
Pericentral ring	-0.11	0.138
Peripheral ring	0.06	0.464
<b>IPL</b>		
Overall	-0.21	0.005
Pericentral ring	-0.26	<0.001
Peripheral ring	-0.26	<0.001
<b>INL</b>		
Overall	-0.25	<0.001
Pericentral ring	-0.26	<0.001
Peripheral ring	-0.22	0.003
<b>OPL</b>		
Overall	-0.23	0.002
center	-0.08	0.325
Pericentral ring	-0.11	0.163
Peripheral ring	-0.28	<0.001

**Continued Table 4.** Associations between magnification corrected DFD and individual intraretinal layers/combined retinal layers, adjusting for axial length, gender, age, magnification corrected disc area, and image quality (Partial correlation test, n=182).

	r	p
<b>ONL</b>		
Overall	-0.17	0.028
center	-0.16	0.040
Pericentral ring	-0.14	0.063
Peripheral ring	-0.17	0.028
<b>IS/OS</b>		
Overall	-0.06	0.419
center	-0.09	0.262
Pericentral ring	-0.11	0.148
Peripheral ring	0.10	0.170
<b>RPE</b>		
Overall	-0.24	0.001
Center	-0.08	0.308
Pericentral ring	-0.06	0.437
Peripheral ring	-0.24	0.001
<b>GCIPL</b>		
Overall	-0.15	0.048
Pericentral ring	-0.18	0.019
Peripheral ring	-0.28	<0.001
<b>GCC</b>		
Overall	-0.38	<0.001
Pericentral ring	-0.27	<0.001
Peripheral ring	-0.34	<0.001
<b>Total macular thickness</b>		
Overall	-0.37	<0.001
Center	-0.30	<0.001
Pericentral ring	-0.33	<0.001
Peripheral ring	-0.31	<0.001

**Table 5.** Associations between DFD and individual intraretinal layers/combined retinal layers, adjusting for axial length, gender, age, disc area, and image quality (Partial correlation test in a subgroup of eyes with axial length ranging from 22.88 to 24.33mm, n=93).

	r	p
<b>NFL</b>		
Overall	-0.58	<0.001
Pericentral ring	-0.36	<0.001
Peripheral ring	-0.53	<0.001
<b>GCL</b>		
Overall	-0.17	0.104
Pericentral ring	-0.20	0.059
Peripheral ring	-0.13	0.230
<b>IPL</b>		
Overall	-0.29	0.005
Pericentral ring	-0.28	0.009
Peripheral ring	-0.31	0.003
<b>INL</b>		
Overall	-0.40	<0.001
Pericentral ring	-0.39	<0.001
Peripheral ring	-0.38	0.003
<b>OPL</b>		
Overall	-0.26	0.016
center	-0.08	0.465
Pericentral ring	-0.09	0.407
Peripheral ring	-0.32	0.003
<b>ONL</b>		
Overall	-0.23	0.033
center	-0.25	0.020
Pericentral ring	-0.22	0.036
Peripheral ring	-0.23	0.031
<b>IS/OS</b>		
Overall	0.06	0.585
center	-0.20	0.057
Pericentral ring	-0.16	0.137
Peripheral ring	0.15	0.179

**Continued Table 5.** Associations between DFD and individual intraretinal layers/combined retinal layers, adjusting for axial length, gender, age, disc area, and image quality (Partial correlation test in a subgroup of eyes with axial length ranging from 22.88 to 24.33mm, n=93).

	r	p
<b>RPE</b>		
Overall	-0.35	0.001
Center	-0.06	0.572
Pericentral ring	-0.13	0.247
Peripheral ring	-0.37	<0.001
<b>GCIPL</b>		
Overall	-0.29	0.006
Pericentral ring	-0.27	0.011
Peripheral ring	-0.42	<0.001
<b>GCC</b>		
Overall	-0.51	<0.001
Pericentral ring	-0.36	<0.001
Peripheral ring	-0.46	<0.001
<b>Total macular thickness</b>		
Overall	-0.49	<0.001
Center	-0.41	<0.001
Pericentral ring	-0.46	<0.001
Peripheral ring	-0.43	<0.001

## References

- Balk LJ, Cruz-Herranz A, Albrecht P, et al. Timing of retinal neuronal and axonal loss in MS: a longitudinal OCT study. *J Neurol*. 2016; 263(7):1323-1331.
- Barham R, El Rami H, Sun JK, Silva PS. Evidence-Based Treatment of Diabetic Macular Edema. *Semin Ophthalmol*. 2017;32(1):56-66.
- Bennett AG, Rudnicka AR, Edgar DF. Improvements on Littmann's method of determining the size of retinal features by fundus photography. *Graefes Arch Clin Exp Ophthalmol*. 1994; 232:361-367.
- Cheng D, Wang Y, Huang S, et al. Macular Inner Retinal Layer Thickening and Outer Retinal Layer Damage Correlate With Visual Acuity During Remission in Behcet's Disease. *Invest Ophthalmol Vis Sci*. 2016; 57(13):5470-5478.
- Cunha LP, Lopes LC, Costa-Cunha LV, et al. Macular Thickness Measurements with Frequency Domain-OCT for Quantification of Retinal Neural Loss and its Correlation with Cognitive Impairment in Alzheimer's Disease. *PLoS One*. 2016; 11(4):e0153830.
- Demirkaya N, van Dijk HW, van Schuppen SM, et al. Effect of age on individual retinal layer thickness in normal eyes as measured with spectral-domain optical coherence tomography. *Invest Ophthalmol Vis Sci*. 2013; 54(7):4934-4940.
- Gabriele ML, Ishikawa H, Wollstein G, et al. Peripapillary nerve fiber layer thickness profile determined with high speed, ultrahigh resolution optical coherence tomography high-density scanning. *Invest Ophthalmol Vis Sci*. 2007; 48(7):3154-3160.
- Gao E, Chen B, Yang J, et al. Comparison of Retinal Thickness Measurements between the Topcon Algorithm and a Graph-Based Algorithm in Normal and Glaucoma Eyes. *PLoS One*. 2015; 10(6):e0128925.
- Garvin MK, Abramoff MD, Kardon R, et al. Intraretinal layer segmentation of macular optical coherence tomography images using optimal 3-D graph search. *IEEE Trans Med Imaging*. 2008; 27:1495-1505.
- Garvin MK, Abramoff MD, Wu X, et al. Automated 3-D intraretinal layer segmentation of macular spectral-domain optical coherence tomography images. *IEEE Trans Med Imaging*. 2009; 28(9):1436-1447.
- Higashide T, Ohkubo S, Hangai M, et al. Influence of Clinical Factors and Magnification Correction on Normal Thickness Profiles of Macular Retinal

Layers Using Optical Coherence Tomography. PLoS One. 2016; 11(1):e0147782.

Hirasawa K, Shoji N. Association between ganglion cell complex and axial length. Jpn J Ophthalmol. 2013; 57(5):429-434.

Hong SW, Ahn MD, Kang SH, Im SK. Analysis of peripapillary retinal nerve fiber distribution in normal young adults. Invest Ophthalmol Vis Sci. 2010; 51(7):3515-3523.

Hood DC. Improving our understanding, and detection, of glaucomatous damage: An approach based upon optical coherence tomography (OCT). Prog Retin Eye Res. 2017; 57:46-75.

Huang S, Chen Q, Ma Q, et al. THREE-DIMENSIONAL CHARACTERISTICS OF FOUR MACULAR INTRARETINAL LAYER THICKNESSES IN SYMPTOMATIC AND ASYMPTOMATIC CARRIERS OF G11778A MUTATION WITH LEBER'S HEREDITARY OPTIC NEUROPATHY. Retina. 2016; 36(12):2409-2418.

Jonas RA, Wang YX, Yang H, et al. Optic Disc-Fovea Distance, Axial Length and Parapapillary Zones. The Beijing Eye Study 2011. PLoS One. 2015; 10(9):e0138701.

Kim NR, Kim JH, Lee J, et al. Determinants of perimacular inner retinal layer thickness in normal eyes measured by Fourier-domain optical coherence tomography. Invest Ophthalmol Vis Sci. 2011; 52(6):3413-3418.

Koh VT, Tham YC, Cheung CY, et al. Determinants of ganglion cell-inner plexiform layer thickness measured by high-definition optical coherence tomography. Invest Ophthalmol Vis Sci. 2012; 53(9):5853-5859.

Lam DS, Leung KS, Mohamed S, et al. Regional variations in the relationship between macular thickness measurements and myopia. Invest Ophthalmol Vis Sci. 2007; 48(1):376-382.

Lee HJ, Kim MS, Jo YJ, Kim JY. Ganglion Cell-Inner Plexiform Layer Thickness in Retinal Diseases: Repeatability Study of Spectral-Domain Optical Coherence Tomography. Am J Ophthalmol. 2015;160(2):283-289.e1.

Leung CK, Cheng AC, Chong KK. Optic disc measurements in myopia with optical coherence tomography and confocal scanning laser ophthalmoscopy. Invest Ophthalmol Vis Sci. 2007; 48:3178-3183.

Liu X, Shen M, Yuan Y, et al. Macular Thickness Profiles of Intraretinal Layers in Myopia Evaluated by Ultrahigh-Resolution Optical Coherence Tomography. Am J Ophthalmol. 2015; 160(1):53-61.e2.

Loduca AL, Zhang C, Zelkha R, Shahidi M. Thickness mapping of retinal layers by spectral-domain optical coherence tomography. *Am J Ophthalmol*. 2010; 150(6):849-855.

Lujan BJ, Wang F, Gregori G, et al. Calibration of fundus images using spectral domain optical coherence tomography. *Ophthalmic Surg Lasers Imaging*. 2008; 39: S15-20.

Mwanza JC, Durbin MK, Budenz DL, et al. Profile and predictors of normal ganglion cell-inner plexiform layer thickness measured with frequency-domain optical coherence tomography. *Invest Ophthalmol Vis Sci*. 2011; 52(11):7872-7879.

Nolan RC, Narayana K, Galetta SL, Balcer LJ. Optical Coherence Tomography for the Neurologist. *Semin Neurol*. 2015; 35(5):564-577.

Ooto S, Hangai M, Tomidokoro A, et al. Effects of age, sex, and axial length on the three-dimensional profile of normal macular layer structures. *Invest Ophthalmol Vis Sci*. 2011; 52(12):8769-8779.

Ruia S, Saxena S, Gemmy Cheung CM, et al. Spectral Domain Optical Coherence Tomography Features and Classification Systems for Diabetic Macular Edema: A Review. *Asia Pac J Ophthalmol (Phila)*. 2016;5(5):360-367.

Shi F, Chen X, Zhao H, et al. Automated 3-D retinal layer segmentation of macular optical coherence tomography images with serous pigment epithelial detachments. *IEEE Trans Med Imaging*. 2015; 34:441-452

Szigeti A, Tátrai E, Varga BE, et al. The Effect of Axial Length on the Thickness of Intraretinal Layers of the Macula. *PLoS One*. 2015; 10(11):e0142383.

Takeyama A, Kita Y, Kita R, Tomita G. Influence of axial length on ganglion cell complex (GCC) thickness and on GCC thickness to retinal thickness ratios in young adults. *Jpn J Ophthalmol*. 2014; 58(1):86-93.

TOPCON CORPORATION. 3D OPTICAL COHERENCE TOMOGRAPHY(3D OCT-2000) INSTRUCTION MANUAL, Tokyo, Japan; 2009 version (2009.11-100TH )

Virgili G, Menchini F, Casazza G, et al. Optical coherence tomography (OCT) for detection of macular oedema in patients with diabetic retinopathy. *Cochrane Database Syst Rev*. 2015 Jan 7;1:CD008081.

Virgili G, Menchini F, Murro V, et al. Optical coherence tomography (OCT) for detection of macular oedema in patients with diabetic retinopathy. *Cochrane Database Syst Rev*. 2011; 6(7):CD008081.

Zhao Z, Jiang C. Effect of myopia on ganglion cell complex and peripapillary retinal nerve fibre layer measurements: a Fourier-domain optical coherence tomography study of young Chinese persons. *Clin Exp Ophthalmol*. 2013; 41(6):561-566.