

University of Groningen

Putting a Hold on the Downward Spiral of Paranoia in the Social World

Collip, Dina; Geschwind, Nicole; Peeters, Frenk; Myin-Germeys, Inez; van Os, Jim; Wichers, Marieke

Published in:
 PLoS ONE

DOI:
[10.1371/journal.pone.0066747](https://doi.org/10.1371/journal.pone.0066747)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Collip, D., Geschwind, N., Peeters, F., Myin-Germeys, I., van Os, J., & Wichers, M. (2013). Putting a Hold on the Downward Spiral of Paranoia in the Social World: A Randomized Controlled Trial of Mindfulness-Based Cognitive Therapy in Individuals with a History of Depression. *PLoS ONE*, 8(6), [e66747]. <https://doi.org/10.1371/journal.pone.0066747>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Putting a Hold on the Downward Spiral of Paranoia in the Social World: A Randomized Controlled Trial of Mindfulness-Based Cognitive Therapy in Individuals with a History of Depression

Dina Collip^{1*}, Nicole Geschwind^{1,2,3}, Frenk Peeters¹, Inez Myin-Germeys¹, Jim van Os^{1,3}, Marieke Wichers¹

1 Department of Psychiatry and Psychology, School of Mental Health and Neuroscience, Maastricht University Medical Centre, Maastricht, The Netherlands, **2** Health Psychology Group, CLEP, University of Leuven, Leuven, Belgium, **3** Department of Psychosis Studies, Institute of Psychiatry, King's Health Partners, King's College London, London, United Kingdom

Abstract

Context: Paranoia embodies altered representation of the social environment, fuelling altered feelings of social acceptance leading to further mistrust. Mindfulness-based cognitive therapy (MBCT) may relieve paranoia and reduce its impact on social acceptance.

Objective: To determine whether MBCT alters momentary feeling of paranoia and social acceptance in daily life.

Design: Randomized controlled trial of daily-life repeated measures (up to 120 per participant) before and after allocation to MBCT or waiting list control.

Participants: Volunteer sample of 130 eligible men and women with residual affective dysregulation after at least one episode of major depressive disorder.

Interventions: Eight weeks of MBCT in groups of 10–15 participants in addition to participants' usual treatment.

Outcome Measures: Daily-life ratings of paranoia and social acceptance. This manuscript concerns additional analyses of the original trial; hypotheses were developed after data collection (focus initially on depressive symptoms) but before data analysis.

Results: Sixty-six participants were assigned to the waiting list control group and 64 to the MBCT intervention group, of whom 66 and 61 respectively were included in the per-protocol analyses. Intention-to-treat analyses revealed a significant group by time interaction in the model of momentary paranoia ($b = -.18$, $p < 0.001$, $d = -0.35$) and social acceptance ($b = .26$, $p < 0.001$, $d = 0.41$). Paranoia levels in the intervention group were significantly reduced ($b = -.11$, $p < 0.001$) and feelings of social acceptance significantly increased ($b = .18$, $p < 0.001$), whereas in the Control condition a significant increase in paranoia ($b = .07$, $p = 0.008$) and a decrease in social acceptance was apparent ($b = -.09$, $p = 0.013$). The detrimental effect of paranoia on social acceptance was significantly reduced in the MBCT, but not the control group (group by time interaction: $b = .12$, $p = 0.022$).

Conclusions: MBCT confers a substantial benefit on subclinical paranoia and may interrupt the *social* processes that maintain and foster paranoia in individuals with residual affective dysregulation.

Trial Registration: Netherlands Trial Register NTR1084

Citation: Collip D, Geschwind N, Peeters F, Myin-Germeys I, van Os J, et al. (2013) Putting a Hold on the Downward Spiral of Paranoia in the Social World: A Randomized Controlled Trial of Mindfulness-Based Cognitive Therapy in Individuals with a History of Depression. PLoS ONE 8(6): e66747. doi:10.1371/journal.pone.0066747

Editor: Eiji Shimizu, Chiba University Graduate School of Medicine, Japan

Received: October 9, 2012; **Accepted:** May 8, 2013; **Published:** June 27, 2013

Copyright: © 2013 Collip et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: MW was supported by the Dutch Organisation for Scientific Research (NWO, VENI Grant Nr. 916.76.147). DC was supported by a Kootstra Talent Fellowship from Maastricht University. NG was supported by K.U. Leuven Center of Excellence on Generalization Research (GRIP*TT; PF/10/005). All above mentioned funding sources had no role in data collection, analysis, or interpretation, trial design, patient recruitment, or any aspect pertinent to the study nor in the writing of the manuscript or the decision to submit it for publication. None of the authors were paid to write this article by a pharmaceutical company or other agency. The authors had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Competing Interests: FP has received financial compensation as an independent symposium speaker from GlaxoSmithKline, Wyeth, Astra Zeneca, Lundbeck, Eli Lilly, Servier and Janssen-Cilag. JVO is/has been an unrestricted research grant holder with or has received financial compensation as an independent symposium speaker from Eli Lilly, BMS, Lundbeck, Organon, Janssen-Cilag, GlaxoSmithKline, Astra Zeneca, Pfizer and Servier, companies that have an interest in the treatment of mental health disorders. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

* E-mail: d.collip@maastrichtuniversity.nl

These authors contributed equally to this work.

Introduction

Humans have always been social animals. The social world is an important source of satisfaction and support for most of us. However, social contact can also be a source of suffering and problems. Individuals with a diagnosis of mental disorder often have difficulties to adapt to the social world and to trust others [1]. Although representations of social mistrust are common also in the general population, they generally take on more severe forms in the context of mental disorders [2]. Two important factors which fuel paranoia are low self-esteem [3], expectation of mistreatment and affective dysregulation [4,5], illustrating the transdiagnostic nature of paranoia. Paranoia consequently is not only commonly observed in individuals with psychosis but also in individuals with affective disorders such as depression, [5,6]. The presence of paranoid symptoms impacts negatively on outcome even when expressed at subclinical level [7], and increases the risk for developing a full-blown psychotic disorder [8]. Paranoia affects representations of the social environment, resulting in unease in social situations and reduced social approach. The resulting social avoidance will further fuel mistrust, giving rise to a downward spiral [9]. Recent research proposes that mindfulness-based approaches, for instance Mindfulness-Based Cognitive Therapy (MBCT) and Acceptance and Commitment Therapy (ACT), are effective in treating different forms of psychopathology. Most studies were conducted in patients with depression or anxiety disorder [10,11], but a few small randomized controlled trials have reported positive effects of ACT and MBCT for patients recovering from acute psychosis [12,13,14,15,16,17,18]. There is little work on possible psychological processes that might be transformed by of mindfulness-based therapies [19,20], particularly in terms of examining alterations in the subtle dynamic interplay of altered representations and social interaction in real life, for example when people interact with others, or when they feel suspicious. Real-time and real-world assessment approaches such as the Experience Sampling Method (ESM) have been optimized for the investigation of dynamic patterns of interaction between experience and behaviour [21].

Mindfulness-based approaches, unlike cognitive therapy, are not aimed at changing the contents of mental experience, but rather target the relationship the individual has with thoughts and feelings [22]. Participants practice to become more aware of moment-to-moment sensations and to recognize negative feelings/thoughts, without being *wrapped up* in them or considering them to be a reflection of reality. This practice may bring about the ability to deal with feelings of insecurity and paranoia.

In a sample of participants with affective dysregulation and increased risk to experience feelings of mistrust, ESM was used to measure feelings of paranoia and appraisal of social interactions during six days before and six days after either MBCT or waiting list control group (hereafter: Control). This study used data pertaining to a published RCT on the effects of MBCT on depressive symptoms [23], to test further hypotheses, developed after completion of the RCT, regarding the effects of MBCT on paranoia. The primary aim of the study was to examine the effects of MBCT on feelings of paranoia and on the appraisal of social situations, measured as they occur in daily life. A secondary aim was to investigate whether MBCT may interrupt the downward spiral between feeling paranoid and feeling less accepted in social situations, using time-lagged analyses.

The hypothesis was that, following MBCT, individuals would feel (i) less paranoid and (ii) more at ease during social interaction. Furthermore, following MBCT, we expected (iii) a reduction of the

association between paranoia and unease during social interactions.

Methods

Participant Characteristics

In the MindMaastricht study (trial number NTR1084, Netherlands Trial Register Trial registration: <http://www.trialregister.nl/trialreg/admin/rctview.asp?TC=1084>) [23], adults with residual depressive symptomatology (≥ 7 on the 17-item Hamilton Depression Rating Scale; HDRS [24]) after at least one episode of Major Depressive Disorder (MDD) and thus at risk of also displaying elevated paranoid ideation [7], were recruited from outpatient mental health care facilities in Maastricht and through posters in public spaces. As the focus was on stable residual symptomatology in the context of a previous affective disorder, individuals were excluded if they met SCID criteria for a current depressive episode, schizophrenia, or any psychotic disorder in the past year, and recent (past four weeks) or upcoming changes in ongoing psychological or pharmacological treatment. Sample sociodemographic and clinical characteristics are displayed in Table 1.

Sampling procedures

All study procedures were approved by the Medical Ethics Committee of Maastricht University Medical Centre, and all participants signed an informed consent form. The protocol for this trial and supporting CONSORT checklist are available as supporting information; see Checklist S1 and Protocol S1. An initial screening was performed by phone to check for availability during the study period and likelihood of meeting inclusion and exclusion criteria. A second screening included administration of the Structured Clinical Interview for DSM IV axis I (SCID-I) [25] and the 17-item HDRS by trained psychologists. Eligible participants were invited for a detailed explanation of the experience sampling procedure, and then took part in the baseline assessment. The baseline assessment consisted of six days of experience sampling in the participant's natural environment (see below). After the baseline assessment, participants were randomized to either MBCT or waiting list (allocation ratio 1:1). After either eight weeks of MBCT (see below) or equivalent waiting time (Control condition), participants again took part in six days of experience sampling. All participants were compensated with gift vouchers worth 50 Euros. Participants in the Control condition had the opportunity to take part in MBCT after the post-intervention assessment.

Randomization

Randomization to treatment condition was stratified according to number of depressive episodes (two or less versus three or more), as at the time of randomization studies suggested a greater benefit for those with three or more previous episodes [26]. An independent researcher not involved in the project generated the randomization sequence in blocks of five (using the sequence generator on www.random.org), and wrote the randomization code into sealed numbered envelopes. After completion of all baseline assessments, the researcher allocated participants to their treatment condition based on the randomization code in the sealed envelope (opened in order of sequence). No masking of treatment condition took place.

Table 1. Baseline demographic and clinical characteristics in each group.

	MBCT (n = 63)	Control (n = 66)
Age (mean, SD)	44.6 (9.7)	43.2 (9.5)
Female gender	79%	73%
Full-/part-time work	62%	68%
Illness/unemployment benefits	19%	23%
Living with partner/own family	64%	64%
Comorbid anxiety disorder (present)	35%	49%
Comorbid anxiety disorder (past)	51%	64%
Current psycho-counseling/-therapy	13%	12%
Current use of antidepressants	32%	38%
(Occasional) use of benzodiazepines	8%	8%

Note. There were no significant differences between groups (at $p < 0.05$). MBCT = Mindfulness-based cognitive therapy; Control = waiting list control condition.

doi:10.1371/journal.pone.0066747.t001

Intervention (MBCT)

Content of MBCT training sessions followed the protocol of Segal, Williams, and Teasdale [27]. Training consisted of eight weekly meetings lasting 2.5 hours and were run in groups of 10–15 participants. Assessment periods for control participants were matched to those of MBCT participants. Sessions included guided meditation, experiential exercises, and discussions. In addition to the weekly group sessions, participants received CDs with guided exercises and were assigned daily homework exercises (30 to 60 minutes daily). Training sessions were given by experienced trainers in a centre specialized in mindfulness trainings. All trainers were supervised by an experienced health care professional who had trained with Teasdale and Williams, the co-developers of MBCT [28].

Experience Sampling Method and Measures

ESM is a momentary assessment method to repeatedly assess participants in their daily living environment. Compared to retrospective questionnaires and interviews, ESM offers several advantages: (a) enhanced ecological validity, (b) minimized retrospective bias, because participants' experiences are assessed as they occur, and (c) enhanced reliability, because participants' experiences are assessed repeatedly [29]. In the current study, participants received a digital wristwatch and a set of ESM self-assessment forms collated in a booklet for each day. The wristwatch was programmed to emit a signal ("beep") at an unpredictable moment in each of ten 90-minute time blocks between 7:30 am and 10:30 pm, on six consecutive days, resulting in a maximum of 60 beeps per study period. After each beep, participants were asked to fill out ESM self-assessment forms, collecting reports of current emotions, cognition and context. All self-assessments were rated on 7-point Likert scales (ranging from *not at all* to *very*). Trained research assistants explained the ESM procedure to the participants during an initial briefing session, and a practice form was completed to confirm that participants understood the 7-point Likert scale. Participants were instructed to complete their reports immediately after the beep, thus minimizing memory distortion, and to record the time at which they completed the form. All reports not filled in within 15 min after the actual beep were excluded from the analysis, since previous work has shown that reports completed after this interval are less

reliable and consequently less valid. For the same reason, participants with less than 20 valid reports at baseline were excluded from the analysis [30]. Previous studies have demonstrated the feasibility, validity, and reliability of ESM in general and patient populations [21].

Momentary paranoid feelings. In accordance with previous work [31], momentary paranoia was measured with the ESM item *I feel suspicious*.

Momentary social acceptance. In case participants were not alone at the time of the beep, they evaluated their social company in terms of social acceptance (*In this company, I feel accepted*) [32].

Statistical Methods

ESM data have a hierarchical structure. Thus, multiple observations (level 1) are clustered within participants (level 2). Note that assessment period (pre/post) is not a separate level but an independent factor, just as in regular regression analyses. Multilevel analyses take the variability associated with each level of nesting into account [33]. Multilevel linear regression analyses, using the XTREG command on a Windows platform in Stata 11.1 were applied to the ESM data.

First, we examined the effects of MBCT on change in paranoia and feelings of social acceptance relative to the Control condition. Thus, in the models of paranoia and feelings of social acceptance, the two-way interaction between study period (baseline vs. post assessment) and treatment group (Control vs. MBCT) was the parameter of interest.

Second, we examined whether MBCT changed the time-lagged association between paranoia and feelings of social acceptance in the flow of daily life compared to the Control condition. To this end, we analysed the three-way interaction between study period (baseline vs. post assessment), treatment group (Control vs. MBCT), and paranoia at time point $t-1$ on feeling accepted at time point t and respectively with feeling accepted at time point $t-1$ on paranoia at time point t . In case of significant interactions, stratified effects were calculated for the four different conditions (MBCT and Control, at baseline and post-intervention). Within each treatment group, statistical significance of differences between baseline and post-assessment was assessed by Wald test [34].

The reported analyses are based on the whole sample (intention to treat). Participants who attended less than four MBCT sessions ($n = 3$) were excluded for the per-protocol analysis (outcomes were similar and are not reported).

Results

Participants

Recruitment of participants started in January 2008 and ended in February 2009, and all post-intervention assessments were completed by August 2009, when the pre-determined number of participants was reached. Sociodemographic and clinical characteristics of MBCT and Control participants are displayed in Table 1. At baseline, there were no large or significant differences between treatment groups with respect to sociodemographic and clinical characteristics. Table 2 shows baseline and post assessment scores of variables used in the analyses, stratified by treatment group. Again, there were no large or significant differences between groups at baseline. The baseline overall mean of momentary paranoia was 1.65 (SD = 1.27, range = 1–7). Participant flow through the study can be found in an earlier publication [35]. No known harms or unintended treatment effects were reported in either group.

Table 2. Baseline and post-assessment means and standard deviations of experience sampling variables used in the analyses, stratified by group.

	MBCT (n = 63)					CONTROL (n = 66)						
	baseline		post		change p-value	Cohen's d	baseline		post		change p-value	Cohen's d
	M	SD	M	SD			M	SD	M	SD		
I feel suspicious	1.63	1.27	1.48	1.04	<0.001	-.13	1.67	1.28	1.72	1.25	0.008	.04
I feel accepted	5.91	1.19	6.09	1.06	<0.001	.16	5.98	1.28	5.88	1.2	0.013	-.08
MBCT sessions attended			7.2	1.5								
Minutes practiced per day			29.7	13.2								

Note. MBCT = Mindfulness-based cognitive therapy plus treatment as usual; Control = waiting list control condition (treatment as usual); M = mean; SD = Standard deviation.

doi:10.1371/journal.pone.0066747.t002

Participants completed 12,453 ESM entries. Of these, 559 (4%) were excluded as invalid entries, because completion times fell outside the pre-determined window of 15 minutes after the beep. On average, participants completed 49 (of 60; *SD* = 7.6) valid entries per ESM assessment period. One participant had completed fewer than 20 valid entries at baseline and was therefore excluded from the analyses.

Effects of MBCT on paranoia and social acceptance

Multilevel analysis revealed a significant interaction between treatment group and study period in the model of momentary paranoia ($b = -.18$; 95%CI $-.25, -.10$, $p < 0.001$; $d = -0.35$); the MBCT group reported significantly reduced feelings of paranoia from pre- to post-assessment ($b = -.11$; 95%CI $-.16, -.05$, $p < 0.001$), whereas in the Control condition a significant increase was apparent ($b = .07$; 95%CI $.02, .12$, $p = 0.008$).

Similarly, significant interactions between treatment group and study period were apparent for the variable *I feel accepted* ($b = .26$; 95%CI $.16, .36$, $p < 0.001$; $d = 0.41$). Thus, MBCT participants felt more accepted after the MBCT training ($b = .18$; 95%CI $.13, .25$, $p < 0.001$), compared to before, whereas Control participants felt less accepted after the waiting period ($b = -.09$; 95%CI $-.16,$

$.02$, $p = 0.013$). For baseline and post-assessment means and effect sizes, see Table 2.

MBCT and time-lagged associations between paranoia and feeling accepted

MBCT compared to Control was associated with a significant change in the effect of paranoia on prospective feelings of social acceptance between pre- and post-treatment ($b = .12$; 95%CI $.02, .21$, $p = 0.022$). Before MBCT, feeling more suspicious was associated with feeling less accepted the next moment ($b = -.06$; 95%CI $-.11, -.02$, $p = 0.002$). After MBCT, the association between feeling suspicious at $t-1$ and feeling less accepted at t disappeared ($b = .03$; 95%CI $-.03, .10$, $p = 0.30$) (Figure 1 and Figure 2), while in the Control condition, the association remained and feeling more suspicious was associated with feeling less accepted the next moment both pre- ($b = -.07$; 95%CI $-.12, -.02$, $p = 0.004$) and post-measurement ($b = -.08$; 95%CI $-.14, -.03$, $p = 0.003$).

No difference was found between MBCT compared to Control with respect to the effect of social acceptance on prospective feelings of paranoia between pre- and post-treatment ($b = .04$; 95%CI $-.06, .14$, $p = 0.43$). Overall, feeling less accepted resulted in feeling more suspicious the next moment ($b = -.06$; 95%CI $-.08, -.03$, $p < 0.001$), illustrating the vicious circle between these two variables (see Figure 1 and Figure 2). All results remained significant and similar when controlling for momentary paranoia at t .

Discussion

This is the first daily life study to show that MBCT changes the way in which individuals with affective dysregulation deal with feelings of insecurity and paranoia in the social world. Our results indicate that MBCT decreases feelings of paranoia in the flow of daily life, while paranoia grew worse in the control group. Moreover, MBCT significantly increased feeling accepted in social situations while controls reported reduced social acceptance from baseline to follow-up. A bi-directional link between feeling paranoid and feeling accepted was apparent, which was attenuated as a result of MBCT, suggesting that MBCT interrupts the vicious circle of feeling paranoid and not feeling accepted (see Figure 1 and Figure 2). This interruption seems to particularly take place between paranoia and social acceptance the next moment, but not between feeling accepted on paranoid feeling the next moment.

Improvements in social interaction (because of their central importance for human beings) may be one of the indirect




Figure 1. Time-lagged associations between feeling paranoid and feeling accepted in the flow of daily life: baseline and post-Mindfulness-based cognitive therapy (MBCT).

doi:10.1371/journal.pone.0066747.g001

MBCT group



Control group



Figure 2. The effect of Mindfulness-based cognitive therapy (MBCT) on feeling accepted, feeling paranoid and their association compared to the control condition: Overview of the main results.

doi:10.1371/journal.pone.0066747.g002

mechanisms underlying the beneficial effects of mindfulness-based interventions in depression. Feeling paranoid and not accepted during social interaction may become detrimental for wellbeing, leading to increased isolation [9]. Earlier work shows that dispositional and state mindfulness predict self-regulatory behaviour and positive emotional states [36]. Thereby, MBCT may prevent individuals from becoming socially isolated and developing additional symptoms, for example social anhedonia. Moreover, the interpretation of events is at the heart of affective dysregulation, according to influential cognitive theories [37]. MBCT may weaken the impact of paranoia on affective dysregulation by promoting a different relationship to intrusive thoughts (aware yet detached; [38]) as well as self-compassion [39]. By affecting paranoia and its impact on a network of are causally linked symptoms [40], MBCT may be able to prevent a cascade of symptoms that would normally result in a worse prognosis and increased risk for need for care in individuals with affective dysregulation. Therefore, MBCT may be particularly suited for individuals suffering from (sub-clinical) paranoia. In early intervention, circular processes that otherwise would take place and induce shifts upward the psychosis continuum may be prevented. ESM is ideally suited to examine these interactions at the most fundamental “micro-level” expression of psychopathology [41] and to examine directionality of momentary processes and the smallest building blocks that may eventually form part of a mental illness. More research is necessary to investigate whether MBCT also improves feeling accepted during social interaction when feelings of paranoia are more pronounced (e.g., as in individuals in

an acute psychotic episode or at ultra-high risk to develop a psychotic disorder).

Only a handful of small RCTs using mindfulness-based therapies in participants with psychotic symptoms exist and it is difficult to compare the current study with earlier work because of differences in sample characteristics (sub-clinical feelings of suspiciousness vs. psychotic patients) and employed methodology (experience sampling vs. retrospective questionnaires). Two small randomized trials showed that ACT significantly reduces distress related to hallucinations [42,43] as well as re-hospitalization for inpatients with psychosis [18], with effects extending to up to 1 year post-discharge [44]. Three other studies in individuals with a psychotic disorder examined the effect of MBCT [14,15,16] compared to waiting list control conditions and found improvements in clinical functioning and mindfulness. Recently, improvements in psychological symptoms, command hallucinations, negative symptoms, depression and crisis contacts were reported after mindfulness-based therapies in psychosis [6,45,46]. Our findings extend these findings to real-life paranoia in the context of major mood disorder, and provide a framework for an understanding of the (psychological) mechanisms underlying these positive clinical changes.

Some strengths and limitations of the current RCT should be mentioned. An important observation is that conclusions are limited to subclinical paranoia. One limitation concerns the absence of an active control group. Moreover, we had no measure of MBCT fidelity with respect to both adherence and competence. For some of the ESM variables used in the current study, single

item measures were used. ESM requires selectivity in choosing one's items, because participants have to answer them so frequently (in this study, up to 120 times). While construct validity may suffer due to these restrictions, ecological validity and reliability of the estimation are strongly boosted [47]. These particular single-item measures items have been used in several ESM studies in our research group (e.g. [31]) and have demonstrated predictive and ecological validity. The momentary paranoia scores of the current sample with residual mood complaints were elevated compared to the general population, but were below the momentary paranoia scores of (paranoid) psychotic patients. Strengths include the large sample size and sufficient power for the analysis of group by time interaction effects. Examining (subclinical) paranoid ideas in a sample with affective dysregulation has the advantage that patterns underlying psychosis remain relative plastic, so that changes in psychological processes related to paranoia may be investigated more easily. The most important asset of the current study is the use of repeated assessment of paranoia and social acceptance in daily life, enabling fine-grained examination of changes in psychological processes.

References

- van Os J, Kenis G, Rutten BPF (2010) The environment and schizophrenia. *Nature* 468: 203–212.
- Freeman D (2007) Suspicious minds: the psychology of persecutory delusions. *Clinical Psychology Review* 27: 425–457.
- Thewissen V, Bentall RP, Lecomte T, van Os J, Myin-Germeys I (2008) Fluctuations in self-esteem and paranoia in the context of daily life. *J Abnorm Psychol* 117: 143–153.
- Freeman D, Garety PA (2003) Connecting neurosis and psychosis: the direct influence of emotion on delusions and hallucinations. *Behaviour Research and Therapy* 41: 923–947.
- Bentall RP, Rowse G, Shryane N, Kinderman P, Howard R, et al. (2009) The cognitive and affective structure of paranoid delusions: a transdiagnostic investigation of patients with schizophrenia spectrum disorders and depression. *Archives of General Psychiatry* 66: 236.
- White R, Gumley A, McTaggart J, Rattrie L, McConville D, et al. (2011) A feasibility study of Acceptance and Commitment Therapy for emotional dysfunction following psychosis. *Behaviour Research and Therapy*.
- Wigman JT, van Nierop M, Vollebergh WAM, Lieb R, Beesdo-Baum K, et al. (2012) Evidence That Psychotic Symptoms Are Prevalent in Disorders of Anxiety and Depression, Impacting on Illness Onset, Risk, and Severity—Implications for Diagnosis and Ultra-High Risk Research. *Schizophrenia Bulletin* 38: 247–257.
- van Rossum I, Lieb R, Wittchen HU, van Os J (2011) Affective dysregulation and reality distortion: a 10-year prospective study of their association and clinical relevance. *Schizophrenia Bulletin* 37: 561–571.
- Green MJ, Phillips ML (2004) Social threat perception and the evolution of paranoia. *Neuroscience & Biobehavioral Reviews* 28: 333–342.
- Hofmann SG, Sawyer AT, Witt AA, Oh D (2010) The effect of mindfulness-based therapy on anxiety and depression: A meta-analytic review. *Journal of Consulting and Clinical Psychology* 78: 169.
- Fjorback L, Arendt M, Ørnbøl E, Fink P, Walach H (2011) Mindfulness-Based Stress Reduction and Mindfulness-Based Cognitive Therapy—a systematic review of randomized controlled trials. *Acta Psychiatrica Scandinavica*.
- Bach PA, Gaudio BA, Pankey J, Herbert JD, Hayes SC (2006) Acceptance, Mindfulness, Values, and Psychosis: Applying Acceptance and Commitment Therapy (ACT) to the Chronically Mentally Ill.
- Pankey J, Hayes SC (2003) Acceptance and commitment therapy for psychosis. *International Journal of Psychology and Psychological Therapy*: 311–328.
- Chadwick P, Taylor KN, Abba N (2005) Mindfulness groups for people with psychosis. *Behavioural and Cognitive Psychotherapy* 33: 351–359.
- Chadwick P, Hughes S, Russell D, Russell I, Dagnan D (2010) Mindfulness groups for distressing voices and paranoia: A replication and randomized feasibility trial. *Behavioural and Cognitive Psychotherapy* 37: 403.
- Langer Á, Cangas AJ, Salcedo E, Fuentes B (2012) Applying Mindfulness Therapy in a Group of Psychotic Individuals: A Controlled Study. *Behavioural and Cognitive Psychotherapy* 40: 105.
- Taylor KN, Harper S, Chadwick P (2009) Impact of mindfulness on cognition and affect in voice hearing: evidence from two case studies. *Behavioural and Cognitive Psychotherapy* 37: 397–402.
- Bach P, Hayes SC (2002) The use of acceptance and commitment therapy to prevent the rehospitalization of psychotic patients: a randomized controlled trial. *Journal of Consulting and Clinical Psychology* 70: 1129–1139.
- Bach P, Gaudio BA, Hayes SC, Herbert JD (2012) Acceptance and commitment therapy for psychosis: intent to treat, hospitalization outcome and mediation by believability.
- Ludwig DS, Kabat-Zinn J (2008) Mindfulness in medicine. *JAMA: the journal of the American Medical Association* 300: 1350–1352.
- Myin-Germeys I, Oorschot M, Collip D, Lataster J, Delespaul P, et al. (2009) Experience sampling research in psychopathology: opening the black box of daily life. *Psychological Medicine*: 1–15.
- Williams JMG, Russell I, Russell D (2008) Mindfulness-based cognitive therapy: Further issues in current evidence and future research.
- Geschwind N, Peeters F, Drukker M, Van Os J, Wichers M (2011) Mindfulness training increases momentary positive emotions and reward experience in adults vulnerable to depression: A randomized controlled trial. *Journal of Consulting and Clinical Psychology* 79: 618.
- Hamilton M (1960) A rating scale for depression. *Journal of neurology, neurosurgery, and psychiatry* 23: 56.
- First MB, Spitzer RL, Gibbon M, Williams JBW (2002) Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version. New York: Biometrics Research, New York State Psychiatric Institute.
- Teasdale JD, Segal ZV, Williams JMG, Ridgeway VA, Soulsby JM, et al. (2000) Prevention of relapse/recurrence in major depression by mindfulness-based cognitive therapy. *Journal of Consulting and Clinical Psychology* 68: 615–623.
- Segal ZV, Williams JMG, Teasdale JD (2002) Mindfulness-Based Cognitive Therapy for Depression: A New Approach to Preventing Relapse. New York: The Guilford Press.
- Teasdale JD, Segal ZV, Williams JMG (1995) How does cognitive therapy prevent depressive relapse and why should attentional control (mindfulness) training help. *Behaviour Research and Therapy* 33: 25–39.
- Csikszentmihalyi M, Larson R (1987) Validity and reliability of the Experience Sampling Method. *Journal of Nervous and Mental Disease* 175: 526–536.
- Delespaul PAEG (1995) Assessing schizophrenia in daily life. The Experience Sampling Method. Maastricht: Maastricht University Press.
- Simons CJ, Wichers M, Derom C, Thiery E, Myin-Germeys I, et al. (2009) Subtle gene-environment interactions driving paranoia in daily life. *Genes Brain Behav* 8: 5–12.
- Myin-Germeys I, van Os J, Schwartz J, Stone A, Delespaul P (2001) Stressreactivity in daily life: a marker for psychosis. *Archives of General Psychiatry* 58: 1137–1144.
- Snijders T, Bosker R (1999) Multilevel analysis: an introduction to basis and advanced multilevel modeling. London: SAGE publications Ltd.
- Clayton D, Hill M (1993) Wald tests. In: Clayton D, Hills M, editors. *Statistical Models in Epidemiology*. Oxford: Oxford Science Publications. pp. 101–102.
- Geschwind N, Peeters F, Huibers M, van Os J, Wichers M (2012) Efficacy of mindfulness-based cognitive therapy in relation to prior history of depression: randomised controlled trial. *The British Journal of Psychiatry*.
- Brown KW, Ryan RM (2003) The benefits of being present: mindfulness and its role in psychological well-being. *J Pers Soc Psychol* 84: 822.
- Harvey DM (1981) Depression and attributional style: Interpretations of important personal events. *J Abnorm Psychol* 90: 134.
- Greeson J, Brantley J (2009) Mindfulness and anxiety disorders: Developing a wise relationship with the inner experience of fear. *Clinical Handbook of Mindfulness*: 171–188.
- Neff KD, Germer CK (2013) A Pilot Study and Randomized Controlled Trial of the Mindful Self-Compassion Program. *Journal of clinical psychology* 69: 28–44.

Supporting Information

Checklist S1 CONSORT Checklist.
(DOC)

Protocol S1 Trial Protocol.
(PDF)

Acknowledgments

We thank our study participants for their time and effort. Furthermore, we thank Rufa Diederer for her help with data collection, and Truda Driesen, Philippe Delespaul, Frieda van Goethem, and the 4D database team for their help with data entry.

Author Contributions

Conceived and designed the experiments: MW. Performed the experiments: NG. Analyzed the data: DC NG. Wrote the paper: DC NG MW FP JvO IM-G. Trial design: NG MW FP JvO.

40. Cramer AOJ, Kendler KS, Borsboom D (2012) A constructionist account of emotional disorders. *Behavioral and Brain Sciences* 35: 146–147.
41. van Os J, Linscott RJ (2012) Introduction: the extended psychosis phenotype—relationship with schizophrenia and with ultrahigh risk status for psychosis. *Schizophrenia Bulletin* 38: 227–230.
42. Gaudiano BA, Herbert JD (2006) Acute treatment of inpatients with psychotic symptoms using Acceptance and Commitment Therapy: Pilot results. *Behaviour Research and Therapy* 44: 415–437.
43. Gaudiano BA, Herbert JD, Hayes SC (2010) Is it the symptom or the relation to it? Investigating potential mediators of change in acceptance and commitment therapy for psychosis. *Behavior Therapy* 41: 543–554.
44. Bach P, Hayes SC, Gallop R (2012) Long-Term Effects of Brief Acceptance and Commitment Therapy for Psychosis. *Behavior Modification* 36: 165–181.
45. van der Valk R, van de Waerdt S, Meijer CJ, van den Hout I, de Haan L (2012) Feasibility of mindfulness-based therapy in patients recovering from a first psychotic episode: a pilot study. *Early Intervention in Psychiatry*.
46. Shawyer F, Farhall J, Mackinnon A, Trauer T, Sims E, et al. (2011) A randomised controlled trial of acceptance-based cognitive behavioural therapy for command hallucinations in psychotic disorders. *Behaviour Research and Therapy*.
47. Delespaul PAEG (1995) *Assessing Schizophrenia in Daily Life*; Vries MWd, editor. Maastricht: Universitaire Pers Maastricht. 374 p.