

University of Groningen

Draft Genome Sequences of Seven Thermophilic Spore-Forming Bacteria Isolated from Foods That Produce Highly Heat-Resistant Spores, Comprising *Geobacillus* spp., *Caldibacillus debilis*, and *Anoxybacillus flavithermus*

Berendsen, Erwin M; Wells-Bennik, Marjon H J; Krawczyk, Antonina O; de Jong, Anne; van Heel, Auke; Holsappel, Siger; Eijlander, Robyn T; Kuipers, Oscar P

Published in:
Microbiology resource announcements

DOI:
[10.1128/genomeA.00105-16](https://doi.org/10.1128/genomeA.00105-16)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Berendsen, E. M., Wells-Bennik, M. H. J., Krawczyk, A. O., de Jong, A., van Heel, A., Holsappel, S., Eijlander, R. T., & Kuipers, O. P. (2016). Draft Genome Sequences of Seven Thermophilic Spore-Forming Bacteria Isolated from Foods That Produce Highly Heat-Resistant Spores, Comprising *Geobacillus* spp., *Caldibacillus debilis*, and *Anoxybacillus flavithermus*. *Microbiology resource announcements*, 4(3), 1-2. [00105-16]. <https://doi.org/10.1128/genomeA.00105-16>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Draft Genome Sequences of Seven Thermophilic Spore-Forming Bacteria Isolated from Foods That Produce Highly Heat-Resistant Spores, Comprising *Geobacillus* spp., *Caldibacillus debilis*, and *Anoxybacillus flavithermus*

Erwin M. Berendsen,^{a,b,c} Marjon H. J. Wells-Bennik,^{b,c} Antonina O. Krawczyk,^{a,c} Anne de Jong,^{a,c} Auke van Heel,^a Siger Holsappel,^a Robyn T. Eijlander,^{a,c} Oscar P. Kuipers^{a,c}

Molecular Genetics, University of Groningen, Groningen, The Netherlands^a; NIZO food research, Ede, The Netherlands^b; Top Institute Food and Nutrition (TIFN), Wageningen, The Netherlands^c

Here, we report the draft genomes of five strains of *Geobacillus* spp., one *Caldibacillus debilis* strain, and one draft genome of *Anoxybacillus flavithermus*, all thermophilic spore-forming Gram-positive bacteria.

Received 27 January 2016 Accepted 17 March 2016 Published 5 May 2016

Citation Berendsen EM, Wells-Bennik MHJ, Krawczyk AO, de Jong A, van Heel A, Holsappel S, Eijlander RT, Kuipers OP. 2016. Draft genome sequences of seven thermophilic spore-forming bacteria isolated from foods that produce highly heat-resistant spores, comprising *Geobacillus* spp., *Caldibacillus debilis*, and *Anoxybacillus flavithermus*. *Genome Announc* 4(3):e00105-16 doi:10.1128/genomeA.00105-16.

Copyright © 2016 Berendsen et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Oscar P. Kuipers, o.p.kuipers@rug.nl.

Thermophilic spore-forming bacteria typically grow at temperatures above 45°C and produce spores that are highly resistant to environmental insults, including high temperatures. Thermophilic spore-formers are encountered in a range of different environments (1) and their spores may be able to survive intense heat treatments, including commercial food sterilization processes used for canning and ultra-high temperature processes applied in the manufacturing of liquid foods. Spores that survive such heat treatments may subsequently germinate, and subsequent growth may result in food spoilage (2–4). Here, we sequenced the genomes of five strains of *Geobacillus* spp. (*G. toebii*, *G. stearothermophilus*, and *Geobacillus* sp.), one strain of *Anoxybacillus flavithermus*, and one strain of *Caldibacillus debilis*. The genome sequences of these strains can provide information on the thermophilic lifestyle of these strains and on the ability of the strains to form spores with high-level heat resistance (5).

The seven strains were grown overnight in 10 ml brain heart infusion broth (Difco) at 55°C. Cultures were 100-fold diluted, cultivated at 55°C, and harvested by centrifugation (5,000 relative centrifugal force) at the exponential growth phase. Total DNA was isolated as described previously (5). The isolated DNA was

sheared to 500-bp fragments in the Covaris (KBioscience) ultrasonication device; a next-generation sequencing library was prepared using the paired-end NEB NExtGen library preparation kit. The libraries were 101-bp paired-end sequenced on an Illumina HiSeq2000 by multiplexing 12 samples per flow cell. Draft genome sequences of all strains were assembled *de novo* using Velvet (6). The genomes were annotated using the RAST server (7). Scaffolds were mapped on the closest genome according to RAST, using CONTIGuator (8). Potential bacteriocin gene clusters were identified using BAGEL3 (9) in strains B4109, B4113, B4114, and B4119. Finally, InterProScan (10) was used to extend protein annotations.

Nucleotide sequence accession numbers. The genome sequences of the seven thermophilic spore-forming strains have been deposited as whole-genome shotgun projects at DDBJ/EMBL/GenBank under the accession numbers listed in Table 1.

ACKNOWLEDGMENTS

We thank the NGS sequence facility of the University Medical Center of Groningen (UMCG) for performing the sequencing of the strains. We thank TIFN for contributing to the funding of the project.

TABLE 1 Genome features and GenBank accession numbers of the strains

Strain	Species	Source of isolation	BioProject no.	Accession no.
B4109	<i>Geobacillus stearothermophilus</i>	Pea soup	PRJNA270597	LQYV00000000
B4110	<i>Geobacillus toebii</i>	Pea soup	PRJNA270597	LQYW00000000
B4113	<i>Geobacillus</i> sp.	Mushroom soup	PRJNA270597	LQYX00000000
B4114	<i>Geobacillus stearothermophilus</i>	Buttermilk powder	PRJNA270597	LQYY00000000
B4119	<i>Geobacillus caldioxysilyticus</i>	Dairy	PRJNA270597	LQYS00000000
B4135	<i>Caldibacillus debilis</i>	Dairy	PRJNA270597	LQYT00000000
B4168	<i>Anoxybacillus flavithermus</i>	Dairy	PRJNA270597	LQYU00000000

REFERENCES

1. Zeigler DR. 2014. The *Geobacillus* paradox: why is a thermophilic bacterial genus so prevalent on a mesophilic planet? *Microbiology* 160:1–11. <http://dx.doi.org/10.1099/mic.0.071696-0>.
2. André S, Zuber F, Remize F. 2013. Thermophilic spore-forming bacteria isolated from spoiled canned food and their heat resistance: results of a French ten-year survey. *Int J Food Microbiol* 165:134–143. <http://dx.doi.org/10.1016/j.ijfoodmicro.2013.04.019>.
3. Burgess SA, Lindsay D, Flint SH. 2010. Thermophilic bacilli and their importance in dairy processing. *Int J Food Microbiol* 144:215–225. <http://dx.doi.org/10.1016/j.ijfoodmicro.2010.09.027>.
4. Eijlander RT, de Jong A, Krawczyk AO, Holsappel S, Kuipers OP. 2014. SporeWeb: an interactive journey through the complete sporulation cycle of *Bacillus subtilis*. *Nucleic Acids Res* 42:D685–D691. <http://dx.doi.org/10.1093/nar/gkt1007>.
5. Scheldeman P, Herman L, Foster S, Heyndrickx M. 2006. *Bacillus sporothermodurans* and other highly heat-resistant spore formers in milk. *J Appl Microbiol* 101:542–555. <http://dx.doi.org/10.1111/j.1365-2672.2006.02964.x>.
6. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
7. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
8. Galardini M, Biondi EG, Bazzicalupo M, Mengoni A. 2011. CONTIGuator: a bacterial genomes finishing tool for structural insights on draft genomes. *Source Code Biol Med* 6:11. <http://dx.doi.org/10.1186/1751-0473-6-11>.
9. van Heel AJ, de Jong A, Montalbán-López M, Kok J, Kuipers OP. 2013. BAGEL3: automated identification of genes encoding bacteriocins and (non-)bactericidal posttranslationally modified peptides. *Nucleic Acids Res* 41:W448–W453. <http://dx.doi.org/10.1093/nar/gkt391>.
10. Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong SY, Lopez R, Hunter S. 2014. InterProScan 5: Genome-scale protein function classification. *Bioinformatics* 30:1236–1240. <http://dx.doi.org/10.1093/bioinformatics/btu031>.