

University of Groningen

The disappearance of a significant other

Lenferink, Lonneke Ingrid Maria

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Lenferink, L. I. M. (2018). *The disappearance of a significant other: Consequences and care*. Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

3

Prolonged grief and post-traumatic stress among relatives of missing persons and homicidally bereaved individuals: A comparative study

Lenferink, L.I.M., van Denderen, M.Y., de Keijser, J., Wessel, I., & Boelen, P.A. (2017). Prolonged grief and post-traumatic stress among relatives of missing persons and homicidally bereaved individuals: A comparative study. *Journal of Affective Disorders*, 209, 1-2.

ABSTRACT

Background

Traumatic loss (e.g., homicide) is associated with elevated prolonged grief disorder (PGD) and posttraumatic stress disorder (PTSD). Several studies comparing relatives of missing persons with homicidally bereaved individuals showed inconsistent results about the difference in PGD- and PTSD-levels between the groups. These studies were conducted in the context of armed conflict, which may confound the results. The current study aims to compare PGD- and PTSD-levels between the groups outside the context of armed conflict.

Methods

Relatives of long-term missing persons ($n = 134$) and homicidally bereaved individuals ($n = 331$) completed self-report measures of PGD and PTSD. Multilevel regression modelling was used to compare symptom scores between the groups.

Results

Homicidally bereaved individuals reported significantly higher levels of PGD ($d = 0.86$) and PTSD ($d = 0.28$) than relatives of missing persons, when taking relevant covariates (i.e., gender, time since loss, and kinship to the disappeared/deceased person) into account.

Limitations

A limitation of this study is the use of self-report measures instead of clinical interviews.

Conclusion

Prior studies among relatives of missing persons and homicidally bereaved individuals in the context of armed conflict may not be generalizable to similar samples outside these contexts. Future research is needed to further explore differences in bereavement-related psychopathology between different groups and correlates and treatment of this psychopathology.

Keywords: Missing persons; Homicidal loss; Grief; Trauma

Boss (2006) claimed that the disappearance of a loved one is the most traumatic loss, because the whereabouts of the missing person are uncertain. It is important to verify this assumption, as it may contribute to psychopathology through stigmatization (Feigelman et al., 2009). Furthermore, insight into different psychological responses to different types of loss may help determining the necessity of tailored interventions.

Studies comparing symptom-levels of prolonged grief disorder (PGD)¹ and posttraumatic stress disorder (PTSD) between relatives of missing persons and homicidally bereaved individuals are scarce and inconclusive. One study showed a significantly higher lifetime-prevalence of PTSD post-homicide (9.0%) than post-disappearance (1.3%), whereas PGD-rates did not differ (Pérez-Sales et al., 2000). Conversely, another study found equal PTSD-levels, but significantly higher PGD-levels post-disappearance than post-homicide ($d = .79$) (Powell et al., 2010). A third study showed no group-differences in PGD and PTSD (Heeke et al., 2015).

These studies were conducted in armed conflict. Additional stressors (e.g., homelessness) may confound the results. Consequently, their generalizability to a context other than armed conflict is questionable. Therefore, we compared PGD- and PTSD-levels of relatives of missing persons and homicidally bereaved individuals outside armed conflict, using data from (on-going) research-projects.

Data were available from 134 relatives of missing persons² who had been missing for more than 3 months and 331 people confronted with the homicidal loss of a loved one at least 6 months earlier. Table 1 summarizes sample characteristics. A local review board provided ethics approval.

The Inventory of Complicated Grief (ICG) assessed PGD-symptoms (Boelen et al., 2003). Participants rated the frequency of 19 grief reactions during the last month on 5-point scales (anchors 0="never" and 4="always"). In keeping with prior research, scores >25 indicated probable PGD.

The 20-item PTSD Checklist for DSM-5 (PCL-5) assessed the DSM-5 PTSD-criteria in relatives of missing persons (Blevins et al., 2015). Participants rated on scales of 0 ("not at all") to 4 ("extremely") to what extent they experienced PTSD symptoms during the preceding month. We considered scores >1 on at least one B- and C-cluster-item and two D- and E-cluster-items as indicative of probable PTSD.

The PTSD Symptom Scale–Self-Report (PSS-SR) assessed the DSM-IV PTSD-criteria in homicidally bereaved individuals (Foa et al., 1993). Respondents briefly described an event

-
1. PGD, also referred to as complicated grief or persistent complex bereavement disorder, is characterized by yearning for the deceased and intense sorrow and is distinguishable from normal grief in terms of higher intensity and longer duration.
 2. A missing person is defined as: "Anyone whose whereabouts is unknown whatever the circumstances of disappearance. They will be considered missing until located and their well-being or otherwise established" (Association of Chief Police Officers, 2010, pp. 15).

that bothered them most during the past month and rated the severity of 17 PTSD-symptoms during the preceding month on scales ranging from 0="not at all" to 3="five or more times per week/almost always". Scores of >1 on at least one re-experiencing, three avoidance, and two hyperarousal symptoms indicated probable PTSD.

Items that referred to "death" (ICG) or "stressful experience" (PCL-5) were replaced by references to the disappearance. Psychometric properties of the measures are adequate (Blevins et al., 2015; Boelen et al., 2003; Foa et al., 1993; $\alpha > .90$ for all measures in the current samples).

Multilevel regression-analysis was used to deal with the nested data (i.e., multiple relatives of the same missing/deceased person completed questionnaires). Type of loss was included as independent variable and PGD- or PTSD-scores as dependent variables. Gender, time since loss, and kinship to the disappeared/deceased relative (i.e., spouse, parent, child, or sibling versus other) were included as covariates. An adapted PTSD score³ for relatives of missing persons was used in the group-comparison. Because the PTSD measure in the homicidally bereaved sample could be related to other events than the homicide, we repeated the analysis for the subgroup of people explicitly referring to the homicidal loss in their event-description ($n = 169$).

3. To be able to compare the PTSD levels between the samples we transformed the 5-point scale (of the PCL-5) into a 4-point scale (in accord with the PSS-SR). Furthermore, items 10 to 12 of the PCL-5 were not covered in the PSS-SR and were therefore excluded. Item 1, 11, and 12 were not sufficiently covered in the PCL-5 and were therefore also assessed in the sample of relatives of missing persons.

Table 1. Characteristics of the participants

	Relatives of missing persons (<i>n</i> = 134)	Homicidally bereaved individuals (<i>n</i> = 331)
Gender, <i>N</i> (%)		
Men	45 (33.6)	113 (34.1)
Women	89 (66.4)	218 (65.9)
Age (years), <i>M</i> (<i>SD</i>)	57.8 (14.2)	52.7 (15.4)
Educational level, <i>N</i> (%)		
Low	32 (23.9)	-
Middle	44 (32.8)	-
High	58 (43.3)	-
Lost relative is, <i>N</i> (%)		
Partner/spouse	18 (13.4)	25 (7.8)
Child	41 (30.6)	157 (48.8)
Parent	14 (10.4)	41 (12.7)
Sibling	31 (23.1)	52 (16.1)
Other family member	28 (20.9)	28 (8.7)
Other	2 (1.5)	19 (5.9)
Number of years since loss, <i>M</i> (<i>SD</i>)	15.5 (17.0)	6.9 (6.5)
Type of disappearance, <i>N</i> (%)		
Criminal act	44 (32.8)	n.a.
Voluntarily	33 (24.6)	n.a.
Accident	33 (24.6)	n.a.
No specific suspicion	24 (17.9)	n.a.
Unique victims	89 (66.4)	254 (76.7)
Recruitment via <i>N</i> (%)		
Editorial office of a Tv-show about missing persons	36 (26.9)	n.a.
Peer support organizations	30 (22.4)	172 (52.0)
Non-governmental support organization	21 (15.7)	136 (41.1)
Family or friends	35 (26.1)	n.a.
Other	12 (9.0)	23 (6.9)
PGD severity, <i>Grand mean</i> (<i>SE</i>)	25.44 (1.40)	39.77 (0.81)
PTSD severity, <i>Grand mean</i> (<i>SE</i>)	13.71 (1.04)	19.96 (0.74)

Note. - = data were not available; n.a. = not applicable; PGD = prolonged grief disorder; PTSD = posttraumatic stress disorder.

In relatives of missing persons, prevalence rates were 47.0% and 23.1% for probable PGD and PTSD, respectively. These rates were 83.1% for PGD and 31.4% for PTSD in homicidally bereaved individuals. Homicidally bereaved individuals scored significantly higher than relatives of missing persons on PGD-symptoms ($F(1, 243.05) = 45.37, p < .001, d = 0.86$) and PTSD-symptoms ($F(1, 291.57) = 5.79, p < .05, d = 0.28$). Similar findings for PTSD-symptoms emerged when including the subgroup of $n = 169$ homicidally bereaved individuals ($F(1, 187.00) = 7.72, p < .01, d = 0.41$).

Our findings that PGD-levels and PTSD-levels are higher post-homicide than post-disappearance contrast with prior studies. This indicates the lack of generalizability of findings from studies in the context of armed conflict to populations outside these contexts. Additional war-related stressors in situations of armed conflict may account for differences. Furthermore, non-Western samples were used in previous studies. Grief may differ across cultures (Rosenblatt, 2008). In contrast to previous studies (Heeke et al., 2015; Pérez-Sales et al., 2000) we used self-reports instead of interviews, which may overestimate symptom-levels.

In contrast to relatives of missing persons, homicidally bereaved individuals need to deal with facts about the violent cause of death and the (presumed) perpetrator, which might be an explanation for the higher PGD- and PTSD-levels post-homicide than post-disappearance. Clearly, before conclusions can be drawn, future research should confirm the present findings. The large difference in PGD-levels suggests that PGD should be more central to treatments for people post-homicide relative to post-disappearance. Future research should further explore differences in psychopathology-symptoms and their correlates after different types of loss to inform intervention development.

REFERENCES

- Association of Chief Police Officers (2010). *Guidance on the management, recording and investigation of missing persons*. http://ec.europa.eu/justice/fundamental-rights/files/missing_persons_sec_edn_2010_en.pdf (accessed 29.07.2016).
- Blevins, C.A., Weathers, F.W., Davis, M.T., Witte, T.K., & Domino J.L. (2015). The Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5): Development and Initial Psychometric Evaluation. *Journal of Traumatic Stress, 28*, 489-98.
- Boelen, P.A., van den Bout, J., de Keijser, J., & Hoijtink, H. (2003). Reliability and validity of the Dutch version of the inventory of traumatic grief (ITG). *Death Studies, 27*, 227-47.
- Boss, P. (2006). Introduction: Loss and ambiguity, in: *Loss, trauma, and resilience: Therapeutic work with ambiguous loss*. W.W. Norton and company, New York, pp. 1-22.
- Feigelman, W., Gorman, B.S., & Jordan, J.R. (2009). Stigmatization and suicide bereavement. *Death Studies, 33*, 591-608.
- Foa, E.B., Riggs, D., Dancu, C., & Rothbaum, B. (1993). Reliability and validity of a brief instrument for assessing post-traumatic stress disorder. *Journal of Traumatic Stress, 6*, 459-474.
- Heeke, C., Stammel, N., & Knaevelsrud, C. (2015). When hope and grief intersect: Rates and risks of prolonged grief disorder among bereaved individuals and relatives of disappeared persons in Colombia. *Journal of Affective Disorders, 173*, 59-64.
- Pérez-Sales, P., Durán-Pérez, T., & Bacic Herzfeld, R. (2000). Long-term psychosocial consequences in first-degree relatives of people detained-disappeared or executed for political reasons in Chile: A study in Mapuce and non-Mapuce persons. *Psicothema, 12*, 109-116.
- Powell, S., Butollo, W., & Hagl, M. (2010). Missing or killed the differential effect on mental health in women in Bosnia and Herzegovina of the confirmed or unconfirmed loss of their husbands. *European Psychologist, 15*, 185-192.
- Rosenblatt, P.C. (2008). Grief across cultures: A review and research agenda, in: Stroebe, M. S. (Eds.), *Handbook of bereavement research and practice: advances in theory and intervention*. American Psychological Association Washington, DC, US, pp. 207-222.

