

University of Groningen

Mortality following elective abdominal aortic aneurysm repair in women

Dutch Society of Vascular Surgery; Tedjawirja, V. N.; Alberga, A. J.; Hof, M. H.P.; Vahl, A. C.; Koelemay, M. J.W.; Balm, R.

Published in:
British Journal of Surgery

DOI:
[10.1093/bjs/znab465](https://doi.org/10.1093/bjs/znab465)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Dutch Society of Vascular Surgery, Tedjawirja, V. N., Alberga, A. J., Hof, M. H. P., Vahl, A. C., Koelemay, M. J. W., & Balm, R. (2022). Mortality following elective abdominal aortic aneurysm repair in women. *British Journal of Surgery*, 109(4), 340-345. <https://doi.org/10.1093/bjs/znab465>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Mortality following elective abdominal aortic aneurysm repair in women

V. N. Tedjawirja¹, A. J. Alberga^{2,3} , M. H. P. Hof⁴ , A. C. Vahl⁵ , M. J. W. Koelemay¹, and R. Balm^{1,*} and in collaboration with the Dutch Society of Vascular Surgery

¹Department of Surgery, Amsterdam Cardiovascular Sciences, Amsterdam UMC, University of Amsterdam, Amsterdam, the Netherlands

²Department of Vascular Surgery, Erasmus University Medical Centre, Rotterdam, the Netherlands

³Dutch Institute of Clinical Auditing, Scientific Bureau, Leiden, the Netherlands

⁴Department of Epidemiology and Data Science, Amsterdam Public Health, Amsterdam UMC, University of Amsterdam, Amsterdam, the Netherlands

⁵Department of Surgery, OLVG, Amsterdam, the Netherlands

*Correspondence to: R. Balm, Amsterdam University Medical Centres, University of Amsterdam, Department of Surgery, Meibergdreef 9, 1105 AZ, Amsterdam, the Netherlands (e-mail: r.balm@amsterdamumc.nl)

Members of the Dutch Society of Vascular Surgery are co-authors of this study and are listed under the heading Collaborators.

Abstract

Background: Previous studies have focused on patient-related risk factors to explain the higher mortality risk in women undergoing elective abdominal aortic aneurysm (AAA) repair. The aim of this study was to evaluate whether hospital-related factors influence outcomes following AAA repair in women.

Methods: Patients undergoing elective AAA repair in 61 hospitals in the Netherlands were identified from the Dutch Surgical Aneurysm Audit registry (2013–2018). A mixed-effects logistic regression analysis was conducted to assess the effect of sex on in-hospital and/or 30-day mortality. This analysis accounted for possible correlation of outcomes among patients who were treated in the same hospital, by adding a hospital-specific random effect to the statistical model. The analysis adjusted for patient-related risk factors and hospital volume of open surgical repair (OSR) and endovascular aneurysm repair (EVAR).

Results: Some 12 034 patients were included in the analysis. The mortality rate was higher in women than among men: 53 of 1780 (3.0 per cent) versus 152 of 10 254 (1.5 per cent) respectively. Female sex was significantly associated with mortality after correction for patient- and hospital-related factors (odds ratio 1.68, 95 per cent c.i. 1.20 to 2.37). OSR volume was associated with lower mortality (OR 0.91 (0.85 to 0.95) per 10-procedure increase) whereas no such relationship was identified with EVAR volume (OR 1.03 (1.01 to 1.05) per 10-procedure increase).

Conclusion: Women are at higher risk of death after abdominal aortic aneurysm repair irrespective of patient- and hospital-related factors.

Introduction

Patients with an abdominal aortic aneurysm (AAA) can be treated electively by open surgical repair (OSR) or endovascular aneurysm repair (EVAR)¹. Previous studies^{2–5} have shown that excess perioperative mortality is evident among women following both types of repair. Well known patient-related risk factors are associated with increased mortality risk, including age, cardiac and pulmonary co-morbidity, and impaired renal function^{6–9}. Despite correction for such factors, female sex has persistently been associated with increased mortality^{2,8}.

Hospital-level factors such as expertise in AAA surgery may influence patient outcomes. Volume can be used as a proxy for expertise and has been found to have an inverse relationship with mortality^{1,10,11}. However, previous studies^{2,8,12} have focused only on patient-related factors. The aim of this study was to establish whether hospital-level factors could explain some of the differences in outcome associated with women after AAA surgery.

Methods

Study design and data source

A retrospective study from the Dutch Surgical Aneurysm Audit (DSAA) was conducted in accordance with the STROBE statement¹³. The DSAA is a nationwide and mandatory quality registry that was initiated in 2013, and obtains data on all patients who undergo surgery for an aortic aneurysm in the Netherlands across 61 hospitals.

Study population

Patients eligible for the present study were women and men registered in the DSAA who underwent primary elective OSR or EVAR for an asymptomatic AAA between January 2013 and December 2018.

Variables and definitions

Patient- and hospital-related factors considered to have an impact on mortality from a clinical point of view and/or known from the literature were assessed before the analysis by means

Received: June 15, 2021. Revised: September 10, 2021. Accepted: December 17, 2021

© The Author(s) 2022. Published by Oxford University Press on behalf of BJS Society Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.




Fig. 1 Study flow diagram

*Used in calculation of hospital volume. AAA, abdominal aortic aneurysm.

of a directed acyclic graph to minimize bias (Table S1). The patient-related risk factors age, AAA diameter, cardiac and pulmonary co-morbidity, serum creatinine levels, and type of repair were extracted from the registry. Cardiac and pulmonary co-morbidities were registered in the DSAA in accordance with POSSUM¹⁴. This score is used to predict 30-day mortality and morbidity rates after surgery, and was designed specifically for surgical audit purposes^{15,16}. The hospital-related factor hospital volume was divided into OSR and EVAR volume, as the separate volumes can be differently associated with mortality¹⁷. The volumes of both types of repair were calculated as the total number of primary elective repairs in each hospital throughout the 6-year study period. The total number of patients who had surgery for an aortic aneurysm per hospital was used to calculate hospital volume, regardless of whether patients had missing values on patient-related risk factors as all of the registered repairs add to the cumulative hospital expertise.

Outcome

The primary outcome of interest for this study was the effect of sex on perioperative mortality, comprising in-hospital mortality during primary admission and 30-day mortality.

Statistical analysis

Baseline characteristics of both the total study cohort and complete cases are reported, along with hospital characteristics including the percentage of women treated with OSR and EVAR per hospital. Continuous variables, stratified by sex, are reported as mean (s.d.) or median (i.q.r.), depending on the distribution. Categorical variables are reported as absolute numbers with percentages.

The data can be regarded to have a clustered structure as they were obtained from 61 hospitals, whereby patients from the same hospital formed a single cluster (group). It is possible that patients treated in the same hospital have correlated outcomes^{18,19}. A wide variety of factors may lead to higher or lower mortality rates in particular hospitals. To deal with possible correlated outcomes, a mixed-effects logistic regression model was used. The following were used as fixed effects in the analysis: OSR volume, EVAR volume, age, sex, AAA diameter, cardiac and pulmonary co-morbidity, serum creatinine levels, and type of repair. The random effect in the statistical model was a hospital-specific offset, which was assumed to follow a normal distribution with a mean of zero. To assess the degree of correlation between

patients treated in the same hospital, the intraclass correlation coefficient (ICC) was calculated¹⁹. The ICC is calculated by dividing the random-effect variance (between-hospital variance) by the total unexplained variance (between-hospital variance and assumed within-hospital variance; fixed value of $\pi^2/3$ in standard logistic distribution)^{19,20}.

The analysis included patients with complete data on patient-related risk factors (complete-case analysis). The association between variables and perioperative mortality was expressed as odds ratios (ORs) with the corresponding 95 per cent confidence intervals. $P < 0.050$ was considered statistically significant. Statistical analyses were conducted with SPSS® 25 (IBM, Armonk, NY, USA) and R studio version 1.3.959 (The R Foundation for Statistical Computing, Vienna, Austria).

Results

Participants and descriptive data

Some 13 091 patients who underwent elective primary AAA repair in 61 hospitals were registered in the DSAA (Fig. 1). These data were used to calculate hospital volume. After exclusion of 30 patients aged 18 years or under, or for whom information on sex or mortality was missing, the total study cohort comprised 13 061 patients. Data were considered to be missing completely at random as patients with missing values on patient-related factors were not treated at specific hospitals. Hence, no hospital was excluded from the analysis. Ultimately, 12 034 patients were included in the complete case-analysis. A total of 2827 patients were treated with OSR (550 women, 19.5 per cent) and 9207 with EVAR (1230 women, 13.4 per cent). Women were older than men at the time of surgery: mean(s.d.) 74.1(7.7) versus 73.1(7.6) years respectively ($P < 0.001$) (Table 1).

Hospital characteristics

The median total number of elective AAA repairs over 6 years was 243 (i.q.r. 187–320) per hospital. The median OSR volume was 55 (39–78) and median EVAR volume was 193 (140–240). Some 18.8 per cent of all patients treated by OSR per hospital, and 12.9 per cent of all those treated by EVAR per hospital, were women (Table 2).

Mortality data

The overall mortality rate was higher in women than men (53 of 1780 (3.0 per cent) versus 152 of 10 254 (1.5 per cent); $P < 0.001$). Mortality rates were higher in women than in men after both

Table 1 Baseline characteristics of patients undergoing primary elective abdominal aortic aneurysm repair in the Netherlands (2013–2018)

	Total study cohort		Complete cases	
	Women (n = 1916)	Men (n = 11 145)	Women (n = 1780)	Men (n = 10 254)
Age (years)*	74.1 (7.7)	73.1 (7.6)	74.1 (7.7)	73.1 (7.6)
AAA diameter (mm)†	55 (52–60)	58 (55–65)	55 (52–60)	58 (55–65)
Missing	30 (1.6)	125 (1.1)		
Preoperative cardiac status				
No cardiac history	820 (43.9)	4702 (43.7)	780 (43.8)	4492 (43.8)
Medication for hypertension, angina pectoris, diuretics or digoxin	883 (47.3)	4889 (45.4)	841 (47.2)	4631 (45.2)
Peripheral oedema, anticoagulation (vitamin K antagonist), borderline cardiomyopathy	144 (7.7)	995 (9.2)	138 (7.8)	957 (9.3)
Increased central venous pressure, cardiomegaly	21 (1.1)	182 (1.7)	21 (1.2)	174 (1.7)
Missing	48 (2.5)	377 (3.4)		
Preoperative pulmonary status				
No dyspnoea	1313 (69.6)	8280 (75.4)	1232 (69.2)	7715 (75.2)
Dyspnoea on exertion	476 (25.2)	2281 (20.8)	457 (25.7)	2145 (20.9)
Disabling dyspnoea	72 (3.8)	318 (2.9)	69 (3.9)	302 (2.9)
Dyspnoea at rest, consolidation, fibrosis	26 (1.4)	100 (0.9)	22 (1.2)	92 (0.90)
Missing	29 (1.5)	166 (1.5)		
Creatinine (µmol/l)†	76 (65–92)	92 (79–110)	76 (64–92)	92 (79–109)
Missing	44 (2.3)	305 (2.7)		

Values in parentheses are percentages unless indicated otherwise; values are *mean (s.d.) and †median (i.q.r.). Baseline characteristics are shown for 13 061 patients who underwent primary elective abdominal aortic aneurysm (AAA) repair after exclusion of those aged 18 years or less, or with missing values on sex or mortality, and for 12 034 patients after excluding those whose data set was incomplete.

Table 2 Type of repair and proportion of women treated per hospital across 61 hospitals in the Netherlands (2013–2018)

	Total study cohort (n = 13 061)	Complete cases (n = 12 034)
% OSR per hospital	23.2 (16.5–30.0)	23.1 (16.3–29.1)
% women treated per hospital	14.8 (12.6–16.2)	14.6 (12.7–16.0)
% women treated by OSR per hospital	18.2 (14.7–21.8)	18.8 (15.2–22.9)
% women treated by EVAR per hospital	12.6 (10.5–15.2)	12.9 (10.7–15.4)

Values are median (interquartile range). OSR, open surgical repair. EVAR, endovascular aneurysm repair.

OSR (38 of 550 (6.9 per cent) versus 104 of 2277 (4.6 per cent); $P = 0.024$) and EVAR (15 of 1230 (1.2 per cent) versus 48 of 7977 (0.6 per cent); $P = 0.014$).

Mixed-effects logistic regression analysis

After adjusting for patient- and hospital-related factors, female sex was significantly associated with perioperative mortality (OR 1.68, 95 per cent c.i. 1.20 to 2.37). Advanced age, cardiac and pulmonary co-morbidity, higher serum creatinine levels, and OSR as type of repair were also associated with an increased mortality risk (Table 3). Higher hospital OSR volume was associated with a lower risk of mortality (OR 0.91 (0.85 to 0.95) per 10-procedure increase), whereas higher hospital EVAR volume was associated with a higher risk of death (OR 1.03 (1.01 to 1.05) per 10-procedure increase).

The estimated hospital-specific offset variance across hospitals was 0.08. An ICC of 0.024 (2.4 per cent) suggested that the outcomes of patients treated in the same hospital were only slightly correlated.

Discussion

The associations between both patient- and hospital-related factors and mortality in AAA surgery have been well reported

previously. In the investigation of the higher mortality rate in women following elective AAA repair, contemporary studies^{2,3,5} have focused foremost on patient-related risk factors. As hospital-level factors can affect outcomes as well, a study combining both factors was conducted to find an explanation beyond patient-related factors for why women are at higher risk. Using nationwide data on aortic aneurysm repair in the Netherlands, the present study found that female sex was associated with mortality after additional correction for interhospital variation.

These findings corroborated those of a recent study²¹ that investigated sex as a modifier in the volume–outcome relationship. The authors concluded that female sex was associated with increased mortality and that hospital volume did not have a consistent effect in women²¹. Another study¹¹ that investigated various hospital-level variables showed that institutional practice patterns had a relatively minor impact on mortality in comparison to patient-level risk factors. These reports suggest that factors at patient level may be more important in explaining the higher mortality risk among women. The patient-level risk factors advanced age, cardiac and pulmonary co-morbidity, high serum creatinine levels, and OSR as type of repair were associated with mortality in the present study, in agreement with previous studies^{6,22,23}. Further in-depth research on other patient-related risk factors, such as anatomical, genetic or biological differences between women and men, are needed to identify potential explanations for the sex-specific mortality risk.

Hospital volume as a measurable parameter at hospital level was used as a proxy to express possible variation in expertise in AAA surgery and other hospital-related processes, such as resources for dealing with postoperative complications. Cumulative number of OSR or EVAR procedures performed over 6 years in each hospital was used as hospital volume, which can be considered to correspond to the average annual volume used in previous studies^{24–26}. The focus of the present analysis was the mortality risk among patients who underwent elective repair. As such, ruptured AAA procedures were not taken into the calculation of hospital volume, which may not have done justice

Table 3 Mixed-effects logistic regression analysis to determine the effect of sex on 30-day and/or in-hospital mortality following elective abdominal aortic aneurysm repair in the Netherlands

	Odds ratio	P
Sex (F versus M)	1.68 (1.20, 2.37)	0.003
Age (per year)	1.07 (1.05, 1.10)	< 0.001
AAA diameter (per 10 mm)	1.04 (0.91, 1.18)	0.577
Cardiac co-morbidity		
No cardiac history	1.00 (reference)	
Medication for hypertension, angina pectoris, diuretics or digoxin	1.31 (0.94, 1.83)	0.108
Peripheral oedema, anticoagulation (vitamin K antagonist), borderline cardiomyopathy	1.86 (1.17, 2.95)	0.009
Increased central venous pressure, cardiomegaly	2.64 (1.12, 6.23)	0.026
Pulmonary co-morbidity		
No dyspnoea	1.00 (reference)	
Dyspnoea on exertion	2.36 (1.73, 3.21)	< 0.001
Disabling dyspnoea	2.12 (1.03, 4.36)	0.042
Dyspnoea at rest, consolidation, fibrosis	8.33 (3.86, 17.99)	< 0.001
Creatinine (per 100-μmol/l increase)	1.61 (1.33, 1.96)	< 0.001
Type of repair (OSR versus EVAR)	12.23 (8.69, 17.23)	< 0.001
Hospital volume OSR (per 10 procedures)	0.91 (0.85, 0.95)	0.002
Hospital volume EVAR (per 10 procedures)	1.03 (1.01, 1.05)	0.017

Values in parentheses are 95 per cent confidence intervals. OSR, open surgical repair; EVAR, endovascular aneurysm repair. These are the results of the analysis investigating the effect of sex on perioperative mortality, with correction for confounders.

to tertiary referral centres that performed more repairs for ruptured AAA than other centres and may potentially have affected outcomes. Although different definitions of volume have been used, higher hospital volume is reported to be associated with lower mortality after AAA repair^{27–29}. However, there seems to be a difference in strength of the association of OSR and EVAR volume with mortality. Previous research¹⁷ has shown that the association between OSR volume and mortality is stronger than that for EVAR volume. Although higher OSR volume was associated with lower mortality as identified previously^{24,30}, EVAR volume was associated with a slightly higher mortality risk in the present study. As EVAR has a relatively low mortality rate and EVAR volume has been reported to have no or little relationship with mortality, a possible explanation for this surprising observation is that selection bias had occurred^{24,25}. EVAR as a less invasive operation is often the procedure of choice in older patients with more co-morbidity, and/or offered to a broader selection of patients^{31–34}. Alternatively, heterogeneity in definitions of hospital volume may also have had an impact on differences in outcomes. For example, some analyses^{24–26} used the average annual hospital volume, whereas another study³⁵ used the annual hospital volume. Although these studies revealed similar outcomes for OSR (volume–outcome association), a minor difference was noted for EVAR (minor volume–outcome association or no association). Notably, the interpretation of the volume–outcome relationship in the present study is different from that in studies that investigated the effect of hospital volume on the mortality risk of patients undergoing OSR or EVAR^{25,26,35}. As such, the hospital volume–outcome relationship can be investigated in various ways, reflecting the complexity of the underlying mechanisms.

The study aimed to control for possible unexplained interhospital variation that may have affected patient outcomes, by accounting for possible correlation between outcomes of patients treated in the same hospital. Hospital parameters that may vary included the concept of heterogeneity in differences in surgical experience with type of repair or differences in experience with postoperative AAA care in women. As a secondary finding, the analysis showed that there was no heterogeneity between hospitals after correction for the fixed effects; all hospitals

performed equally. Although the study aimed to capture these unmeasured hospital-related parameters, there is a possibility that the authors could not have accounted for all such factors.

The study showed that female sex is associated with high mortality after elective AAA repair. The high mortality risk in women may in part be due to a minor delay in treatment, reflected by a median AAA diameter of 5.5 cm, with potentially more advanced AAA disease and need for complex repair. However, as women are at higher risk of perioperative mortality, perhaps the trade-off of treating women with surgery should be re-evaluated. The threshold for treating AAA in women is currently set at an aortic diameter of 5.0 cm, which is lower than the threshold of 5.5 cm in men, possibly because women have a higher risk of aneurysm rupture than men^{1,36}. Yet, perhaps the perioperative mortality risk exceeds the rupture risk at the lower AAA diameter threshold. As further studies are warranted to investigate this trade-off, a more dynamic approach to treatment may be suggested meanwhile. For women undergoing open repair, the threshold should perhaps be increased until as-yet unidentified risk factors for mortality have been elucidated, whereas a lower threshold may be indicated for EVAR considering the low mortality risk. It is clear that a tailor-made decision is required, by incorporating the patient's preference into shared decision-making³⁷.

There were some limitations to this study. First, potential risk factors that were not registered in the DSAA could not be taken into account. These include both patient- and hospital-related factors; the former include AAA parameters such as aneurysm anatomy and operative complexity, and the latter surgeon volume (number of procedures performed per surgeon) which has been proposed to be associated with mortality^{38,39}. Social factors such as caregiver status may also influence outcomes, which could not be taken into account in the present analysis. Second, this retrospective study used data from a quality registry that was not primarily designed for research and could have missing values. The percentage of missing values for each co-morbidity was less than 4 per cent and the incomplete data were distributed over approximately 8 per cent of the patients. The information bias of the extracted variables was therefore considered to be acceptable.

Collaborators

P.J. van den Akker, G.J. Akkersdijk, G.P. Akkersdijk, W.L. Akkersdijk, M.G. van Andringa de Kempnaer, C.H.P. Arts, J.A.M. Avontuur, O.J. Bakker, R. Balm, W.B. Barendregt, J.A. Bekken, M.H. Bender, B.L. Bendermacher, M. van den Berg, P. Berger, R.J. Beuk, J.D. Blankensteijn, R.J. Bleker, J.J. Blok, A.S. Bode, M.E. Bodegom, K.E. van der Bogt, A.P.M. Boll, M.H. Booster, B.L. Borger van der Burg, G.J. de Borst, W.T.G.J. Bos- van Rossum, J. Bosma, J.M.J. Botman, L.H. Bouwman, V. Brehm, M.T. de Bruijn, J.L. de Bruin, P. Brummel, J.P. van Brussel, S.E. Buijk, M.A. Buijs, M.G. Buimer, D.H. Burger, H.C. Buscher, E. Cancrinus, P.H. Castenmiller, G. Cazander, A.M. Coester, P.H. Cuypers, J.H. Daemen, I. Dawson, J.E. Dierikx, M.L. Dijkstra, J. Diks, M.K. Dinkelman, M. Dirven, D.E. Dolmans, R.C. van Doorn, L.M. van Dortmont, J.W. Drouven, M.M. van der Eb, D. Eefting, G.J. van Eijck, J.W. Elshof, B.H. Elsman, A. van der Elst, M.I. van Engeland, R.G. van Eps, M.J. Faber, W.M. de Fijter, B. Fioole, T.M. Fokkema, F.A. Frans, W.M. Fritschy, P.H. Fung Kon Jin, R.H. Geelkerken, W.B. van Gent, G.J. Glade, B. Govaert, R.P. Groenendijk, H.G. de Groot, R.F. van den Haak, E.F. de Haan, G.F. Hajer, J.F. Hamming, E.S. van Hattum, C.E. Hazenberg, P.P. Hedeman Joosten, J.N. Helleman, L.G. van der Hem, J.M. Hendriks, J.A. van Herwaarden, J.M. Heyligers, J.W. Hinnen, R.J. Hissink, G.H. Ho, P.T. den Hoed, M.T. Hoedt, F. van Hoek, R. Hoencamp, W.H. Hoffmann, W. Hogendoorn, A.W. Hoksbergen, E.J. Hollander, M. Hommes, C.J. Hopmans, L.C. Huisman, R.G. Hulsebos, K.M. Huntjens, M.M. Idu, M.J. Jacobs, M.F. van der Jagt, J.R. Jansbeken, R.J. Janssen, H.H. Jiang, S.C. de Jong, T.A. Jongbloed-Winkel, V. Jongkind, M.R. Kapma, B.P. Keller, A. Khodadade Jahrome, J.K. Kievit, P.L. Klemm, P. Klinkert, N.A. Koedam, M.J. Koelemaij, J.L. Kolkert, G.G. Koning, O.H. Koning, R. Konings, A.G. Krasznai, R.M. Krol, R.H. Kropman, R.R. Kruse, L. van der Laan, M.J. van der Laan, J.H. van Laanen, G.W. van Lammeren, D.A. Lamprou, J.H. Lardenoye, G.J. Lauret, B.J. Leenders, D.A. Legemate, V.J. Leijdekkers, M.S. Lemson, M.M. Lensvelt, M.A. Lijkwan, R.C. Lind, F.T. van der Linden, P.F. Liqui Lung, M.J. Loos, M.C. Loubert, K.M. van de Luijngaarden, D.E. Mahmoud, C.G. Manshanden, E.C. Mattens, R. Meerwaldt, B.M. Mees, G.C. von Meijenfheldt, T.P. Menting, R. Metz, R.C. Minnee, J.C. de Mol van Otterloo, M.J. Molegraaf, Y.C. Montauban van Swijndregt, M.J. Morak, R.H. van de Mortel, W. Mulder, S.K. Nagesser, C.C. Naves, J.H. Nederhoed, A.M. Nevenzal-Putters, A.J. de Nie, D.H. Nieuwenhuis, J. Nieuwenhuizen, R.C. van Nieuwenhuizen, D. Nio, V.J. Noyez, A.P. Oomen, B.I. Oranen, J. Oskam, H.W. Palamba, A.G. Peppelenbosch, A.S. van Petersen, B.J. Petri, M.E. Pierie, A.J. Ploeg, R.A. Pol, E.D. Ponfoort, I.C. Post, P.P. Poyck, A. Prent, S. ten Raa, J.T. Raymakers, M. Reichart, B.L. Reichmann, M.M. Reijnen, J.A. de Ridder, A. Rijbroek, M.J. van Rijn, R.A. de Roo, E.V. Rouwet, B.R. Saleem, P.B. Salemans, M.R. van Sambeek, M.G. Samyn, H.P. van 't Sant, J. van Schaik, P.M. van Schaik, D.M. Scharn, M.R. Scheltinga, A. Schepers, P.M. Schlejen, F.J. Schlosser, F.P. Schol, V.P. Scholtes, O. Schouten, M.A. Schreve, G.W. Schurink, C.J. Sikkink, A. te Slaa, H.J. Smeets, L. Smeets, R.R. Smeets, A.A. de Smet, P.C. Smit, T.M. Smits, M.G. Snoeijs, A.O. Sondakh, M.J. Speijers, T.J. van der Steenhoven, S.M. van Sterkenburg, D.A. Stigter, R.A. Stokmans, R.P. Strating, G.N. Stultiëns, J.E. Sybrandy, J.A. Teijink, B.J. Telgenkamp, M. Teraa, M.J. Testroote, T. Tha-In, R.M. The, W.J. Thijsse, I. Thomassen, I.F. Tielliu, R.B. van Tongeren, R.J. Toorop, E. Tournioij, M. Truijers, K. Türkcan, R.P. Tutein Nolthenius, Ç. Ünlü, R.H. Vaes, A.A. Vafi, A.C. Vahl, E.J. Veen, H.T. Veger, M.G. Veldman, S. Velthuis, H.J. Verhagen, B.A.

Verhoeven, C.F. Vermeulen, E.G. Vermeulen, B.P. Vierhout, R.J. van der Vijver-Coppen, M.J. Visser, J.A. van der Vliet, C.J. Vlijmen - van Keulen, R. Voorhoeve, J.R. van der Vorst, A.W. Vos, B. de Vos, C.G. Vos, G.A. Vos, M.T. Voute, B.H. Vriens, P.W. Vriens, A.C. de Vries, D.K. de Vries, J.P. de Vries, M. de Vries, C. van der Waal, E.J. Waasdorp, B.M. Wallis de Vries, L.A. van Walraven, J.L. van Wanroij, M.C. Warlé, W. van de Water, V. van Weel, A.M. van Well, G.M. Welten, R.J. Welten, J.J. Wever, A.M. Wiersema, O.R. Wikkeling, W.I. Willaert, J. Wille, M.C. Willems, E.M. Willigendael, E.D. Wilschut, W. Wisselink, M.E. Witte, C.H. Wittens, C.Y. Wong, R. Wouda, O. Yazar, K.K. Yeung, C.J. Zeebregts, M.L. van Zeeland.

Acknowledgements

No preregistration exists for the study reported in this article.

Funding

V.N. Tedjawirja was funded by the AMC Foundation.

Disclosure. The authors declare no conflict of interest.

Supplementary material

[Supplementary material](#) is available at *BJS* online.

References

1. Wanhainen A, Verzini F, Van Herzelee I, Allaire E, Bown M, Cohnert T *et al.* Editor's choice—European Society for Vascular Surgery (ESVS) 2019 clinical practice guidelines on the management of abdominal aorto-iliac artery aneurysms. *Eur J Vasc Endovasc Surg* 2019;**57**:8–93
2. Sidloff DA, Saratzis A, Sweeting MJ, Michaels J, Powell JT, Thompson SG *et al.* Sex differences in mortality after abdominal aortic aneurysm repair in the UK. *Br J Surg* 2017;**104**:1656–1664
3. Deery SE, Soden PA, Zettervall SL, Shean KE, Bodewes TCF, Pothof AB *et al.* Sex differences in mortality and morbidity following repair of intact abdominal aortic aneurysms. *J Vasc Surg* 2017;**65**:1006–1013
4. Desai M, Choke E, Sayers RD, Nath M, Bown MJ. Sex-related trends in mortality after elective abdominal aortic aneurysm surgery between 2002 and 2013 at National Health Service hospitals in England: less benefit for women compared with men. *Eur Heart J* 2016;**37**:3452–3460
5. Lo RC, Bensley RP, Hamdan AD, Wyers M, Adams JE, Schermerhorn ML *et al.* Gender differences in abdominal aortic aneurysm presentation, repair, and mortality in the Vascular Study Group of New England. *J Vasc Surg* 2013;**57**:1261–1268, 1268.e1–5
6. Grant SW, Grayson AD, Purkayastha D, Wilson SD, McCollum C. Logistic risk model for mortality following elective abdominal aortic aneurysm repair. *Br J Surg* 2011;**98**:652–658
7. Chaikof EL, Dalman RL, Eskandari MK, Jackson BM, Lee WA, Mansour MA *et al.* The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *J Vasc Surg* 2018;**67**:2–77.e2
8. Mehta M, Byrne WJ, Robinson H, Roddy SP, Paty PS, Kreienberg PB *et al.* Women derive less benefit from elective endovascular aneurysm repair than men. *J Vasc Surg* 2012;**55**:906–913
9. Nathan DP, Brinster CJ, Jackson BM, Wang GJ, Carpenter JP, Fairman RM *et al.* Predictors of decreased short- and long-term

- survival following open abdominal aortic aneurysm repair. *J Vasc Surg* 2011;**54**:1237–1243
10. Holt PJE, Poloniecki JD, Gerrard D, Loftus IM, Thompson MM. Meta-analysis and systematic review of the relationship between volume and outcome in abdominal aortic aneurysm surgery. *Br J Surg* 2007;**94**:395–403
 11. Hicks CW, Canner JK, Arhuidese I, Obeid T, Black JH III, Malas MB. Comprehensive assessment of factors associated with in-hospital mortality after elective abdominal aortic aneurysm repair. *JAMA Surg* 2016;**151**:838–845
 12. Indrakusuma R, Jalalzadeh H, Vahl AC, Koelemay MJW, Balm R. Editor's choice—sex related differences in peri-operative mortality after elective repair of an asymptomatic abdominal aortic aneurysm in the Netherlands: a retrospective analysis of 2013 to 2018. *Eur J Vasc Endovasc Surg* 2019;**58**:813–820
 13. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Int J Surg* 2014;**12**:1495–1499
 14. DICA. Jaarrapportage 2014 DSAA Aneurysma Chirurgie. <https://dica.nl/jaarrapportage-2014/dsaa.html> (accessed 7 October 2020)
 15. Copeland GP. The POSSUM system of surgical audit. *Arch Surg* 2002;**137**:15–19
 16. Copeland GP, Jones D, Walters M. POSSUM: a scoring system for surgical audit. *Br J Surg* 2005;**78**:355–360
 17. Landon BE, O'Malley AJ, Giles K, Cotterill P, Schermerhorn ML. Volume–outcome relationships and abdominal aortic aneurysm repair. *Circulation* 2010;**122**:1290–1297
 18. Berlin JA, Kimmel SE, Ten Have TR, Sammel MD. An empirical comparison of several clustered data approaches under confounding due to cluster effects in the analysis of complications of coronary angioplasty. *Biometrics* 1999;**55**:470–476
 19. Austin PC, Merlo J. Intermediate and advanced topics in multi-level logistic regression analysis. *Stat Med* 2017;**36**:3257–3277
 20. Wu S, Crespi CM, Wong WK. Comparison of methods for estimating the intraclass correlation coefficient for binary responses in cancer prevention cluster randomized trials. *Contemp Clin Trials* 2012;**33**:869–880
 21. Trenner M, Salvermoser M, Busch A, Reutersberg B, Eckstein HH, Kuehnl A. Effect modification of sex and age for the hospital volume–outcome relationship in abdominal aortic aneurysm treatment: secondary data analysis of the nationwide German diagnosis related groups statistics from 2005 to 2014. *J Am Heart Assoc* 2020;**9**:e014534
 22. Becquemin JP, Chemla E, Chatellier G, Allaire E, Mellièrè D, Desgranges P. Perioperative factors influencing the outcome of elective abdominal aorta aneurysm repair. *Eur J Vasc Endovasc Surg* 2000;**20**:84–89
 23. Steffen M, Schmitz-Rixen T, Jung G, Böckler D, Grundmann RT. [The DIGG risk score: a risk predictive model of perioperative mortality after elective treatment of intact abdominal aortic aneurysms in the DIGG register.] *Chirurg* 2019;**90**:913–920
 24. Scali ST, Beck AW, Sedrakyan A, Mao J, Venermo M, Faizer R et al. Hospital volume association with abdominal aortic aneurysm repair mortality: analysis of the international consortium of vascular registries. *Circulation* 2019;**140**:1285–1287
 25. Zettervall SL, Schermerhorn ML, Soden PA, McCallum JC, Shean KE, Deery SE et al. The effect of surgeon and hospital volume on mortality after open and endovascular repair of abdominal aortic aneurysms. *J Vasc Surg* 2017;**65**:626–634
 26. Gray WK, Day J, Horrocks M. Editor's choice—volume–outcome relationships in elective abdominal aortic aneurysm surgery: analysis of the UK Hospital Episodes Statistics database for the Getting It Right First Time (GIRFT) programme. *Eur J Vasc Endovasc Surg* 2020;**60**:509–517
 27. Trenner M, Kuehnl A, Salvermoser M, Reutersberg B, Geisbuesch S, Schmid V et al. Editor's choice—high annual hospital volume is associated with decreased in hospital mortality and complication rates following treatment of abdominal aortic aneurysms: secondary data analysis of the nationwide German DRG statistics from 2005 to 2013. *Eur J Vasc Endovasc Surg* 2018;**55**:185–194
 28. Phillips P, Poku E, Essat M, Woods HB, Goka EA, Kaltenthaler EC et al. Procedure volume and the association with short-term mortality following abdominal aortic aneurysm repair in European populations: a systematic review. *Eur J Vasc Endovasc Surg* 2017;**53**:77–88
 29. Holt PJE, Poloniecki JD, Khalid U, Hinchliffe RJ, Loftus IM, Thompson MM. Effect of endovascular aneurysm repair on the volume–outcome relationship in aneurysm repair. *Circ Cardiovasc Qual Outcomes* 2009;**2**:624–632
 30. Scali ST, Beck A, Sedrakyan A, Mao J, Behrendt CA, Boyle JR et al. Editor's choice—optimal threshold for the volume–outcome relationship after open AAA repair in the endovascular era: analysis of the International Consortium of Vascular Registries. *Eur J Vasc Endovasc Surg* 2021;**61**:747–755
 31. Anderson PL, Arons RR, Moskowitz AJ, Gelijns A, Magnell C, Faries PL et al. A statewide experience with endovascular abdominal aortic aneurysm repair: rapid diffusion with excellent early results. *J Vasc Surg* 2004;**39**:10–19
 32. Greenhalgh RM, Brown LC, Powell JT, Thompson SG, Epstein D. Endovascular repair of aortic aneurysm in patients physically ineligible for open repair. *N Engl J Med* 2010;**362**:1872–1880
 33. Dias-Neto M, Norton L, Sousa-Nunes F, Silva JR, Rocha-Neves J, Teixeira JF et al. Impact of gradual adoption of EVAR in elective repair of abdominal aortic aneurysm: a retrospective cohort study from 2009 to 2015. *Ann Vasc Surg* 2021;**70**:411–424
 34. Dias-Neto M, Mani K, Leite-Moreira A, Freitas A, Sampaio S. Nationwide analysis of intact abdominal aortic aneurysm repair in Portugal from 2000 to 2015. *Ann Vasc Surg* 2020;**66**:54–64.e1
 35. Tong T, Aber A, Chilcott J, Thokala P, Walters SJ, Maheswaran R et al. Volume–outcome relationships in open and endovascular repair of abdominal aortic aneurysm: administrative data 2006–2018. *Br J Surg* 2021;**108**:521–527
 36. Skibba AA, Evans JR, Hopkins SP, Yoon HR, Katras T, Kalbfleisch JH et al. Reconsidering gender relative to risk of rupture in the contemporary management of abdominal aortic aneurysms. *J Vasc Surg* 2015;**62**:1429–1436
 37. Ubbink DT, Koelemay MJW. Shared decision making in vascular surgery. why would you? *Eur J Vasc Endovasc Surg* 2018;**56**:749–750
 38. Arnaoutakis DJ, Scali ST, Neal D, Giles KA, Huber TS, Powell RJ et al. Surgeon experience association with patient selection and outcomes after open abdominal aortic aneurysm repair. *J Vasc Surg* 2020;**72**:1325–1336.e2
 39. Scali ST, Arnaoutakis DJ, Neal D, Giles KA, Goodney PP, Suckow BD et al. Association between surgeon case volume and years of practice experience with open abdominal aortic aneurysm repair outcomes. *J Vasc Surg* 2021;**73**:1213–1226.e2