

University of Groningen

Surveying the management of Achilles tendon ruptures in the Netherlands

Dams, Olivier C; van den Akker-Scheek, Inge; Diercks, Ron L; Wendt, Klaus W; Zwerver, Johannes; Reininga, Inge H F

Published in:
Knee Surgery, Sports Traumatology, Arthroscopy

DOI:
[10.1007/s00167-018-5049-5](https://doi.org/10.1007/s00167-018-5049-5)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Dams, O. C., van den Akker-Scheek, I., Diercks, R. L., Wendt, K. W., Zwerver, J., & Reininga, I. H. F. (2019). Surveying the management of Achilles tendon ruptures in the Netherlands: lack of consensus and need for treatment guidelines. *Knee Surgery, Sports Traumatology, Arthroscopy*, 27(9), 2754-2764. <https://doi.org/10.1007/s00167-018-5049-5>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Surveying the management of Achilles tendon ruptures in the Netherlands: lack of consensus and need for treatment guidelines

Olivier C. Dams¹ · Inge van den Akker-Scheek^{1,2} · Ron L. Diercks² · Klaus W. Wendt³ · Johannes Zwerver¹ · Inge H. F. Reininga³

Received: 2 February 2018 / Accepted: 29 June 2018
© The Author(s) 2018

Abstract

Purpose This study aimed to describe and analyse usual care of Achilles tendon ruptures (ATRs) by orthopaedic surgeons and trauma surgeons in the Netherlands.

Methods A nationwide online survey of ATR management was sent to all consultant orthopaedic and trauma surgeons in the Netherlands, requesting participation of those involved in ATR management. Data on individual characteristics and the entire ATR management (from diagnosis to rehabilitation) were gathered. Consensus was defined as $\geq 70\%$ agreement on an answer.

Results A total of 91 responses (70 orthopaedic surgeons and 21 trauma surgeons) were analysed. There was consensus on the importance of the physical examination in terms of diagnosis ($> 90\%$) and a lack of consensus on diagnostic imaging (ultrasound/MRI). There was consensus that non-surgical treatment is preferred for sedentary and systemically diseased patients and surgery for patients who are younger and athletic and present with larger tendon gap sizes. There was consensus on most of the non-surgical methods used: initial immobilisation in plaster cast with the foot in equinus position (90%) and its gradual regression (82%) every 2 weeks (85%). Only length of immobilisation lacked consensus. Surgery was generally preferred, but there was a lack of consensus on the entire followed protocol. Orthopaedic and trauma surgeons differed significantly on their surgical ($p=0.001$) and suturing techniques ($p=0.002$) and methods of postoperative immobilisation ($p<0.001$). Orthopaedic surgeons employed open repair and Bunnell sutures more often, whereas trauma surgeons used minimally invasive approaches and bone anchors. Rehabilitation methods and advised time until weight-bearing and return to sport varied. Orthopaedic surgeons advised a significantly longer time until return to sport after both non-surgical treatment ($p=0.001$) and surgery ($p=0.002$) than trauma surgeons.

Conclusion This is the first study to describe the entire ATR management. The results show a lack of consensus and wide variation in management of ATRs in the Netherlands. This study shows that especially the methods of the perioperative and rehabilitation phases were inconclusive and differed between orthopaedic and trauma surgeons. Further research into optimal ATR management regimens is recommended. In addition, to achieve uniformity in management more multidisciplinary collaboration between Dutch and international surgeons treating ATRs is needed.

Level of evidence Cross-sectional survey, Level V.

Keywords Achilles · Rupture · Epidemiology · Ankle injuries · Treatment · Diagnostics · Clinical protocol · Traumatology · Orthopaedics · Surgical procedures · Rehabilitation

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00167-018-5049-5>) contains supplementary material, which is available to authorized users.

✉ Olivier C. Dams
O.C.Dams@umcg.nl

Extended author information available on the last page of the article

Introduction

Although the Achilles tendon is capable of tolerating heavy loading, it can be susceptible to injury, as the most frequently ruptured tendon [1]. The incidence of Achilles tendon rupture (ATR) has steadily increased in the Western world over the past years [2–5] and is expected to rise further, especially among the middle-aged, likely

due to higher rates of recreational sports participation and obesity [6–9]. In addition to the rising incidence, ATRs can significantly burden patients, with more than half showing functional deficits and/or reporting pain even 12 months after injury, and many unable to return to their pre-injury level of activity [2, 10–12].

Despite these figures, a clear international management consensus for the treatment of ATRs is lacking. The only available guidelines are those of the American Academy of Orthopedic Surgeons (AAOS) and have limited or inconclusive recommendations for the role of imaging, choice of treatment (non-surgical/ surgical) and rehabilitation methods as a result of either lacking or unconvincing scientific evidence [13]. In addition, neither the UK's National Institute for Health and Care Excellence (NICE), the British Orthopaedic Association, the Nordic Orthopaedic Federation, the Dutch Society for Orthopaedic Surgery (NOV) nor the Dutch Society for Trauma Surgery (NVT) has published evidence-based guidelines for the management of ATRs. Because of this lack of evidence, we hypothesise that all phases of ATR management in the Netherlands lack consensus; this leads to a divergent clinical protocol and potentially variation in outcome for patients.

Two studies have described multiple phases of reported ATR management in the United Kingdom (UK) and Scandinavia, concluding that there is a wide variation in treatment options that practitioners adhere to and that clear consensus is lacking [14, 15]. However, no such studies have been conducted in the Netherlands where ATRs are treated by both trauma and orthopaedic surgeons.

The aim of this study was to describe and analyse usual ATR care in the Netherlands, to create transparency about how this injury is being managed without clinical guidelines. Differences within and between the two treating specialisms were also compared and contrasted. This is the first study to describe the entire state of practice and the first to do so in the Netherlands. The analysis will highlight the barriers to achieving clinical guidelines, guide future research directions and allow clinicians to reflect on and compare their ATR treatment.

Materials and methods

Study design

Design of this study was a cross-sectional survey of practice. The local ethics committee of the University Medical Center Groningen judged the methods employed and waived further need for approval (METc #2016.475).

Study population

All consultant orthopaedic surgeons ($n = 601$) and trauma surgeons ($n = 230$) registered in their respective Dutch specialist societies [the Dutch Societies for Orthopaedic Surgery (NOV) and Trauma Surgery (NVT)] were approached as potential study participants. Within these groups, those surgeons actually treating ATRs in their respective hospitals were asked to participate by filling in an online survey about ATR management. We estimate that the study population of surgeons actually treating ATRs consists of approximately 150 orthopaedic surgeons and 50 trauma surgeons, based on one or two surgeons per hospital treating ATRs.

The survey

The survey was developed by epidemiologists, orthopaedic surgeons, trauma surgeons and sport and exercise medicine physicians with relevant clinical and methodological expertise (OCD, IvdAS, RLD, KWW, JZ, IHFR). It was designed to assess responders' characteristics and the complete applied/preferred management decisions from diagnosis to rehabilitation. In designing the questions, attention was paid to recommendations made in the AAOS guidelines [13]. The survey consisted of a minimum of 21 and a maximum of 35 items, depending on the type of ATR treatment (non-surgical, surgical or both) responders use in their practice. Most questions were presented in multiple-choice format. The Appendix describes the survey questions.

Items in the survey were divided into four major sections: (1) individual characteristics, (2) diagnostic tools used, (3) preferred/applied primary treatment of ATRs, and (4) rehabilitation methods and advised time until return to sport (RTS). In each section responders were asked questions about:

1. Their medical specialism, years of experience, practice setting and number of ATRs treated per year.
2. Use of diagnostic tools [physical examination: palpation of tendon gap, Thompson test [16]; imaging: ultrasound (US), MRI, X-ray and CT] for which responders were asked to select all they apply for diagnosis and treatment-planning.
3. Type of preferred primary treatment (non-surgical or surgical) in the context of the individual patient, dependent on clinical factors such as age, American Society of Anesthesiologists (ASA) physical status, body mass index (BMI), activity level, tendon gap size and time between presentation and injury. Responders had to

report whether they had a general preference for non-surgical or surgical treatment or whether they treated one way or the other exclusively. Subsequently, they were presented with an online pathway asking about the non-surgical and/or surgical treatment methods used.

4. Methods of rehabilitation, referral to physiotherapists, time advised for commencement of return to sport (RTS) and a selection of multiple outcomes (patient follow-up, questionnaires, physical tests, imaging) deemed relevant when monitoring recovery. For this study, the start of the rehabilitation phase was defined as the period after initial immobilisation in non-surgical patients and 6 weeks after surgery in surgical patients. Rehabilitation methods and advised time to RTS were asked about after both non-surgical and surgical treatment.

Survey administration

Consultant surgeons were approached through the regular electronic newsletters of their Dutch specialist associations, NOV and NVT. The newsletters contained a brief description and a hyperlink to the survey. Although the survey was sent to all consultant surgeon members, it asked exclusively for participation of surgeons actually treating ATRs. The survey was sent out once to the NOV in December 2016 and twice to the NVT in December 2016 and January 2017.

In the online survey, environment responders were presented with a distinct pathway concerning only the respective treatment methods (non-surgical and/or surgical) they use. This pathway was generated by the question about the general preferred treatment, for which responders could indicate if they only treated surgically or only treated non-surgically. Data of the survey were stored on the server of the University Medical Center Groningen.

Statistical analysis

Data were recorded on a tabulated form on Microsoft Excel 2010. All the required variables were converted to IBM SPSS Statistics for Windows software (Version 23.0, Armonk, NY: IBM Corp.) for statistical analysis. A p value < 0.05 was considered statistically significant in all analyses.

Descriptive statistics n (%) were calculated for the categorical survey data on all individual responder characteristics and applied/preferred reported management. Group comparisons were performed using chi-squared tests for nominal data and Kruskal–Wallis/Mann–Whitney U tests for ordinal data.

Odds ratios (ORs) including 95% confidence intervals (CIs) were calculated for dichotomous categorical variables using binary logistic regression analysis with specialism (comparing trauma surgeons with orthopaedic surgeons) as

independent variable and the response (e.g. yes vs. no or surgical vs. non-surgical) as dependent variable.

Consistent with the definition employed by Sumsion regarding the Delphi survey technique, consensus was defined as $\geq 70\%$ of agreement on a topic [17].

Due to the descriptive nature of this study, no formal sample size calculation was performed prior to data collection. It was aimed to include as many ATR-treating surgeons as possible to accurately describe the current state of practice.

Results

Survey responders

A total of 91 medical specialists completed the survey: 70 orthopaedic surgeons and 21 trauma surgeons. The response rate amounts to a total of 47% (70/150) of orthopaedic surgeons and 42% (21/50) of trauma surgeons based on the estimated number of eligible participants. Table 1 shows the responders' characteristics. No statistically significant differences were found in terms of experience or practice setting between orthopaedic and trauma surgeon responders. Individually, trauma surgeons treated significantly more ATRs per year ($p < 0.001$) than orthopaedic surgeons.

Diagnosis

Responders used physical examination, palpation of tendon gap (92%) or the Thompson test (91%) as primary diagnostic tool. Concerning imaging, US was used by 45% of

Table 1 Characteristics of survey responders

Characteristic	n (%)
Experience (years)	
0–5	18 (20)
5–10	21 (23)
10–15	15 (17)
15–20	15 (17)
> 20	22 (24)
Practice setting	
Non-academic hospital without residents	27 (30)
Non-academic hospital with residents	43 (47)
Academic hospital	16 (18)
Private clinic	4 (4)
Combination of academic and private practice	1 (1)
Number of ATRs treated/year	
< 5	29 (32)
5–15	46 (51)
15–25	11 (12)
> 25	5 (6)

responders, one responder used X-rays, one MRI, and no responders used CT to diagnose an ATR.

Subgroup analyses based on specialism, experience, practice setting or number of ATRs treated showed no significant differences in the diagnostic modalities chosen.

Primary treatment

General preferred primary treatment

Surgery was the reported preferred treatment among all responders (52%). Non-surgical treatment was preferred by 34%, and 14% had no preference. One responder treated all ATRs surgically, none treated only non-surgically. Although not statistically significant, trauma surgeons tended to prefer surgical treatment more often than orthopaedic surgeons (71 vs. 45%, $p=0.12$).

Subgroup analyses based on experience, practice setting or number of ATRs treated showed no significant differences in preferred treatment.

Preferred primary treatment per patient factor

Table 2 shows the preferred primary treatment based on specific clinical factors. There was consensus among responders that non-surgical treatment was preferred for patients presenting with an ASA status >3 and patients with a sedentary lifestyle. Surgical treatment was preferred for patients who were younger (age <40) and athletic, and when a gap size >1 cm was present. There was no consensus on a preferred primary treatment based on BMI or chronicity of rupture. Only seven responders (8%) considered other patient-related factors such as smoking, diabetes mellitus, impaired arterial circulation of the lower extremity and rupture location (distal or midportion) for their treatment choice.

Orthopaedic and trauma surgeons differed significantly in their preferred primary treatment of patients older than 40 ($p=0.02$) and with ASA status <3 ($p=0.04$); trauma surgeons preferred surgical treatment more often for both (OR 3.19, 95% CI 1.14–8.90 and OR 3.38, 95% CI 1.03–11.07). Although not statistically significant, more trauma surgeons seemed to prefer surgical treatment for patients with a higher BMI ($p=0.08$) and gap size <1 cm ($p=0.09$) than orthopaedic surgeons.

Subgroup analyses based on experience or number of ATRs treated showed no significant differences by clinical factors in preferred primary treatment.

Non-surgical treatment methods

The non-surgical treatment methods are presented in Table 3. There was consensus that the initial immobilisation should consist of placing the foot in a plaster cast in equinus

Table 2 Treatment preference by clinical factors

Factor	Treatment	n (%)
Age <40 years	Surgical	67 (74)
	Non-surgical	24 (26)
Age >40 years	Surgical	41 (45)
	Non-surgical	50 (55)
ASA <3	Surgical	56 (62)
	Non-surgical	35 (38)
ASA >3	Surgical	5 (5)
	Non-surgical	86 (95)
Athletic	Surgical	72 (79)
	Non-surgical	19 (21)
Sedentary	Surgical	6 (6)
	Non-surgical	85 (94)
BMI <30 kg/m ²	Surgical	58 (64)
	Non-surgical	33 (36)
BMI >30 kg/m ²	Surgical	33 (36)
	Non-surgical	58 (64)
Gap size <1 cm	Surgical	46 (51)
	Non-surgical	45 (49)
Gap size >1 cm	Surgical	68 (75)
	Non-surgical	23 (25)
Injury <6 weeks old	Surgical	58 (64)
	Non-surgical	33 (36)
Injury >6 weeks old	Surgical	38 (42)
	Non-surgical	53 (58)

Bold indicates consensus was reached among responders, ($>70\%$) agreed on an answer

position (90%). No consensus was found on the exact length of initial immobilisation or the time at which weight-bearing was allowed; although there was consensus for weight-bearing within 6 weeks (89%), the chosen period ranged from 2 to 12 weeks. The two most chosen durations of immobilisation were either 2 weeks (46%) or 6 weeks (38%). There was consensus that the foot position should be changed (82%) and that this should be done every 2 weeks (85%).

Orthopaedic surgeons and trauma surgeons differed significantly on their methods of initial immobilisation ($p=0.001$), with trauma surgeons prescribing methods other than a plaster cast more often (24 vs. 6%). Although not statistically significant, trauma surgeons tended to prescribe a shorter length of initial immobilisation ($p=0.07$), with 72% of trauma surgeons immobilising for ≤ 4 weeks compared to 51% of orthopaedic surgeons. Responders with more experience recommended a longer period until the patient could bear weight ($p=0.002$), with 72% of responders with ≥ 15 years' experience recommending weight-bearing after 6 weeks compared to 40% of those with <15 years' experience.

Table 3 Methods of non-surgical and surgical treatment

Treatment method	Surgical <i>n</i> (%)	Non-surgical ^a <i>n</i> (%)
Preoperative antibiotics		
Yes	37 (41)	
No	54 (59)	
Preoperative anticoagulants		
Yes	38 (42)	
No	53 (58)	
Surgical technique		
Open repair	59 (65)	
Augmented repair	5 (5)	
Percutaneous	9 (10)	
Combined mini-open	17 (19)	
Other not specified	1 (1)	
Tunneling through the calcaneus		
Yes	12 (13)	
No	79 (87)	
Suturing technique		
Bunnell	50 (55)	
Kessler	19 (21)	
Epitendinous	2 (2)	
Mitek-anchors	7 (8)	
Krackow	2 (2)	
Other	11 (12)	
Type of sutures		
Absorbable	61 (67)	
Non-absorbable	30 (33)	
Initial immobilisation		
Equinus and plaster cast	54 (59)	81 (90)
Non-equinus and plaster cast	24 (26)	2 (2)
Boot/brace	9 (10)	5 (6)
Tape	4 (4)	2 (2)
Length of initial immobilisation (weeks)		
2		41 (46)
3		3 (33)
4		6 (7)
5		1 (1)
6		34 (38)
> 6		5 (6)
Change foot position during immobilisation		
Yes	61 (67)	74 (82)
No	30 (33)	16 (18)
Frequency of foot position change ^b		
Every week	7 (11)	7 (9)
Every 2 weeks	52 (85)	63 (85)
Every 3 weeks	2 (3)	3 (4)
Every 4 weeks	0 (0)	1 (1)
When can the patient bear weight (weeks)		
2	35 (38)	30 (33)

Table 3 (continued)

Treatment method	Surgical <i>n</i> (%)	Non-surgical ^a <i>n</i> (%)
4	14 (15)	12 (13)
6	34 (37)	39 (43)
8	4 (4)	4 (4)
10	2 (2)	2 (2)
12	1 (1)	3 (3)

Bold indicates consensus was reached among responders, (> 70%) agreed on an answer

^aOne orthopaedic surgeon did not complete the questions as he/she never treats non-surgically

^bEight trauma surgeons and 22 orthopaedic surgeons (surgical) and three trauma surgeons and 13 orthopaedic surgeons (non-surgical) did not answer this question as they did not change the foot position during immobilisation

Subgroup analyses based on practice setting or number of ATRs treated showed no significant differences in non-surgical treatment methods.

Surgical treatment methods

The surgical treatment methods are presented in Table 3. There was no consensus on the administration of preoperative antibiotics and anticoagulants, surgical technique, suturing type and methods or postsurgical immobilisation methods. Open repair (65%) was the most common surgical technique and Bunnell sutures (55%) were mostly applied. The postoperative immobilisation method was the same as that chosen for non-surgical: plaster cast with the foot in equinus position (59%).

Trauma surgeons and orthopaedic surgeons differed significantly on surgical technique ($p=0.001$), suturing technique ($p=0.002$) and initial immobilisation methods ($p<0.001$). The most frequent choices of trauma surgeons were minimally invasive techniques [combined mini-open (48%) and percutaneous (14%)], with orthopaedic surgeons gravitating towards open repair techniques (73%). Trauma surgeons used bone anchors more often (24 vs. 3%), and orthopaedic surgeons Bunnell sutures (61 vs. 33%). Trauma surgeons prescribed a brace as initial postoperative immobilisation method more often than orthopaedic surgeons (29 vs. 4%).

Subgroup analyses based on practice setting or number of ATRs treated showed no significant differences in surgical treatment methods.

Table 4 Rehabilitation methods

Rehabilitation method	Surgical <i>n</i> (%)	Non-surgical ^a <i>n</i> (%)
Applied protection		
Walking boot	23 (25)	27 (30)
Heel-lift	31 (34)	28 (31)
Brace	4 (4)	15 (17)
Tape	14 (15)	14 (16)
None	19 (21)	5 (6)
Not specified		1 (1)
Referral to physiotherapist		
Yes	79 (87)	79 (88)
No	12 (13)	11 (12)
Monitoring modality		
Patient follow-up	85 (93)	84 (92)
Questionnaires	5 (6)	4 (4)
Heel-rise test	17 (19)	17 (19)
Tolerated load on tendon	16 (18)	15 (17)
US	0 (0)	1 (1)
MRI	0 (0)	0 (0)

Bold indicates consensus was reached among responders, (> 70%) agreed on an answer

^aOne orthopaedic surgeon did not answer the questions as he/she never treats non-surgically

Rehabilitation

Table 4 shows that responders differed greatly on rehabilitation methods after surgical and non-surgical treatment. The type of protection varied, with heel lifts and walking boots as the most commonly used. Although there was consensus among responders to refer their patients to physiotherapists, trauma surgeons were significantly less likely to refer patients to physiotherapy after surgery than orthopaedic surgeons (OR 0.23; 95% CI 0.06–0.83). Tendon healing was primarily monitored via patient follow-up, one responder used US and none used MRI to monitor healing. There were

no significant differences in rehabilitation methods after surgical and non-surgical treatment.

Subgroup analyses based on experience, practice setting or number of ATRs treated showed no significant differences in methods of rehabilitation after non-surgical treatment or surgery.

The advised time to commence RTS after both surgery and non-surgical treatment ranged from 2 to 6 weeks to > 26 weeks after initial injury, with 14–18 weeks as most responded answer. There was no significant difference in advised time to RTS after surgical or non-surgical treatment. Figure 1 shows trauma and orthopaedic surgeons differed significantly on advised time to RTS after surgical ($p=0.002$) and non-surgical treatment ($p=0.001$). Orthopaedic surgeons recommended a longer period until patients could RTS.

Subgroup analyses based on experience, practice setting or number of ATRs treated showed no significant differences in advised time to return to sport.

Discussion

The most important finding of this present study was the variation in care of ATRs in the Netherlands provided by orthopaedic and trauma surgeons. To our knowledge this is the first study to gather such information and to describe the entire ATR management applied from diagnosis to rehabilitation.

This description showed considerable practice variation and a lack of consensus on ATR management, particularly concerning management in the perioperative and rehabilitation phases. There are also significant differences in applied/preferred management of ATRs between orthopaedic and trauma surgeons. Orthopaedic surgeons tended to prefer non-surgical treatment more often, followed a different perioperative protocol and advised a longer period before patients could return to sport. Lastly, management—especially concerning time to weight-bearing as well as preferred primary




Fig. 1 Advised time to RTS after **a** surgical, **b** non-surgical treatment

treatment for specific patient groups—is not in concordance with recent scientific evidence and clinical guidelines.

Although only an estimated response rate was provided, we consider the sample of this study a good representation of ATR-treating orthopaedic surgeons and trauma surgeons in the Netherlands. Due to subspecialisations in the Dutch healthcare system, usually one or two surgeons per hospital or department treat ATRs (foot/ankle and/or sport-injury specialists in their respective practices). Given the total number of eight academic hospitals and 83 general hospitals in the Netherlands [18], we estimate that the ATR-treating specialists in at least 60% of all practice settings are represented in the results. Multiplying the median number of ATRs treated by the responders (Table 1) amounts to roughly 900 ATRs per year. This is equivalent to approximately 80% of all ATRs presenting in the Netherlands. We, therefore, consider the 91 responses that we analysed to accurately reflect current ATR management in the Netherlands. This is a higher absolute response number than described by the two other studies that surveyed individual specialists (in the UK) on acute ATR management [15, 19].

Current practice

Diagnostics

There is consensus on the diagnosis and the diagnostic tools used in ATRs: responders relied on the physical examination rather than imaging. This low dependence on imaging is likely due to the ease and high diagnostic accuracy of clinical tests [20] as well as the ‘typical’ ATR presentation involving sports trauma, an audible snap and/or a feeling of being kicked. In addition, recent surveys show that Dutch orthopaedic surgeons see no additional value in diagnostic musculoskeletal imaging [21]. This preference adheres to the recommendations made by the AAOS and a recent systematic review for a comprehensive physical examination of ATR patients [13, 22].

Primary treatment

In terms of primary treatment, responders agreed on non-surgical management for systemically diseased (ASA > 3) and sedentary patients with an ATR. The recommendation made in the AAOS guidelines to ‘cautiously approach surgery in the sedentary, obese and systemically diseased’ was adhered to [13]. In this study, obesity (BMI > 30 kg/m²) was not reported to be a clear indication for non-surgical treatment. Obesity is a significant risk factor for developing an ATR [7, 23] as well as for infection and other complications following orthopaedic surgery [24]. The lack of consensus on the treatment of obese patients requires additional

awareness of the AAOS guideline and careful consideration by surgeons treating ATRs.

There was consensus that surgery is the preferred treatment for athletic patients, as is also preferred by Scandinavian and British specialists and supported by the literature [14, 15, 25, 26]. The agreement among the responders for surgical treatment of ATRs with a larger tendon gap size (> 1 cm) adheres to recent scientific evidence showing non-surgical managed patients with a gap size of > 1 cm to have a worse (functional) outcome [27]. Future research, however, should establish at which specific gap size non-surgical treatment should be considered, as prospective studies have employed various ultrasonographically measured gap sizes (5 mm and 1 cm) without sufficient evidence to support a specific cutoff point [25, 28, 29].

Literature has defined ATRs older than 4 weeks as ‘chronic’ [30]. These undiagnosed or untreated ruptures cause considerable morbidity and require special attention. The results showed that there is no consensus on management of these ruptures (> 6 weeks old), as responders supported both surgical and non-surgical treatments. This inconclusiveness is in line with the literature, supporting both treatments [31, 32]. To date there is insufficient evidence for clear treatment indications; future research should examine both treatment modalities for chronic ruptures.

In contrast to practice in other countries, Dutch surgeons did not generally prefer non-surgical treatment [33, 34], yet consensus was found on most of its primary methods. The protocol for non-surgical ATR treatment was similar to the SMART protocol proposed by Hutchison et al. [28], who also emphasised initial immobilisation in equinus position with casting for 2 weeks followed by gradual regression of foot position and physiotherapy. The SMART protocol resulted in the lowest rate of re-ruptures to date, as well as excellent patient-reported outcomes [28]. Despite these promising results, Dutch surgeons seem not to adhere to this protocol entirely and prescribed either 2 or 6 weeks of initial immobilisation. This prescribed period is dissimilar to the 1-week period applied in recent Dutch papers [35, 36]. We recommend further adherence to this non-surgical treatment protocol with short casting (maximum 2 weeks), followed by careful weight-bearing.

Responders generally preferred surgical treatment, yet there was a lack of consensus on perioperative methods. This might be due to a dearth of scientific evidence: a systematic review yields that only four trials have compared open and less invasive techniques, showing no significant difference in outcome and insufficient evidence comparing suturing methods [37]. We hypothesise that the preference for surgery among responders is due to the influence of several Dutch studies showing the reliable results of a minimally invasive technique [38, 39] and its superiority to non-surgical treatment in terms of complications [36]. Nonetheless, we

recommend that Dutch surgeons re-evaluate their general preference for surgical treatment, as recent evidence establishing that non-surgical treatment that shows similar results has produced an international rise (in Scandinavia and North America) in the use of non-surgical ATR treatment [3, 33, 34, 40]. There is consensus on postoperative immobilisation after ATR repair, probably based on the assumption that early mobilisation puts patients at risk for re-ruptures. Several systematic reviews, however, have determined immobilisation may not be necessary or helpful at all, and that early dynamic rehabilitation using a brace results in safe results with higher patient satisfaction [41]; superior, more rapid recovery [42]; and a quicker return to sporting activities [43]. A trend towards minimal immobilisation (2 weeks) is also advocated after peroneal tendon repair [44]. Similarly to non-surgical treatment we therefore recommend weight-bearing as early as possible after surgery, as already proposed by other Dutch authors [36, 45].

Rehabilitation

Opinions and usual care concerning the rehabilitation phase varied, and practice seemed to be based on the individual decisions. This is in line with previous research on ATR rehabilitation practice in the UK [46]. Although early weight-bearing (<2 weeks) after surgery is associated with fewer complications and better functional recovery [27, 41, 42, 47–49], only 39% of specialists allow for weight-bearing within 2 weeks of surgery and 33% after non-surgical treatment. This study showed that many Dutch surgeons (49%) do not adhere to the AAOS recommendations of early mobilisation (2–4 weeks). This contrasts the practice of Scandinavian surgeons, 83–100% of whom allow weight-bearing within 4 weeks [14] as well as the weight-bearing protocols proposed in Dutch publications [36, 38, 39, 45].

The protection measures used during rehabilitation varied. This was also reported in the UK [46] and is in line with the lack of evidence in the literature. With the availability of novel orthoses and increasing use of functional rehabilitation more high-quality research on rehabilitation protection is required.

There was consensus to refer patients to physiotherapists. Qualitative research has shown that trauma patients report physiotherapy as beneficial to recovery [50], implying that this referral potentially increases patient satisfaction. Nonetheless, the Royal Dutch Society for Physiotherapy (KNGF) has no recommendations or guideline statements for physiotherapists managing rehabilitation after ATR, which further impairs uniformity in ATR rehabilitation methods in the Netherlands.

A recent systematic review on RTS following ATR reported a range of 2.9–10.4 months [51] and a mere 80% return to their pre-injury levels of physical activity at ATR

[51]. Likewise, in this survey no consensus on the topic was found. Because many ATR patients are physically active and ultimately sustain their rupture during sports [9], we believe a minimum period until RTS should be determined to optimise return yet avoid performance deficits and/or re-injury. We recommend this minimum period takes into consideration the vulnerable phase in which most re-ruptures occur (6–12 weeks) [52]. Ultimately, in clinical practice RTS should be tailored to each individual patient, while also considering other factors such as their fear of re-injury and motivation for RTS, among other things. This study, therefore, concludes that a multifaceted approach involving cooperation between surgeons, sports and exercise medicine physicians, physiotherapists and sport psychologists is required.

Responder comparison

Nearly all of the differences in ATR management between responders were based on their specialism (trauma vs. orthopaedic surgery) as opposed to experience, practice setting, or number of ATRs treated per year. Non-surgical treatment and different immobilisation and surgical methods were more likely to be used by orthopaedic surgeons. Orthopaedic surgeons generally employed more traditional methods of ATR management such as immobilisation via plaster casts and open repair with Bunnell sutures, whereas trauma surgeons were more likely to use various orthoses and minimally invasive surgical techniques. Orthopaedic surgeons were also more conservative in their advised period of return to sport, advising a longer wait.

At first glance, these differences in usual practice between trauma and orthopaedic surgeons could be explained by the structure of the Dutch healthcare system, where orthopaedic surgery and trauma surgery are different specialisms following distinct medical training, yet both treating trauma patients. As a prior survey on the treatment of ankle fractures in the Netherlands showed few or no differences between orthopaedic and trauma surgeons [53], the lack of guidelines is most probably what underlies differences in ATR management between the two treating specialisms. Perhaps each specialism has separate guidelines written by individual experts with a limited following, hence the differences in management.

The clinical relevance of this study is that it allows clinicians in the Netherlands and abroad to reflect on and compare their current ATR practice, guide treatment decisions and define future (research) directions. The findings suggest that to develop clinical guidelines, multidisciplinary collaboration is required; this warrants the cooperation of orthopaedic surgeons and trauma surgeons in the Netherlands.

Conclusion

This study provides insight into current management of ATR in the Netherlands. The diagnosis of ATR and indications for primary non-surgical or surgical treatment based on clinical factors are only partially in concordance with the limited available scientific evidence and guidelines. There is a general lack of consensus among individual specialists as well as significant differences in preferred and applied management between Dutch orthopaedic surgeons and trauma surgeons. The use of existing evidence, the development of clinical guidelines for primary treatment of different patient populations and the application of evidence-based rehabilitation principles should be encouraged.

Acknowledgements We would like to thank Pieter Davids from the University Medical Center Groningen Department of Communications for the technical assistance in developing the survey, the NOV and NVT for their cooperation in distributing the survey and all those who took the time to respond to this survey.

Funding This study was not funded.

Compliance with ethical standards

Conflict of interest The authors declare that they have no competing interest.

Ethical approval Approved by the local ethics committee of the University Medical Center Groningen (METc #2016.475).

Informed consent Informed consent is not required for this type of article.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Pedowitz D, Kirwan G (2013) Achilles tendon ruptures. *Curr Rev Musculoskelet Med* 6:285–293
- Holm C, Kjaer M, Eliasson P (2014) Achilles tendon rupture - treatment and complications: a systematic review. *Scand J Med Sci Sports* 25:e1–e10. <https://doi.org/10.1111/sms.12209>
- Huttunen TT, Kannus P, Rolf C, Fellander-Tsai L, Mattila VM (2014) Acute achilles tendon ruptures: incidence of injury and surgery in Sweden between 2001 and 2012. *Am J Sports Med* 42:2419–2423
- Lanto I, Heikkinen J, Flinkkilä T, Ohtonen P, Leppilähti J (2015) Epidemiology of Achilles tendon ruptures: increasing incidence over a 33-year period. *Scand J Med Sci Sport* 25:e133–e138. <https://doi.org/10.1111/sms.12253>
- Nyyssönen T, Lüthje P, Kröger H (2008) The increasing incidence and difference in sex distribution of Achilles tendon rupture in Finland in 1987–1999. *Scand J Surg* 97:272–275
- Erickson BJ, Cvetanovich GL, Nwachukwu BU, Villarreal LD, Lin JL, Bach BR, McCormick FM (2014) Trends in the management of Achilles tendon ruptures in the United States medicare population, 2005–2011. *Orthop J Sport Med* 2:2325967114549948
- Gaida JE, Ashe MC, Bass SL, Cook JL (2009) Is adiposity an under-recognized risk factor for tendinopathy? A systematic review. *Arthritis Care Res* 61:840–849
- Ho G, Tantigate D, Kirschenbaum J, Greisberg JK, Vosseller JT (2017) Increasing age in Achilles rupture patients over time. *Injury* 48:1701–1709
- Raikin SM, Garras DN, Krapchev PV (2013) Achilles tendon injuries in a United States population. *Foot Ankle Int* 34:475–480
- Fox G, Gabbe BJ, Richardson M, Oppy A, Page R, Edwards ER, Hau R, Ekegren CL (2016) Twelve-month outcomes following surgical repair of the Achilles tendon. *Injury* 47:2370–2374
- Rosso C, Buckland DM, Polzer C, Sadoghi P, Schuh R, Weisskopf L, Vavken P, Valderrabano V (2015) Long-term biomechanical outcomes after Achilles tendon ruptures. *Knee Surg Sports Traumatol Arthrosc* 23:890–898
- Trofa DP, Miller JC, Jang ES, Woode DR, Greisberg JK, Vosseller JT (2017) Professional Athletes' return to play and performance after operative repair of an Achilles tendon rupture. *Am J Sports Med* 45:2864–2871
- Chiodo CP, Glazebrook M, Bluman EM, Cohen BE, Femino JE, Giza E, Watters WC 3rd, Goldberg MJ, Keith M, Haralson RH 3rd, Turkelson CM, Wies JL, Raymond L, Anderson S, Boyer K, Sluka P, American Academy of Orthopaedic Surgeons (2010) Diagnosis and treatment of acute Achilles tendon rupture. *J Am Acad Orthop Surg* 18:503–510
- Barfod KW, Nielsen F, Helander KN, Mattila VM, Tingby O, Boesen A, Troelsen A (2013) Treatment of acute achilles tendon rupture in scandinavia does not adhere to evidence-based guidelines: a cross-sectional questionnaire-based study of 138 departments. *J Foot Ankle Surg* 52:629–633
- Worth N, Ghosh S, Maffulli N (2007) Management of acute Achilles tendon ruptures in the United Kingdom. *J Orthop Surg (Hong Kong)* 15:311–314
- Thompson TC (2003) A test for rupture of the tendo achillis. *Acta Orthop Scand* 32:461–465
- Sumsion T (1998) The delphi technique: an adaptive research tool. *Br J Occup Ther* 61:153–156
- Dutch Hospital Data (2016) Kengetallen Nederlandse Ziekenhuizen 2014. Utrecht
- Osarumwense D, Wright J, Gardner K, James L (2013) Conservative treatment for acute Achilles tendon rupture: survey of current practice. *J Orthop Surg (Hong Kong)* 21:44–46
- Maffulli N (1998) The clinical diagnosis of subcutaneous tear of the Achilles tendon. A prospective study in 174 patients. *Am J Sports Med* 26:266–270
- Scholten-Peters GGM, Franken N, Beumer A, Verhagen AP (2014) The opinion and experiences of Dutch orthopedic surgeons and radiologists about diagnostic musculoskeletal ultrasound imaging in primary care: a survey. *Man Ther* 19:109–113
- Dams OC, Reininga IHF, Gielen JL, van den Akker-Scheek I, Zwerver J (2017) Imaging modalities in the diagnosis and monitoring of Achilles tendon ruptures: a systematic review. *Injury* 48:2383–2399
- Claessen FMAP, de Vos RJ, Reijman M, Meuffels DE (2014) Predictors of primary Achilles tendon ruptures. *Sports Med* 44:1241–1259
- Yuan K, Chen HL (2013) Obesity and surgical site infections risk in orthopedics: a meta-analysis. *Int J Surg* 11:383–388

25. Guss D, Smith JT, Chiodo CP (2015) Acute achilles tendon rupture: a critical analysis review. *JBJS Rev.* <https://doi.org/10.2106/JBJS.RVW.N.00015>
26. Maffulli N, Longo UG, Maffulli GD, Khanna A, Denaro V (2011) Achilles tendon ruptures in elite athletes. *Foot Ankle Int* 32:9–15
27. Lawrence JE, Nasr P, Fountain DM, Berman L, Robinson AHN (2017) Functional outcomes of conservatively managed acute ruptures of the Achilles tendon. *Bone Joint J* 99–B:87–93
28. Hutchison AM, Topliss C, Beard D, Evans RM, Williams P (2015) The treatment of a rupture of the achilles tendon using a dedicated management programme. *Bone Joint J* 97–B:510–515
29. Kotnis R (2006) Dynamic ultrasound as a selection tool for reducing Achilles tendon reruptures. *Am J Sports Med* 34:1395–1400
30. Flint JH, Wade AM, Giuliani J, Rue J-P (2014) Defining the terms acute and chronic in orthopaedic sports injuries: a systematic review. *Am J Sports Med* 42:235–241
31. Maffulli N, Ajis A (2008) Management of chronic ruptures of the Achilles tendon. *J Bone Joint Surg Am* 90:1348–1360
32. Maffulli N, Via AG, Oliva F (2015) Chronic Achilles tendon disorders: tendinopathy and chronic rupture. *Clin Sports Med* 34:607–624
33. Ganestam A, Kallemose T, Troelsen A, Barfod KW (2016) Increasing incidence of acute Achilles tendon rupture and a noticeable decline in surgical treatment from 1994 to 2013. A nationwide registry study of 33,160 patients. *Knee Surg Sports Traumatol Arthrosc* 24:3730–3737
34. Sheth U, Wasserstein D, Jenkinson R, Moineddin R, Kreder H, Jaglal S (2017) Practice patterns in the care of acute Achilles tendon ruptures. *Bone Joint J* 99–B:1629–1636
35. Metz R, van der Heijden GJMG, Verleisdonk E-JMM, Tamminga R, van der Werken C (2009) Recovery of calf muscle strength following acute achilles tendon rupture treatment: a comparison between minimally invasive surgery and conservative treatment. *Foot Ankle Spec* 2:219–226
36. Metz R, Verleisdonk E-JMM, van der Heijden GJ-M-G, Clevers G-J, Hammacher ER, Verhofstad MHJ, van der Werken C (2008) Acute Achilles tendon rupture: minimally invasive surgery versus nonoperative treatment with immediate full weightbearing—a randomized controlled trial. *Am J Sports Med* 36:1688–1694
37. Khan RJ, Carey Smith RL (2010) Surgical interventions for treating acute Achilles tendon ruptures. *Cochrane database Syst Rev* 9:CD003674. <https://doi.org/10.1002/14651858.CD003674.pub4>
38. Groetelaers RPTGC, Janssen L, van der Velden J, Wieland AWJ, Amendt AGFM, Geelen PHJ, Janzing HMJ (2014) Functional treatment or cast immobilization after minimally invasive repair of an acute Achilles tendon rupture: prospective, randomized trial. *Foot Ankle Int* 35:771–778
39. Lansdaal JR, Goslings JC, Reichart M, Govaert GAM, van Scherpenzeel KM, Haverlag R, Ponsen KJ (2007) The results of 163 Achilles tendon ruptures treated by a minimally invasive surgical technique and functional after treatment. *Injury* 38:839–844
40. Sheth U, Wasserstein D, Jenkinson R, Moineddin R, Kreder H, Jaglal SB (2017) The epidemiology and trends in management of acute Achilles tendon ruptures in Ontario, Canada. *Bone Joint J* 99–B:78–86
41. McCormack R, Bovard J, McCormack RG (2015) Early functional rehabilitation or cast immobilisation for the postoperative management of acute Achilles tendon rupture? A systematic review and meta-analysis of randomised controlled trials. *Br J Sport Med* 49:1329–1335
42. Huang J, Wang C, Ma X, Wang X, Zhang C, Chen L (2015) Rehabilitation regimen after surgical treatment of acute Achilles tendon ruptures: a systematic review with meta-analysis. *Am J Sport Med* 43:1008–1016
43. Mark-Christensen T, Troelsen A, Kallemose T, Barfod KW (2016) Functional rehabilitation of patients with acute Achilles tendon rupture: a meta-analysis of current evidence. *Knee Surg Sports Traumatol Arthrosc* 24:1852–1859
44. van Dijk PAD, Lubberts B, Verheul C, DiGiovanni CW, Kerkhoffs GMMJ (2016) Rehabilitation after surgical treatment of peroneal tendon tears and ruptures. *Knee Surg Sports Traumatol Arthrosc* 24:1165–1174
45. Kerkhoffs GMMJ, Struijs PAA, Raaymakers ELFB, Marti RK (2002) Functional treatment after surgical repair of acute Achilles tendon rupture: wrap vs walking cast. *Arch Orthop Trauma Surg* 122:102–105
46. Kearney RS, Parsons N, Underwood M, Costa ML (2015) Achilles tendon rupture rehabilitation: a mixed methods investigation of current practice among orthopaedic surgeons in the United Kingdom. *Bone Joint Res* 4:65–69
47. Braunstein M, Baumbach SF, Boecker W, Carmont MR, Polzer H (2015) Development of an accelerated functional rehabilitation protocol following minimal invasive Achilles tendon repair. *Knee Surg Sports Traumatol Arthrosc* 26:846–853
48. Brumann M, Baumbach SF, Mutschler W, Polzer H (2014) Accelerated rehabilitation following Achilles tendon repair after acute rupture—development of an evidence-based treatment protocol. *Injury* 45:1782–1790
49. Valkering KP, Aufwerber S, Ranuccio F, Lunini E, Edman G, Ackermann PW (2017) Functional weight-bearing mobilization after Achilles tendon rupture enhances early healing response: a single-blinded randomized controlled trial. *Knee Surg Sports Traumatol Arthrosc* 25:1807–1816
50. Sleney J, Christie N, Earthy S, Lyons RA, Kendrick D, Towner E (2014) Improving recovery—Learning from patients’ experiences after injury: a qualitative study. *Injury* 45:312–319
51. Zellers JA, Carmont MR, Grävare Silbernagel K (2016) Return to play post-Achilles tendon rupture: a systematic review and meta-analysis of rate and measures of return to play. *Br J Sports Med* 50:1325–1332
52. Ecker TM, Bremer AK, Krause FG, Müller T, Weber M (2016) Prospective use of a standardized nonoperative early weightbearing protocol for Achilles tendon rupture: 17 years of experience. *Am J Sports Med* 44:1004–1010
53. Schepers T, Van Zuuren WJ, Van Den Bekerom MPJ, Vogels LMM, Van Lieshout EMM (2012) The management of acute distal tibio-fibular syndesmotic injuries: results of a nationwide survey. *Injury* 43:1718–1723

Affiliations

Olivier C. Dams¹ · Inge van den Akker-Scheek^{1,2} · Ron L. Diercks² · Klaus W. Wendt³ · Johannes Zwerver¹ · Inge H. F. Reininga³

¹ Department of Sport and Exercise Medicine, University of Groningen, University Medical Center Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands

² Department of Orthopaedics, University of Groningen, University Medical Center Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands

³ Department of Trauma Surgery, University of Groningen, University Medical Center Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands