

University of Groningen

**Presence of retinopathy and incident kidney and cardiovascular events in type 2 diabetes with normoalbuminuria – A post-hoc analysis of the PRIORITY randomized clinical trial**

on the behalf of the PRIORITY Study Group; Rotbain Curovic, Viktor; Tofte, Nete; Lindhardt, Morten; Adamova, Katarina; Bakker, Stephan J.L.; Beige, Joachim; Beulens, Joline W.J.; Birkenfeld, Andreas L.; Currie, Gemma

*Published in:*  
 JOURNAL OF DIABETES AND ITS COMPLICATIONS

*DOI:*  
[10.1016/j.jdiacomp.2023.108433](https://doi.org/10.1016/j.jdiacomp.2023.108433)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
 Publisher's PDF, also known as Version of record

*Publication date:*  
 2023

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

on the behalf of the PRIORITY Study Group, Rotbain Curovic, V., Tofte, N., Lindhardt, M., Adamova, K., Bakker, S. J. L., Beige, J., Beulens, J. W. J., Birkenfeld, A. L., Currie, G., Delles, C., Dimos, I., Francová, L., Frimodt-Møller, M., Girman, P., Göke, R., Hansen, T. W., Havrdova, T., Kooy, A., ... Rossing, P. (2023). Presence of retinopathy and incident kidney and cardiovascular events in type 2 diabetes with normoalbuminuria – A post-hoc analysis of the PRIORITY randomized clinical trial. *JOURNAL OF DIABETES AND ITS COMPLICATIONS*, 37(4), Article 108433.  
<https://doi.org/10.1016/j.jdiacomp.2023.108433>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Presence of retinopathy and incident kidney and cardiovascular events in type 2 diabetes with normoalbuminuria – a post-hoc analysis of the PRIORITY randomized clinical trial

**Viktor Curovic** (✉ [viktor.rotbain.curovic@regionh.dk](mailto:viktor.rotbain.curovic@regionh.dk))

Steno Diabetes Center Copenhagen

**Nete Tofte**

Steno Diabetes Center Copenhagen

**Morten Lindhardt**

Steno Diabetes Center Copenhagen

**Katarina Adamova**

University Clinic of Endocrinology

**Stephan J.L. Bakker**

University of Groningen

**Joachim Beige**

Hospital St Georg

**Joline W.J. Beulens**

Amsterdam UMC, location Vrije Universiteit Amsterdam

**Andreas L. Birkenfeld**

University Hospital Tübingen

**Gemma Currie**

University of Glasgow

**Christian Delles**

University of Glasgow

**Ingo Dimos**

Diabetespraxis

**Lidmila Francová**

Charles University

**Marie Frimodt-Møller**

Steno Diabetes Center Copenhagen

**Peter Girman**

Diabetes Center Institute for Clinical and Experimental Medicine

**Rüdiger Göke**

Diabetologen Hessen

**Tine W. Hansen**

Steno Diabetes Center Copenhagen

**Tereza Havrdova**

Diabetes Center Institute for Clinical and Experimental Medicine

**Adriaan Kooy**

Bethesda Diabetes Research Center

**Gozewijn D. Laverman**

Ziekenhuisgroep Twente Hospital

**Harald Mischak**

Mosaiques Diagnostics GmbH

**Gerjan Navis**

University of Groningen

**Giel Nijpels**

Department of General Practice and Elderly Care

**Marina Noutsou**

National and Kapodistrian University of Athens, Hippokratio General Hospital

**Alberto Ortiz**

Instituto de Investigacion Sanitaria de la Fundacion Jiménez Díaz UAM

**Aneliya Parvanova**

Istituto di Ricerche Farmacologiche Mario Negri IRCCS

**Frederik Persson**

Steno Diabetes Center Copenhagen

**John R. Petrie**

University of Glasgow

**Piero L. Ruggenenti**

Istituto di Ricerche Farmacologiche Mario Negri IRCCS

**Femke Rutters**

Amsterdam UMC, location Vrije Universiteit Amsterdam

**Ivan Rychlík**

Charles University

**Justyna Siwy**

Mosaiques Diagnostics GmbH

**Goce Spasovski**

Cyril and Methodius University in Skopje

**Marijn Speeckaert**

Ghent University Hospital

**Matias Trillini**

Istituto di Ricerche Farmacologiche Mario Negri IRCCS

**Petra Züribig**

Mosaiques Diagnostics GmbH

**Heiko von der Leyen**

Orgenesis Germany GmbH

**Peter Rossing**

Steno Diabetes Center Copenhagen

---

## Research Article

**Keywords:** Type 2 diabetes, macrovascular disease, cardiovascular disease, microvascular disease, diabetic retinopathy, chronic kidney disease, albuminuria, diabetic kidney disease, risk stratification, clinical medicine

**Posted Date:** November 14th, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-2255769/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

## Background

Diabetic retinopathy (DR) is a microvascular complication of diabetes highly associated to cardiovascular disease and diabetic kidney disease. However, these associations are not thoroughly investigated at an early type 2 diabetes disease stage. This study therefore evaluated the association between baseline DR status and development of cardiovascular events (CVEs), microalbuminuria, and kidney function decline and in persons with type 2 diabetes and normal urinary albumin excretion.

## Methods

Post-hoc analysis of the PRIORITY study including 1758 persons with type 2 diabetes and normoalbuminuria followed for a median of 2.5 (IQR: 2.0–3.0) years. The study was originally designed to investigate a urinary proteomic risk classifier predictor of microalbuminuria development. DR at baseline was defined as non-proliferative and proliferative abnormalities, macular oedema, or history of laser treatment. Cox models were fitted to investigate the association of DR status with development of 1) a CVE composite defined as non-fatal myocardial infarction, stroke, coronary artery bypass graft, percutaneous coronary intervention, hospitalization for heart failure, or all-cause mortality; 2) persistent microalbuminuria (urinary albumin-creatinine ratio > 30mg/g); and 3) chronic kidney disease (CKD) G3 (eGFR < 60 mL/min/1.73m<sup>2</sup>). Models were adjusted for relevant risk factors.

## Results

At baseline, 304 (17.3%) had DR. Compared to persons without DR, they were older (mean ± SD: 62.7 ± 7.7 vs 61.4 ± 8.3 years,  $p = 0.019$ ), had longer diabetes duration (17.9 ± 8.4 vs. 10.6 ± 7.0 years,  $p < 0.001$ ), and higher HbA<sub>1c</sub> (62 ± 13 vs. 56 ± 12 mmol/mol,  $p < 0.001$ ). The adjusted hazard ratios of DR at baseline for development of CVE ( $n = 64$ ), microalbuminuria ( $n = 197$ ), and CKD ( $n = 166$ ) were: 2.61 (95%CI: 1.44, 4.72), 1.50 (95%CI: 1.07, 2.11), and 0.87 (95%CI: 0.56, 1.34), and, compared to without DR. Baseline levels of the urinary proteomics classifier did not influence the results.

## Conclusions

Presence of DR in normoalbuminuric type 2 diabetes was associated with an increased risk of developing CVE and microalbuminuria, but not with kidney function decline.

## Background

One of the main concerns for a person diagnosed with type 2 diabetes is the considerable risk of developing debilitating and potentially fatal diabetic complications over time (1). Improvements in pharmacological agents, multifactorial treatment, and diabetes care have radically progressed, reducing the incidence of diabetes-related complications (2). However, despite this success a residual risk remains. While increased blood glucose is the main driving factor for the development of diabetes complications in type 1 and type 2 diabetes alike, the interplay between the various complications is still not fully understood. Diabetic retinopathy (DR), a microvascular complication leading to proliferation in retinal vessels, leaky vessels with oedema formation, and potentially blindness if left untreated, is one of the most feared complications for an individual with diabetes and a leading cause of blindness in adults in developed countries (3). Another microvascular diabetes complication, chronic kidney disease (CKD), is the single largest cause of kidney failure and dialysis in developed countries (4). Furthermore, cardiovascular events (CVE) are frequent macrovascular complications and the leading cause of death in type 2 diabetes (5).

DR and CKD in diabetes specifically affect microvascular tissue and are commonly present in long-term diabetes, to the extent that they have been considered as different manifestations of the same vascular complication (6, 7). In type 1 diabetes, DR almost invariably precedes CKD, a pattern that can also be found for type 2 diabetes, albeit to a lower degree (8). While an association between DR and CKD has been shown in prospective studies, only a few of them have investigated the association between DR and the development of CKD in otherwise uncomplicated diabetes, none in normoalbuminuric type 2 diabetes, and none simultaneously investigating the association between DR and CVE (9).

Recent studies have suggested that DR and CKD in diabetes might lack a common molecular interplay and physiology (10, 11). Therefore, we have investigated their relationship, utilizing data from the PRIORITY study; a prospective multi-center observational study with an embedded randomized clinical trial, including 1775 individuals with type 2 diabetes and normoalbuminuria. The study demonstrated that the urinary proteomic classifier – CKD273 – predicts onset of microalbuminuria and CKD in type 2 diabetes (12). In the present study we aimed to evaluate whether, and to which extent, the presence of DR at baseline is associated with the onset of microalbuminuria, kidney function decline, and development of CVE in individuals with type 2 diabetes and normoalbuminuria.

## Methods

### Study design and participants

The details of the PRIORITY study design and population have been previously described (12, 13). In short, individuals aged 18–75 with type 2 diabetes, preserved kidney function, and normoalbuminuria were recruited from 15 highly specialized diabetes medical centers in 10 European countries. The clinical study protocol and informed consent documents were reviewed and approved by the respective local independent ethics committees and competent authorities, respectively. Every patient gave written informed consent prior to the conduct of any study-related procedures. This study was registered

(EudraCT 20120-004523-4; ClinicalTrials.gov NCT02040441) and is completed. The main inclusion criteria were urinary albumin-creatinine ratio (UACR) < 30 mg/g of at least two out of three consecutive first-morning void urine samples and an estimated glomerular filtration rate (eGFR) > 45 mL/min/1.73m<sup>2</sup>. Historical UACR status was not referred. The main exclusion criteria were treatment with dual renin-angiotensin-aldosterone system blockade or mineralocorticoid receptor antagonist, or heart failure requiring treatment with a mineralocorticoid receptor antagonist. The complete list of inclusion and exclusion criteria can be found in Supplementary Table 1. In the original study, all participants were stratified by low- ( $\leq 0.154$ ) or high-risk ( $> 0.154$ ) urinary proteomic classifier (CKD273) status. The classifier is based on 273 peptides differentially present in urine in people with vs without CKD (14). High-risk participants were included in a nested randomized clinical trial and subsequently randomized to receive daily spironolactone 25 mg or placebo. Low-risk participants were seen once yearly, and high-risk participants once quarterly for the study duration of median 2.5 (IQR: 2.0–3.0) years. At all visits, local blood and urine sampling was performed and vital parameters were measured, and adverse events were recorded by the interviewing investigator using subject-reported information and electronic medical records. The primary outcome was the development of persistent microalbuminuria (moderately increased albuminuria: UACR > 30 mg/g in two out of three consecutive measurements).

The present study aimed to assess baseline DR status with: 1) the primary outcome – development of confirmed microalbuminuria (UACR > 30 mg/g) in at least two of three first morning voids with a 30% increase (geometric mean) in UACR from ‘run-in-phase’ samples, or > 40 mg/g (geometric mean), 2) development of CKD grade 3 (G3), defined as eGFR < 60 mL/min/1.73m<sup>2</sup> based on two consecutive serum creatinine measurements; 3) development of a composite endpoint of CVE defined as non-fatal myocardial infarction, stroke, coronary artery bypass graft, percutaneous coronary intervention, hospitalization for heart failure, or all-cause mortality; and 4) yearly eGFR decline. The CKD G3 endpoint was only applicable for participants with a baseline eGFR > 60 mL/min/1.73m<sup>2</sup>; Therefore, a total of 111 participants were excluded from this endpoint calculation due to baseline eGFR < 60 mL/min/1.73m<sup>2</sup>.

## Procedures

Demographics including age, sex, diabetes duration, smoking, medical history, and concomitant medical treatment were collected at the baseline visit by the interviewing investigator. Likewise, the interviewing investigator evaluated DR status at baseline based on participant information and the participant’s electronic medical record. Participants were included from highly specialized diabetes medical centers in which participants were subjected to regular retinal photo assessments graded by trained specialized staff according to local protocols. UACR was measured centrally (at Steno Diabetes Center Copenhagen, Gentofte, Denmark) at all visits using a Vitros 5600 MicroSlide (Orto Clinical Diagnostics, Raritan, NJ, USA). eGFR was determined using serum creatinine values measured centrally using the 2009 Chronic Kidney Disease Epidemiology Collaboration Eq. (15). HbA1c and potassium were analyzed at the participating center’s local routine laboratory using standardized methods. Urinary proteomics were performed by capillary electrophoresis mass spectrometry by Mosaiques Diagnostics (Hannover,

Germany) using previously described methods (12), assessing a predefined renal risk profile based on 273 peptide fragments (CKD273).

## Statistical analysis

Baseline values are presented as mean  $\pm$  standard deviation (SD), median (inter-quartile range) if non-normally distributed, and n (%) if categorical. Comparisons between groups were performed using one-way analysis of variance, Kruskal-Wallis rank sum test, and  $\chi^2$ -test, respectively. Cox proportional hazard models were used to investigate associations between baseline presence of DR, alone or stratified by CKD273 risk profile, and time to development of persistent microalbuminuria, CKD G3, or the composite cardiovascular endpoint. Results are presented as hazard ratios (HR) with 95% confidence intervals (95% CI) with participants without DR at baseline being the reference. Decline in eGFR was derived using all available eGFR measurements for all participants (yearly measurements for low-risk participants and quarterly for high-risk) and calculating an estimated yearly decline using individual linear regression models. To calculate the estimated yearly decline, participants were required to have a minimum of three eGFR measurements across a minimum of 6 months to acquire an acceptable estimate. All adjusted models included the following covariates: sex, baseline age, diabetes duration, HbA<sub>1c</sub>, systolic blood pressure, eGFR, UACR, and CKD273 risk status. Adjustments in the analysis of the CVE endpoint also included baseline LDL cholesterol, body mass index, and history of ischemic heart disease and stroke. Baseline DR, alone or stratified by CKD273 risk profile was investigated categorically using linear regression models, in association with the calculated yearly eGFR decline. Models stratified by DR and CKD273 risk profiles were subjected to a limited adjustment including only HbA<sub>1c</sub> and diabetes duration due to low event rates and significant risk of over-adjustment. Cumulative incidence plots were drawn, and differences between groups are in these figures calculated using log-rank test. Sensitivity analyses were performed, first by further adjustment of all models with renin-angiotensin-aldosterone system inhibitor, statin, aspirin, glucagon like peptide 1 receptor agonist, and sodium-glucose co-transporter 2 inhibitor treatment; and second by substratification of baseline DR into non-proliferative and proliferative DR. Furthermore, a sensitivity analysis was performed investigating DR status at baseline in association with difference in eGFR slope in which only values from month 3 (first visit after randomization) and forward were included, only in the high-risk group. This to address the possible acute influence spironolactone administration might have had on eGFR. All statistical analyses and data visualizations were performed using R v. 4.1.0 (R Core Team, 2021) and RStudio v. 1.4.1 (RStudio Team, 2021).

## Results

Original enrollment of the PRIORITY study occurred between March 25, 2014, and Sept 30, 2018. Of the 1775 participants included in the PRIORITY study, 1758 had baseline information regarding DR status while the status was unknown for 17 (Supplementary Fig. 1). Of the 1758 participants, 304 (17.3%) had DR at baseline, with 231 (76.0%) classified as non-proliferative DR (NPDR), 56 (18.4%), as proliferative DR (PDR), and 17 (5.6%) as background or non-classified DR (only maculopathy or history of laser treatment). Furthermore, 79 (26.8%) participants had maculopathy at baseline, and 76 (25.0%) had a



history of laser therapy. 1454 participants did not have DR at baseline. Participants with DR at baseline had, on average, longer diabetes duration ( $17.7 \pm 8.4$  vs  $10.5 \pm 7.0$  years), higher HbA1c ( $62 \pm 13$  vs.  $56 \pm 12$  mmol/mol ( $7.8 \pm 1.2$  vs.  $7.3 \pm 1.1\%$ )), lower eGFR ( $85 \pm 16$  vs.  $87 \pm 15$  ml/min/1.73m<sup>2</sup>), lower diastolic blood pressure ( $76 \pm 10$  vs.  $79 \pm 8$  mmHg), and lower LDL cholesterol ( $2.22 \pm 0.83$  vs.  $2.45 \pm 0.94$  mmol/l) than participants without DR. In addition, history of ischemic heart disease was higher in participants with DR than in those without DR (18% vs. 11%,  $p < 0.001$ ). However, the proportion with a history of stroke was identical (4% vs. 4%,  $p = 1.000$ ). Participants with DR at baseline were also more likely to be treated with insulin, glucagon like peptide 1 receptor agonists, or sodium-glucose co-transporter 2 inhibitors compared to those without DR (Table 1). These overall associations were intact when stratifying baseline characteristics by both presence of DR and urinary proteomic risk classifier status (Supplementary Table 1). Baseline characteristics were also largely identical for the subset of participants with eGFR  $> 60$  mL/min/1.73m<sup>2</sup> at baseline, in which development of CKD G3 was assessed (Supplementary Table 2).

Table 1

Baseline characteristics stratified by presence of diabetic retinopathy (DR) at baseline. Data are presented as mean (standard deviation), median [interquartile range], or n (%). P-values were calculated using Student's t-test, Kruskal-Wallis rank sum test, and the  $\chi^2$ -test for continuous, non-normal continuous, and categorical variables, respectively. HbA1c: glycated hemoglobin; eGFR: estimated glomerular filtration rate; UACR: urinary albumin-creatinine rate; CKD273: urinary proteomics classifier; BP: blood pressure, BMI: body mass index, RAAS: renin-angiotensin-aldosterone system; GLP1-RA: glucagon like peptide 1 receptor agonist; SGLT2: sodium-glucose co-transporter 2; IHD: ischemic heart disease.

Variable	No DR	DR	p
N	1454	304	
Age, years	61.4 (8.4)	62.7 (7.7)	0.010
Male, n (%)	899 (61.8)	198 (65.1)	0.310
Non-White	46 (3.2)	14 (4.6)	0.278
Diabetes duration, years	10.5 (7.0)	17.7 (8.4)	< 0.001
Low-risk CKD273 pattern, n (%)	1279 (88.0)	263 (86.5)	0.545
Retinopathy grade, n (%)			
Non-proliferative	-	231 (76.5)	
Proliferative	-	56 (18.5)	
Maculopathy, n (%)	-	79 (26.8)	
History of laser therapy, n (%)	-	76 (25.8)	
HbA1c, mmol/mol	56 (12)	62 (13)	< 0.001
HbA1c, %	7.3 (1.1)	7.8 (1.2)	
eGFR, mL/min/1.73m <sup>2</sup>	87 (15)	85 (16)	0.010
UACR, mg/g	5 [3, 8]	6 [4, 10]	< 0.001
CKD273, arbitrary unit	-0.34 (0.42)	-0.26 (0.38)	0.002
Systolic BP, mmHg	133 (12)	134 (12)	0.250
Diastolic BP, mmHg	79 (8)	76 (10)	< 0.001
BMI, kg/m <sup>2</sup>	30.3 (5.0)	30.6 (5.2)	0.487
LDL, mmol/l	2.45 (0.94)	2.22 (0.83)	< 0.001
Smoker, n (%)	656 (45.2)	126 (41.7)	0.291
RAAS-inhibitor treatment, n (%)	909 (62.5)	232 (76.3)	< 0.001

Variable	No DR	DR	p
Metformin treatment, n (%)	1092 (78.6)	240 (79.7)	0.709
Insulin treatment, n (%)	365 (26.3)	167 (55.5)	< 0.001
GLP1-RA treatment, n (%)	200 (14.4)	71 (23.6)	< 0.001
SGLT2-inhibitor treatment, n (%)	91 (6.5)	30 (10.0)	0.050
History of IHD, n (%)	157 (11)	58 (18)	< 0.001
History of stroke, n (%)	62 (4)	13 (4)	1.000

Median follow-up was 2.5 (IQR: 2.0–3.0) years for all outcomes. During this time, 198 (11.3%) participants progressed to microalbuminuria, 166 (9.4% of the 1675 participants with eGFR > 60 mL/min/1.73m<sup>2</sup> at baseline) to CKD G3, and 64 (3.6%) had a CVE. Results from the primary Cox proportional hazards models can be seen in Table 2 and Fig. 1. Presence of DR at baseline was significantly associated with progression to persistent microalbuminuria in both unadjusted (HR: 1.81 (95% CI: 1.32, 2.48), p < 0.001) and adjusted (1.50 (1.07, 2.11), p = 0.018) models compared to participants without DR. No association was found with progression to CKD G3, neither in unadjusted (0.95 (0.63, 1.42), p = 0.801) nor in adjusted (0.87 (0.56, 1.34), p = 0.55) models. The risk of developing CVE was more than 2.6-fold increased for participants with DR compared to those without, irrespective of adjustment (unadjusted: 2.74 (1.64, 4.57), p < 0.001; adjusted: 2.61 (1.44, 4.72), p = 0.002). The cumulative incidence for all dichotomous endpoints, stratified by the presence of DR at baseline, can be seen in Fig. 2, showing a higher event rate for microalbuminuria and CVE in participants with DR. When sub-stratifying the baseline DR, similar associations were found for NPDR, both before and after adjustment (adjusted HR (95% CI): microalbuminuria: 1.53 (1.06, 2.22), p = 0.024; CKD G3: 0.69 (0.41, 1.17), p = 0.173; and CVE: 2.49 (1.33, 4.67), p = 0.004) when compared to no DR. Although having similar HRs, PDR was not associated with endpoints when compared to no DR, in any of the adjusted models, presumably due to the low number of participants with PDR at baseline (n = 56) (Supplementary Table 3). Furthermore, we subdivided participants into strata by the combined presence of DR and CKD273 risk status at baseline. Results were largely confirmatory with the primary analyses; compared with no DR and low-risk status, no DR and high-risk status at baseline was associated with development of all endpoints but not with a steeper eGFR slope. Presence of DR and low-risk status at baseline were associated with increased risk of microalbuminuria and CVE, but not CKD G3 and eGFR slope. Finally, the presence of DR and high-risk status compared to no DR and low-risk status at baseline was associated with all outcomes with magnitude higher HRs for the development of microalbuminuria and CVE (Supplementary Table 4, Supplementary Fig. 2).

Table 2

Presence of diabetic retinopathy (DR) with the development of microalbuminuria, chronic kidney disease (CKD) G3, or cardiovascular events (CVE), and difference in yearly estimated glomerular filtration (eGFR) slope, compared to no diabetic retinopathy. Associations to dichotomous outcomes were estimated using Cox proportional hazards models and association to yearly eGFR slope with linear regression. Adjustments included sex, baseline age, diabetes duration, HbA1c, systolic blood pressure, eGFR, urinary albumin-creatinine rate, and urinary proteomic risk classifier status. Adjustments for the CVE endpoint also included baseline LDL cholesterol, body mass index, and ischemic heart disease and stroke history.

Outcome	Presence of DR	p	Presence of DR	p
	Unadjusted HR (95% CI)		Adjusted HR (95% CI)	
Microalbuminuria (n = 198)	1.81 (1.32, 2.48)	< 0.001	1.50 (1.07, 2.11)	0.018
CKD G3 (eGFR < 60 mL/min/1.73m <sup>2</sup> ) (n = 166)	0.95 (0.63, 1.42)	0.801	0.87 (0.56, 1.34)	0.553
CVE (n = 64)	2.74 (1.64, 4.57)	< 0.001	2.61 (1.44, 4.72)	0.002
	Presence of DR Unadjusted mL/min/1.73m <sup>2</sup> /year		Presence of DR Adjusted mL/min/1.73m <sup>2</sup> /year	
eGFR slope difference	-0.13 (-0.45, 0.70)	0.669	-0.01 (-0.61, 0.58)	0.968

Further investigation of the relationship between baseline DR status and yearly eGFR change (mean for entire population: -1.01 (-1.27, -0.76) mL/min/1.73 m<sup>2</sup>/year) showed no difference in yearly eGFR slope of individuals with DR at baseline compared to those without (adjusted  $\beta$  estimate: -0.01 (-0.61, 0.58),  $p = 0.968$ ), Table 2). To account for a possible acute influence spironolactone administration might have had on eGFR, a sensitivity analysis was performed including only measurements from month 3 and forward, in the high-risk group. Results were unchanged compared to the main analysis. When sub-stratifying baseline DR, no differences in yearly eGFR slopes were observed neither in NPDR nor in PDR (Supplementary Table 3). Likewise, when the analysis considered the population stratified by both baseline presence of DR and CKD273 risk status, no significant associations were observed in any combination of baseline presence of DR or risk status compared to no DR and low-risk status (Supplementary Table 4).

Additional adjustment for renin-angiotensin-aldosterone system inhibitors, statins, aspirin treatment, sodium-glucose co-transporter 2 inhibitor treatment, glucagon like peptide 1 receptor agonist treatment, or

randomization group (for the nested trial) did not affect any of the above-described results significantly.

## **Sub-analysis combined microalbuminuria and CKD G3 development**

Of 198 (microalbuminuria) and 166 (CKD G3) events, 27 participants experienced both endpoints. The association between presence of DR at baseline and the development of the combined microalbuminuria and CKD G3 endpoint did not result in significant associations neither in crude nor in adjusted models. We also performed two further analyses, including only participants who had developed either CKD G3 or microalbuminuria. We assessed the relationship between baseline DR and the development of both microalbuminuria and CKD G3. In both subsets, no significant associations were found between DR and the development of the combined endpoint neither crude nor adjusted models (Supplementary Table 5).

## **Discussion**

We have demonstrated that the presence of DR in people with type 2 diabetes and normoalbuminuria was associated with increased risk of development of microalbuminuria and CVE across a median follow-up of 2.5 years. Interestingly, DR was not associated with an increased risk of developing impaired kidney function in this relatively short observation period. CKD in individuals with diabetes, and type 1 diabetes in particular, has long been considered closely associated with the presence of DR (6). This relationship is so deeply established, that clinical diabetic kidney disease can solely be diagnosed in persons with diabetes and macro- (severe) albuminuria, presence of DR, and absence of clinical or laboratory evidence of other kidney diseases. Previous studies, however, have primarily focused on macroalbuminuric CKD in diabetes, thereby highlighting a severe form of kidney impairment that was not included in this study and with which DR has been strongly associated. In addition, our results showed no association between the presence of DR and a steeper yearly eGFR slope, irrespective of the progression of the albuminuria grade during follow-up.

In our study, participants with DR at baseline had longer diabetes duration, but similar age and were more likely to have a history of ischemic heart disease. These factors did, however, not affect the risk effect size or significance, indicating certain robustness of the findings. We have previously demonstrated that participants identified at high risk based on the CKD273 classifier had a higher risk for development of microalbuminuria or CKD G3. In this study, we demonstrate that the combination of DR and a high risk CKD273 profile indicates an additive effect on the risk for development of all outcomes, compared to the presence of DR or a high-risk CKD273 profile alone. Given the much fewer individuals in each group, confidence intervals were large, and no adjustments were performed; thus, no firm conclusion should be drawn from these analyses. However, as we have previously shown, urinary proteomics, and CKD273 specifically, were not associated with baseline DR grade or progression in DR in both type 1 and type 2 diabetes (10). These results may reflect two pathogeneses leading to the development of microalbuminuria.

Our results beg the question of how specific microalbuminuria is to CKD in type 2 diabetes or if it instead is a phenotype of general vascular damage. This has been suggested before, as it has been shown that more than 30% of people with type 2 diabetes and  $eGFR < 60 \text{ mL/min/1.73m}^2$  had non-albuminuric CKD (16, 17). On a molecular level, recent studies have supported the distinction between DR and CKD (10, 11). A study by Hong et al., which investigated the relationship between baseline DR and development of kidney impairment or cardiovascular disease over 14 years, found a significantly higher risk of cardiovascular disease compared to CKD (18), suggesting that DR may be primarily associated with cardiovascular disease, rather than kidney disease. Even in type 1 diabetes, few metabolite similarities were discovered using exploratory omics-based methods in two studies that used the same cohort and metabolomics platform (19, 20). Ribitol and ribonic acid, both derivatives of ribose, which in turn is active in the pentose phosphate pathway and highly influenced by hyperglycemia (21), were positively correlated to presence of DR (19), a finding also identified in type 2 diabetes (22). Similarly, ribonic acid was significantly associated with future kidney function decline after adjustment for clinical covariates and multiple testing (20), but no metabolites independent of hyperglycemia influence were identified. Furthermore, another recent study investigating proteomic biomarkers for risk of kidney disease found very little association with prevalence, incidence, or progression of DR in a population of 958 individuals included in the Fremantle Diabetes Study (23).

While the current study does not provide a comprehensive answer to the relationship between DR and diabetic kidney disease and vice versa, it does support the idea that a distinction between these two microvascular complications may be indicated. Alternatively, albuminuric CKD and non-albuminuric CKD in type 2 diabetes may be separate diseases and should be targeted differently, especially at an early stage. The sooner this can be fully elucidated, and the respective pathophysiologies charted, the sooner more targeted solutions might be proposed and applied in the treatment and prevention of DR and kidney disease. It is also important to note that our findings do not invalidate previous research linking DR and CKD in diabetes, but rather that the complete picture may be more complex than previously assumed while emphasizing that our study is limited to normoalbuminuric type 2 diabetes

Our study also has its limitations. Mainly, no standardized retinal photography was performed at baseline to grade DR and thus the presence of DR at baseline was assessed by the interviewing physician at the participating center. This is somewhat ameliorated by the fact that all participating centers and investigators specialized in diabetes treatment and care, leading to a supposed higher validity of the DR diagnosis. Notwithstanding, a formal, homogenous, retinal examination across the entire cohort would be preferred. The study is a large European multi-center cohort of individuals with type 2 diabetes and normoalbuminuria which increases the generalizability of the results across a western population, albeit the homogenous racial composition of the participants as well as the strict inclusion and exclusion criteria inherently limits the applicability of the study in other populations.

## Conclusions

We have demonstrated that the presence of DR in a population of individuals with type 2 diabetes and normoalbuminuria is mainly a risk factor for the development of microalbuminuria and cardiovascular disease. At the same time, it is not associated with kidney function decline, implying that DR is an indicator of microvasculopathy in general and appears to be a prognostic factor. Our results may lead to a more precise risk assessment of individuals with type 2 diabetes. It furthers the notion that DR and impaired kidney function may not be markedly interlinked.

## List Of Abbreviations

DR

diabetic retinopathy

CKD

chronic kidney disease

CVE

cardiovascular events

UACR

urinary albumin-creatinine rate

eGFR

estimated glomerular filtration rate

CKD273

urinary proteomic renal risk classifier based on 273 peptide fragments

G3

grade 3

NPDR

non-proliferative diabetic retinopathy

PDR

proliferative diabetic retinopathy

## Declarations

### *Ethics approval and consent to participate*

The clinical study protocol and informed consent documents were reviewed and approved by the respective local independent ethics committees and competent authorities at each participating study site. Every patient gave written informed consent prior to the conduct of any study-related procedures. This study was registered (EudraCT 20120-004523-4; ClinicalTrials.gov NCT02040441) and is completed. The study complied Good Clinical Practice and with the Declaration of Helsinki.

### *Availability of data and materials*

The datasets generated and analyzed during the current study are not publicly available due to Danish GDPR legislation. Scrambled, anonymized datasets could be available from the corresponding author on reasonable request.

### *Competing interests*

NT is a full-time employee of Novo Nordisk A/S. TWH has equity in Novo Nordisk A/S. ML has received research support from Boehringer Ingelheim, Bayer and Merck Sharp & Dohme, and lecture fees from AstraZeneca, Bayer and Boehringer Ingelheim. All fees are given to Holbaek hospital. FP has served as a consultant, on advisory boards or as educator for AstraZeneca, Novo Nordisk, Boehringer Ingelheim, Sanofi, Mundipharma, MSD, Novartis, Amgen and has received research grants to institution from Novo Nordisk, Boehringer Ingelheim, Amgen and AstraZeneca.

AO has received grants from Sanofi and consultancy or speaker fees or travel support from Advicciene, Astellas, Astrazeneca, Amicus, Amgen, Fresenius Medical Care, GSK, Bayer, Sanofi-Genzyme, Menarini, Mundipharma, Kyowa Kirin, Alexion, Freeline, Idorsia, Chiesi, Otsuka, Novo-Nordisk, Sysmex and Vifor Fresenius Medical Care Renal Pharma and is Director of the Catedra Mundipharma-UAM of diabetic kidney disease and the Catedra Astrazeneca-UAM of chronic kidney disease and electrolytes. HM is the cofounder and co-owner of Mosaiques Diagnostics GmbH. JS and PZ are employees of Mosaiques Diagnostics GmbH. MF-M has received speaker fees or support for medical meetings and research grants from AstraZeneca and Novo Nordisk and speaking fees from Boehringer Ingelheim, Eli Lilly, Novo Nordisk, Novartis, Baxter, and Sanofi-Aventis. IR has received honoraria for lectures of AstraZenec, Bayer, Boehringer Ingelheim, and Mundipharma. PR has received research support and personal fees from AstraZeneca, Bayer and Novo Nordisk, and personal fees from Astellas Pharma Inc., Bayer, Boehringer Ingelheim, Eli Lilly and Company, Gilead, Merck, Merck Sharp & Dohme, Mundipharma, Sanofi, and Vifor Pharma. All fees are given to Steno Diabetes Center Copenhagen. All other authors declare no competing interests that could be perceived as affecting the impartiality of the reported results.

### *Funding*

The PRIORITY study was funded by the European Union Seventh Framework Programme (FP7/20072-013) under grant agreement number 279277.

### *Author contributions*

VRC, TWH, and PR designed the study. All authors contributed to data collection. VRC performed the statistical analysis and wrote the first draft of the manuscript. All authors contributed to the interpretation of the results. All authors contributed to critical revision and review of the manuscript. All authors approved the final version of the manuscript.

### *Acknowledgments*




We would like to thank the individuals who participated in the study for their time and commitment. Furthermore, we also thank the investigators, laboratory technicians, and study nurses for their invaluable efforts and contributions.

## References

1. Young-Hyman D, de Groot M, Hill-Briggs F, Gonzalez JS, Hood K, Peyrot M. Psychosocial Care for People With Diabetes: A Position Statement of the American Diabetes Association. *Diabetes Care*. 2016;39(12):2126–40.
2. Association AD. 11. Microvascular Complications and Foot Care: Standards of Medical Care in Diabetes—2021. *Diabetes Care*. 2020;44(Supplement\_1):151-S67.
3. Solomon SD, Chew E, Duh EJ, Sobrin L, Sun JK, VanderBeek BL, et al. Diabetic Retinopathy: A Position Statement by the American Diabetes Association. *Diabetes Care*. 2017;40(3):412–8.
4. Tuttle KR, Bakris GL, Bilous RW, Chiang JL, de Boer IH, Goldstein-Fuchs J, et al. Diabetic kidney disease: a report from an ADA Consensus Conference. *Diabetes care*. 2014;37(10):2864-83.
5. American Diabetes A 10. Cardiovascular Disease and Risk Management: Standards of Medical Care in Diabetes—2021. *Diabetes Care*. 2020;44(Supplement\_1):S125-S50.
6. Afkarian M, Zelnick LR, Hall YN, Heagerty PJ, Tuttle K, Weiss NS, et al. Clinical Manifestations of Kidney Disease Among US Adults With Diabetes, 1988–2014. *JAMA*. 2016;316(6):602–10.
7. Estacio RO, McFarling E, Biggerstaff S, Jeffers BW, Johnson D, Schrier RW. Overt albuminuria predicts diabetic retinopathy in Hispanics with NIDDM. *Am J Kidney Dis*. 1998;31(6):947–53.
8. He F, Xia X, Wu XF, Yu XQ, Huang FX. Diabetic retinopathy in predicting diabetic nephropathy in patients with type 2 diabetes and renal disease: a meta-analysis. *Diabetologia*. 2013;56(3):457–66.
9. Pearce I, Simó R, Lövestam-Adrian M, Wong DT, Evans M. Association between diabetic eye disease and other complications of diabetes: Implications for care. A systematic review. *Diabetes Obes Metab*. 2019;21(3):467–78.
10. Rotbain Curovic V, Magalhães P, He T, Hansen TW, Mischak H, Rossing P. Urinary peptidome and diabetic retinopathy in the DIRECT-Protect 1 and 2 trials. *Diabet Med*. 2021;38(9):e14634.
11. Lye WK, Paterson E, Patterson CC, Maxwell AP, Binte Mohammed Abdul RB, Tai ES, et al. A systematic review and participant-level meta-analysis found little association of retinal microvascular caliber with reduced kidney function. *Kidney Int*. 2021;99(3):696–706.
12. Toft N, Lindhardt M, Adamova K, Bakker SJL, Beige J, Beulens JWJ, et al. Early detection of diabetic kidney disease by urinary proteomics and subsequent intervention with spironolactone to delay progression (PRIORITY): a prospective observational study and embedded randomised placebo-controlled trial. *Lancet Diabetes Endocrinol*. 2020;8(4):301–12.
13. Lindhardt M, Persson F, Currie G, Pontillo C, Beige J, Delles C, et al. Proteomic prediction and Renin angiotensin aldosterone system Inhibition prevention Of early diabetic nephropathy in Type 2


- diabetic patients with normoalbuminuria (PRIORITY): essential study design and rationale of a randomised clinical multicentre trial. *BMJ Open*. 2016;6(3):e010310.
14. Good DM, Zürbig P, Argilés A, Bauer HW, Behrens G, Coon JJ, et al. Naturally occurring human urinary peptides for use in diagnosis of chronic kidney disease. *Mol Cell Proteomics*. 2010;9(11):2424–37.
  15. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF 3rd, Feldman HI, et al. A new equation to estimate glomerular filtration rate. *Ann Intern Med*. 2009;150(9):604–12.
  16. Kramer HJ, Nguyen QD, Curhan G, Hsu C-y. Renal Insufficiency in the Absence of Albuminuria and Retinopathy Among Adults With Type 2 Diabetes Mellitus. *JAMA*. 2003;289(24):3273–7.
  17. Penno G, Solini A, Orsi E, Bonora E, Fondelli C, Trevisan R, et al. Non-albuminuric renal impairment is a strong predictor of mortality in individuals with type 2 diabetes: the Renal Insufficiency And Cardiovascular Events (RIACE) Italian multicentre study. *Diabetologia*. 2018;61(11):2277–89.
  18. Hong J, Surapaneni A, Daya N, Selvin E, Coresh J, Grams ME, et al. Retinopathy and Risk of Kidney Disease in Persons With Diabetes. *Kidney Med*. 2021;3(5):808 – 15.e1.
  19. Curovic VR, Suvitaival T, Mattila I, Ahonen L, Trošt K, Theilade S, et al. Circulating Metabolites and Lipids Are Associated to Diabetic Retinopathy in Individuals With Type 1 Diabetes. *Diabetes*. 2020;69(10):2217–26.
  20. Tofte N, Suvitaival T, Trost K, Mattila IM, Theilade S, Winther SA, et al. Metabolomic Assessment Reveals Alteration in Polyols and Branched Chain Amino Acids Associated With Present and Future Renal Impairment in a Discovery Cohort of 637 Persons With Type 1 Diabetes. *Frontiers in Endocrinology*. 2019;10.
  21. Kinoshita JH. A thirty year journey in the polyol pathway. *Exp Eye Res*. 1990;50(6):567–73.
  22. Chen L, Cheng C-Y, Choi H, Ikram MK, Sabanayagam C, Tan GSW, et al. Plasma Metabonomic Profiling of Diabetic Retinopathy *Diabetes*. 2016;65(4):1099–108.
  23. Drinkwater JJ, Peters K, Davis WA, Turner AW, Bringans SD, Lipscombe RJ, et al. Assessment of biomarkers associated with rapid renal decline in the detection of retinopathy and its progression in type 2 diabetes: The Fremantle Diabetes Study Phase II. *J Diabetes Complications*. 2021;35(4):107853.

## Figures



**Figure 1**

Forest plot illustrating the unadjusted and adjusted hazard ratios of baseline presence of diabetic retinopathy with the development of microalbuminuria, chronic kidney disease (CKD) G3, and cardiovascular events (CVE) compared to no diabetic retinopathy. Adjustments included sex, baseline age, diabetes duration, HbA1c, systolic blood pressure, eGFR, urinary albumin-creatinine rate, and urinary proteomic risk classifier status. Adjustments for the CVE endpoint also included baseline LDL cholesterol, body mass index, an ischemic heart disease and stroke history.



**Figure 2**

Cumulative event plot illustrating the event rate of microalbuminuria, chronic kidney disease G3, and cardiovascular events stratified by presence of diabetic retinopathy (DR) at baseline. Each plot is presented with a complete y-axis and a minor embedded plot with a broken y-axis.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementalinformation091122.docx](#)