

## University of Groningen

### AdS-Carroll branes

Clark, T. E.; ter Veldhuis, T.

*Published in:*  
Journal of Mathematical Physics

*DOI:*  
[10.1063/1.4967969](https://doi.org/10.1063/1.4967969)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2016

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*  
Clark, T. E., & ter Veldhuis, T. (2016). AdS-Carroll branes. *Journal of Mathematical Physics*, 57(11), Article 112303. <https://doi.org/10.1063/1.4967969>

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## AdS-Carroll branes

T. E. Clark, and T. ter Veldhuis

Citation: [Journal of Mathematical Physics](#) **57**, 112303 (2016); doi: 10.1063/1.4967969

View online: <https://doi.org/10.1063/1.4967969>

View Table of Contents: <http://aip.scitation.org/toc/jmp/57/11>

Published by the [American Institute of Physics](#)

---

### Articles you may be interested in

[Oscillator versus prefundamental representations](#)

[Journal of Mathematical Physics](#) **57**, 111702 (2016); 10.1063/1.4966925

[Inverse eigenvalue problems](#)

[Journal of Mathematical Physics](#) **57**, 112102 (2016); 10.1063/1.4964390

[All-order bounds for correlation functions of gauge-invariant operators in Yang-Mills theory](#)

[Journal of Mathematical Physics](#) **57**, 122301 (2016); 10.1063/1.4967747

[Exactly solvable Hermite, Laguerre, and Jacobi type quantum parametric oscillators](#)

[Journal of Mathematical Physics](#) **57**, 122107 (2016); 10.1063/1.4972293

[Bender-Wu singularities](#)

[Journal of Mathematical Physics](#) **57**, 122109 (2016); 10.1063/1.4972290

[Cosmic anisotropic doomsday in Bianchi type I universes](#)

[Journal of Mathematical Physics](#) **57**, 112502 (2016); 10.1063/1.4967954

---

PHYSICS TODAY

WHITEPAPERS

### MANAGER'S GUIDE

Accelerate R&D with  
Multiphysics Simulation

READ NOW

PRESENTED BY  
 COMSOL

## AdS-Carroll branes

T. E. Clark<sup>1,a)</sup> and T. ter Veldhuis<sup>2,b)</sup>

<sup>1</sup>*Department of Physics and Astronomy, Purdue University, West Lafayette, Indiana 47907-2036, USA*

<sup>2</sup>*Van Swinderen Institute for Particle Physics and Gravity, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands and Department of Physics and Astronomy, Macalester College, Saint Paul, Minnesota 55105-1899, USA*

(Received 18 May 2016; accepted 6 November 2016; published online 30 November 2016)

Coset methods are used to determine the action of a co-dimension one brane (domain wall) embedded in  $(d + 1)$ -dimensional AdS space in the Carroll limit in which the speed of light goes to zero. The action is invariant under the non-linearly realized symmetries of the AdS-Carroll spacetime. The Nambu-Goldstone field exhibits a static spatial distribution for the brane with a time varying momentum density related to the brane's spatial shape as well as the AdS-C geometry. The AdS-C vector field dual theory is obtained. *Published by AIP Publishing.* [<http://dx.doi.org/10.1063/1.4967969>]

### I. INTRODUCTION

The symmetries of spacetime delimit the form of the action for fields on it. The familiar case of Poincaré symmetric spacetime results in particle motion being restricted to the local forward lightcone. This lightcone opens up to be the forward time half space in the Galilean limit in which the speed of light  $c \rightarrow \infty$  and instantaneous interaction is possible. On the other hand, as the speed of light vanishes,  $c \rightarrow 0$ , the causal lightcone closes to be just the forward time half-line. Such a contraction of spacetime is known as Carroll spacetime with symmetries generated by the Wigner-Inönü contracted Poincaré algebra,  $c \rightarrow 0$ , to the Carroll algebra.<sup>1,2</sup> A particle in such a spacetime must remain stationary as the time axis is the lightcone. This lack of motion can be found by considering the  $c \rightarrow 0$  limit of its Poincaré geodesic action. For a free particle moving in  $1 + 1$  dimensional Minkowski spacetime its action is given by

$$\begin{aligned} \Gamma &= -mc^2 \int d\tau = -mc^2 \int \sqrt{dt^2 - dx^2/c^2} \\ &= -mc^2 \int dt \sqrt{1 - \dot{x}(t)^2/c^2}. \end{aligned} \quad (1.1)$$

Introducing a Lagrange multiplier auxiliary velocity  $v(t)$

$$\dot{x}(t)/c = \tanh v(t), \quad (1.2)$$

the action becomes

$$\Gamma = -mc^2 \int dt \cosh v(t) \left[ 1 - \frac{1}{c} \dot{x}(t) \tanh v(t) \right]. \quad (1.3)$$

In order to take the Carroll limit  $c \rightarrow 0$ , the velocity is scaled by the speed of light  $v(t) = 2cw(t)$  yielding the action

$$\Gamma = -mc^2 \int dt \cosh 2cw(t) \left[ 1 - \frac{1}{c} \dot{x}(t) \tanh 2cw(t) \right]. \quad (1.4)$$

<sup>a)</sup>e-mail address: [clarkt@purdue.edu](mailto:clarkt@purdue.edu)

<sup>b)</sup>e-mail address: [terveldhuis@macalester.edu](mailto:terveldhuis@macalester.edu)

Letting  $c \rightarrow 0$  the Carroll limit for the action  $\Gamma_C = \Gamma/mc^2$  is obtained

$$\Gamma_C = - \int dt [1 - 2w(t)\dot{x}(t)]. \tag{1.5}$$

As expected there is no causal relation between different events along the particle’s trajectory and it remains stationary  $\dot{x}(t) = 0 = \dot{w}(t)$ .<sup>3,4</sup>

Extending the limiting procedure to membranes inserted into  $(d + 1)$ -dimensional Minkowski spacetime, the  $c \rightarrow 0$  contraction yields the Carrollian Nambu-Goto action for the brane. Such a limit occurs in the case of effective field theory of tachyon brane condensation in which the tachyon field rolls to the Carrollian limit.<sup>5</sup> The Carroll brane action can be obtained from the contraction of the Nambu-Goto action for a one-codimensional brane

$$\begin{aligned} \Gamma_{\text{NG}} &= -\sigma \int d^d x \sqrt{-(-1)^d \det g} = -\sigma \int d^d x \sqrt{-(-1)^d \det(\eta_{\mu\nu} - \partial_\mu \phi \partial_\nu \phi)} \\ &= -\sigma \int d^d x \sqrt{1 - \partial_\mu \phi \partial^\mu \phi}, \end{aligned} \tag{1.6}$$

where the  $(d + 1)$ -dimensional spacetime has been spontaneously broken to that of a  $d$ -dimensional world volume by the formation of a domain wall in the additional dimension. These Poincaré symmetries are compactly described by the invariant interval  $ds^2 = dx^\mu \eta_{\mu\nu} dx^\nu - dz^2$  with  $z$  denoting the one-dimensional covolume coordinate and  $x^\mu$  with  $\mu = 0, 1, \dots, p = (d - 1)$  denoting the  $d$ -dimensional world volume coordinates. Replacing  $z = \phi(x)$  it is obtained that  $ds^2 = dx^\mu g_{\mu\nu} dx^\nu = dx^\mu (\eta_{\mu\nu} - \partial_\mu \phi \partial_\nu \phi) dx^\nu$  resulting in the  $(d + 1)$  dimensional space-time invariance of the Nambu-Goto action (1.6) with brane tension  $\sigma$ .

In order to implement the Carrollian contraction, Lagrange multiplier auxiliary fields  $V_\mu(x)$  are introduced so that

$$\partial^\mu \phi = -\eta^{\mu\nu} V_\nu \frac{\tanh \sqrt{V^2}}{\sqrt{V^2}}, \tag{1.7}$$

with  $V^2 = V_\mu \eta^{\mu\nu} V_\nu$  and  $x_\mu = (ct, x_m)$  while  $\partial^\mu = \frac{\partial}{\partial x_\mu}$ . The action becomes

$$\Gamma_{\text{NG}} = -c\sigma \int dt d^p x \cosh \sqrt{V^2} \left[ 1 + \left( V_\mu \frac{\tanh \sqrt{V^2}}{\sqrt{V^2}} \right) \partial^\mu \phi \right]. \tag{1.8}$$

Making the speed of light explicit  $V_0 = 2cw$  and  $V_m = 2v_m$  for  $m = 1, 2, \dots, p = (d - 1)$ , the Carrollian limit  $c \rightarrow 0$  of the action  $\Gamma_C = -(1/c\sigma)\Gamma_{\text{NG}} = \int dt d^p x \mathcal{L}_C$  is obtained

$$\Gamma_C = \int dt d^p x \mathcal{L}_C = \int dt d^p x \cos \sqrt{4v^2} \left[ 1 + \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} (2w\dot{\phi} + 2v_m \partial^m \phi) \right], \tag{1.9}$$

where, after separation of space and time coordinates, the spatial metric is just  $\delta^{mn}$ . Only subscripts  $x_m, v_m$  and superscript derivatives  $\partial^m = \frac{\partial}{\partial x_m}$  will be used, with  $\dot{\phi} = \frac{\partial}{\partial t} \phi$ . In the case of a Goldstone field, as is  $\phi$ , the leading term in the derivative expansion of the action (thin wall limit) is uniquely fixed. The field equations are found to be

$$\begin{aligned}
0 &= \frac{\delta\Gamma_C}{\delta w(x,t)} = 2\dot{\phi} \left[ \frac{\sin\sqrt{4v^2}}{\sqrt{4v^2}} \right], \\
0 &= \frac{\delta\Gamma_C}{\delta v_m(x,t)} = 2\cos\sqrt{4v^2} \left( \left[ 2\delta^{nr}v_r \frac{\tan\sqrt{4v^2}}{\sqrt{4v^2}} - \partial^n\phi \right] \times \right. \\
&\quad \times \left. \left[ \frac{\tan\sqrt{4v^2}}{\sqrt{4v^2}} P_{Tns}(v) + P_{Lns}(v) \right] \delta^{sm} \right) \\
&\quad + 2\dot{\phi} \frac{wv_s\delta^{sm}}{v^2} \cos\sqrt{4v^2} \left( \frac{\tan\sqrt{4v^2}}{\sqrt{4v^2}} - 1 \right), \\
0 &= \frac{\delta\Gamma_C}{\delta\phi(x,t)} = -\frac{\partial}{\partial t} \left[ 2w \frac{\sin\sqrt{4v^2}}{\sqrt{4v^2}} \right] - \frac{\partial}{\partial x_m} \left[ 2v_m \frac{\sin\sqrt{4v^2}}{\sqrt{4v^2}} \right], \tag{1.10}
\end{aligned}$$

where the transverse  $P_{Tmn}(v)$  and longitudinal  $P_{Lmn}(v)$  projection matrices are defined as

$$\begin{aligned}
P_{Tmn}(v) &= \delta_{mn} - \frac{v_m v_n}{v^2}, \\
P_{Lmn}(v) &= \frac{v_m v_n}{v^2}. \tag{1.11}
\end{aligned}$$

Although informally obtained in the Introduction, these results also follow from the coset method of Section II for the brane embedded in  $AdS - C$  space when the flat (Minkowski) space-time limit  $m^2 \rightarrow 0$  is taken and  $AdS - C \xrightarrow{m^2 \rightarrow 0} C$ , Carroll spacetime.

The field equations reflect the Carroll spacetime symmetries yielding  $\dot{\phi} = 0$  from the  $w$  equation of motion,  $\delta\Gamma_C/\delta w = 0$  so that the brane's initial spatial shape does not evolve as expected from the collapse of the light cone in this limit. Having set  $\dot{\phi} = 0$ , the spatial components of  $v_m$  obey the constraint  $2v_m \frac{\tan\sqrt{4v^2}}{\sqrt{4v^2}} = \partial^m\phi$  as dictated by the  $v_m$  field equation,  $\delta\Gamma_C/\delta v_m = 0$ . Both field equations are consistent with the initial auxiliary velocity field equation (1.7) for  $c \rightarrow 0$ . Finally the broken space translation symmetry in the  $(p + 1)$  direction yields the time variation of the momentum density as given by the  $\phi$  field equation  $\delta\Gamma_C/\delta\phi = 0$ . The momentum density  $\Pi$  is defined by

$$\Pi \equiv \partial\mathcal{L}_C/\partial\dot{\phi} = 2w \frac{\sin\sqrt{4v^2}}{\sqrt{4v^2}}, \tag{1.12}$$

while the derivatives of  $\mathcal{L}_C$  with respect to the spatial derivatives of  $\phi$  are denoted

$$\Pi_m = \partial\mathcal{L}_C/\partial\partial^m\phi = 2v_m \frac{\sin\sqrt{4v^2}}{\sqrt{4v^2}}. \tag{1.13}$$

Thus the  $\phi$ -field equation has the form of a current conservation equation. Indeed, the field equation is the spontaneously broken translation current conservation equation for the Carroll spacetime. The corresponding Noether current has the conserved form as above

$$\frac{\partial}{\partial t}\Pi + \frac{\partial}{\partial x_m}\Pi_m = 0. \tag{1.14}$$

The action, Equation (1.9), is invariant under the Carroll transformations, obtained by contracting the Poincaré transformations as  $c \rightarrow 0$ , of the  $(d + 1)$ -dimensional Carroll spacetime which include the unbroken  $d$ -dimensional worldvolume time and space translations, space rotations and boosts with respective parameters  $\epsilon$ ,  $a_m$ ,  $\alpha_{mn}$ ,  $\beta_m$  and additionally, now non-linearly realized, the broken space translation of the covolume which is just a  $\phi$  shift symmetry, boosts in that direction and rotations in a worldvolume-covolume plane with respective parameters  $\zeta$ ,  $\lambda$ ,  $\kappa_m$ . Exploiting the invariance of the Minkowski interval  $ds^2 = dx^\mu\eta_{\mu\nu}dx^\nu - dz^2$  under  $(d + 1)$ -dimensional Poincaré transformations

$$x'_M = x_M + \lambda_{MN}\eta^{NP}x_P + a_M, \tag{1.15}$$

where  $x_M = (ct, x_m, z)$  with  $z = \phi(x)$  and  $\lambda_{MN} = -\lambda_{NM}$ , the coordinate and field Poincaré transformations have the form

$$\begin{aligned} t' &= t - \frac{1}{c}\lambda_{0n}x_n - \frac{1}{c}\lambda_{0z}\phi + \frac{1}{c}a_0, \\ x'_m &= x_m + c\lambda_{0m}t - \lambda_{mn}x_n - \lambda_{mz}\phi + a_m, \\ \phi'(x', t') &= \phi(x, t) + c\lambda_{z0}t - \lambda_{zm}x_m + a_z. \end{aligned} \quad (1.16)$$

Contracting the Poincaré symmetry transformations to those of the Carroll symmetries requires a rescaling of the time components of the transformation parameters so that

$$a_0 = c\epsilon, \quad \lambda_{0m} = c\beta_m, \quad \lambda_{0z} = c2\lambda, \quad (1.17)$$

while the purely spatial components are unchanged and are denoted as

$$\zeta = a_z, \quad \alpha_{mn} = -\lambda_{mn}, \quad 2\kappa_m = \lambda_{mz}. \quad (1.18)$$

The new parameters denote the Carroll transformation parameters. The  $c \rightarrow 0$  Carroll spacetime transformations of the coordinates and field  $\phi$  are thus obtained and have the non-linear form (see [Appendix A](#) for the coset method derivation and the  $AdS - C$  to Carroll spacetime  $C$  limit to obtain Equations (1.19) and (1.21))

$$\begin{aligned} t' &= t + \epsilon - 2\lambda\phi - \beta_mx_m, \\ x'_m &= x_m + a_m + \alpha_{mn}x_n - 2\phi\kappa_m, \\ \phi'(x', t') &= \phi(x, t) + \zeta + 2\kappa_mx_m. \end{aligned} \quad (1.19)$$

Applying these transformations to the auxiliary Lagrange multiplier field definition, Equation (1.7), so that

$$\partial'_\mu\phi'(x', t') = -V'_\mu(x', t')\frac{\tanh\sqrt{V'^2}}{\sqrt{V'^2}}, \quad (1.20)$$

with  $V'_\mu(x', t') = V_\mu(x, t) + \Delta V_\mu(x, t)$ , where  $\Delta V_\mu = (2c\Delta w, 2\Delta v_m)$  yields the  $w$  and  $v_m$  auxiliary fields' Carroll transformations

$$\begin{aligned} w'(x', t') &= w(x, t) \left[ 1 + \frac{v_m\kappa_m}{v^2} \left( 1 - \sqrt{4v^2} \cot\sqrt{4v^2} \right) \right] \\ &\quad - \beta_mv_m + \lambda\sqrt{4v^2} \cot\sqrt{4v^2}, \\ v'_m(x', t') &= v_m(x, t) + \alpha_{mn}v_n + \left( \sqrt{4v^2} \cot\sqrt{4v^2} P_{Tmn}(v) + P_{Lmn}(v) \right) \kappa_n. \end{aligned} \quad (1.21)$$

Since the time and space transformations involve functions thereof, the differential form of Equation (1.19) yields the general coordinate transformation  $G = \partial(t', x')/\partial(t, x)$ . That is, recombining  $t$  and  $x_m$  in the matrix  $X_M = (t, x_m)$  where now  $M, N = 0, 1, \dots, p$ , the transformations are given by

$$\begin{aligned} dX'_M &= (dt', dx'_m) \\ &= dX_N G^N_M = (dtG^0_0 + dx_n G^n_0, dtG^0_m + dx_n G^n_m), \end{aligned} \quad (1.22)$$

with

$$\begin{aligned} G^N_M &= \begin{pmatrix} \frac{\partial t'}{\partial t} & \frac{\partial x'_m}{\partial t} \\ \frac{\partial t'}{\partial x_n} & \frac{\partial x'_m}{\partial x_n} \end{pmatrix}_M^N \\ &= \begin{pmatrix} 1 - 2\lambda\dot{\phi} & -2\dot{\phi}\kappa_m \\ (-\beta_n - 2\lambda\partial^n\phi) & (\delta_{nm} + \alpha_{nm} - 2\partial^n\phi\kappa_m) \end{pmatrix}_M^N. \end{aligned} \quad (1.23)$$

The spacetime transformation Jacobian is  $dt'd^p x' = dt d^p x \det G$ . On the other hand, the action  $\Gamma_C$  is invariant under the Carrollian symmetry transformations, thus

$$\Gamma'_C = \int dt' d^p x' \mathcal{L}'_C(x', t') = \int dt d^p x \det G \mathcal{L}'_C(x', t')$$

TABLE I. Carroll spacetime transformations and Noether currents.

Transformation	Parameter	Noether currents
Time translations	$\epsilon$	$\mathcal{H} = \Pi\dot{\phi} - \mathcal{L}_C$ $= \cos\sqrt{4v^2 + \Pi_m\dot{\theta}^m}\phi$
Space translations	$a_n$	$h_m = \Pi_m\dot{\phi}$ $T^n = \Pi\dot{\theta}^n\phi$ $T_m^n = \frac{\partial\mathcal{L}_C}{\partial\dot{\theta}^m\phi}\dot{\theta}^n\phi - \delta_m^n\mathcal{L}_C$ $= \delta_m^n[\cos\sqrt{4v^2 + \Pi\dot{\phi}} + \Pi_r\dot{\theta}^r\phi] - \Pi_m\dot{\theta}^n\phi$
Broken space translations	$\zeta$	$z = \Pi$ $z_m = \Pi_m$
Broken boosts	$\lambda$	$l = \phi\mathcal{H}$ $l_m = \phi h_m$
Broken rotations	$\kappa_n$	$K^n = 2\phi T^n + 2\delta^{nr}x_r\Pi$ $K_m^n = 2\phi T_m^n + 2\Pi_m\delta^{nr}x_r$
Unbroken rotations	$\alpha_{rs}$	$\mathcal{M}^{rs} = x_r T^s - x_s T^r$ $\mathcal{M}_m^{rs} = x_r T_m^s - x_s T_m^r$
Unbroken boosts	$\beta_n$	$B^n = \delta^{nr}x_r\mathcal{H}$ $B_m^n = \delta^{nr}x_r h_m$

$$= \int dt d^p x \mathcal{L}_C(x, t) = \Gamma_C, \quad (1.24)$$

so that

$$\mathcal{L}'_C(x', t') = \det G^{-1} \mathcal{L}_C(x, t). \quad (1.25)$$

For these Carrollian transformations, the Noether currents take the couplet form of time and spatial component currents

$$J_M = \begin{pmatrix} \Pi\delta\phi + \delta t \mathcal{L}_C \\ \Pi_m\delta\phi + \delta x_m \mathcal{L}_C \end{pmatrix}_{M=(0,m)} \quad (1.26)$$

with  $\delta t = t' - t$  and  $\delta x_m = x'_m - x_m$ , where the intrinsic transformation  $\delta\phi$  is defined as

$$\delta\phi \equiv \phi'(x, t) - \phi(x, t) = \Delta\phi(x, t) - \delta t \dot{\phi} - \delta x_m \dot{\theta}^m \phi \quad (1.27)$$

with the total variation given by

$$\Delta\phi(x, t) \equiv \phi'(x', t') - \phi(x, t). \quad (1.28)$$

Thus Noether's theorem is ( $\varphi_i = \{\phi, w, v_m\}$ )

$$\begin{aligned} \Delta\mathcal{L}_C &= \mathcal{L}'_C(x', t') - \mathcal{L}_C(x, t) \\ &= \frac{\partial}{\partial t} J_0 + \frac{\partial}{\partial x_m} J_m - \mathcal{L}_C \frac{\partial\delta t}{\partial t} - \mathcal{L}_C \frac{\partial\delta x_m}{\partial x_m} + \frac{\delta\Gamma_C}{\delta\varphi_i} \delta\varphi_i, \end{aligned} \quad (1.29)$$

where the last term vanishes by the field equations  $\delta\Gamma_C/\delta\varphi_i = 0$ . The conserved currents (before use of the field equation constraints) are given by the pairs displayed in Table I. The action is invariant as reflected by the vanishing or explicit cancellation of the  $\Delta\mathcal{L}_C + \mathcal{L}_C \left( \frac{\partial\delta t}{\partial t} + \frac{\partial\delta x_m}{\partial x_m} \right) = 0$  terms in Noether's theorem for each of the Carrollian symmetry transformations. Hence  $\partial^M J_M = \frac{\partial}{\partial t} J_0 + \frac{\partial}{\partial x_m} J_m = -\frac{\delta\Gamma_C}{\delta\varphi_i} \delta\varphi_i$ . The associated conserved charges are given by  $Q = \int d^p x J_0$ , where  $\dot{Q} = -\int_{S \rightarrow \infty} \vec{J} \cdot d\vec{S} \rightarrow 0$ . Once again the time evolution of the momentum density is contained in the broken translation current conservation equation and the invariance of the Lagrangian

$$0 = \dot{\Pi} + \partial^m \Pi_m = -\frac{\delta\Gamma_C}{\delta\phi}. \quad (1.30)$$

In summary, the  $w$ -field equation of motion, Equation (1.10), yields the frozen spatial distribution of the domain wall as expected from the collapse of the lightcone to the positive time half-line

in the Carrollian  $c \rightarrow 0$  limit:  $\frac{\partial}{\partial t} \phi(x, t) = 0$ . Along with this the Lagrange multiplier  $v_m$ -field equation of motion simply reproduces the constraint of the “inverse Higgs mechanism”<sup>13</sup>

$$2v_m \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} = \partial^m \phi. \quad (1.31)$$

These field equation constraints also follow directly from the Lagrange multiplier equation (1.7) in the  $c \rightarrow 0$  limit. Finally, although the brane is stationary, the momentum must vary in time in order to balance the tension due to the domain wall’s local spatial shape where using the Lagrange multiplier constraints so that  $\dot{\phi} = 0$  and  $2v_m \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} = \partial^m \phi$ , it is found that

$$\dot{\Pi} = -\partial^m \left[ \frac{\partial^m \phi}{\sqrt{1 + \partial^s \phi \partial^s \phi}} \right]. \quad (1.32)$$

The purpose of this paper is to determine the Carrollian limit for branes in  $AdS$  spacetime.<sup>6,7</sup> The  $D = d + 1$  dimensional  $AdS$  spacetime symmetry algebra is contracted in the Carroll limit,  $c \rightarrow 0$ . For a p-brane action in the alternate string Carrollian limit of Minkowski space see Ref. 8. Application of the Carrollian limit to gravity and electromagnetism is discussed in Refs. 9 and 10.

In Section II coset methods<sup>11-13</sup> are applied to the  $AdS - C$  algebra for the case of an embedded co-dimension one p-brane (domain wall). The induced vielbeine, covariant derivatives, and spin connections are determined using the Maurer-Cartan one-form associated with the p-brane coset element. The action is constructed and shown to be invariant under the non-linearly realized  $AdS - C_{d+1}$  broken to  $AdS - C_d$  symmetries by the brane embedding. The symmetry transformations are detailed in Appendix A. Alternatively, the  $AdS - C$  action can be obtained by making the speed of light  $c$  dependence explicit in the  $AdS_{d+1} \rightarrow AdS_d$  case and taking the  $c \rightarrow 0$  limit. Using the results of Ref. 6, this approach is demonstrated in Appendix B.

From the action the field equations are determined. As expected due to the collapse of the forward light cone to the positive time half-line, the spatial shape of the brane is stationary. However, the spatial shape of the brane as well as the  $AdS - C$  geometry requires its conjugate momentum density to be time dependent. Finally Noether’s theorem is applied to the broken space translation symmetry in order to calculate the current and its conservation equation. Section III presents the action in terms of a product of the background  $AdS - C_d$  world volume vielbein and the Nambu-Goto-Carrollian vielbein. This is then used to express the action in terms of its dual vector theory. The results of the brane embedding are reviewed in Section IV.

## II. AdS-CARROLL SPACE AND THE COSET METHOD

The AdS-Carroll spacetime is defined by the Wigner-Inönü contraction of the AdS symmetry algebra for the speed of light vanishing,  $c \rightarrow 0$ . The isometry group of the  $D$ -dimensional  $AdS$  space is given by the  $SO(2, D - 1)$  algebra of symmetry generators with the commutation relations

$$\begin{aligned} [M^{MN}, M^{RS}] &= -i(\eta^{MR} M^{NS} - \eta^{MS} M^{NR} + \eta^{NS} M^{MR} - \eta^{NR} M^{MS}), \\ [M^{MN}, P^L] &= i(P^M \eta^{NL} - P^N \eta^{ML}), \\ [P^M, P^N] &= -im^2 M^{MN}, \end{aligned} \quad (2.1)$$

where  $L, M, N, R, S = 0, 1, 2, \dots, D - 1$  and the metric  $\eta_{MN} = (+1, -1, -1, \dots, -1)$  with  $m^2 = 1/R^2$  and  $R$  the curvature of the AdS hyperboloid.

Introducing the explicit factors of the speed of light for the time related components, the time component involved generators  $H$  and  $B^A$  are defined as

$$\begin{aligned} P^0 &= \frac{1}{c} H, \\ M^{A0} &= \frac{1}{c} B^A, \end{aligned} \quad (2.2)$$



while the spatial components remain unscaled  $P^A, M^{AB}$ ,

$$\begin{aligned} P^A &= P^A, \\ M^{AB} &= M^{AB}, \end{aligned} \quad (2.3)$$

for  $A, B, C, D = 1, 2, \dots, (D-1)$  denoting the spatial indices. The  $SO(2, D-1)$  algebra contracts to the AdS-Carroll algebra in the  $c \rightarrow 0$  limit

$$\begin{aligned} [M^{AB}, M^{CD}] &= +i(\delta^{AC}M^{BD} - \delta^{AD}M^{BC} + \delta^{BD}M^{AC} - \delta^{BC}M^{AD}), \\ [M^{AB}, B^C] &= -i(B^A\delta^{BC} - B^B\delta^{AC}), \\ [M^{AB}, P^C] &= -i(P^A\delta^{BC} - P^B\delta^{AC}), \\ [B^A, P^B] &= +i\delta^{AB}H, \\ [H, P^A] &= +im^2B^A, \\ [P^A, P^B] &= -im^2M^{AB}, \end{aligned} \quad (2.4)$$

with remaining commutators vanishing.

A brane embedded in this AdS-Carroll spacetime will break its  $AdS - C_D$  symmetries down to those of the  $d$ -dimensional worldvolume  $AdS - C_d$  and its complementary covolume with the remaining symmetries being spontaneously broken. In the case considered here, the insertion of a domain wall results in a  $d = (1+p) = (D-1)$ -dimensional worldvolume and 1-dimensional covolume. Choosing the  $(p+1)^{\text{th}}$  spatial direction as the broken translation symmetry direction, the AdS-Carroll algebra can be expressed in terms of broken and unbroken generators with the generators  $H, P^m, M^{mn}, B^m$ , with  $m, n = 1, 2, \dots, p$ , as unbroken generators and  $P^{p+1} \equiv Z, M^{p+1}, m \equiv \frac{1}{2}K^m, B^{p+1} \equiv \frac{1}{2}L$  as the broken generators.  $M^{mn}$  are the  $SO(p)$  worldvolume spatial rotation generators while the worldvolume spatial translation generators  $P^m$  form an  $SO(p)$  vector with time translations generated by  $H$ . The  $SO(p)$  vector  $B^m$  generates worldvolume boosts in the  $m$ -direction. The translations in the covolume spatial direction are generated by  $Z$  while boosts in that direction are generated by  $L$ . Finally broken rotations in the covolume- $m$  worldvolume plane are generated by the  $SO(p)$  vector  $K^m$ . Consequently the  $AdS - C_{D=d+1}$  algebra can be expressed in terms of these worldvolume and domain wall charges. The  $AdS - C_{d=p+1}$  worldvolume isometries are given by the  $H, P^m, M^{mn}, B^m$  algebra (only nontrivial commutators listed)

$$\begin{aligned} [M^{mn}, M^{rs}] &= +i(\delta^{mr}M^{ns} - \delta^{ms}M^{nr} + \delta^{ns}M^{mr} - \delta^{nr}M^{ms}), \\ [M^{mn}, B^l] &= -i(B^m\delta^{nl} - B^n\delta^{ml}), \\ [M^{mn}, P^l] &= -i(P^m\delta^{nl} - P^n\delta^{ml}), \\ [B^m, P^n] &= +i\delta^{mn}H, \\ [H, P^m] &= +im^2B^m, \\ [P^m, P^n] &= -im^2M^{mn}. \end{aligned} \quad (2.5)$$

The broken symmetry generators  $Z, L, K^m$  commute with the unbroken generators above according to their unbroken subgroup representation

$$\begin{aligned} [M^{mn}, Z] &= 0, \\ [M^{mn}, L] &= 0, \\ [M^{mn}, K^l] &= -i(K^m\delta^{nl} - K^n\delta^{ml}), \\ [B^m, Z] &= 0, \\ [B^m, L] &= 0, \\ [B^m, K^n] &= +i\delta^{mn}L, \\ [P^m, Z] &= \frac{i}{2}m^2K^m, \\ [P^m, L] &= 0, \\ [P^m, K^n] &= +2i\delta^{mn}Z, \end{aligned}$$

$$\begin{aligned}
[H, Z] &= \frac{i}{2}m^2L, \\
[H, L] &= 0, \\
[H, K^n] &= 0.
\end{aligned} \tag{2.6}$$

Finally the broken charges  $Z, L, K^m$  commute amongst themselves to yield the charges of the unbroken subalgebra

$$\begin{aligned}
[Z, L] &= -2iH, \\
[Z, K^m] &= -2iP^m, \\
[L, K^m] &= -4iB^m, \\
[K^m, K^n] &= 4iM^{mn}.
\end{aligned} \tag{2.7}$$

The domain wall spontaneously breaks the  $AdS - C_D$  spacetime symmetries down to those of the  $AdS - C_d$  worldvolume. As a derivative expansion, the leading form of the brane action is uniquely determined. The Goldstone boson fields  $\phi(x, t)$  corresponding to the long wavelength oscillations of the domain wall parameterize the coset coordinates along with the fields associated with the broken boost and rotations,  $w(x, t)$  and  $v_m(x, t)$ , respectively. The geometry of the underlying  $AdS - C_d$  worldvolume spacetime is described by the time  $t$  and space  $x_m$ ,  $m = 1, 2, \dots, p = (d - 1)$ , coordinate group elements. Overall these fields and spacetime coordinates parameterize the  $AdS - C_D/ISO(p)$  coset element  $\Omega$  (note generators are defined with superscripts)

$$\Omega \equiv e^{itH + ix_m P^m} e^{i\phi(x, t)Z} e^{iw(x, t)L + iv_m(x, t)K^m}, \tag{2.8}$$

where  $ISO(p)$  is the unbroken subgroup with generators  $M^{mn}$  and  $B^m$ . The background worldvolume coset  $\bar{\Omega} \in AdS - C_d/ISO(p)$

$$\bar{\Omega} \equiv e^{itH + ix_m P^m} \tag{2.9}$$

is used to determine the  $AdS - C_d$  background vielbeine and spin connections via the Maurer-Cartan 1-form  $\tilde{\omega}$

$$\tilde{\omega} \equiv -i\bar{\Omega}^{-1}d\bar{\Omega} = \bar{\omega}_H H + \bar{\omega}_{P_a} P^a + \frac{1}{2}\bar{\omega}_{ab} M^{ab} + \bar{\omega}_{B_a} B^a. \tag{2.10}$$

Expanding the 1-forms in terms of the coordinate differentials, the Maurer-Cartan 1-form becomes (with tangent space indices denoted  $a, b = 1, 2, \dots, p$  and world volume indices denoted  $m, n = 1, 2, \dots, p$ )

$$\begin{aligned}
\tilde{\omega} &= (dt\bar{e}_0^0 + dx_m\bar{e}_0^m)H + (dt\bar{e}_a^0 + dx_n\bar{e}_a^n)P^a \\
&\quad + \frac{1}{2}(dt\bar{\omega}_{ab}^t + dx_r\bar{\omega}_{ab}^r)M^{ab} + (dt\bar{\omega}_{0a}^t + dx_r\bar{\omega}_{0a}^r)B^a,
\end{aligned} \tag{2.11}$$

where the background vielbeine are found to be

$$\begin{aligned}
\bar{e}_0^0 &= \frac{\sinh\sqrt{m^2x^2}}{\sqrt{m^2x^2}}, \\
\bar{e}_0^m &= \frac{x_m t}{x^2} \left( 1 - \frac{\sinh\sqrt{m^2x^2}}{\sqrt{m^2x^2}} \right), \\
\bar{e}_a^n &= \left( \frac{\sinh\sqrt{m^2x^2}}{\sqrt{m^2x^2}} \right) P_{Tna}(x) + P_{Lna}(x), \\
\bar{e}_a^0 &= 0,
\end{aligned} \tag{2.12}$$

with  $x^2 = x_m \delta^{mn} x_n = x_m x_m$ . The background spin connections are also obtained as

$$\begin{aligned}
\bar{\omega}_{ab}^t &= 0, \\
\bar{\omega}_{ab}^r &= \left( \frac{1 - \cosh\sqrt{m^2x^2}}{x^2} \right) (\delta_a^s \delta_b^r - \delta_b^s \delta_a^r) x_s,
\end{aligned}$$

$$\begin{aligned}\tilde{\omega}_{0a}^t &= \left( \frac{1 - \cosh \sqrt{m^2 x^2}}{x^2} \right) x_a, \\ \tilde{\omega}_{0a}^r &= - \left( \frac{1 - \cosh \sqrt{m^2 x^2}}{x^2} \right) t \delta_a^r.\end{aligned}\tag{2.13}$$

The  $AdS - C_d$  background vielbein  $\tilde{E}_A^M$  is defined as the matrix relating the coordinate differentials  $dX_N = (dt, dx_n)$ , with  $M, N = 0, 1, \dots, p$ , to the covariant coordinate differentials  $\tilde{\omega}_A = (\tilde{\omega}_H, \tilde{\omega}_{Pa})$ , with  $A, B = 0, 1, \dots, p$  as well as  $\tilde{\omega}_0 = \tilde{\omega}_H$  and  $\tilde{\omega}_a = \tilde{\omega}_{Pa}$ , thus

$$\begin{aligned}\tilde{\omega}_A &= (\tilde{\omega}_H \ \tilde{\omega}_{Pa}) = dX_M \tilde{E}_A^M \\ &= (dt \ dx_m) \begin{pmatrix} \tilde{e}_0^0 & 0 \\ \tilde{e}_0^m & \tilde{e}_a^m \end{pmatrix},\end{aligned}\tag{2.14}$$

that is

$$\tilde{E}_A^M = \begin{pmatrix} \tilde{e}_0^0 & \tilde{e}_a^0 = 0 \\ \tilde{e}_0^m & \tilde{e}_a^m \end{pmatrix} = \begin{pmatrix} \tilde{E}_0^0 & \tilde{E}_a^0 = 0 \\ \tilde{E}_0^m & \tilde{E}_a^m \end{pmatrix},\tag{2.15}$$

with  $\det \tilde{E} = \tilde{e}_0^0 \det \tilde{e}_a^m$ , where  $a, b, m, n = 1, 2, \dots, p$ .

On the other hand, the Maurer-Cartan 1-form for the domain wall breakdown of  $AdS - C_D \rightarrow AdS - C_d$  can be constructed using the coset element  $\Omega$

$$\begin{aligned}\tilde{\omega} &= -i\Omega^{-1}d\Omega \\ &= \omega_H H + \omega_{Pa} P^a + \omega_Z Z + \omega_L L + \omega_{Ka} K^a + \frac{1}{2} \omega_{Mab} M^{ab} + \omega_{Ba} B^a.\end{aligned}\tag{2.16}$$

This yields the vielbeine, covariant derivatives of the fields, and spin connections. The vielbeine are given in terms of the coordinate differentials according to

$$\begin{aligned}\omega_H &= dt e_0^0 + dx_m e_0^m, \\ \omega_{Pa} &= dt e_a^0 + dx_m e_a^m,\end{aligned}\tag{2.17}$$

with  $v^2 = v_a \delta^{ab} v_b = v_a v_a$  and

$$\begin{aligned}e_0^0 &= \tilde{e}_0^0 \cosh \sqrt{m^2 \phi^2} + 2w\phi \left( \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} \right), \\ e_0^m &= \tilde{e}_0^m \cosh \sqrt{m^2 \phi^2} + 2w\partial^m \phi \left( \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} \right) \\ &\quad + \cosh \sqrt{m^2 \phi^2} \left( \frac{\cos \sqrt{4v^2} - 1}{v^2} \right) wv_a \tilde{e}_a^m, \\ e_a^0 &= 2\phi v_a \left( \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} \right), \\ e_a^m &= 2\partial^m \phi v_a \left( \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} \right) \\ &\quad + \cosh \sqrt{m^2 \phi^2} [P_{Tab}(v) + (\cos \sqrt{4v^2}) P_{Lab}(v)] \tilde{e}_b^m.\end{aligned}\tag{2.18}$$

The  $AdS - C_D$  spacetime vielbein  $E_A^M$  is defined as the matrix relating the coordinate differentials  $dX_M = (dt, dx_m)$  to the covariant coordinate differentials  $\tilde{\omega}_A = (\omega_H, \omega_{Pa})$ , that is  $\tilde{\omega}_0 = \omega_H$  and  $\tilde{\omega}_a = \omega_{Pa}$ , thus

$$\begin{aligned}\tilde{\omega}_A &= (\omega_H \ \omega_{Pa}) = dX_M E_A^M \\ &= (dt \ dx_m) \begin{pmatrix} e_0^0 & e_a^0 \\ e_0^m & e_a^m \end{pmatrix},\end{aligned}\tag{2.19}$$

that is

$$E^M_A = \begin{pmatrix} E^0_0 & E^0_a \\ E^m_0 & E^m_a \end{pmatrix} = \begin{pmatrix} e^0_0 & e^0_a \\ e^m_0 & e^m_a \end{pmatrix}. \quad (2.20)$$

The brane field's covariant derivatives,  $\nabla^t \phi$  and  $\nabla^m \phi$ , are given by the  $\omega_Z$  one-form

$$\omega_Z = dt \nabla^t \phi + dx_m \nabla^m \phi, \quad (2.21)$$

where

$$\begin{aligned} \nabla^t \phi &= \dot{\phi} \cos \sqrt{4v^2}, \\ \nabla^m \phi &= \bar{e}^m_a \cosh \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} \left( \frac{\bar{e}^{-1a}_n \frac{\partial}{\partial x_n} \phi}{\cosh \sqrt{m^2 \phi^2}} - 2\delta^{ab} v_b \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) \right). \end{aligned} \quad (2.22)$$

Likewise the auxiliary fields  $w$  and  $v_a$  have covariant derivatives determined by  $\omega_L$  and  $\omega_{K_a}$ ,

$$\begin{aligned} \omega_L &= dt \nabla^t w + dx_m \nabla^m w, \\ \omega_{K_a} &= dt \nabla^t v_a + dx_m \nabla^m v_a, \end{aligned} \quad (2.23)$$

where the derivatives are found to be

$$\begin{aligned} \nabla^t w &= \dot{w} + \left( \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} - 1 \right) \left( \dot{w} - w \frac{v_a \dot{v}_a}{v^2} \right) - \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} \bar{\omega}^t_{0a} v_a \\ &\quad - \frac{1}{2} m^2 \phi \frac{\sinh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \cos \sqrt{4v^2} \bar{e}^0_0, \\ \nabla^m w &= \partial^m w + \left( \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} - 1 \right) \left( \partial^m w - w \frac{v_a \partial^m v_a}{v^2} \right) - \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} \bar{\omega}^m_{0a} v_a \\ &\quad - \frac{1}{2} m^2 \phi \frac{\sinh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \left[ \cos \sqrt{4v^2} \bar{e}^m_0 + \frac{(1 - \cos \sqrt{4v^2})}{v^2} w v_a \bar{e}^m_a \right], \\ \nabla^t v_a &= \dot{v}_a + \left( \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} - 1 \right) P_{Tab}(v) \dot{v}_b + \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} \bar{\omega}^t_{ab} v_b, \\ \nabla^m v_a &= \partial^m v_a + \left( \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} - 1 \right) P_{Tab}(v) \partial^m v_b + \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} \bar{\omega}^s_{ab} v_b \\ &\quad - \frac{1}{2} m^2 \phi \frac{\sinh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \left[ \cos \sqrt{4v^2} P_{Tab}(v) + P_{Lab}(v) \right] \bar{e}^m_b. \end{aligned} \quad (2.24)$$

Finally the spin connections are obtained from  $\omega_M$  and  $\omega_B$

$$\begin{aligned} \omega_{Mab} &= dt \omega^t_{ab} + dx_m \omega^m_{ab} \\ &= \bar{\omega}_{ab} + m^2 \phi \frac{\sinh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} \\ &\quad + \left( 1 - \cos \sqrt{4v^2} \right) \left[ P_{Lac}(v) \bar{\omega}_{bc} - P_{Lbc}(v) \bar{\omega}_{ac} - \left( \frac{dv_a v_b - dv_b v_a}{v^2} \right) \right], \\ \omega_{Ba} &= dt \omega^t_a + dx_m \omega^m_a \\ &= \bar{\omega}_{Ba} + \left( \frac{1 - \cos \sqrt{4v^2}}{v^2} \right) [dw v_a - w dv_a - \bar{\omega}_{ab} w v_b - \bar{\omega}_{Bb} v_b v_a] \\ &\quad + 2m^2 \phi \frac{\sinh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} [w \bar{e}^m_a dx_m - v_a \bar{e}^0_0 dt - v_a dx_m \bar{e}^m_0]. \end{aligned} \quad (2.25)$$

The  $AdS - C_D$  invariant action is constructed in terms of the vielbein  $E$

$$\Gamma_{AdS-C_D} = \int dt d^p x \mathcal{L}_{AdS-C_D}(x, t) = \int dt d^p x \det E. \quad (2.26)$$

The  $AdS - C_D$  transformations are non-linearly realized according to the group multiplication properties involving the coset  $\Omega$  as detailed in [Appendix A](#). The invariance of the action follows from the transformation properties of the vielbein  $E$ . The Maurer-Cartan one-forms transform according to which representation of the local  $ISO(p)$  tangent space transformations that the associated operator belongs given by the unbroken subgroup element  $h(x, t)$  obtained in [Appendix A](#). Using the coset transformation law  $g\Omega(x, t) = \Omega'(x', t')h(x, t)$ , the one-forms transform as  $\omega'(x', t') = h(x, t)\omega(x, t)h^{-1}(x, t) - ih(x, t)dh^{-1}(x, t)$ , yielding

$$\begin{aligned} \omega' &= \omega'_H H + \omega'_{P_a} P^a + \omega'_Z Z + \omega'_L L + \omega'_{K_a} K^a + \frac{1}{2} \omega'_{M_{ab}} M^{ab} + \omega'_{B_a} B^a \\ &= h\omega h^{-1} - ih dh^{-1} \\ &= \omega_H h H h^{-1} + \omega_{P_a} h P^a h^{-1} + \omega_Z h Z h^{-1} + \omega_L h L h^{-1} + \omega_{K_a} h K^a h^{-1} \\ &\quad + \frac{1}{2} \omega_{M_{ab}} h M^{ab} h^{-1} + \omega_{B_a} h B^a h^{-1} - \frac{1}{2} d\theta_{ab} M^{ab} - d\theta_a B^a. \end{aligned} \quad (2.27)$$

Hence the one-forms' variations are obtained

$$\begin{aligned} \omega'_H &= \omega_H - \omega_{P_a} \theta_a, \\ \omega'_{P_a} &= \omega_{P_b} (\delta_{ba} - \theta_{ba}) \equiv \omega_{P_b} R_{ba}^{-1}, \\ \omega'_Z &= \omega_Z, \\ \omega'_L &= \omega_L - \omega_{K_a} \theta_a, \\ \omega'_{K_a} &= \omega_{K_b} R_{ba}^{-1}, \\ \omega'_{M_{cd}} &= \omega_{M_{ab}} R_{ac}^{-1} R_{db}^{-1} - d\theta_{cd}, \\ \omega'_{B_a} &= \omega_{B_b} R_{ba}^{-1} - d\theta_a + \frac{1}{2} \omega_{M_{cd}} (\theta_c \delta_{da} - \theta_d \delta_{ca}). \end{aligned} \quad (2.28)$$

The covariant coordinate differentials and vielbeine transform as

$$\begin{aligned} \tilde{\omega}'_A &= (\omega'_H \ \omega'_{P_a}) = (\omega_H \ \omega_{P_b}) \begin{pmatrix} 1 & 0 \\ -\theta_b & R_{ba}^{-1} \end{pmatrix} \\ &= dX'_M E'^M_A = (dt' \ dx'_m) \begin{pmatrix} e'^0_0 & e'^0_a \\ e'^m_0 & e'^m_a \end{pmatrix} \\ &= (dt \ dx_r) \begin{pmatrix} e^0_0 & e^0_b \\ e^r_0 & e^r_b \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -\theta_b & R_{ba}^{-1} \end{pmatrix} = dX_R E^R_B \Lambda^B_A, \end{aligned} \quad (2.29)$$

with letters from the beginning of the alphabet denoting the tangent space transformation properties. From [Appendix A](#) the coordinate differentials transform according to the general coordinate transformation

$$dX'_M = (dt' \ dx'_m) = (dt \ dx_n) \begin{pmatrix} G^0_0 & G^0_m \\ G^n_0 & G^n_m \end{pmatrix} = dX_N G^N_M, \quad (2.30)$$

where the complicated general coordinate transformation matrix is denoted by  $G^N_M$ , with letters in the middle of the alphabet indicating world volume coordinate transformations. The spacetime differentials have the Jacobian  $dt' d^p x' = dt d^p x \det G$ . Thus the vielbein  $E^M_A$  transforms as

$$E'^M_A = G^{-1M}_N E^N_B \Lambda^B_A, \quad (2.31)$$

where the tangent space transformations have been denoted by

$$\Lambda^B_A = \begin{pmatrix} 1 & 0 \\ -\theta_b & R^{-1}_{ba} \end{pmatrix}. \tag{2.32}$$

Noting that  $\det \Lambda = 1$  so that  $\det E' = \det G^{-1} \det E$ , the action is invariant

$$\begin{aligned} \Gamma'_{AdS-C_D} &= \int dt' d^p x' \det E' = \int dt d^p x \det G \det G^{-1} \det E \\ &= \int dt d^p x \det E = \Gamma_{AdS-C_D}. \end{aligned} \tag{2.33}$$

With the vielbeine in Equation (2.18) the  $AdS - C_D$  invariant action is found

$$\Gamma_{AdS-C_D} = \int dt d^p x \det E = \int dt d^p x \det(e^m_a) [e^0_0 - e^0_a e^{-1a}_n e^n_0], \tag{2.34}$$

with (noting that  $\bar{e}^{-1a}_m \bar{e}^m_0 = \delta^a_m \bar{e}^m_0 = \bar{e}^a_0$ )

$$\begin{aligned} \det E &= \det \bar{E} \cosh^p \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} \left\{ \cosh \sqrt{m^2 \phi^2} + \mathcal{D}^a \phi 2v_a \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right. \\ &\quad \left. + 2\bar{e}^0_0^{-1} \dot{\phi} (w - v_a \bar{e}^a_0) \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right\}. \end{aligned} \tag{2.35}$$

The background  $AdS - C_d$  spacetime measure is given by  $\det \bar{E} = \bar{e}^0_0 \det \bar{e}^m_a$  and the partially covariant spatial derivative  $\mathcal{D}^a \phi = \bar{e}^{-1a}_m \frac{\partial}{\partial x_m} \phi$ . Note that the covariant derivatives of  $\phi$ , Equation (2.22), are given by

$$\begin{aligned} \nabla^t \phi &= \dot{\phi} \cos \sqrt{4v^2}, \\ \nabla^m \phi &= \bar{e}^m_a \cosh \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} \left( \frac{\mathcal{D}^a \phi}{\cosh \sqrt{m^2 \phi^2}} - 2\delta^{ab} v_b \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) \right), \end{aligned} \tag{2.36}$$

and can be used to covariantly constrain (as  $\omega'_Z = \omega_Z$  is invariant, Equation (2.28)) the field  $v_a$  equivalent to the constraint obtained from the  $v_a$  field equation as well as constrain  $\phi$  to be static as obtained from the  $w$  field equation.

Indeed the field equations are obtained directly from the  $AdS - C_D$  invariant action. The  $w$ -equation of motion is obtained as

$$\begin{aligned} 0 &= \frac{\delta}{\delta w(x,t)} \Gamma_{AdS-C_D} \\ &= (\det \bar{e}^m_a) \cosh^p \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} \left[ 2 \left( \frac{\partial}{\partial t} \phi \right) \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) \right]. \end{aligned} \tag{2.37}$$

The  $v_a$ -field equation yields

$$\begin{aligned} 0 &= \frac{\delta}{\delta v_a(x,t)} \Gamma_{AdS-C_D} \\ &= 2 \det \bar{E} \cosh^{p+1} \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} \left( \frac{\mathcal{D}^b \phi}{\cosh \sqrt{m^2 \phi^2}} - 2\delta^{bd} v_d \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) \right) \times \\ &\quad \times \left[ -2v_b \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) \frac{\mathcal{D}^a \phi}{\cosh \sqrt{m^2 \phi^2}} \right. \\ &\quad \left. + \left( \frac{\mathcal{D}^c \phi}{\cosh \sqrt{m^2 \phi^2}} 2v_c \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) - \tan^2 \sqrt{4v^2} + \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) P_{Tba}(v) \right. \\ &\quad \left. + \left( \frac{\mathcal{D}^c \phi}{\cosh \sqrt{m^2 \phi^2}} 2v_c \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) + 1 \right) P_{Lba}(v) \right] \end{aligned}$$

$$\begin{aligned}
 &+2 \left( \frac{\partial}{\partial t} \phi \right) (\det \bar{e}^n_b) \cosh^p \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} \left[ -\bar{e}^a_0 \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) \right. \\
 &\qquad \qquad \qquad \left. + (w - v_f \bar{e}^f_0) \frac{v_a}{v^2} \left( 1 - \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) \right]. \tag{2.38}
 \end{aligned}$$

Finally the  $\phi$ -equation of motion is obtained

$$\begin{aligned}
 0 &= \frac{\delta}{\delta \phi(x,t)} \Gamma_{AdS-CD} \\
 &= \det \bar{E} \cosh^{p+1} \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} (1+p) m^2 \phi \left( \frac{\tanh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \right) \\
 &\quad - \det \bar{E} \cosh^p \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} (2 \bar{e}^{-1a}_m \partial^m v_b) \left[ \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) P_{Tba}(v) + P_{Lba}(v) \right] \\
 &\quad - [\partial^m (\det \bar{E} \bar{e}^{-1a}_m)] \cosh^p \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} \left[ 2v_a \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) \right] \\
 &\quad - (\det \bar{e}^m_c) \cosh^p \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} 2 (\dot{w} - v_a \dot{\bar{e}}^a_0) \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) \\
 &\quad + (\det \bar{e}^m_c) \cosh^p \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} (2 \dot{v}_a \bar{e}^a_0) \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) \\
 &\quad - (\det \bar{e}^m_c) \cosh^p \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} \left( \frac{v_a \dot{v}_a}{v^2} \right) 2 (w - v_b \bar{e}^b_0) \left[ 1 - \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right) \right]. \tag{2.39}
 \end{aligned}$$

Introducing the momentum density  $\Pi(x,t)$  as

$$\Pi(x,t) \equiv \frac{\partial \det E}{\partial \dot{\phi}} = \det \bar{E} \bar{e}^{0-1} \cosh^p \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} 2 \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} (w - v_a \bar{e}^a_0), \tag{2.40}$$

and likewise defining the derivative of the Lagrangian with respect to the spatial derivatives of  $\phi$  as  $\Pi_m(x,t)$

$$\Pi_m(x,t) \equiv \frac{\partial \det E}{\partial \partial^m \phi} = \det \bar{E} \cosh^p \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} 2v_a \bar{e}^{-1a}_m, \tag{2.41}$$

the  $\phi$ -field equation is expressed as

$$\begin{aligned}
 0 &= \frac{\delta \Gamma_{AdS-CD}}{\delta \phi(x,t)} \\
 &= -\dot{\Pi} - \partial^m \Pi_m + \det \bar{E} \cosh^p \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} \left[ \frac{\sinh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} m^2 \phi \right] \times \\
 &\quad \times \left\{ p + 1 + p \left[ \frac{\mathcal{D}^a \phi}{\cosh \sqrt{m^2 \phi^2}} 2v_a \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} + \frac{\bar{e}^{0-1} \phi}{\cosh \sqrt{m^2 \phi^2}} 2(w - v_a \bar{e}^a_0) \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right] \right\}. \tag{2.42}
 \end{aligned}$$

Applying the  $w$ -field equation implies that  $\dot{\phi} = 0$ . As expected there is no causal connection so the field has a static spatial distribution. Applying this to the  $v_m$ -field equation yields the ‘‘inverse Higgs mechanism’’<sup>13</sup> for the (spatial) components of the  $SO(p)$  vector field  $v_a$

$$\frac{\mathcal{D}^a \phi}{\cosh \sqrt{m^2 \phi^2}} = 2 \delta^{ab} v_b \left( \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right). \tag{2.43}$$

Since the  $\phi$  covariant derivatives have the same form, Equation (2.22), the static nature of the spatial distribution of the  $\phi$  field and the inverse Higgs constraint for the spatial vector field  $v_a$

could equivalently be covariantly imposed on the fields by  $\omega_Z = 0$ . Finally applying the first two field equations to the  $\phi$ -field equation the momentum density time dependence is obtained

$$\begin{aligned} \dot{\Pi} + \partial^m \Pi_m &= \frac{\det \bar{E} \cosh^{(p+1)} \sqrt{m^2 \phi^2}}{\sqrt{\cosh^2 \sqrt{m^2 \phi^2} + \mathcal{D}^a \phi \mathcal{D}^a \phi}} \left[ m^2 \phi \frac{\sinh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \right] \times \\ &\quad \times \left\{ p + 1 + p \left[ \frac{\mathcal{D}^b \phi \mathcal{D}^b \phi}{\cosh^2 \sqrt{m^2 \phi^2}} \right] \right\}, \end{aligned} \quad (2.44)$$

with the spatial momentum  $\Pi_m$  becoming

$$\Pi_m = \frac{\det \bar{E} \cosh^{(p+1)} \sqrt{m^2 \phi^2}}{\sqrt{\cosh^2 \sqrt{m^2 \phi^2} + \mathcal{D}^b \phi \mathcal{D}^b \phi}} \bar{e}^{-1a}_m \mathcal{D}^a \phi. \quad (2.45)$$

The Noether current for the broken translation symmetry with parameter  $\zeta$  is more complex in the background  $AdS - C_d$  case and no longer simply produces the  $\phi$  field equation as in the Carroll space case but involves time variation of a related composite operator. Noether's theorem has the same form as Equation (1.29) (with  $\mathcal{L}_{AdS-C_D}$  replacing  $\mathcal{L}_C$ ) with the broken space translation variations given in Appendix A. In addition to these variations, the induced general coordinate transformation matrix  $G = \partial(t', x') / \partial(t, x)$  is obtained and hence the  $\det G^{-1}$

$$\det G^{-1} = 1 + \left( 1 + p + t \frac{\partial}{\partial t} + x_m \partial^m \right) \left[ m^2 \zeta \phi \frac{\tanh \sqrt{m^2 \phi^2} \sinh \sqrt{m^2 x^2}}{\sqrt{m^2 \phi^2} \sqrt{m^2 x^2}} \right]. \quad (2.46)$$

This yields the relation  $\Delta \mathcal{L}_{AdS-C} + \mathcal{L}_{AdS-C}(\frac{\partial}{\partial t} \delta t + \partial^m \delta x_m) = 0$  and Noether's theorem is obtained

$$0 = \dot{D} + \partial^m D_m, \quad (2.47)$$

where

$$\begin{aligned} D &= \Pi \delta \phi + \delta t \mathcal{L}_{AdS-C_D} \\ &= \Pi \cosh \sqrt{m^2 x^2} + m^2 \phi \frac{\tanh \sqrt{m^2 \phi^2} \sinh \sqrt{m^2 x^2}}{\sqrt{m^2 \phi^2} \sqrt{m^2 x^2}} \left\{ \Pi x_m \partial^m \phi - t \Pi_m \partial^m \phi \right. \\ &\quad \left. + t \det \bar{E} \cosh^{(p+1)} \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} \right\}, \\ D_m &= \Pi_m \delta \phi + \delta x_m \mathcal{L}_{AdS-C_D} \\ &= \Pi_m \cosh \sqrt{m^2 x^2} + m^2 \phi \frac{\tanh \sqrt{m^2 \phi^2} \sinh \sqrt{m^2 x^2}}{\sqrt{m^2 \phi^2} \sqrt{m^2 x^2}} \left\{ t \Pi_m \phi - x_m \Pi \phi \right. \\ &\quad \left. + (\Pi_m x_n - \Pi_n x_m) \partial^n \phi - x_m \det \bar{E} \cosh^{(p+1)} \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} \right\}. \end{aligned} \quad (2.48)$$

Applying the stationary constraint,  $\dot{\phi} = 0$ , the broken translation symmetry currents become

$$\begin{aligned} D &= \Pi \left[ \cosh \sqrt{m^2 x^2} + m^2 \phi \frac{\tanh \sqrt{m^2 \phi^2} \sinh \sqrt{m^2 x^2}}{\sqrt{m^2 \phi^2} \sqrt{m^2 x^2}} x_m \partial^m \phi \right] \\ &\quad + t m^2 \phi \frac{\tanh \sqrt{m^2 \phi^2} \sinh \sqrt{m^2 x^2}}{\sqrt{m^2 \phi^2} \sqrt{m^2 x^2}} \times \\ &\quad \times \left( \det \bar{E} \cosh^{(p+1)} \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} - \Pi_m \partial^m \phi \right), \\ D_m &= \Pi_m \left[ \cosh \sqrt{m^2 x^2} + m^2 \phi \frac{\tanh \sqrt{m^2 \phi^2} \sinh \sqrt{m^2 x^2}}{\sqrt{m^2 \phi^2} \sqrt{m^2 x^2}} x_n \partial^n \phi \right] \end{aligned}$$



$$\begin{aligned}
& -x_m m^2 \phi \frac{\tanh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \frac{\sinh \sqrt{m^2 x^2}}{\sqrt{m^2 x^2}} \times \\
& \times \left( \det \bar{E} \cosh^{(p+1)} \sqrt{m^2 \phi^2} \cos \sqrt{4v^2} - \Pi_n \partial^n \phi \right). \quad (2.49)
\end{aligned}$$

The non-linear broken translation symmetry of  $AdS - C_D$  space no longer results in the  $\phi$  field equation directly but involves a composite operator current. In the Carroll spacetime limit  $m^2 \rightarrow 0$ , as applied to the above currents,  $D \rightarrow \Pi$  and  $D_m \rightarrow \Pi_m$  and the broken translation current conservation equation is just the  $\phi$  field equation as the broken translation transformation is a simple  $\phi$  field shift symmetry (1.19).

Similarly combining the  $L$  and  $K^a$  one-forms in matrix notation as was done for the  $H$  and  $P^a$  one-forms, Equations (2.19) and (2.20), the covariant derivatives are given as  $\nabla^M V_A$  with  $V_A = (w, v_a)$  and

$$\tilde{\omega}_{KA} = (\omega_L \quad \omega_{Ka}) = dX_M \nabla^M V_A = \begin{pmatrix} dt & dx_m \end{pmatrix} \begin{pmatrix} \nabla^t w & \nabla^t v_a \\ \nabla^m w & \nabla^m v_a \end{pmatrix}. \quad (2.50)$$

Noting that under the non-linear transformations (2.30) of the coordinates and the transformations of the covariant derivative one-forms with Equation (2.32) so that

$$\tilde{\omega}'_{KA} = (\omega'_L \quad \omega'_{Ka}) = (\omega_L \quad \omega_{Kb}) \begin{pmatrix} 1 & 0 \\ -\theta_b & R_{ba}^{-1} \end{pmatrix} = \tilde{\omega}_{KB} \Lambda^B_A \quad (2.51)$$

implying that

$$(\nabla^M V_A)' = G^{-1M}_N (\nabla^N V_B) \Lambda^B_A. \quad (2.52)$$

Combining the vielbeine into the matrix vielbein  $E$  as in Equation (2.19), its variation involves  $G$  and  $\Lambda$  as shown in Equation (2.31). Thus calculating the trace of  $\nabla V$  using  $E^{-1}$

$$(E^{-1A}_M \nabla^M V_B)' = \Lambda^{-1A}_D (E^{-1D}_M \nabla^M V_C) \Lambda^C_B, \quad (2.53)$$

an invariant is obtained

$$(\text{Tr}[E^{-1} \nabla V])' = \text{Tr}[E^{-1} \nabla V]. \quad (2.54)$$

Applying the  $w$  and  $v_a$  field equation constraints,  $\dot{\phi} = 0 = \dot{v}_a$  and implicitly the Equation (2.43), the remaining  $\phi$  field equation (2.39) can be shown to be given by the trace of the covariant derivative of  $w$  and  $v_a$ . The inverse vielbein with the  $\dot{\phi} = 0$  constraint explicit so that  $e^0_0 = 0$  is found to be

$$E^{-1A}_M = \begin{pmatrix} e^{0-1}_0 & 0 \\ -e^{-1a}_n e^n_0 e^{0-1}_0 & e^{-1a}_m \end{pmatrix}, \quad (2.55)$$

where  $e^{-10}_0 = e^{0-1}_0$  with  $e^0_0 = \bar{e}^0_0 \cosh \sqrt{m^2 \phi^2}$  and the inverse submatrix  $e^{-1a}_m$  is found to be

$$e^{-1a}_m = \frac{1}{\cosh \sqrt{m^2 \phi^2}} \left[ P_{Tab}(v) + \cos \sqrt{4v^2} P_{Lab}(v) \right] \bar{e}^{-1b}_m, \quad (2.56)$$

both with the field constraints having been used. Multiplying these matrices together yields the vector field covariant derivative trace

$$\text{Tr}[E^{-1} \nabla V] = e^{0-1}_0 \nabla^t w + e^{-1a}_m \nabla^m v_a, \quad (2.57)$$

where again the constraints have been used, particularly  $\dot{v}_a = 0$ . Applying the covariant derivatives which are recalled in Equations (2.24) with again the constraints applied finally yields the trace

$$\begin{aligned}
\text{Tr}[E^{-1} \nabla V] = & -\frac{1}{2} m^2 \phi (1+p) \cos \sqrt{4v^2} \frac{\tanh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \\
& + \frac{1}{\cosh \sqrt{m^2 \phi^2}} \left( \bar{e}^{0-1}_0 (\dot{w} - p \bar{\omega}_{0a}^t v_a) \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} \right. \\
& \left. + \bar{e}^{-1b}_n \partial^n v_a \left[ \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} P_{Tab}(v) + \cos \sqrt{4v^2} P_{Lab}(v) \right] \right). \quad (2.58)
\end{aligned}$$

Using the background one-forms, Equation (2.12), and background spin connections, Equation (2.13), it is found that

$$\partial^m [\det \bar{e}_0^0 \bar{e}^{-1a}] = \det \bar{e} [\dot{\bar{e}}_0^a - p \bar{\omega}_{0b}^t \delta^{ba}], \quad (2.59)$$

with  $\bar{e}_0^a = \bar{e}^{-1a}_r \bar{e}^r_0$  so that  $\dot{\bar{e}}_0^a = \bar{e}^{-1a}_r \dot{\bar{e}}^r_0$ . Applying this and the field constraints to the  $\phi$  field Equation (2.39) and exploiting the background spin connections  $\bar{\omega}_{ab}^n$  and  $\bar{\omega}_{0a}^t$ , it is found that (2.39) (with  $\dot{\phi} = 0 = \dot{v}_a$ ) can be written as

$$0 = \frac{\delta \Gamma_{AdS-CD}}{\delta \phi} = -2 \det \bar{E} \cosh^{(1+p)} \sqrt{m^2 \phi^2} \text{Tr}[E^{-1} \nabla V]. \quad (2.60)$$

### III. THE NAMBU-GOTO CARROLLIAN VIELBEIN AND THE DUAL VECTOR ACTION

Returning to the unconstrained fields and combining the vielbeine in matrix form  $E^M_A$ , it and the Lagrangian,  $\det E$ , can be factorized into a product of the background  $AdS_d$  world volume vielbein,  $\bar{E}^M_A$ , times the Nambu-Goto Carrollian vielbein,  $N^B_A$ , as such

$$E^M_A = \bar{E}^M_B N^B_A, \quad (3.1)$$

with Equation (2.20)

$$E^M_A = \begin{pmatrix} e^0_0 & e^0_a \\ e^m_0 & e^m_a \end{pmatrix} \quad (3.2)$$

and background  $AdS_d$  vielbein

$$\bar{E}^M_A = \begin{pmatrix} \bar{e}^0_0 & \bar{e}^0_a = 0 \\ \bar{e}^m_0 & \bar{e}^m_a \end{pmatrix} \quad (3.3)$$

and correspondingly the inverse background world volume vielbein

$$\bar{E}^{-1A}_M = \begin{pmatrix} \bar{e}^{0-1}_0 & 0 \\ -\bar{e}^{-1a}_r \bar{e}^r_0 \bar{e}^{0-1}_0 & \bar{e}^{-1a}_m \end{pmatrix}. \quad (3.4)$$

This yields the Nambu-Goto Carrollian vielbein  $N^B_A = \bar{E}^{-1B}_M E^M_A$

$$N^B_A = \begin{pmatrix} N^0_0 & N^0_a \\ N^b_0 & N^b_a \end{pmatrix} \quad (3.5)$$

with

$$\begin{aligned} N^0_0 &= \bar{e}^{0-1}_0 e^0_0 \\ &= e^A \left[ 1 + 2w e^{-A} \mathcal{D}^t \phi \frac{\sin 2v}{2v} \right], \\ N^b_0 &= \bar{e}^{-1b}_n [e^n_0 - \bar{e}^n_0 \bar{e}^{0-1}_0 e^0_0] \\ &= 2w \frac{\sin 2v}{2v} [\mathcal{D}^b \phi - \bar{e}^b_0 \mathcal{D}^t \phi] + e^A w v_b \left( \frac{\cos 2v - 1}{v^2} \right), \\ N^0_a &= \bar{e}^{0-1}_0 e^0_a \\ &= 2 \mathcal{D}^t \phi v_a \frac{\sin 2v}{2v}, \\ N^b_a &= \bar{e}^{-1b}_m [e^m_a - \bar{e}^m_0 \bar{e}^{0-1}_0 e^0_a] \\ &= 2v_a \frac{\sin 2v}{2v} [\mathcal{D}^b \phi - \bar{e}^b_0 \mathcal{D}^t \phi] + e^A [P_{Tba}(v) + \cos 2v P_{Lba}(v)], \end{aligned} \quad (3.6)$$

where the notation uses the simplifying expressions

$$e^A = e^{\ln \cosh \sqrt{m^2 \phi^2}} = \cosh \sqrt{m^2 \phi^2} = \cosh m\phi,$$

$$\begin{aligned}
 \mathcal{D}^t \phi &= \bar{e}_0^{0-1} \frac{\partial}{\partial t} \phi = \bar{e}_0^{0-1} \dot{\phi}, \\
 \mathcal{D}^a \phi &= \bar{e}_m^{-1a} \frac{\partial}{\partial x_m} \phi = \bar{e}_m^{-1a} \partial^m \phi, \\
 x^2 &= x_m \delta^{mn} x_n = x_m x_m, \\
 v &= \sqrt{v^2} = \sqrt{v_a v_a},
 \end{aligned} \tag{3.7}$$

where recall the shorthand expressions such as  $v_a = \delta^{ab} v_b$  and likewise  $x_m = \delta^{mn} x_n$  are used and recall the background component vielbeine

$$\begin{aligned}
 \bar{e}_m^{-1a} \bar{e}_0^m &= \frac{x_a t}{x^2} (1 - \bar{e}_0^0) = \bar{e}_0^a, \\
 \bar{e}_m^{-1a} &= \bar{e}_0^{0-1} P_{Tam}(x) + P_{Lam}(x), \\
 \bar{e}_0^0 &= \frac{\sinh \sqrt{m^2 x^2}}{\sqrt{m^2 x^2}}.
 \end{aligned} \tag{3.8}$$

Hence the  $AdS - C_D$  action is given by

$$\Gamma_{AdS-C_D} = \int dt d^p x \det E = \int dt d^p x \det \bar{E} \det N, \tag{3.9}$$

where  $\det \bar{E} = \bar{e}_0^0 \det \bar{e}_a^m$  and

$$\det N = (\det N_a^c) [N_0^0 - N_a^0 N^{-1a} N_b^0]. \tag{3.10}$$

Letting  $N_b^a = u_a v_b + e^A (P_{Tab}(v) + \cos 2v P_{Lab}(v))$  the inverse submatrix  $N^{-1a}$  is

$$N^{-1a} = \alpha u_a v_b + e^{-A} [P_{Tab}(v) + \beta P_{Lab}(v)], \tag{3.11}$$

where

$$\begin{aligned}
 \alpha &= \frac{-e^{-A}}{[u_a v_a + e^A \cos 2v]}, \\
 \beta &= \frac{(e^A + u_b v_b)}{[u_a v_a + e^A \cos 2v]}.
 \end{aligned} \tag{3.12}$$

For the case at hand

$$u_a = 2 (\mathcal{D}^a \phi - \bar{e}_0^a \mathcal{D}^t \phi) \frac{\sin 2v}{2v}, \tag{3.13}$$

and so  $u_a v_a = 2 (\mathcal{D}^a \phi - \bar{e}_0^a \mathcal{D}^t \phi) v_a \frac{\sin 2v}{2v}$ . The submatrix determinant,  $\det N_b^a$ , is also determined to be

$$\det N_b^a = e^{pA} \cos 2v \left[ 1 + 2e^{-A} (\mathcal{D}^a \phi - \bar{e}_0^a \mathcal{D}^t \phi) v_a \frac{\tan 2v}{2v} \right]. \tag{3.14}$$

Putting all these expressions together yields the Nambu-Goto Carrollian determinant

$$\det N = \cosh^p m \phi \left[ \cosh m \phi \cos 2v + 2 \left( \mathcal{D}^a \phi v_a \frac{\sin 2v}{2v} \right) + 2 (w - v_a \bar{e}_m^{-1a} \bar{e}_0^m) \mathcal{D}^t \phi \frac{\sin 2v}{2v} \right]. \tag{3.15}$$

Thus with  $\det \bar{E} = \bar{e}_0^0 \det \bar{e}_a^m = (\bar{e}_0^0)^p$ , the  $\det E = \det \bar{E} \det N$  Lagrangian is in agreement with the Lagrangian obtained in Equation (2.35). The invariant action is obtained

$$\begin{aligned}
 \Gamma_{AdS-C_D} &= \int dt d^p x \det E = \int dt d^p x \det \bar{E} \det N \\
 &= \int dt d^p x \bar{e}_0^0 (\det \bar{e}) e^{pA} \left[ e^A \cos 2v + 2 \left( \mathcal{D}^a \phi v_a \frac{\sin 2v}{2v} \right) \right. \\
 &\quad \left. + 2 (w - v_a \bar{e}_m^{-1a} \bar{e}_0^m) \mathcal{D}^t \phi \frac{\sin 2v}{2v} \right].
 \end{aligned} \tag{3.16}$$

As in the *AdS* case<sup>6</sup> a vector field dual formulation of the *AdS* –  $C_D$  action can be obtained by introducing a vector field  $F_M$ , with  $M = 0, 1, 2, \dots, p$

$$F_M \equiv 2 \det \bar{E} \frac{\sin 2v}{2v} V_A \bar{E}^{-1A}_M, \quad (3.17)$$

with space-time component fields

$$V_A = (w, v_a) \quad (3.18)$$

and

$$F_M = (f, f_m). \quad (3.19)$$

It is useful to introduce the singular tangent space metric

$$h^{AB} = \begin{pmatrix} 0 & 0 \\ 0 & \delta^{ab} \end{pmatrix} \quad (3.20)$$

as well as the singular background *AdS* –  $C_d$  metric

$$\begin{aligned} \bar{G}^{MN} &\equiv \bar{E}^M_A h^{AB} \bar{E}^N_B \\ &= \begin{pmatrix} 0 & 0 \\ 0 & \bar{E}^m_a \delta^{ab} \bar{E}^n_b \end{pmatrix} \\ &\equiv \begin{pmatrix} 0 & 0 \\ 0 & \bar{g}^{mn} \end{pmatrix}. \end{aligned} \quad (3.21)$$

Note that  $\det \bar{G} = 0$  and hence has no inverse. Using the singular metric it is found that

$$F_M \bar{G}^{MN} F_N = f_m \bar{g}^{mn} f_n. \quad (3.22)$$

In terms of component fields it is obtained that

$$F_M = \begin{pmatrix} f \\ f_m \end{pmatrix} = 2 \det \bar{E} \frac{\sin 2v}{2v} \begin{pmatrix} (w - v_a \bar{e}^{-1a}_r \bar{e}^r_0) \bar{e}^{0-1}_0 \\ v_a \bar{e}^{-1a}_m \end{pmatrix}, \quad (3.23)$$

Inverting Equation (3.17)

$$V_A = \left( \frac{1}{2 \det \bar{E} \frac{\sin 2v}{2v}} \right) F_M \bar{E}^M_A, \quad (3.24)$$

so that the component fields are related by

$$V_A = \begin{pmatrix} w \\ v_a \end{pmatrix} = \frac{1}{2 \det \bar{E} \frac{\sin 2v}{2v}} \begin{pmatrix} f \bar{e}^0_0 + f_m \bar{e}^m_0 \\ f_m \bar{e}^m_a \end{pmatrix}. \quad (3.25)$$

Hence the useful expressions are found

$$\begin{aligned} v^2 &= v_a \delta^{ab} v_b = V_A h^{AB} V_B = \frac{1}{(2 \det \bar{E} \frac{\sin 2v}{2v})^2} F_M \bar{G}^{MN} F_N \\ &= \frac{1}{(2 \det \bar{E} \frac{\sin 2v}{2v})^2} f_m \bar{g}^{mn} f_n, \end{aligned} \quad (3.26)$$

while

$$\det \bar{E} \cos 2v = \sqrt{(\det \bar{E})^2 - F_M \bar{G}^{MN} F_N} \quad (3.27)$$

and

$$\begin{aligned} V_A \bar{E}^{-1A}_M \partial^M \phi &= (w - v_a \bar{e}^{-1a}_m \bar{e}^m_0) \mathcal{D}^t \phi + v_a \mathcal{D}^a \phi \\ &\equiv V_A \hat{\mathcal{D}}^A \phi, \end{aligned} \quad (3.28)$$

where  $\hat{\mathcal{D}}^A \phi = \bar{E}^{-1A} \partial^M \phi$  with  $\partial^M = (\partial^t, \partial^m)$  and the partial covariant derivatives  $\mathcal{D}^t \phi = \bar{e}^{0-1} \dot{\phi}$  and  $\mathcal{D}^a \phi = \bar{e}^{-1a} \partial^m \phi$ .

The determinant of the Nambu-Goto Carrollian vielbein, Equation (3.15), can be written in terms of  $\phi$  and  $F_M$  as

$$\det \bar{E} \det N = e^{pA} \left( e^A \sqrt{(\det \bar{E})^2 - F_M \bar{G}^{MN} F_N} + F_M \partial^M \phi \right). \tag{3.29}$$

Following Ref. 6, introduce the function  $h(\phi)$  so that

$$\partial^M \phi e^{pA(\phi)} \equiv \partial^M h(\phi) = \frac{dh}{d\phi} \partial^M \phi \tag{3.30}$$

and hence  $\frac{dh(\phi)}{d\phi} = e^{pA(\phi)}$ , the  $AdS - C_D$  action now has the form, after integration by parts,

$$\Gamma_{AdS-C_D} = \int dt d^p x \left[ e^{(p+1)A(\phi)} \sqrt{(\det \bar{E})^2 - F_M \bar{G}^{MN} F_N} - h(\phi) \partial^M F_M \right]. \tag{3.31}$$

The scalar field  $\phi$  equation of motion follows ( $d = (1 + p)$ )

$$\frac{\delta \Gamma_{AdS-C_D}}{\delta \phi} = 0 = \frac{dh}{d\phi} \left[ dm^2 \phi \frac{\sinh m\phi}{m\phi} \sqrt{(\det \bar{E})^2 - F_M \bar{G}^{MN} F_N} - \partial^M F_M \right]. \tag{3.32}$$

The  $\phi$  equation of motion can be enforced by introducing the Lagrange multiplier field  $L$  yielding the action

$$\Gamma_{AdS-C_D} = \int dt d^p x \left[ \sqrt{(\det \bar{E})^2 - F_M \bar{G}^{MN} F_N} \left( T(\phi) + L dm^2 \phi \frac{\sinh m\phi}{m\phi} \right) - L \partial^M F_M \right], \tag{3.33}$$

with

$$T(\phi) = e^{(1+p)A(\phi)} - h(\phi) dm^2 \phi \frac{\sinh m\phi}{m\phi}. \tag{3.34}$$

Thus we see that the previous  $\phi$  equation of motion, now coming from the  $L$  field equation, results in the  $L$  dependent terms cancelling and the  $\phi$  field itself being expressed in terms of  $F_M$ . Thus using this equation of motion

$$\begin{aligned} \sinh m\phi &= \frac{\partial^M F_M}{dm \sqrt{(\det \bar{E})^2 - F_R \bar{G}^{RS} F_S}} \\ &= \frac{\dot{f} + \partial^m f_m}{dm \sqrt{(\det \bar{E})^2 - f_r \bar{g}^{rs} f_s}}, \end{aligned} \tag{3.35}$$

the  $T(\phi)$  can be written in terms of  $F_M$ , so adopting the notation  $T(\phi) = T(\phi(F^M)) \rightarrow T(F)$ , the dual vector form of the action is

$$\begin{aligned} \Gamma_{AdS-C_D} &= \int dt d^p x T(F) \sqrt{(\det \bar{E})^2 - F_M \bar{G}^{MN} F_N} \\ &= \int dt d^p x T(f, f_m) \sqrt{(\det \bar{E})^2 - f_m \bar{g}^{mn} f_n} \end{aligned} \tag{3.36}$$

along with Equations (3.35) and (3.34) to determine  $\phi = \phi(F)$  and  $T(F)$ .

The equivalence runs in reverse as well, introducing a Lagrange multiplier  $L$  to enforce Equation (3.35) yields the action of Equation (3.33) where  $T(\phi)$  is given in Equation (3.34). The fields  $\phi$ ,  $F_M$ , and  $L$  are independent. The  $\phi$  equation of motion allows  $L$  to be eliminated as  $\delta \Gamma_{AdS-C_D} / \delta \phi = 0$  implies

$$L = -\frac{1}{dm^2} e^{-A} \frac{dT}{d\phi} = h(\phi). \tag{3.37}$$

Upon substitution into  $\Gamma_{AdS-C_D}$ , Equation (3.33), yields Equation (3.31). The definition of  $F_M$  in terms of  $V_A$ , Equation (3.17), can be applied along with (recalling that  $V^2 = V_A h^{AB} V_B = v^2$ )

$$\cos 2V = \cos 2v = \frac{1}{(\det \bar{E})} \sqrt{(\det \bar{E})^2 - F_M \bar{G}^{MN} F_N}. \quad (3.38)$$

Thus substituting this into Equation (3.31) and integrating by parts results in the Nambu-Goto-Carrollian action, note Equation (B58),

$$\Gamma_{AdS-C_D} = \int dt d^p x \det \bar{E} e^{pA(\phi)} \left[ e^{A(\phi)} \cos 2V + \frac{\sin 2V}{2V} 2V_A \bar{E}^{-1A} \partial^M \phi \right]. \quad (3.39)$$

This can be expanded in terms of component fields  $w$  and  $v_a$  to obtain the Nambu-Goto Carrollian action of Equation (3.16).

#### IV. CONCLUSION

A  $p$ -brane with codimension one was embedded in  $D = d + 1$ -dimensional AdS-Carroll space by means of the coset method. The vanishing speed of light,  $c \rightarrow 0$ , Wigner-Inönü contraction of the AdS space  $SO(2, D - 1)$  symmetry algebra was obtained and the  $AdS - C_D/ISO(p)$  coset element, Equation (2.8), was formed along with the unbroken background  $AdS - C_d/ISO(p)$  coset element Equation (2.9). The non-linearly realized spontaneously broken  $AdS - C_D \rightarrow AdS - C_d$  symmetry transformations were obtained in Appendix A. The invariant brane action was found using the Maurer-Cartan one-forms. Expanding the one-forms in terms of coordinate differentials, the vielbeine and background vielbeine were obtained. These component vielbeine were re-assembled as a Carroll spacetime matrix vielbein, Equations (2.14) and (2.19). The  $AdS - C_D SO(2, D - 1)$  invariant action was shown to be given by the determinant of this matrix vielbein

$$\Gamma_{AdS-C_D} = \int dt d^p x \det E \quad (4.1)$$

with  $\det E$  expressed in terms of the component fields in Equation (2.35).

The  $w$ ,  $v_a$ , and  $\phi$  field equations followed directly from the action. The  $w$  and  $v_a$  equations of motion implied the inverse Higgs mechanism constraints yielding the static nature of the spatial shape of the  $\phi$  field,  $\dot{\phi} = 0$ , as expected from the contraction of the light cone to the time axis as  $c \rightarrow 0$ , and the auxiliary vector field  $v_a$  and the spatial derivatives of  $\phi$  were related, Equation (2.43). Both of these constraints can alternatively be imposed by the invariant  $\phi$ -covariant derivative constraint  $\omega_Z = 0$ . The canonical momentum density defined in Equation (2.40) on the other hand exhibits time variation related to the shape of the brane,  $\phi(x_m)$ , as well as the  $AdS - C$  geometry. Finally the  $\phi$ -field equation can be written in terms of the covariant derivatives of the auxiliary fields  $w$  and  $v_a$  as Equation (2.60).

In the flat Minkowski space limit,  $m^2 \rightarrow 0$ , the  $AdS - C_D$  results describe the  $p$ -brane embedded in a Carroll spacetime,  $AdS - C_D \rightarrow C_D$ . These results agree with those of the more informally derived results discussed in the Introduction. In addition the broken translation symmetry Noether current in the  $AdS - C$  case, Equation (2.47), with currents Equation (2.49), goes over to the Carroll space currents  $z = \Pi$ ,  $z_m = \Pi_m$ , found in Table I, and the Carrollian component vielbeine can be obtained as the  $m^2 \rightarrow 0$  limit of Equation (2.18) and correspondingly the  $m^2 \rightarrow 0$  action  $\Gamma_C = \int dt d^p x \mathcal{L}_C$  with  $\det E \xrightarrow{c \rightarrow 0} \mathcal{L}_C$  and action  $\Gamma_C$  of Equation (1.9).

The  $AdS - C_D$  vielbein  $E$  has the product form of the  $AdS - C_d$  background vielbein  $\bar{E}$  times the Nambu-Goto-Carroll vielbein  $N$ ,  $E = \bar{E}N$ , as expressed in Equations (3.1)-(3.7). The  $p$ -brane action can be re-formulated in terms of its dual vector field  $F_M$  action, Equation (3.36), with functions  $\phi = \phi(F)$ , Equation (3.35), and  $T(F)$ , Equation (3.34). Likewise the dual action can be reformulated to yield the brane Nambu-Goto-Carrollian action Equation (3.39) with component form Equation (3.16).

An equivalent approach to obtain the  $p$ -brane action is to expose the speed of light in the already known  $AdS_{(d+1)} \rightarrow AdS_d$  brane action results and take the  $c \rightarrow 0$  Carrollian limit thereof.

This method was presented in [Appendix B](#) where the Maurer-Cartan one-forms and Nambu-Goto vielbein of Ref. [6](#) were used to obtain the  $AdS$  action with the speed of light parameter, Equation ([B58](#)). The Carrollian limit was then taken to obtain the coset method component action ([3.16](#)) results.

## ACKNOWLEDGMENTS

The work of T.t.V was supported in part by the NSF under Grant No. PHY- 1102585. T.t.V gratefully acknowledges the hospitality of E. Bergshoeff and the Van Swinderen Institute for Particle Physics and Gravity at the University of Groningen while on sabbatical leave from Macalester College. T.t.V also thanks Joaquim Gomis for useful discussions.

## APPENDIX A: $AdS - C$ TRANSFORMATIONS

Using the group multiplication laws as applied to the coset  $\Omega$ , the non-linearly realized  $AdS - C_D$  transformations are determined from

$$g\Omega(x, t) = \Omega'(x', t')h(x, t), \quad (A1)$$

where the infinitesimal  $AdS - C_D$  transformations form the group elements

$$g = e^{i\epsilon H} e^{ia_m P^m} e^{i\zeta Z} e^{i\lambda L} e^{i\kappa_m K^m} e^{\frac{i}{2}\alpha_{mn} M^{mn}} e^{i\beta_m B^m}, \quad (A2)$$

while the transformed coset element is given by

$$\Omega'(x', t') = e^{it'H + ix'_m P^m} e^{i\phi'(x', t')Z} e^{iw'(x', t')L + v'_m(x', t')K^m}. \quad (A3)$$

The  $h(x, t)$  is an element of the invariant  $ISO(p)$  subgroup

$$h(x, t) = e^{\frac{i}{2}\theta_{mn}(x, t)M^{mn}} e^{i\theta_m(x, t)B^m} \quad (A4)$$

with parameters  $\theta_{mn}$  and  $\theta_m$  that also depend on  $g$ . The transformations of the spacetime coordinates and fields are found to be non-linearly realized

$$\begin{aligned} t' &= t \left[ 1 + \frac{a_m x_m}{x^2} \left( 1 - \sqrt{m^2 x^2} \coth \sqrt{m^2 x^2} \right) \right. \\ &\quad - m^2 \zeta \phi \frac{\tanh \sqrt{m^2 \phi^2} \sinh \sqrt{m^2 x^2}}{\sqrt{m^2 \phi^2} \sqrt{m^2 x^2}} \\ &\quad \left. - \frac{\kappa_m x_m}{x^2} 2\phi \frac{\tanh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \left( \cosh \sqrt{m^2 x^2} - \frac{\sqrt{m^2 x^2}}{\sinh \sqrt{m^2 x^2}} \right) \right] \\ &\quad + \epsilon \sqrt{m^2 x^2} \coth \sqrt{m^2 x^2} - \lambda 2\phi \frac{\tanh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \frac{\sqrt{m^2 x^2}}{\sinh \sqrt{m^2 x^2}} - \beta_m x_m, \\ x'_m &= x_m \left[ 1 - m^2 \zeta \phi \frac{\tanh \sqrt{m^2 \phi^2} \sinh \sqrt{m^2 x^2}}{\sqrt{m^2 \phi^2} \sqrt{m^2 x^2}} \right] + \alpha_{mn} x_n \\ &\quad + \left( \sqrt{m^2 x^2} \coth \sqrt{m^2 x^2} P_{Tmn}(x) + P_{Lmn}(x) \right) a_n \\ &\quad - 2\phi \frac{\tanh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \left[ \frac{\sqrt{m^2 x^2}}{\sinh \sqrt{m^2 x^2}} P_{Tmn}(x) + \cosh \sqrt{m^2 x^2} P_{Lmn}(x) \right] \kappa_n, \\ \phi'(x', t') &= \phi(x, t) + \zeta \cosh \sqrt{m^2 x^2} + 2\kappa_m x_m \frac{\sinh \sqrt{m^2 x^2}}{\sqrt{m^2 x^2}}, \\ w'(x', t') &= w(x, t) + \epsilon m^2 x_m v_m \frac{\tanh \sqrt{m^2 x^2}/2}{\sqrt{m^2 x^2}} - \beta_m v_m \end{aligned}$$

$$\begin{aligned}
 & -m^2 t \frac{\tanh \sqrt{m^2 x^2} / 2}{\sqrt{m^2 x^2}} a_m v_m + \lambda \frac{\sqrt{4v^2} \cot \sqrt{4v^2}}{\cosh \sqrt{m^2 \phi^2}} \\
 & + \lambda 2 \phi \frac{v_m x_m}{x^2} \left( \sqrt{m^2 x^2} \tanh \sqrt{m^2 x^2} / 2 \right) \left( \frac{\tanh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \right) \\
 & + \frac{1}{2} m^2 \zeta \frac{1}{\cosh \sqrt{m^2 \phi^2}} \frac{\sinh \sqrt{m^2 x^2}}{\sqrt{m^2 x^2}} \left[ w \frac{x_m v_m}{v^2} + \sqrt{4v^2} \cot \sqrt{4v^2} \left( t - w \frac{x_m v_m}{v^2} \right) \right] \\
 & + w \frac{1}{\cosh \sqrt{m^2 \phi^2}} \frac{[1 - \sqrt{4v^2} \cot \sqrt{4v^2}]}{v^2} v_m [P_{Tmn}(x) + \cosh \sqrt{m^2 x^2} P_{Lmn}(x)] \kappa_n \\
 & - m^2 t \kappa_m v_m \phi \left( \frac{\tanh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \right) \left( \frac{\tanh \sqrt{m^2 x^2} / 2}{\sqrt{m^2 x^2} / 2} \right) \\
 & + t \kappa_m x_m \left( \frac{\sqrt{4v^2} \cot \sqrt{4v^2}}{\cosh \sqrt{m^2 \phi^2}} \right) \left( \frac{\cosh \sqrt{m^2 x^2} - 1}{x^2} \right), \\
 v'_m(x', t') = & v_m(x, t) + \alpha_{mn} v_n + \frac{m^2}{2} (a_m x_n - a_n x_m) \frac{2}{\sqrt{m^2 x^2}} \tanh \sqrt{m^2 x^2} / 2 v_n \\
 & + \frac{m^2}{2} \zeta \frac{1}{\cosh \sqrt{m^2 \phi^2}} \frac{\sinh \sqrt{m^2 x^2}}{\sqrt{m^2 x^2}} \left[ \sqrt{4v^2} \cot \sqrt{4v^2} P_{Tmn}(v) + P_{Lmn}(v) \right] x_n \\
 & - m^2 \phi \frac{\tanh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \frac{2}{\sqrt{m^2 x^2}} \tanh \sqrt{m^2 x^2} / 2 (x_m (\kappa_n v_n) - \kappa_m (x_n v_n)) \\
 & + \left( \sqrt{4v^2} \cot \sqrt{4v^2} P_{Tmn}(v) + P_{Lmn}(v) \right) \frac{1}{\cosh \sqrt{m^2 \phi^2}} \\
 & \left( P_{Tnr}(x) + \cosh \sqrt{m^2 x^2} P_{Lnr}(x) \right) \kappa_r. \tag{A5}
 \end{aligned}$$

The invariant  $ISO(p)$  subgroup parameters  $\theta_{mn}$  and  $\theta_m$  are also obtained

$$\begin{aligned}
 \theta_{mn} = & \alpha_{mn} + \frac{m^2}{2} (a_m x_n - a_n x_m) \frac{2}{\sqrt{m^2 x^2}} \tanh \sqrt{m^2 x^2} / 2 \\
 & - m^2 \zeta \frac{1}{\cosh \sqrt{m^2 \phi^2}} \frac{\sinh \sqrt{m^2 x^2}}{\sqrt{m^2 x^2}} (x_m v_n - x_n v_m) \frac{\tan \sqrt{v^2}}{\sqrt{v^2}} \\
 & - m^2 \phi \frac{\tanh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \frac{2}{\sqrt{m^2 x^2}} \tanh \sqrt{m^2 x^2} / 2 (x_m \kappa_n - x_n \kappa_m) \\
 & - 2 \frac{\tan \sqrt{v^2}}{\sqrt{v^2}} \frac{1}{\cosh \sqrt{m^2 \phi^2}} \left[ \left( P_{Tmr}(x) + \cosh \sqrt{m^2 x^2} P_{Lmr}(x) \right) \kappa_r v_n \right. \\
 & \quad \left. - \left( P_{Tnr}(x) + \cosh \sqrt{m^2 x^2} P_{Lnr}(x) \right) \kappa_r v_m \right], \\
 \theta_m = & \beta_m - \epsilon m^2 x_m \frac{\tanh \sqrt{m^2 x^2} / 2}{\sqrt{m^2 x^2}} + m^2 t a_m \frac{\tanh \sqrt{m^2 x^2} / 2}{\sqrt{m^2 x^2}} \\
 & + m^2 \zeta \frac{\tan \sqrt{v^2}}{\sqrt{v^2}} \frac{1}{\cosh \sqrt{m^2 \phi^2}} \frac{\sinh \sqrt{m^2 x^2}}{\sqrt{m^2 x^2}} (t v_m - x_m w) \\
 & - \lambda m^2 x_m \phi \frac{\tanh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \frac{2}{\sqrt{m^2 x^2}} \tanh \sqrt{m^2 x^2} / 2 + 2 \lambda v_m \frac{\tan \sqrt{v^2}}{\sqrt{v^2}} \frac{1}{\cosh \sqrt{m^2 \phi^2}} \\
 & + m^2 t \phi \frac{\tanh \sqrt{m^2 \phi^2}}{\sqrt{m^2 \phi^2}} \frac{2}{\sqrt{m^2 x^2}} \tanh \sqrt{m^2 x^2} / 2 \kappa_m
 \end{aligned}$$



$$\begin{aligned}
& +2tv_m(\kappa_n x_n) \frac{\tan \sqrt{v^2}}{\sqrt{v^2}} \frac{1}{\cosh \sqrt{m^2 \phi^2}} \frac{\cosh \sqrt{m^2 x^2} - 1}{x^2} \\
& -2w \frac{\tan \sqrt{v^2}}{\sqrt{v^2}} \frac{1}{\cosh \sqrt{m^2 \phi^2}} \left( P_{Tmn}(x) + \cosh \sqrt{m^2 x^2} P_{Lmn}(x) \right) \kappa_n.
\end{aligned} \tag{A6}$$

The symmetry transformations for  $D = d + 1$  Carrollian spacetime as given in Equations (1.19) and (1.21) and as well as the induced local rotations and boosts

$$\Lambda = \begin{pmatrix} 1 & 0 \\ -\theta_n & R_{nm}^{-1} \end{pmatrix} \tag{A7}$$

with  $R_{nm}^{-1} = \delta_{nm} - \theta_{nm}$ , where the induced infinitesimal rotation has parameter  $\theta_{mn}$ ,

$$\theta_{nm} = \alpha_{nm} - 2 \frac{\tan \sqrt{v^2}}{\sqrt{v^2}} (\kappa_n v_m - \kappa_m v_n), \tag{A8}$$

while the unbroken induced boosts have parameter  $\theta_n$ ,

$$\theta_n = \beta_n + 2 (\lambda v_n - w \kappa_n) \frac{\tan \sqrt{v^2}}{\sqrt{v^2}} \tag{A9}$$

are obtained as the  $m^2 \rightarrow 0$  limit of these  $AdS - C_D \rightarrow AdS - C_d$  transformations.

## APPENDIX B: $AdS \xrightarrow{c \rightarrow 0} AdS - C$

The purpose of this appendix is to make the speed of light  $c$  explicit in the  $AdS_{d+1} \rightarrow AdS_d$  isometry algebra and associated coset elements in order to implement the  $c \rightarrow 0$  limit directly, reproducing the action of Sections II and III. Returning to the  $SO(2, d)$  symmetry algebra for  $AdS_{d+1}$ , Equation (2.1), where now the  $SO(2, d) \rightarrow SO(2, d - 1)$  isometry algebra for  $AdS_{d+1} \rightarrow AdS_d$  is denoted with hatted operators so that

$$\begin{aligned}
P^M &= \hat{P}^M \text{ for } M = 0, 1, 2, \dots, p, \\
P^{p+1} &= -\hat{Z}
\end{aligned} \tag{B1}$$

and

$$\begin{aligned}
M^{MN} &= \hat{M}^{MN} \text{ for } M, N = 0, 1, 2, \dots, p, \\
M^{p+1M} &= \hat{K}^M \text{ for } M = 0, 1, 2, \dots, p,
\end{aligned} \tag{B2}$$

where now  $M, N = 0, 1, 2, \dots, p$  labelling only the  $AdS_d$  components while the  $(p + 1)^{th}$  components are separated into  $\hat{Z}$  and  $\hat{K}^M$ . The  $SO(2, d)$  algebra becomes that used in Equation (B.5) of Ref. 6

$$\begin{aligned}
[\hat{M}^{MN}, \hat{M}^{RS}] &= -i (\eta^{MR} \hat{M}^{NS} - \eta^{MS} \hat{M}^{NR} + \eta^{NS} \hat{M}^{MR} - \eta^{NR} \hat{M}^{MS}), \\
[\hat{M}^{MN}, \hat{P}^L] &= i (\hat{P}^M \eta^{NL} - \hat{P}^N \eta^{ML}), \\
[\hat{M}^{MN}, \hat{K}^L] &= i (\hat{K}^M \eta^{NL} - \hat{K}^N \eta^{ML}), \\
[\hat{M}^{MN}, \hat{Z}] &= 0, & [\hat{P}^M, \hat{K}^N] &= i \eta^{MN} \hat{Z}, \\
[\hat{P}^M, \hat{P}^N] &= -im^2 \hat{M}^{MN}, & [\hat{P}^M, \hat{Z}] &= -im^2 \hat{K}^M, \\
[\hat{K}^M, \hat{K}^N] &= i \hat{M}^{MN}, & [\hat{Z}, \hat{K}^M] &= i \hat{P}^M.
\end{aligned} \tag{B3}$$

To make the speed of light explicit introduce the generators

$$\begin{aligned}
\hat{P}^0 &= \frac{1}{c} H, \\
\hat{P}^m &= P^m, \\
\hat{K}^0 &= \frac{1}{2c} L,
\end{aligned}$$

$$\begin{aligned}
\hat{K}^m &= \frac{1}{2}K^m, \\
\hat{Z} &= -Z, \\
\hat{M}^{m0} &= \frac{1}{c}B^m, \\
\hat{M}^{mn} &= M^{mn},
\end{aligned} \tag{B4}$$

where the spatial indices are labelled by  $m, n = 1, 2, \dots, p = (d - 1)$ . Hence the  $SO(2, d)$  algebra of Equation (B3) is as given in Equations (2.5)-(2.7) except for the four commutators involving the explicit factor of the speed of light, which are now

$$\begin{aligned}
[B^m, B^n] &= -ic^2 M^{mn}, & [B^m, L] &= ic^2 K^m, \\
[B^m, H] &= ic^2 P^m, & [H, L] &= -2ic^2 Z.
\end{aligned} \tag{B5}$$

Define the operators  $\mathcal{P}^M$ ,  $\mathcal{M}^{MN}$ ,  $\mathcal{Z}$ , and  $\mathcal{K}^M$  with the explicit speed of light factors removed as

$$\begin{aligned}
\mathcal{P}^M &= \begin{pmatrix} H \\ P^m \end{pmatrix}, \\
\mathcal{K}^M &= \begin{pmatrix} L \\ K^m \end{pmatrix}, \\
\mathcal{Z} &= Z, \\
\mathcal{M}^{MN} &= \begin{pmatrix} 0 & -B^n \\ B^m & M^{mn} \end{pmatrix}.
\end{aligned} \tag{B6}$$

The relation to the hatted operators is given succinctly by

$$\begin{aligned}
\mathcal{P}^M &= C_N^M \hat{P}^M, \\
\mathcal{K}^M &= 2C_N^M \hat{K}^M, \\
\mathcal{Z} &= -\hat{Z}, \\
\mathcal{M}^{MN} &= C_R^M \hat{M}^{RS} C_S^N,
\end{aligned} \tag{B7}$$

with

$$C_N^M = \begin{pmatrix} c & 0 \\ 0 & \delta_{mn} \end{pmatrix}. \tag{B8}$$

In terms of these operators, the  $SO(2, d)$  algebra of Equation (B3) becomes

$$\begin{aligned}
[\mathcal{M}^{MN}, \mathcal{M}^{RS}] &= -i(n^{MR}\mathcal{M}^{NS} - n^{MS}\mathcal{M}^{NR} + n^{NS}\mathcal{M}^{MR} - n^{NR}\mathcal{M}^{MS}), \\
[\mathcal{M}^{MN}, \mathcal{P}^L] &= i(\mathcal{P}^M n^{NL} - \mathcal{P}^N n^{ML}), \\
[\mathcal{M}^{MN}, \mathcal{K}^L] &= i(\mathcal{K}^M n^{NL} - \mathcal{K}^N n^{ML}), \\
[\mathcal{M}^{MN}, \mathcal{Z}] &= 0, & [\mathcal{P}^M, \mathcal{K}^N] &= -2in^{MN}\mathcal{Z}, \\
[\mathcal{P}^M, \mathcal{P}^N] &= -im^2\mathcal{M}^{MN}, & [\mathcal{P}^M, \mathcal{Z}] &= +\frac{i}{2}m^2\mathcal{K}^M, \\
[\mathcal{K}^M, \mathcal{K}^N] &= 4i\mathcal{M}^{MN}, & [\mathcal{Z}, \mathcal{K}^M] &= -2i\mathcal{P}^M,
\end{aligned} \tag{B9}$$

where the metric has the form of a  $(p + 1) \times (p + 1)$  diagonal matrix denoted  $n^{MN}$

$$n^{MN} \equiv C_R^M \eta^{RS} C_S^N = \begin{pmatrix} c^2 & 0 \\ 0 & -\delta_{mn} \end{pmatrix}. \tag{B10}$$

Rather than use the coset method directly with this form of the algebra, the Maurer-Cartan one-forms found using the hatted form of the algebra can be converted to one-forms with the explicit powers of  $c$  exhibited and then the  $c \rightarrow 0$  limit performed. First the coset elements for the

two sets of operators are identified. Consider the coordinates

$$\hat{x}_M \equiv (x_0, x_m) = (ct, x_m) \tag{B11}$$

and

$$X_M \equiv (t, x_m) \tag{B12}$$

that is  $X_M = \hat{x}_N C^{-1N}_M$ . Hence the coset elements

$$e^{i\hat{x}_M \hat{P}^M} = e^{iX_M \mathcal{P}^M}. \tag{B13}$$

Likewise let  $\hat{\phi}(\hat{x})\hat{Z} = \phi(X)\mathcal{Z}$  so that

$$e^{i\hat{\phi}\hat{Z}} = e^{i\phi\mathcal{Z}}, \tag{B14}$$

and  $\phi = -\hat{\phi}$ . Also define the components of  $\hat{v}_M$  as

$$\hat{v}_M = (\hat{v}_0, \hat{v}_m) = (2cw, 2v_m) \tag{B15}$$

and those of  $V_M$  as

$$V_M = (w, v_m). \tag{B16}$$

Thus  $V_M = \frac{1}{2}\hat{v}_N C^{-1N}_M$  so that

$$e^{i\hat{v}_M \hat{K}^M} = e^{iV_M \mathcal{K}^M}. \tag{B17}$$

Finally equating the unbroken subgroup operators  $\hat{\theta}_{MN}\hat{M}^{MN} = \Theta_{MN}\mathcal{M}^{MN}$  with

$$\Theta_{RS} = C^{-1M}_R \hat{\theta}_{MN} C^{-1N}_S \tag{B18}$$

so that

$$\Theta_{RS} = \begin{pmatrix} 0 & -\theta_s \\ \theta_r & \theta_{rs} \end{pmatrix} \tag{B19}$$

while

$$\hat{\theta}_{MN} = \begin{pmatrix} 0 & -\hat{\theta}_n \\ \hat{\theta}_m & \hat{\theta}_{mn} \end{pmatrix} = \begin{pmatrix} 0 & -c\theta_n \\ c\theta_m & \theta_{mn} \end{pmatrix} \tag{B20}$$

and the subgroup elements are equal

$$e^{\frac{i}{2}\hat{\theta}_{MN}\hat{M}^{MN}} = e^{\frac{i}{2}\Theta_{MN}\mathcal{M}^{MN}}. \tag{B21}$$

These coset elements so identified,

$$\hat{\Omega} = e^{i\hat{x}_M \hat{P}^M} e^{i\hat{\phi}\hat{Z}} e^{i\hat{v}_M \hat{K}^M} = \Omega = e^{iX_M \mathcal{P}^M} e^{i\phi\mathcal{Z}} e^{iV_M \mathcal{K}^M}, \tag{B22}$$

allow their respective Maurer-Cartan one-forms to be related, recalling the one-forms  $\omega = -i\Omega^{-1}d\Omega$  and  $\hat{\omega} = -i\hat{\Omega}^{-1}d\hat{\Omega}$  with  $\hat{d} = d\hat{x}_M \frac{\partial}{\partial \hat{x}_M} = dt \frac{\partial}{\partial t} + dx_m \frac{\partial}{\partial x_m} = dX_M \frac{\partial}{\partial X_M} = d$ , the one-forms are equal  $\omega = \hat{\omega}$ . Expanding them in terms of the generators with tangent space indices  $A, B = 0, 1, \dots, p$  (recall world indices  $M, N = 0, 1, \dots, p$  also)

$$\omega = \omega_A \mathcal{P}^A + \omega_Z \mathcal{Z} + \omega_{KA} \mathcal{K}^A + \frac{1}{2}\omega_{AB} \mathcal{M}^{AB} \tag{B23}$$

and

$$\hat{\omega} = \hat{\omega}_A \hat{P}^A + \hat{\omega}_Z \hat{Z} + \hat{\omega}_{KA} \hat{K}^A + \frac{1}{2}\hat{\omega}_{AB} \hat{M}^{AB}, \tag{B24}$$

and utilizing Equation (B7) the Maurer-Cartan one-forms are related

$$\begin{aligned} \omega_A &= \hat{\omega}_B C^{-1B}_A, \\ \omega_Z &= -\hat{\omega}_Z, \end{aligned}$$

$$\begin{aligned}\omega_{KA} &= \frac{1}{2}\hat{\omega}_{KB}C^{-1B}_A, \\ \omega_{AB} &= \hat{\omega}_{CD}C^{-1C}_A C^{-1D}_B.\end{aligned}\quad (\text{B25})$$

These yield the relations for the component one-forms and the eventual  $c \rightarrow 0$  relation to the one-forms of Section II. The explicit factors of  $c$  relating the component one-forms are, with  $a, b = 1, 2, \dots, p$ ,

$$\begin{aligned}\omega_0 &= \frac{1}{c}\hat{\omega}_0, \\ \omega_a &= \hat{\omega}_a, \\ \omega_Z &= -\hat{\omega}_Z, \\ \omega_{K0} &= \frac{1}{2c}\hat{\omega}_{K0}, \\ \omega_{Ka} &= \frac{1}{2}\hat{\omega}_{Ka}, \\ \omega_{a0} &= \frac{1}{c}\hat{\omega}_{a0}, \\ \omega_{ab} &= \hat{\omega}_{ab}.\end{aligned}\quad (\text{B26})$$

The relation to the  $AdS - C$  one-forms of Section II is found in the  $c \rightarrow 0$  limit of the above, for example,  $\omega_0 = \frac{1}{c}\hat{\omega}_0 \xrightarrow{c \rightarrow 0} \omega_H$ .

Similarly for the background one-forms for which  $\hat{\Omega} = e^{i\hat{x}_A \hat{P}^A} = \tilde{\Omega} = e^{iX_A \mathcal{P}^A}$  and so  $\bar{\omega} = \hat{\omega}$ . Expanding in terms of the generators

$$\bar{\omega} = \bar{\omega}_A \mathcal{P}^A + \frac{1}{2}\bar{\omega}_{AB} \mathcal{M}^{AB} \quad (\text{B27})$$

and

$$\hat{\omega} = \hat{\omega}_A \hat{P}^A + \frac{1}{2}\hat{\omega}_{AB} \hat{M}^{AB}, \quad (\text{B28})$$

and using the relations for the one-forms

$$\begin{aligned}\bar{\omega}_A &= \hat{\omega}_B C^{-1B}_A, \\ \bar{\omega}_{AB} &= \hat{\omega}_{CD} C^{-1C}_A C^{-1D}_B,\end{aligned}\quad (\text{B29})$$

these yield the component background one-form equalities

$$\begin{aligned}\bar{\omega}_0 &= \frac{1}{c}\hat{\omega}_0, \\ \bar{\omega}_a &= \hat{\omega}_a, \\ \bar{\omega}_{a0} &= \frac{1}{c}\hat{\omega}_{a0}, \\ \bar{\omega}_{ab} &= \hat{\omega}_{ab},\end{aligned}\quad (\text{B30})$$

with the  $AdS - C$  background one-forms of Section II found in the  $c \rightarrow 0$  limit, for example  $\bar{\omega}_0 = \frac{1}{c}\hat{\omega}_0 \xrightarrow{c \rightarrow 0} \bar{\omega}_H$ .

Applying these  $c$ -factor conversions to the Maurer-Cartan one-form  $\hat{\omega}_A$  found in Equation (2.10) of Ref. 6 for charges defined with upper indices, as is the convention here,

$$\hat{\omega}_A = -\frac{\sinh \sqrt{\hat{v}^2}}{\sqrt{\hat{v}^2}} \hat{v}_A \hat{d}\hat{\phi} + \cosh \sqrt{m^2 \hat{\phi}^2} \left[ P_{TAB}(\hat{v}) + \cosh \sqrt{\hat{v}^2} P_{LAB}(\hat{v}) \right] \eta^{BC} \hat{\omega}_C, \quad (\text{B31})$$

with the corresponding background one-form  $\hat{\omega}_A$  of Ref. 6

$$\hat{\omega}_C = \left[ \frac{\sin \sqrt{m^2 \hat{x}^2}}{\sqrt{m^2 \hat{x}^2}} P_{TC D}(\hat{x}) + P_{LCD}(\hat{x}) \right] \eta^{DE} d\hat{x}_E, \quad (\text{B32})$$

yields the resulting  $\omega_A$  one-form  $\omega_A = \hat{\omega}_B C^{-1B}_A$

$$\omega_A = \frac{\sinh \sqrt{4V^2}}{\sqrt{4V^2}} 2V_A d\phi + \cosh \sqrt{m^2 \phi^2} \left[ \mathcal{P}_{TA}^B(V) + \cosh \sqrt{4V^2} \mathcal{P}_{LA}^B(V) \right] \bar{\omega}_B, \quad (\text{B33})$$

with the projection operators

$$\begin{aligned} \mathcal{P}_{LA}^B(V) &= P_{LAC}(V) n^{CB} = \frac{V_A V_C n^{CB}}{V_D n^{DE} V_E}, \\ \mathcal{P}_{TA}^B(V) &= \delta_A^B - \mathcal{P}_{LA}^B(V), \end{aligned} \quad (\text{B34})$$

and where  $\hat{v}^2 = \hat{v}_A \eta^{AB} \hat{v}_B = 4V_D n^{DE} V_E \equiv 4V^2$ . In analogous fashion the background one-form is derived  $\bar{\omega}_A = \hat{\omega}_B C^{-1B}_A$

$$\bar{\omega}_A = \left[ \frac{\sin \sqrt{m^2 X^2}}{\sqrt{m^2 X^2}} \mathcal{P}_{TA}^M(X) + \mathcal{P}_{LA}^M(X) \right] dX_M, \quad (\text{B35})$$

with  $\hat{x}^2 = \hat{x}_M \eta^{MN} \hat{x}_N = X_M n^{MN} X_N \equiv X^2$ . Likewise from Equation (2.10) of Ref. 6

$$\hat{\omega}_Z = \cosh \sqrt{\hat{v}^2} \left[ \hat{d}\hat{\phi} - \cosh \sqrt{m^2 \hat{\phi}^2} \hat{\omega}_A \eta^{AB} \hat{v}_B \frac{\tanh \sqrt{\hat{v}^2}}{\sqrt{\hat{v}^2}} \right], \quad (\text{B36})$$

from which it is found that

$$\omega_Z = -\hat{\omega}_Z = \cosh \sqrt{4V^2} \left[ d\phi + \cosh \sqrt{m^2 \phi^2} \bar{\omega}_A n^{AB} 2V_B \frac{\tanh \sqrt{4V^2}}{\sqrt{4V^2}} \right]. \quad (\text{B37})$$

The vielbein  $\mathcal{E}_A^M$  is defined by relating the covariant differentials  $\omega_A$  to the coordinate differentials  $dX_M$

$$\omega_A \equiv dX_M \mathcal{E}_A^M. \quad (\text{B38})$$

Likewise

$$\hat{\omega}_B \equiv d\hat{x}_M \hat{\mathcal{E}}_B^M \quad (\text{B39})$$

and so the vielbeine are related through the one-forms  $\omega_A = \hat{\omega}_B C^{-1B}_A$  as

$$\mathcal{E}_A^M = C_N^M \hat{\mathcal{E}}_B^N C^{-1B}_A. \quad (\text{B40})$$

Similarly for the background one-forms

$$\begin{aligned} \bar{\omega}_A &= dX_M \bar{\mathcal{E}}_A^M, \\ \hat{\omega}_A &= d\hat{x}_M \hat{\mathcal{E}}_A^M, \end{aligned} \quad (\text{B41})$$

and hence the related vielbeine

$$\bar{\mathcal{E}}_A^M = C_N^M \hat{\mathcal{E}}_B^N C^{-1B}_A. \quad (\text{B42})$$

Since the one-forms  $\omega_A$  and  $\bar{\omega}_A$  are already obtained, the vielbeine can be read off from their forms. From Equation (B35) the background vielbein  $\bar{\mathcal{E}}_A^M$  is seen to be equal to

$$\bar{\mathcal{E}}_A^M = \frac{\sin \sqrt{m^2 X^2}}{\sqrt{m^2 X^2}} \mathcal{P}_{TA}^M(X) + \mathcal{P}_{LA}^M(X). \quad (\text{B43})$$

Equation (B33) with  $d\phi = dX_M \frac{\partial}{\partial X_M} \phi = dX_M \partial^M \phi$  and  $\bar{\omega}_B = dX_M \bar{\mathcal{E}}_B^M$  provides the vielbein  $\mathcal{E}_A^M$

$$\mathcal{E}_A^M = \frac{\sinh \sqrt{4V^2}}{\sqrt{4V^2}} 2V_A \frac{\partial}{\partial X_M} \phi + \cosh \sqrt{m^2 \phi^2} \left[ \mathcal{P}_{TA}^B(V) + \cosh \sqrt{4V^2} \mathcal{P}_{LA}^B(V) \right] \bar{\mathcal{E}}_B^M. \quad (\text{B44})$$

The speed of light can be taken to zero to obtain the results of Sections II and III. Displaying the component one-forms and vielbeine, it is found for the background one-forms and vielbeine

that

$$\begin{aligned}
\bar{\omega}_0 &= dt\bar{\mathcal{E}}_0^0 + dx_m\bar{\mathcal{E}}_0^m \\
&\xrightarrow{c \rightarrow 0} dt \left( \frac{\sinh \sqrt{m^2 x^2}}{\sqrt{m^2 x^2}} \right) + dx_m \left( \frac{tx_m}{x^2} \left( 1 - \frac{\sinh \sqrt{m^2 x^2}}{\sqrt{m^2 x^2}} \right) \right) \\
&= \bar{\omega}_H = dt\bar{e}_0^0 + dx_m\bar{e}_0^m, \\
\bar{\omega}_a &= dt\bar{\mathcal{E}}_a^0 + dx_m\bar{\mathcal{E}}_a^m \\
&\xrightarrow{c \rightarrow 0} dx_m \left( \frac{\sinh \sqrt{m^2 x^2}}{\sqrt{m^2 x^2}} P_{Tam}(x) + P_{Lam}(x) \right) \\
&= \bar{\omega}_{Pa} = dt\bar{e}_a^0 + dx_m\bar{e}_a^m.
\end{aligned} \tag{B45}$$

Thus the vielbeine components of Section II, Equations (2.12) and (2.14), have been obtained. In short this  $c \rightarrow 0$  limit is

$$\bar{\mathcal{E}}_A^M \xrightarrow{c \rightarrow 0} \bar{E}_A^M. \tag{B46}$$

Proceeding in a similar manner for the one-forms and vielbeine their  $c \rightarrow 0$  limits are obtained as those of Section II

$$\begin{aligned}
\omega_0 &= dt\mathcal{E}_0^0 + dx_m\mathcal{E}_0^m \\
&\xrightarrow{c \rightarrow 0} dt \left[ \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} 2w\partial^t \phi + \cosh \sqrt{m^2 \phi^2} \bar{\mathcal{E}}_0^0 \right] \\
&\quad + dx_m \left[ \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} 2w\partial^m \phi + \cosh \sqrt{m^2 \phi^2} \bar{\mathcal{E}}_0^m \right. \\
&\quad \quad \left. + \cosh \sqrt{m^2 \phi^2} (\cos \sqrt{4v^2} - 1) \frac{wv_b}{v^2} \bar{\mathcal{E}}_b^m \right] \\
&= \omega_H = dtE_0^0 + dx_mE_0^m, \\
\omega_a &= dt\mathcal{E}_a^0 + dx_m\mathcal{E}_a^m \\
&\xrightarrow{c \rightarrow 0} dt \left[ \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} 2v_a\partial^t \phi \right] + dx_m \left( \frac{\sin \sqrt{4v^2}}{\sqrt{4v^2}} 2v_a\partial^m \phi \right. \\
&\quad \left. + \cosh \sqrt{m^2 \phi^2} [P_{Tab}(v) + \cos \sqrt{4v^2} P_{Lab}(v)] \bar{\mathcal{E}}_b^m \right) \\
&= \omega_{Pa} = dtE_a^0 + dx_mE_a^m.
\end{aligned} \tag{B47}$$

Thus the vielbeine components of Section II, Equations (2.18)-(2.20), have been obtained. In short this  $c \rightarrow 0$  limit is

$$\mathcal{E}_A^M \xrightarrow{c \rightarrow 0} E_A^M. \tag{B48}$$

The  $c \rightarrow 0$  limit of the  $\phi$  covariant derivatives is found from the  $\omega_Z$  one-form Equation (B37)

$$\begin{aligned}
\omega_Z &\equiv dt\tilde{\nabla}^t \phi + dx_m\tilde{\nabla}^m \phi \\
&\xrightarrow{c \rightarrow 0} dt \left( \phi \cos \sqrt{4v^2} \right) \\
&\quad + dx_m \cos \sqrt{4v^2} \left[ \partial^m \phi - \cosh \sqrt{m^2 \phi^2} 2v_a \bar{e}_a^m \frac{\tan \sqrt{4v^2}}{\sqrt{4v^2}} \right] \\
&= \omega_Z = dt\nabla^t \phi + dx_m\nabla^m \phi,
\end{aligned} \tag{B49}$$

which agree with the  $\phi$  covariant derivatives in Section II Equation (2.22). Thus the same  $AdS - C$  results are obtained as in the use of the coset method.

The Nambu-Goto vielbein,  $\hat{N}_A^B$ , is defined by factoring the background vielbein from  $\hat{E}_A^M$  so that

$$\hat{E}_A^M = \hat{E}_B^M \hat{N}_A^B \tag{B50}$$

and so

$$\hat{N}_A^B = \hat{E}^{-1B}_M \hat{E}_A^M. \tag{B51}$$

Correspondingly  $\mathcal{E}_A^M = \bar{\mathcal{E}}^M_B \mathcal{N}_A^B$  and  $\mathcal{N}_A^B = \bar{\mathcal{E}}^{-1B}_M \mathcal{E}_A^M$ . Exploiting the relations between  $\hat{E}$ ,  $\hat{\hat{E}}$  and  $\mathcal{E}$ ,  $\bar{\mathcal{E}}$ , it is obtained that

$$\mathcal{N}_A^B = C^B_D \hat{N}^D_C C^{-1C}_A. \tag{B52}$$

Thus  $\det \mathcal{N} = \det \hat{N}$  and from Equations (B40) and (B42)  $\det \mathcal{E} = \det \hat{E}$  as well as  $\det \bar{\mathcal{E}} = \det \hat{\hat{E}}$ . Consequently the *AdS* invariant action, Equation (3.18) of Ref. 6 rescaled by  $-\sigma/c$ ,  $\Gamma_{AdS} \equiv \int dt d^p x \det \hat{E}$ , is written in terms of  $\det \mathcal{E}$  as

$$\det \hat{E} = \det \hat{\hat{E}} \det \hat{N} = \det \bar{\mathcal{E}} \det \mathcal{N} = \det \mathcal{E}. \tag{B53}$$

Utilizing Equation (3.19) or (3.20) of Ref. 6 for the  $\det \hat{N}$

$$\det \hat{N} = \cosh^d \sqrt{m^2 \hat{\phi}^2} \cosh \sqrt{\hat{\delta}^2} \left[ 1 - \left( \hat{\nu}_A \frac{\tanh \sqrt{\hat{\delta}^2}}{\sqrt{\hat{\delta}^2}} \right) \left( \frac{\hat{\hat{E}}^{-1A}_M \hat{\delta}^M \hat{\phi}}{\cosh \sqrt{m^2 \hat{\phi}^2}} \right) \right], \tag{B54}$$

and converting  $\hat{\phi}$ ,  $\hat{\nu}_A$  and  $\hat{x}_M$  to  $\phi$ ,  $V_A$  and  $X_M$  as well as using the relation

$$\bar{\mathcal{E}}^{-1A}_M = C^A_B \hat{\hat{E}}^{-1B}_N C^{-1N}_M \tag{B55}$$

in order to find that

$$C^B_A \hat{\hat{E}}^{-1A}_M \frac{\partial}{\partial \hat{x}_M} \hat{\phi} = -\bar{\mathcal{E}}^{-1B}_M \frac{\partial}{\partial X_M} \phi, \tag{B56}$$

the  $\det \mathcal{N}$  is found

$$\det \mathcal{N} = \cosh^d \sqrt{m^2 \phi^2} \cosh \sqrt{4V^2} \left[ 1 + \left( 2V_A \frac{\tanh \sqrt{4V^2}}{\sqrt{4V^2}} \right) \left( \frac{\bar{\mathcal{E}}^{-1A}_M \partial^M \phi}{\cosh \sqrt{m^2 \phi^2}} \right) \right]. \tag{B57}$$

Thus the  $AdS_{d+1} \rightarrow AdS_d$  brane embedded action  $\Gamma_{AdS}$  is obtained (note Equation (3.20) of Ref. 6)

$$\begin{aligned} \Gamma_{AdS} &= \int d^d X \det \bar{\mathcal{E}} \det \mathcal{N} \\ &= \int d^d X \det \bar{\mathcal{E}} \cosh^d \sqrt{m^2 \phi^2} \cosh \sqrt{4V^2} \left[ 1 + \left( 2V_A \frac{\tanh \sqrt{4V^2}}{\sqrt{4V^2}} \right) \right. \\ &\quad \left. \left( \frac{1}{\cosh \sqrt{m^2 \phi^2}} \bar{\mathcal{E}}^{-1A}_M \frac{\partial}{\partial X_M} \phi \right) \right], \end{aligned} \tag{B58}$$

in which the explicit factors of  $c$  are in the background vielbein  $\bar{\mathcal{E}}^M_A$  and its inverse  $\bar{\mathcal{E}}^{-1A}_M$  (note the form of Equation (3.39) in which the  $c \rightarrow 0$  limit is already taken). Further, taking the  $c \rightarrow 0$  limit, the explicit component fields can be exhibited from Equations (B12) and (B16) to obtain Equation (3.15) of Section III for  $\det \mathcal{N} \xrightarrow{c \rightarrow 0} \det N$  and likewise  $\det \bar{\mathcal{E}} \xrightarrow{c \rightarrow 0} \det \bar{E} = \bar{e}^0_a \det \bar{e}^m_a$ .

Thus  $\Gamma_{AdS} \xrightarrow{c \rightarrow 0} \Gamma_{AdS-C_D}$  and Equation (3.16) is obtained.

<sup>1</sup> J.-M. Levy-Leblond, Ann. I.H.P.: Phys. Theor. **3**, 1 (1965), <http://eudml.org/doc/75509>.

<sup>2</sup> H. Bacry and J. Levy-Leblond, J. Math. Phys. **9**, 1605 (1968).

<sup>3</sup> E. Bergshoeff, J. Gomis, and G. Longhi, Classical Quantum Gravity **31**(20), 205009 (2014); e-print [arXiv:1405.2264](https://arxiv.org/abs/1405.2264) [hep-th].

- <sup>4</sup> E. Bergshoeff, J. Gomis, and L. Parra, “The Symmetries of the Carroll Superparticle,” *J. Phys. A* **49**(18), 185402 (2016).
- <sup>5</sup> G. Gibbons, K. Hashimoto, and P. Yi, *J. High Energy Phys.* **2002**, 061; e-print [arXiv:hep-th/0209034](https://arxiv.org/abs/hep-th/0209034).
- <sup>6</sup> T. E. Clark, S. T. Love, M. Nitta, and T. ter Veldhuis, *J. Math. Phys.* **46**, 102304 (2005); e-print [arXiv:hep-th/0501241](https://arxiv.org/abs/hep-th/0501241).
- <sup>7</sup> T. E. Clark, S. T. Love, M. Nitta, T. ter Veldhuis, and C. Xiong, *Phys. Rev. D* **76**, 105014 (2007); e-print [arXiv:hep-th/0703179](https://arxiv.org/abs/hep-th/0703179) [HEP-TH].
- <sup>8</sup> B. Cardona, J. Gomis, and J. M. Pons, “Dynamics of Carroll Strings,” *JHEP* **1607**, 050 (2016).
- <sup>9</sup> J. Hartong, *J. High Energy Phys.* **1508**, 069 (2015); e-print [arXiv:1505.05011](https://arxiv.org/abs/1505.05011) [hep-th].
- <sup>10</sup> C. Duval, G. W. Gibbons, P. A. Horvathy, and P. M. Zhang, *Classical Quantum Gravity* **31**, 085016 (2014); e-print [arXiv:1402.0657](https://arxiv.org/abs/1402.0657) [gr-qc].
- <sup>11</sup> S. R. Coleman, J. Wess, and B. Zumino, *Phys. Rev.* **177**, 2239 (1969); C. G. Callan, Jr., S. R. Coleman, J. Wess, and B. Zumino, *Phys. Rev.* **177**, 2247 (1969).
- <sup>12</sup> D. V. Volkov, *Sov. J. Part. Nucl.* **4**, 3 (1973); V. I. Ogievetsky, in *Proceedings of the X-th Winter School of Theoretical Physics in Karpacz* (Wroclaw, 1974), Vol. 1, p. 227.
- <sup>13</sup> E. A. Ivanov and V. I. Ogievetsky, *Teor. Mat. Fiz.* **25**, 164 (1975).