

University of Groningen

Vanadium (beta-(dimethylamino)ethyl)cyclopentadienyl complexes with diphenylacetylene ligands

Liu, Guohua; Lu, Xiaoquan; Gagliardo, Marcella; Beetstra, Dirk J.; Meetsma, Auke; Hessen, Bart; Liu, Gaifen

Published in:
 Organometallics

DOI:
[10.1021/om8000718](https://doi.org/10.1021/om8000718)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2008

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Liu, G., Lu, X., Gagliardo, M., Beetstra, D. J., Meetsma, A., Hessen, B., & Liu, G. (2008). Vanadium (beta-(dimethylamino)ethyl)cyclopentadienyl complexes with diphenylacetylene ligands. *Organometallics*, 27(10), 2316-2320. <https://doi.org/10.1021/om8000718>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Vanadium (β -(Dimethylamino)ethyl)cyclopentadienyl Complexes with Diphenylacetylene Ligands

Guohua Liu,^{*,†} Xiaoquan Lu,[†] Marcella Gagliardo,[‡] Dirk J. Beetstra,[‡] Auke Meetsma,[‡] and Bart Hessen^{*,‡}

Department of Chemistry, College of Life and Environmental Science, Shanghai Normal University, Shanghai 200234, People's Republic of China, and Center for Catalytic Olefin Polymerization, Stratingh Institute for Chemistry and Chemical Engineering, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

Received January 26, 2008

Reduction of the V(III) (β -(dimethylamino)ethyl)cyclopentadienyl dichloride complex $[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{VCl}_2(\text{PMe}_3)$ (**1**) with 1 equiv of Na/Hg yielded the V(II) dimer $\{[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{V}(\mu\text{-Cl})\}_2$ (**2**). This compound reacted with diphenylacetylene in THF to give the V(II) alkyne adduct $[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{VCl}(\eta^2\text{-PhC}\equiv\text{CPh})$ (**3**). Further reduction of **2** with Mg in the presence of diphenylacetylene resulted in oxidative coupling of two diphenylacetylene groups to yield the diamagnetic, formally V(V), bent metallacyclopentatriene complex $[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{V}(\text{C}_4\text{Ph}_4)$ (**4**).

Amino-functionalized cyclopentadienyl transition-metal complexes have attracted much attention, owing to their dramatic effect on catalytic function compared to the case for the corresponding parent complexes.¹ Playing a major role in this area are titanium and chromium complexes, which exhibit good activity in ethene and propene polymerization.² However, there are relatively few reports concerning vanadium complexes of this type.³ This is mainly due to the fact that such compounds are extremely air-sensitive and paramagnetic, due to the inherent instability of monocyclopentadienyl vanadium analogues. The limiting step in the development of this chemistry has been the absence of suitable organometallic vanadium starting materials. Amino-functionalized cyclopentadienyl ligands with additional pendant Lewis basic functionalities have been used to enhance the stability of metal complexes through the chelate effect, thus leading to interesting products. It has been recognized that such ligands can exhibit hemilabile behavior, in which the pendant

functionality can reversibly dissociate from the metal center. This behavior can strongly affect the reactivity of such complexes: for instance, in catalytic conversions.⁴ Recently, we described the chemistry of the vanadium(III) complex $(\eta^5:\eta^1\text{-C}_5\text{H}_4\text{CH}_2\text{CH}_2\text{NMe}_2)\text{VCl}_2(\text{PMe}_3)$,⁵ containing a (β -(dimethylamino)ethyl)cyclopentadienyl ligand, which seemed to us to be a suitable starting material for the development of new organovanadium chemistry.⁶ Also, we observed that the tendency of the pendant amino group to bind to or dissociate from the vanadium center depends strongly on the nature of the other ligands bound to the vanadium atom.

In this contribution, we present the chemistry of the dimeric vanadium(II) (β -(dimethylamino)ethyl)cyclopentadienyl complex $\{[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{V}(\mu\text{-Cl})\}_2$ (**2**). It has been found that the reaction of **2** with diphenylacetylene produces the V(II) alkyne adduct $[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{VCl}(\eta^2\text{-PhC}\equiv\text{CPh})$ (**3**) and reduction of **2** with Mg in the presence of diphenylacetylene results in the formation of the bent V(V) metallacyclopentatriene complex $[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{V}(\text{C}_4\text{Ph}_4)$ (**4**), in which the Lewis basic amino group can bind to the vanadium center through the chelate effect.

Results and Discussion

Synthesis and Molecular Structure of $\{[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{V}(\mu\text{-Cl})\}_2$ (2**).** The vanadium (β -(dimethylamino)ethyl)cyclopentadienyl complex $[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{VCl}_2(\text{PMe}_3)$ (**1**)⁵ was prepared in high yield by a straightforward reaction between $\text{VCl}_3(\text{PMe}_3)_2$ and $\text{Li}[\text{C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]$ in THF. One-electron reduction of the V(III) complex **1** with 1 equiv of Na/Hg in THF afforded the red-violet dinuclear V(II) chloride-bridged complex $\{[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{V}(\mu\text{-Cl})\}_2$ (**2**; eq 1) in 56% isolated yield. Due to its paramagnetism, the ¹H NMR spectrum of **2** only shows a very broad resonance for

* To whom correspondence should be addressed. Tel: +86-21-64321819. Fax: +86-21-64322511. E-mail: ghliu@shnu.edu.cn.

[†] Shanghai Normal University.

[‡] University of Groningen.

(1) (a) Baretta, A.; Chong, K. S.; Cloke, F. G. N.; Feigenbaum, A.; Green, M. L. H. *J. Chem. Soc., Dalton Trans.* **1983**, 86, 1–864. (b) Jutzi, P.; Redeker, T. *Eur. J. Inorg. Chem.* **1998**, 663–674. (c) Britovsek, G. J. P.; Gibson, V. C.; Wass, D. F. *Angew. Chem., Int. Ed.* **1999**, 38, 428–447. (d) Arndt, S.; Okuda, J. *Chem. Rev.* **2002**, 102, 1953–1976. (e) Schrock, R. R. *Chem. Rev.* **2002**, 102, 145–179.

(2) (a) Flores, J. C.; Chien, J. C. W.; Rausch, M. D. *Organometallics* **1994**, 13, 4140–4142. (b) Flores, J. C.; Chien, J. C. W.; Rausch, M. D. *Macromolecules* **1996**, 29, 8030–8035. (c) Blais, M. S.; Chien, J. C. W.; Rausch, M. D. *Organometallics* **1998**, 17, 3775–3783. (d) van Beek, J. A. M.; van Doremaele, G. H. J.; Gruter, G. J. M.; Arts, H. J.; Eggels, G. H. M. R. *WO 96/13529*, 1995, issued to DSM NV. (e) Ypey, E. G.; van Beek, J. A. M.; Gruter, G. J. M. *WO 97/42157*, 1997, issued to DSM NV. (f) van Beek, J. A. M.; Gruter, G. J. M. *EP 0805142 A1*, 1997, issued to DSM NV. (g) Herberich, G. E.; Schmidt, B.; Schmitz, A.; van Beek, J. A. M. *WO 97/23493*, 1997, issued to DSM NV. (h) van Beek, J. A. M.; van Doremaele, G. H. J.; Gruter, G. J. M.; Arts, H. J.; Eggels, G. H. M. R. *U.S. Patent 5,986,029*, 1999, issued to DSM NV. (i) Jolly, P. W.; Joans, K.; Verhovnik, G. P. J.; Döhning, A.; Göhre, J.; Weber, J. C. *WO 98/04570*, 1998, issued to Studiengesellschaft Kohle MBH. (k) Jolly, P. W.; Joans, K.; Verhovnik, G. P. J. *DE 19630580*, 1998, assigned to Studiengesellschaft Kohle mbH.

(3) Bradley, S.; Camm, K. D.; Furtado, S. J.; Gott, A. L.; McGowan, P. C.; Podesta, T. J.; Thornton-Pett, M. *Organometallics* **2002**, 21, 3443–3453.

(4) (a) Deckers, P. J. W.; Hessen, B.; Teuben, J. H. *Angew. Chem., Int. Ed.* **2001**, 40, 2516–2519. (b) Deckers, P. J. W.; Hessen, B.; Teuben, J. H. *Organometallics* **2002**, 21, 5122–5135. (c) de Bruin, T. J. M.; Magna, L.; Raybaud, P.; Toulhoat, H. *Organometallics* **2003**, 22, 3404–3413.

(5) Liu, G. H.; Beetstra, D. J.; Meetsma, A.; Hessen, B. *Organometallics* **2004**, 23, 3914–3920.





Figure 1. Molecular structure of $\{[\eta^5\text{-}\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{V}(\mu\text{-Cl})_2\}_2$ (**2**). Thermal ellipsoids are drawn at the 50% probability level. Hydrogen atoms are omitted for clarity.

Table 1. Selected Bond Lengths (Å) and Angles (deg) for **2**


V(1)–Cl(1)	2.443(2)	V(1)–C(1)	2.277(7)
V(1)–Cl(2)	2.434(2)	V(1)–C(2)	2.308(7)
V(2)–Cl(1)	2.430(2)	V(1)–C(3)	2.326(7)
V(2)–Cl(2)	2.454(2)	V(1)–C(4)	2.290(7)
V(1)–N(1)	2.251(5)	V(1)–C(5)	2.253(7)
V(2)–N(2)	2.258(7)		
Cl(1)–V(1)–Cl(2)	92.78(7)	Cl(1)–V(2)–N(2)	100.22(18)
Cl(1)–V(1)–N(1)	93.43(15)	Cl(2)–V(2)–N(2)	91.21(17)
Cl(2)–V(1)–N(1)	93.81(14)	V(1)–Cl(1)–V(2)	76.43(6)
Cl(1)–V(2)–Cl(2)	92.60(7)	V(1)–Cl(2)–V(2)	76.16(6)

all protons. However, it is clear that **2** is a phosphine-free complex, as confirmed by the disappearance of the PMe_3 proton resonances in the ^1H NMR spectrum. A crystal structure determination of **2** (Figure 1, with selected bond lengths and angles given in Table 1) shows a puckered V_2Cl_2 core with the cyclopentadienyl ligands in a cis arrangement. It strongly resembles the dimeric V(II) monochloride triethylphosphine complex $[\text{Cp}(\text{Et}_3\text{P})\text{V}(\mu\text{-Cl})_2]$ reported previously,^{6g,h} with very similar V–Cl distances in the puckered V_2Cl_2 unit, which is essentially equilateral. The V–Cl distances (2.443(2), 2.434(2), 2.430(2), and 2.454(2) Å) are comparable to those observed in other chloride-bridged dimeric vanadium complexes (2.4128(15) and 2.5365(15) Å in $[\text{V}(=\text{NAr})\text{Cl}_2(\text{dppm})_2]$ ⁷ and 2.459(2) and 2.373(2) Å in $\{[(\text{Me}_3\text{Si})\text{NCH}_2\text{CH}_2]_2\text{N}(\text{Me}_3\text{Si})_2\text{V}_2(\mu\text{-Cl}_2)\}$,⁸ although there are slight differences. Furthermore, the Cl–V–Cl angles of 92.78(7) and 93.43(15)° in the dimeric complex **2** are obviously larger than those observed in a closely related dimeric titanium complex (77.11(5), 78.21(7), and 78.63(7)° in $(\text{C}_5\text{H}_4)_2\text{TiCl}_2$),⁹ indicating the steric nature of the β -aminoethyl-functionalized cyclopentadienyl ligand. Apparently the (Cp-ethylamino)VCl fragment prefers to form dimeric **2** rather than

to bind the PMe_3 ligand. This behavior was also observed for the $\text{CpVCl}(\text{PR}_3)$ system (R = Et, Me), although for R = Me it was seen that the equilibrium may be shifted to the side of $\text{CpVCl}(\text{PMe}_3)_2$ when an excess of PMe_3 is added.^{6g}



Reaction of 2 with Diphenylacetylene: Synthesis and Molecular Structure of $[\eta^5\text{-}\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{VCl}(\eta^2\text{-PhC}\equiv\text{CPh})$ (3**).** Reaction of metal chloride complexes with alkynes can result in highly interesting derivatives.¹⁰ However, for vanadium compounds, only several examples have been reported.¹¹ In this case, when a toluene solution of **2** was treated with 1 equiv of phenylacetylene at ambient temperature, no reaction was observed and **2** could be recovered unchanged. However, when the same reaction was performed in THF solution, the V(II) diphenylacetylene adduct $[\eta^5\text{-}\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{VCl}(\eta^2\text{-PhC}\equiv\text{CPh})$ (**3**) was isolated as red crystals in 57% yield after recrystallization from pentane. Apparently, the coordination of the alkyne to the V(II) center is thermodynamically favorable, but diphenylacetylene is kinetically unable to cleave the $(\mu\text{-Cl})_2$ bridge in dinuclear **2**. Although the low-valent metal center is expected to have a relatively low affinity for THF, the ether apparently is kinetically competent to cleave **2** to give a transient mononuclear THF adduct, from which the THF subsequently is displaced by the alkyne (eq 2). The alkyne adduct **3** was characterized by single-crystal X-ray diffraction, and its structure is shown in Figure 2 (selected bond lengths and angles are given in Table 2). Its structure is geometrically similar to that of the V(I) complex $\text{CpV}(\text{PMe}_3)_2(\eta^2\text{-PhC}\equiv\text{CPh})$.^{6b} In the latter, the alkyne $\text{C}\equiv\text{C}$ bond lies approximately in the same plane as one of the V–P bonds. In **3** the alkyne is similarly oriented relative to the V–N bond. As was observed in $\text{CpV}(\text{PMe}_3)_2(\eta^2\text{-PhC}\equiv\text{CPh})$, the bonding of the cyclopentadienyl moiety to vanadium in **3** is noticeably distorted from the regular η^5 mode, with the longest V–C distances to C(3) and C(4) (2.35–2.36 Å) and the shortest to C(1) (2.24 Å). A closer look at the coordinated alkyne reveals that both the $\text{C}\equiv\text{C}$ distance of 1.312(3) Å and the C–C–C(Ph) angles of 139° are indications of a somewhat lesser extent of π -back-donation in the V(II) complex **3** than in the V(I) complex $\text{CpV}(\text{PMe}_3)_2(\eta^2\text{-PhC}\equiv\text{CPh})$, where the related parameters are 1.328(3) Å and 136°.



Reduction of 2 in the Presence of Diphenylacetylene: Synthesis and Molecular Structure of $[\eta^5\text{-}\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{V}(\text{C}_4\text{Ph}_4)$ (4**).** Further reduction of the V(II) complex **2** by Mg in THF in the presence of diphenylacetylene (performed

(6) (a) Nieman, J.; Scholtens, H.; Teuben, J. H. *J. Organomet. Chem.* **1980**, *186*, C12–C14. (b) Hessen, B.; Meetsma, A.; Van Bolhuis, F.; Teuben, J. H.; Helgesson, G.; Jagner, S. *Organometallics* **1990**, *9*, 1925–1936. (c) Hessen, B.; Teuben, J. H.; Lemmen, T. H.; Huffman, J. C.; Caulton, K. G. *Organometallics* **1985**, *4*, 946–948. (d) Hessen, B.; Lemmen, T. H.; Lutikhedde, H. J. G.; Teuben, J. H.; Petersen, J. L.; Jagner, S.; Huffman, J. C.; Caulton, K. G. *Organometallics* **1987**, *6*, 2354–2362. (e) Hessen, B.; Buijink, J. K. F.; Meetsma, A.; Teuben, J. H.; Helgesson, G.; Hakansson, M.; Jagner, S.; Spek, A. L. *Organometallics* **1993**, *12*, 2268–2276. (f) Hessen, B.; Van Bolhuis, F.; Teuben, J. H.; Petersen, J. L. *J. Am. Chem. Soc.* **1988**, *110*, 295–296. (g) Nieman, J.; Teuben, J. H. *Organometallics* **1986**, *5*, 1149–1153. (h) Nieman, J.; Teuben, J. H.; Hulsbergen, F. B.; de Graaff, R. A. G.; Reedijk, J. *Inorg. Chem.* **1987**, *26*, 2376. (i) Hessen, B.; Meetsma, A.; Teuben, J. H. *J. Am. Chem. Soc.* **1988**, *110*, 4860–4861.

(7) Feghali, K.; Harding, D. J. H.; Reardon, D.; Gambarotta, S.; Yap, G.; Wang, Q. *Organometallics* **2002**, *21*, 968–976.

(8) Lorber, C.; Choukroun, R.; Donnadiou, B. *Inorg. Chem.* **2003**, *42*, 673–675.

(9) Jungst, R.; Dekutowshi, D.; Davis, J.; Luly, M.; Stucky, G. *Inorg. Chem.* **1977**, *7*, 1645–1655.





Figure 2. Molecular structure of $[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{VCl}(\eta^2\text{-PhC}\equiv\text{CPh})$ (**3**). Thermal ellipsoids are drawn at the 50% probability level. Hydrogen atoms are omitted for clarity.

Table 2. Selected Bond Lengths (Å) and Angles (deg) for **3**

V(1)–Cl(1)	2.3366(6)	V(1)–C(1)	2.239(3)
V(1)–N(1)	2.276(3)	V(1)–C(2)	2.284(3)
V(1)–C(16)	1.969(3)	V(1)–C(3)	2.351(2)
V(1)–C(17)	2.003(2)	V(1)–C(4)	2.360(3)
C(16)–C(17)	1.312(3)	V(1)–C(5)	2.305(3)
C(16)–V(1)–C(17)	38.56(9)	N(1)–V(1)–C(17)	88.66(9)
Cl(1)–V(1)–C(16)	103.43(7)	N(1)–V(1)–Cl(1)	90.69(5)
Cl(1)–V(1)–C(17)	109.09(7)	C(15)–C(16)–C(17)	139.1(2)
N(1)–V(1)–C(16)	127.19(9)	C(16)–C(17)–C(18)	138.2(3)

at low temperature, -30 to -5 °C) resulted in the isolation of a diamagnetic red crystalline compound that was characterized by single-crystal X-ray diffraction as the bent metallacyclopentatriene complex $[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2]\text{V}(\text{C}_4\text{Ph}_4)$ (**4**; eq 3). It is likely to be formed by reduction of the vanadium to V(I) and coordination of two alkyne molecules to the metal center followed by an oxidative coupling of the diphenylacetylene ligands to yield a metallacycle. It was observed previously that the metallacycle of the formula $\text{CpV}(\text{C}_4\text{R}_4)(\text{PMe}_3)$ takes on a bent metallacyclopentatriene structure rather than the more common planar metallacyclopentadiene structure.^{6b} The crystal structure of **4** (Figure 3, with selected bond lengths and angles given in Table 3) shows two short V–C bond distances of 1.888(5) and 1.895(4) Å, which are shorter than that of 1.922 Å in a benzylidene complex.¹³ Such short V–C bond distances



In conclusion, the vanadium(III) (β -(dimethylamino)ethyl)-cyclopentadienyl dichloride complex $(\eta^5:\eta^1\text{-C}_5\text{H}_4\text{CH}_2\text{CH}_2\text{NMe}_2)\text{-VCl}_2(\text{PMe}_3)$ is a convenient precursor for synthesis of a range of organometallic vanadium derivatives. It has also been recognized that amino-functionalized cyclopentadienyl ligands with additional Lewis basic functionalities can enhance the stability of the vanadium complexes through the chelate effect, thus resulting in novel complexes.

Experimental Section

General Considerations. All manipulations were performed under an inert nitrogen atmosphere, using standard Schlenk or glovebox techniques. Pentane (Aldrich, anhydrous, 99.8%) was passed over columns of Al_2O_3 (Fluka), BASF R3-11-supported Cu oxygen scavenger, and molecular sieves (Aldrich, 4 Å). Diethyl ether and THF (Aldrich, anhydrous, 99.8%) were dried over Al_2O_3 (Fluka). All solvents were degassed prior to use and stored under nitrogen. Deuterated solvents (C_6D_6 , $\text{THF-}d_8$; Aldrich) were vacuum-transferred from Na/K alloy prior to use. Starting materials: $(\text{C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2)\text{VCl}_2(\text{PMe}_3)$ was prepared according to the reported method.⁶ ^1H NMR spectra were recorded on Varian VXR-300 (300 MHz) spectrometers in NMR tubes sealed with a Teflon

(15) Ahmed, K. J.; Chisholm, M. H.; Huffman, J. C. *Organometallics* **1985**, *4*, 1168–1174.

(16) (a) Basuli, F.; Kilgore, U. J.; Hu, X. L.; Meyer, K.; Pink, M.; Huffman, J. C.; Mindiola, D. J. *Angew. Chem., Int. Ed.* **2004**, *43*, 3156–3159. (b) Mindiola, D. J. *Acc. Chem. Res.* **2006**, *39*, 813–821.

(17) Hirpo, W.; Curtis, M. D. *J. Am. Chem. Soc.* **1988**, *110*, 5218–5221.

(10) (a) Marschner, C. *Angew. Chem., Int. Ed.* **2007**, *46*, 6770–6771. (b) Rosenthal, U.; Burlakov, V. V.; Arndt, P.; Baumann, W.; Spannenberg, A. *Organometallics* **2003**, *22*, 884–900. (c) Rosenthal, U.; Burlakov, V. V.; Arndt, P.; Baumann, W.; Spannenberg, A. *Organometallics* **2005**, *24*, 456–471. (d) Beveries, T.; Burlakov, V. V.; Bach, M. A.; Arndt, P.; Baumann, W.; Spannenberg, A.; Rosenthal, U. *Organometallics* **2007**, *26*, 247–249.

(11) (a) Köhler, F. H.; Hofmann, P.; Prössdorf, W. *J. Am. Chem. Soc.* **1981**, *103*, 6359–6367. (b) Köhler, F. H.; Prössdorf, W.; Schubert, U. *Inorg. Chem.* **1981**, *20*, 4096–4101. (c) Buijink, J. K. F.; Kloetstra, K. R.; Meetsma, A.; Teuben, J. H.; Smeets, W. J. J.; Spek, A. L. *Organometallics* **1996**, *15*, 2523–2533. (d) Evans, W. J.; Bloom, I.; Doedens, R. J. *J. Organomet. Chem.* **1984**, *265*, 249–255. (e) Fachinetti, G.; Floriani, C.; Chiesi-Villa, A.; Guastini, C. *Inorg. Chem.* **1979**, *18*, 2282–2287.

(12) Johnson, S. A.; Liu, F. Q.; Suh, M. C.; Surcher, S.; Haufe, M.; Mao, S. S. H.; Tilley, T. D. *J. Am. Chem. Soc.* **2003**, *125*, 4199–4211.

(13) Buijink, J.-K. F.; Teuben, J. H.; Kooijman, H.; Spek, A. L. *Organometallics* **1994**, *13*, 2922–2924.

(14) Moore, M.; Gambarotta, S.; Yap, G.; Liable-Sands, L. M.; Rheingold, A. L. *Chem. Commun.* **1997**, 643–644.




Figure 3. Molecular structure of the cation of $[\eta^5:\eta^1\text{-C}_5\text{H}_4\text{-(CH}_2\text{)}_2\text{NMe}_2\text{]V(C}_4\text{Ph}_4\text{)}$ (**4**). Thermal ellipsoids are drawn at the 50% probability level. Hydrogen atoms have been omitted for clarity.

Table 3. Selected Bond Lengths (Å) and Angles (deg) for **4**

V(1)–C(10)	1.888(5)	C(10)–C(17)	1.433(6)
V(1)–C(31)	1.895(4)	C(24)–C(31)	1.438(6)
V(1)–C(17)	2.360(5)	C(17)–C(24)	1.417(5)
V(1)–C(24)	2.339(5)	V(1)–N(1)	2.254(4)
C(10)–V(1)–C(31)	92.7(2)	N(1)–V(1)–C(31)	116.32(15)
C(10)–V(1)–C(17)	37.40(15)	C(24)–C(31)–V(1)	88.0(3)
C(17)–V(1)–C(24)	35.11(14)	V(1)–C(10)–C(17)	89.5(3)
C(24)–V(1)–C(31)	37.91(16)	C(17)–C(24)–C(31)	118.0(4)
N(1)–V(1)–C(10)	112.60(15)	C(10)–C(17)–C(24)	116.8(4)

Table 4. Crystallographic Data for **2–4**

	2	3	4
mol formula	C ₁₈ H ₂₈ Cl ₂ N ₂ V ₂	C ₂₃ H ₂₄ ClNV	C ₃₇ H ₃₄ NV
fw	445.20	400.82	543.59
diffractometer	SMART APEX CCD	SMART APEX CCD	SMART APEX CCD
temp (K)	100(1)	100(1)	100(1)
cryst syst	monoclinic	trigonal	monoclinic
space group	<i>P</i> ₂ / <i>1</i> / <i>c</i>	<i>R</i> ³	<i>P</i> ₂ / <i>1</i> / <i>n</i>
<i>a</i> (Å)	7.736(2)	31.948(2)	9.5142(9)
<i>b</i> (Å)	16.663(3)	31.948(2)	30.774(3)
<i>c</i> (Å)	16.225(3)	11.0765(7)	10.501(1)
β (deg)	99.529(3)		116.330(2)
<i>V</i> (Å ³)	2062.6(8)	9790.9(11)	2755.6(5)
<i>Z</i>	4	18	4
<i>d</i> _{calcd} (g cm ⁻³)	1.434	1.224	1.310
F(000)	920	3672	1144
ν (Mo K α), cm ⁻¹	11.67	5.84	3.87
θ range (deg)	2.44, 26.02	2.21, 28.28	2.41, 24.73
<i>R</i> _w (<i>F</i> ²)	0.2044	0.1257	0.1612
no. of indep rflns	4062	5403	4842
no. of params	221	331	354
<i>R</i> (<i>F</i>) for <i>F</i> _o > 4.0 σ (<i>F</i> _o)	0.0736	0.0391	0.0678
GOF	1.012	1.158	0.986
largest diff peak/hole (e Å ⁻³)	1.2(1), -0.7(1)	0.43(10), -0.28	1.12(10), -0.43

(Young) stopcock. IR spectra were recorded on a Mattson-4020 Galaxy FT-IR spectrometer from Nujol mulls between KBr disks unless stated otherwise. Elemental analyses were performed by Kolbe Analytical Laboratories, Mülheim a.d. Ruhr, Germany.

Preparation of $[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2\text{]V(Cl)}_2$ (2**).** Na sand (0.090 g, 3.90 mmol) was added to 45 g of frozen Hg and carefully dissolved by thawing out the Hg. When the Na/Hg was at room temperature, it was added to a solution of complex **1** (1.30

g, 3.90 mmol) in 30 mL of dry THF. The deep purple solution turned violet over 2 h. After it had been stirred overnight, the violet THF solution was transferred into a new Schlenk flask and the residual Hg was washed twice with 5 mL of THF. All the THF solutions were combined, the volatiles were removed in vacuo, and the resulting violet solid was stripped twice with 15 mL of pentane. The violet-red solid was repeatedly extracted with 30 mL portions of pentane. The violet-red extracts were filtered and concentrated to 10 mL. Cooling to -30°C produced violet-red crystals of **2** (0.98 g; 2.2 mmol; 56%). IR (Nujol mull): 635, 678, 754, 772, 786, 817, 920, 953, 996, 1022, 1046, 1098, 1117, 1167, 1210, 1236, 1267, 1323, 1377, 1402, 1461, 2831, 2887, 2910, 2942, 2963 cm⁻¹. ¹H NMR (benzene-*d*₆, 20 °C, 300 MHz): δ 50.69 (s), 47.68 (s), 35.89 ($\Delta\nu_{1/2}$ = 1240 Hz), 31.19 ($\Delta\nu_{1/2}$ = 749 Hz), 21.58 ($\Delta\nu_{1/2}$ = 480 Hz), 12.19 ($\Delta\nu_{1/2}$ = 429 Hz), 11.18 ($\Delta\nu_{1/2}$ = 342 Hz), -2.39 ($\Delta\nu_{1/2}$ = 146 Hz), -4.59 ($\Delta\nu_{1/2}$ = 240 Hz). Anal. Calcd for C₁₈H₂₈Cl₂N₂V₂: C, 48.56; H, 6.34; N, 6.29. Found: C, 48.53; H, 6.33; N, 6.09.

Preparation of $[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2\text{]VCl}(\eta^2\text{-PhC}\equiv\text{CPh})$ (3**).** A solution of **2** (148 mg, 0.33 mmol) together with PhC \equiv CPh (118 mg, 0.66 mmol) in 5 mL of THF was stirred overnight at room temperature. The solvents were removed in vacuo, and the resulting solid was stripped with two portions of 5 mL of ether. The red solid was repeatedly extracted with 30 mL portions of ether. The red extracts were filtered and concentrated to 5 mL. Cooling to -30°C produced red crystals of **3** (152 mg, 0.38 mmol, 57%). IR (Nujol mull): 689, 722, 754, 773, 802, 912, 921, 1001, 1024, 1044, 168, 1098, 1260, 1377, 1461, 1498, 1587, 1603, 1636, 2854, 2924, 2954 cm⁻¹. ¹H NMR (benzene-*d*₆, 20 °C, 300 MHz): δ 5.12 ($\Delta\nu_{1/2}$ = 11 Hz), 5.07 ($\Delta\nu_{1/2}$ = 18 Hz), 4.89 (s, 2H, Ph), 4.87 (s, Ph), 4.86 (s, Ph), 4.66 ($\Delta\nu_{1/2}$ = 12 Hz), 4.61 (s), 4.46 (s). Anal. Calcd for C₂₃H₂₄ClNV: C, 68.92; H, 6.04; N, 3.49. Found: C, 69.11; H, 5.89; N, 3.36.

Preparation of $[\eta^5:\eta^1\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{NMe}_2\text{]V(C}_4\text{Ph}_4)$ (4**).** To 0.3 g of activated Mg (12.3 mmol) was added a solution of **2** (124 mg, 0.28 mmol) together with PhC \equiv CPh (200 mg, 1.12 mmol) in 5 mL of THF at -30°C . After 20 min, the solution changed from violet to deep red. The solution was warmed to -5°C over another 40 min. The solvent was removed in vacuo and the residue stripped with two 5 mL portions of pentane. The brown-red solid was repeatedly extracted with 30 mL of pentane. The extracts were filtered and concentrated to 5 mL. Cooling to -30°C produced brown-red crystals of **4** (193 mg; 0.36 mmol; 59.6%). IR (Nujol mull): 695, 721, 753, 773, 784, 828, 842, 925, 957, 995, 1023, 1071, 1097, 1113, 1152, 1262, 1326, 1377, 1461, 1484, 1584, 2853, 2923, 2951 cm⁻¹. ¹H NMR (benzene-*d*₆, 20 °C, 300 MHz): δ 7.61, 7.59 (d, 4 H, Ph), 7.00–6.88 (m, 12 H, Ph), 6.37 (t, 2 H, *J* = 2.1 Hz, Cp), 4.45 (t, 2 H, *J* = 2.1 Hz, Cp), 1.79 (t, 2H, *J* = 6.3 Hz, CpCH₂), 1.44 (t, 2H, *J* = 6.3 Hz, CH₂N), 1.28 (s, 6 H, NMe₂). ¹³C NMR (benzene-*d*₆, 20 °C, 75.4 MHz): δ 25.45 (t, CpCH₂), 48.52 (q, NMe₂), 69.70 (t, NCH₂), 94.93 (b, C=C), 104.42 (b, Cp C), 123.70, 124.02, 125.14, 127.14, 127.39, 127.61, 133.91, 141.55, 150.92 (all, b, Ph C), 263.64 (b, V=C). Anal. Calcd for C₃₇H₃₄NV: C, 81.75; H, 6.30; N, 2.58. Found: C, 81.75; H, 6.38; N, 2.50.

Structure Determinations. Suitable crystals for single-crystal X-ray diffraction were obtained by cooling solutions of the compounds in pentane (**2** and **4**) and diethyl ether (**3**). Crystals were mounted on a glass fiber inside a drybox and transferred under an inert atmosphere to the cold nitrogen stream of a Bruker SMART APEX CCD diffractometer. Intensity data were collected with Mo K α radiation (λ = 0.710 73 Å). Intensity data were corrected for Lorentz and polarization effects. A semiempirical absorption correction was applied, based on the intensities of symmetry-related reflections measured at different angular settings (SADABS¹⁸). The structures were solved by Patterson methods, and extension of the

(18) Sheldrick, G. M. SHELXL-97 Program for the Refinement of Crystal Structures; University of Göttingen, Göttingen, Germany, 1997.

models was accomplished by direct methods applied to difference structure factors using the program DIRDIF.¹⁹ Hydrogen atom coordinates and isotropic thermal parameters were refined freely unless mentioned otherwise. All refinements and geometry calculations were performed with the program packages SHELXL and PLATON. Crystallographic data and details of the data collections and structure refinements are given in Table 4.

(19) Spek, A. L. PLATON Program for the Automated Analysis of Molecular Geometry, Version April 2000; University of Utrecht, Utrecht, The Netherlands, 2000.

Acknowledgment. We are grateful to the Shanghai Sciences and Technologies Development Fund (No. 071005119) and China National Natural Science Foundation (No. 20673072) for financial support.

Supporting Information Available: CIF files giving details of the structure determinations of **2–4**, including crystal data, positional and thermal parameters, and interatomic distances and angles. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM8000718