

University of Groningen

Local sex ratio affects the cost of reproduction

Nicolaus, Marion; Michler, Stephanie P. M.; Ubels, Richard; van der Velde, Marco; Bouwman, Karen M.; Both, Christiaan; Tinbergen, Joost M.

Published in:
Journal of Animal Ecology

DOI:
[10.1111/j.1365-2656.2011.01933.x](https://doi.org/10.1111/j.1365-2656.2011.01933.x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Nicolaus, M., Michler, S. P. M., Ubels, R., van der Velde, M., Bouwman, K. M., Both, C., & Tinbergen, J. M. (2012). Local sex ratio affects the cost of reproduction. *Journal of Animal Ecology*, 81(3), 564-572. <https://doi.org/10.1111/j.1365-2656.2011.01933.x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

1 **Appendices**

2 Local sex ratio affects the cost of reproduction

3 Marion Nicolaus, Stephanie P. M. Michler, Richard Ubels, Marco van der Velde,

4 Karen M. Bouwman, Christiaan Both and Joost M. Tinbergen

5

6 **Appendix S1. Collection of provisioning data**

7 In 2006, additional provisioning data of parents were collected on 94 broods using a
8 Radio Frequency Identification system (RFID). At day 5, an antenna was placed
9 around the nest-box entrance to let the parents habituate to the presence of a new
10 device. At day 10, the antenna was connected to a reader (Trovan, Ltd., Douglas, UK)
11 and the number of visits of PIT tagged parents was automatically recorded for 24
12 hours (for details on the PIT tags used see Nicolaus *et al.* 2008). At day 11 (\pm 25
13 hours later), data were collected and the RFID device removed. We defined a visit as
14 being a reading recorded with a minimum time interval of 3 seconds from the
15 previous reading. Because parental visits to the nest were recorded when going in and
16 out the next-box, we divided the total number of visits per individual by two. Video
17 data collected simultaneously outside the nest-boxes revealed that the number of visits
18 observed and automatically recorded were highly significantly and positively
19 correlated ($r=0.98$, $p<0.01$ for males, $r=0.99$, $p<0.01$ for females, $n=20$ individuals of
20 each sex).

21

22 Literature cited

23 Nicolaus M., K. M. Bouwman, & N. J. Dingemanse. 2008. Effect of PIT tags on the
24 survival and recruitment of Great Tits *Parus major*. *Ardea*, 96:286-292.

25

26 **Appendix S2. Overview of the plot manipulations and breeding parameters of**
27 **the Lauwersmeer great tit population in the Netherlands.** A density
28 ('decreased'/'increased') and a sex ratio ('female/balanced/male') treatment were
29 assigned to each of the 12 plots. The mean plot density (total number of nestlings per
30 plot) and mean plot sex ratio (proportion of male nestlings in a plot) are presented
31 before and after manipulation at day 6 (n=number of plots). Mean brood size, sex
32 ratio, probability of producing a second clutch and adult survival are presented with
33 their standard deviation (SD) and their sample size for the three study years 2005,
34 2006 and 2007 (n=number of nests or individuals).
35

parameters	2005			2006			2007		
	mean	SD	n	mean	SD	n	mean	SD	n
plot traits									
decreased before	156.83	28.27	6	134.33	48.73	6	166.83	19.57	6
decreased after	137.83	26.35	6	119.67	48.90	6	143.00	22.02	6
increased before	161.50	16.28	6	118.00	27.62	6	144.83	34.50	6
increased after	181.17	17.33	6	132.67	32.67	6	168.67	42.97	6
female before	0.48	0.02	4	0.47	0.06	4	0.46	0.01	4
female after	0.24	0.00	4	0.24	0.02	4	0.25	0.02	4
balanced before	0.47	0.02	4	0.50	0.02	4	0.51	0.04	4
balanced after	0.49	0.01	4	0.49	0.01	4	0.50	0.03	4
male before	0.50	0.03	4	0.49	0.04	4	0.52	0.06	4
male after	0.74	0.02	4	0.79	0.03	4	0.76	0.02	4
brood traits									
brood size	7.67	1.72	201	8.96	1.27	107	7.35	1.58	198
brood sex ratio	0.49	0.18	201	0.48	0.17	107	0.49	0.18	198
p(second brood)	0.09	0.29	201	0.32	0.47	107	0.12	0.32	198
adult survival									
female survival	0.15	0.36	201	0.39	0.49	107	0.33	0.47	198
male survival	0.15	0.36	189	0.26	0.44	115	0.34	0.47	203

Appendix S3. Annual variation in the probability of producing a second clutch

Model summary of hierarchical models examining the probability of producing a second clutch in relation to the plot social environment (density, 'D' and sex ratio, 'SR') and date in a great tit population in the Netherlands for three study years (2005, 2006 and 2007). In these models, density treatment ('decreased/increased') was fitted as a factor with 'decreased' chosen as reference category. Date (in April days), natural density and natural plot sex ratio centred around population mean, were fitted as fixed effects. Nests and plots were fitted as nested random effects.

Parameters	level	β	s.e. (β)	χ^2	df	P
2005 (n=224 broods)						
intercept		-2.623	0.617			
date	<i>nest</i>	-0.228	0.067	11.56	1	<0.001
2006 (n=158 broods)						
intercept		-0.82	0.192			
natural D	<i>plot</i>	-0.016	0.005	10.21	1	0.001
natural plot SR	<i>plot</i>	12.855	5.485	5.49	2	0.019
date	<i>nest</i>	-0.273	0.07	15.02	1	<0.001
2007 (n=242 broods)						
intercept		-1.602	0.269			
natural D	<i>plot</i>	-0.032	0.009	12.09	1	<0.001
D	<i>plot</i>	-1.575	0.536	8.64	1	0.003
natural plot SR	<i>plot</i>	11.043	4.861	5.16	1	0.023
date	<i>nest</i>	-0.105	0.05	4.3	1	0.038