

University of Groningen

Enantioselective synthesis of almorexant via iridium-catalysed intramolecular allylic amidation

Fananas Mastral, Martin; Teichert, Johannes F.; Fernandez-Salas, Jose Antonio; Heijnen, Dorus; Feringa, Ben L.

Published in:
Organic & Biomolecular Chemistry

DOI:
[10.1039/c3ob40655e](https://doi.org/10.1039/c3ob40655e)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Fananas Mastral, M., Teichert, J. F., Fernandez-Salas, J. A., Heijnen, D., & Feringa, B. L. (2013). Enantioselective synthesis of almorexant via iridium-catalysed intramolecular allylic amidation. *Organic & Biomolecular Chemistry*, 11(27), 4521-4525. <https://doi.org/10.1039/c3ob40655e>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.


ELECTRONIC SUPPLEMENTARY INFORMATION

Enantioselective Synthesis of Almorexant via Iridium-catalysed Intramolecular Allylic Amidation


Martín Fañanás-Mastral, Johannes F. Teichert, José Antonio Fernández-Salas, Dorus Heijnen and Ben L.
Feringa*


*Stratingh Institute for Chemistry, University of Groningen, Nijenborgh 4, 9747 AG, Groningen, The
Netherlands.*

NMR Spectra:





¹H NMR







APT







¹⁹F NMR







¹H NMR







¹H NMR







^{13}C NMR






¹⁹F NMR






Almorexant 1


¹H NMR






Almorexant 1

¹³C NMR





Almorexant 1

¹⁹F NMR

