

University of Groningen

Cuticular hydrocarbon divergence in the jewel wasp *Nasonia*

Buellesbach, J.; Gadau, J.; Beukeboom, L. W.; Echinger, F.; Raychoudhury, R.; Werren, J. H.; Schmitt, T.

Published in:
Journal of Evolutionary Biology

DOI:
[10.1111/jeb.12242](https://doi.org/10.1111/jeb.12242)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Buellesbach, J., Gadau, J., Beukeboom, L. W., Echinger, F., Raychoudhury, R., Werren, J. H., & Schmitt, T. (2013). Cuticular hydrocarbon divergence in the jewel wasp *Nasonia*: Evolutionary shifts in chemical communication channels? *Journal of Evolutionary Biology*, 26(11), 2467-2478.
<https://doi.org/10.1111/jeb.12242>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.




Fig. S2: Comparison of average CHC ratios per sex and species relative to the total peak area sums (percentages). The CHC compounds are categorized and summarized by their respective compound classes (see Table S1). Unknown Hydrocarbons (in white) are placed at the end of the bars. NV = *N. vitripennis* ($N_f = 19$, $N_m = 13$), NG = *N. giraulti* ($N_f = 18$, $N_m = 14$), NO = *N. oneida* ($N_f = 17$, $N_m = 11$), NL = *N. longicornis* ($N_f = 17$, $N_m = 11$), TS = *Trichomalopsis sarcophagae* ($N_f = 14$, $N_m = 12$). N_f refers to the sample size for females, N_m to the sample size for males, 150 CHC profiles analyzed in total for calculating the ratios per compound class.