

University of Groningen

Bronchoscopic Lung Volume Reduction with Endobronchial Valves Exclusively of the Middle Lobe in Patients with Emphysema

Klooster, Karin; Van Dijk, Marlies; Koster, T. David; Hartman, Jorine E.; Slebos, Dirk Jan

Published in:
Journal of Bronchology and Interventional Pulmonology

DOI:
[10.1097/LBR.0000000000000906](https://doi.org/10.1097/LBR.0000000000000906)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2023

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Klooster, K., Van Dijk, M., Koster, T. D., Hartman, J. E., & Slebos, D. J. (2023). Bronchoscopic Lung Volume Reduction with Endobronchial Valves Exclusively of the Middle Lobe in Patients with Emphysema. *Journal of Bronchology and Interventional Pulmonology*, 30(2), 192-195.
<https://doi.org/10.1097/LBR.0000000000000906>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Bronchoscopic Lung Volume Reduction With Endobronchial Valves Exclusively of the Middle Lobe in Patients With Emphysema

Karin Klooster, PhD, Marlies van Dijk, MD, T. David Koster, MD,
Jorine E. Hartman, PhD, and Dirk-Jan Slebos, MD, PhD

Key Words: endobronchial valves, emphysema, COPD, lung volume reduction, middle lobe, hyperinflation

(*J Bronchol Intervent Pulmonol* 2023;30:192–195)

Bronchoscopic lung volume reduction (BLVR) treatment using 1-way endobronchial valves has been shown to significantly improve pulmonary function, exercise tolerance, and quality of life, and also may lead to a survival benefit, in carefully selected chronic obstructive pulmonary disease (COPD) patients with severe emphysema and hyperinflation.^{1–7}

The 1-way endobronchial valves are placed in the most emphysematous lobe. In previous randomized controlled trials, the middle lobe was almost always treated in combination with the right upper lobe.

Only 1 single case has been published before on middle lobe treatment in an emphysema patient, with in that same paper a summary of 5 additional patients treated for a bulla in the middle lobe.⁸ However, to our knowledge, this is the first analysis of patients with emphysema who underwent BLVR with Zephyr endobronchial valves (PulmonX) exclusively in the right middle lobe. Our aim was to investigate the

efficacy of BLVR with endobronchial valves exclusively of the middle lobe in patients with emphysema.

From January 2008 till February 2021, 15 patients (53% male) with COPD underwent a middle lobe treatment at our center (~3% of the total number of patients treated with valves). All patients were included in the analysis and all provided written informed consent for the scientific use of their data.

At baseline and after endobronchial valve treatment, quantitative analysis was performed on high-resolution computed tomography (HRCT) scans (LunQ; Thirona). Furthermore, pulmonary function tests, quality of life questionnaires (Saint George Respiratory Questionnaire), and 6-minute walk tests were performed at baseline and 6 months after treatment. To test for differences between baseline and follow-up, a Wilcoxon signed-rank test was used. Data is described as median (minimum to maximum). All statistical analyses were performed using IBM SPSS Statistics, version 23 (IBM Corp.), and *P* values <0.05 were considered statistically significant.

Baseline characteristics were: median age 63 years (range: 36 to 74 y), forced expiratory volume in 1 second 24 (16 to 51) % of predicted, residual volume of 228 (162 to 392) % of predicted, 6-minute walk distance 323 m (165 to 539 m), and SGRQ total score of 46 points (37 to 78 points). Target lobe volume was 908 mL (436 to 2510 mL), and 67% had a >95% fissure integrity of the right middle lobe fissure [median fissure score of 99% (54% to 100%)]. The destruction of the middle lobe, measured on the HRCT scan was 58% (46% to 84%) on voxel density –950 HU.

Valve placement was performed under general anesthesia in 12 patients, using deep conscious sedation in 2 patients, and under local anesthesia 1 patient. Before valve placement,

Received for publication July 5, 2022; accepted October 4, 2022.
From the Department of Pulmonary Diseases, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands.

K.K.: helped in conception and design of the study, acquisition of data, analysis, and interpretation of data analysis, writing original draft preparation. M.v.D., T.D.K., J.E.H., and D.-J.S.: helped in conceptualization, interpretation of data analysis, and reviewing and editing of the draft. All the authors meet the definition of an author as stated by the International Committee of Medical Journal Editors, and all have seen and approved the final manuscript.

Disclosure: K.K. received travel reimbursement from PulmonX Inc. D.-J.S. is a physician advisor for PulmonX Corp. and a PI for clinical studies funded by PulmonX Corp. The remaining authors declare no conflict of interest or other disclosures.

Correspondence: Karin Klooster, PhD, Department of Pulmonary Diseases, AA11, University Medical Center Groningen, P.O. Box 30.001, Groningen 9700 RB, The Netherlands (e-mail: k.klooster@umcg.nl).
Copyright © 2022 Wolters Kluwer Health, Inc. All rights reserved.
DOI: 10.1097/LBR.0000000000000906

Chartis assessment was performed in 6 patients (40%), in 3 patients the outcome was no collateral ventilation and in 3 patients the outcome was undeterminable due to no flow output. The median procedure time was 3 minutes (1 to 7 min) and in every patient, one endobronchial valve size 5.5 was placed in the entrance of the combined segment RB4/5. The median hospital stay was 1 night (1 to 4 nights) after treatment.

At 6 months follow-up, outcomes significantly improved and were clinically meaningful (Table 1). One pneumothorax occurred 5 days posttreatment and was treated with chest drainage for 9 days. There were no other serious adverse events and no patients died.

Our results showed that BLVR treatment with endobronchial valves exclusively in the middle lobe, leads to a statistically significant and clinically meaningful reduction in target lobar volume, lung function, exercise tolerance, and quality of life (Fig. 1). The results of our study are consistent with the results of 6 previously published case reports, where there was clinical improvement in 5 of 6 patients.⁸

The target lobe with the most pronounced destruction is most often an upper or lower lobe, and therefore chosen as target lobe for lung volume reduction with valves. The right middle lobe is rarely the most destroyed lobe and the lobe volume is usually much smaller compared with the upper and lower lobes. Therefore, closing of the middle lobe with endobronchial valves may have too little effect. However, in this study, we showed that if the middle lobe shows the most pronounced destruction, that treatment of a relatively small volume [middle lobe volume 908 (436 to 2510 mL)] results in clinically relevant outcomes. Furthermore, our analysis demonstrates that occlusion of the middle lobe

leads to significant target lobar volume reduction, with 12 of 14 patients (86%) reaching the minimal clinically important difference of > -22.4% for target lobar volume reduction.⁹

The absence of collateral ventilation between target lobe and ipsilateral lobe is key for inducing an atelectasis after endobronchial valve placement. However, measurement of collateral ventilation in the middle lobe is challenging, due to a “no flow” output. Fissure integrity is a surrogate for the absence or presence of collateral ventilation.¹⁰ In our cohort, 67% had a fissure completeness score > 95%, suggesting absence of collateral ventilation.¹¹

The first cases were performed with doing a Chartis measurement. At that time, we did not have the quantitative computed tomography (CT) analysis reports available, and therefore no information about the fissure integrity. Since we had the quantitative CT and the fissure was above 95%, we decided not to perform Chartis. Other reasons not to perform a Chartis are: in all cases, the middle lobe was very much expanded and overinflated, so the change that there will be collateral ventilation is really low. Furthermore, also to get a reliable flow output from such an overexpanded lobe, was very difficult. And the time and costs of the Chartis assessment was another argument to just place the valve. By placing one valve, the air cannot directly go to the middle lobe, so the lobe will be less hyperinflated and the patient can still have the benefit of the treatment. In our experience over the years, it became clear that at least an 80% middle lobe fissure integrity score can still be worthwhile to schedule for a Chartis assessment.

Posttreatment CT scan showed significant target lobar volume reduction with a high

TABLE 1. Effectiveness Outcomes: Change From Baseline to 6 Months After Endobronchial Valve Treatment Exclusively in the Middle Lobe (N=15)

	Baseline	Absolute Change	Relative Change
TLV	908 mL (436-2510 mL)	-835 mL (-2510 to -68 mL)	-98% (-100% to -16%)
FEV ₁	0.77 L (0.38-2.28 L)	+60 mL (-130 to +730)	+9.1% (-11.5% to +98.6%)
FVC	2.54 L (1.20-7.55 L)	+310 mL (-210 to +2240 mL)	+11.5% (-4.7% to 102.3%)
RV	4.99 L (3.69-7.58 L)	-305 mL (-1770 to +30 mL)	-6.35% (-38% to 0%)
TLC	7.94 L (6.17-13.74 L)	-140 mL (-1110 to +170 mL)	-1.4% (-11% to +2.7%)
SGRQ	46 points (37-78 points)	-6.5 points (-35 to +6 points)	
6MWD	323 m (165-539 m)	+40 m (-16 to +155 m)	

Values are presented in median (minimum-maximum). All outcomes were statistically significant compared with baseline ($P < 0.05$).

FEV₁ (N = 15) indicates forced expiratory volume in 1 second; FVC, forced vital capacity; 6MWD (N = 11), 6-minute walk distance; RV (N = 14), residual volume; SGRQ (N = 13), Saint George Respiratory Questionnaire total score; TLC (N = 15), total lung capacity; TLV, target lobe volume; TLVR (N = 14), target lobe volume reduction.




FIGURE 1. Minimal clinically important difference (MCID) responder results at 6 months after endobronchial valve treatment exclusively in the middle lobe. The MCID is shown in the graph as a black horizontal line. Response rates were calculated by counting the number of patients for whom the change at 6 months met or exceeded the MCID. Individual outcomes per patient in change from baseline. FEV₁ (N=15) indicates forced expiratory volume in 1 second (MCID > +10%); FVC, forced vital capacity; 6MWD (N=11), 6-minute walk distance (MCID >26 m); RV (N=14), residual volume (MCID >-310 mL); SGRQ (N=13), Saint George Respiratory Questionnaire total score (MCID >-7.1 points); TLC (N=15), total lung capacity (MCID RV/TLC >-2.8%); TLVR (N=14), target lobe volume reduction (MCID TLVR >-22.4%); TLV, target lobe volume.

responder rate. In 4 patients, the fissure completeness score was <80%. In 2 of these patients, only slight volume reduction was obtained after treatment. However, 2 patients did show significant volume reduction (-61% and -100% target lobe volume reduction).

Despite the fact that our cohort included only a small sample size, the significant target lobar volume reduction of the middle lobe suggests that with careful patient selection treatment benefit can be achieved. All patients in our cohort had advanced COPD, remained highly symptomatic despite receiving optimal medical, had severe hyperinflation [residual volume of 228 (162 to 392) % of predicted], and a “clear” treatment target lobe with having pronounced destruction of lung tissue in the middle lobe.

To conclude, the benefit of BLVR treatment with Zephyr endobronchial valves is not limited

to only patients with upper or lower lobe predominant emphysema. Endobronchial valve treatment exclusively in the middle lobe in patients with emphysema can improve lung function, exercise tolerance, and quality of life. However, careful selection of patients with hyperinflation and a clear treatment target lobe identified on HRCT is key.

REFERENCES

1. Klooster K, ten Hacken NHT, Hartman JE, et al. Endobronchial valves for emphysema without interlobar collateral ventilation. *N Engl J Med.* 2015;373:2325–2335.
2. Criner GJ, Sue R, Wright S, et al. A multicenter RCT of Zephyr(R) Endobronchial Valve Treatment in Heterogeneous Emphysema (LIBERATE). *Am J Respir Crit Care Med.* 2018;198:1151–1164.
3. Kemp SV, Slebos DJ, Kirk A, et al. A multicenter, prospective, randomized, controlled trial of endobronchial valve treatment vs standard of care in

Downloaded from http://journals.lww.com/bronchology by BMDM56PHKav1zEoun1tIQIN4a+kLLHEZgbsIH04XMIOHC ywCX1AMwvQpIIQIHd3D00DRyTTSF14C13VC1y0abgqZXdwmKZBYtws= on 06/26/2023

- heterogeneous emphysema (transform). *Am J Respir Crit Care Med.* 2017;196:1535–1543.
4. Valipour A, Slebos DJ, Herth F, et al. Endobronchial valve therapy in patients with homogeneous emphysema results from the IMPACT study. *Am J Respir Crit Care Med.* 2016;194:1073–1082.
 5. Criner GJ, Delage A, Voelker K, et al. Improving lung function in severe heterogenous Emphysema with the Spiration Valve System (EMPROVE) a multicenter, open-label randomized controlled clinical trial. *Am J Respir Crit Care Med.* 2019;200:1354–1362.
 6. Klooster K, Hartman JE, Ten Hacken NHT, et al. Improved predictors of survival after endobronchial valve treatment in patients with severe emphysema. *Am J Respir Crit Care Med.* 2017;195:1272–1274.
 7. Hartman JE, Welling JBA, Klooster K, et al. Survival in COPD patients treated with bronchosco-
pic lung volume reduction. *Respir Med.* 2022;196:106825.
 8. Villeneuve T, Fumat R, Héluain V, et al. Response to exclusive right middle lobe treatment with endobronchial valves: a case report. *Breathe.* 2021;17:210108.
 9. Welling JBA, Hartman JE, van Rikxoort EM, et al. Minimal important difference of target lobar volume reduction after endobronchial valve treatment for emphysema. *Respirology.* 2018;23:306–310.
 10. Gesierich W, Samitas K, Reichenberger F, et al. Collapse phenomenon during Chartis collateral ventilation assessment. *Eur Respir J.* 2016;47:1657–1667.
 11. Klooster K, Koster TD, Ruwwe-Glösenkamp C, et al. An integrative approach of the fissure completeness score and chartis assessment in endobronchial valve treatment for emphysema. *Int J Chron Obstruct Pulmon Dis.* 2020;15:1325–1334.