

University of Groningen

## Age-related differences in women's foot shape

Ansuategui Echeita, Jone ; Hijmans, Juha M.; Smits, Sharon; Van der Woude, Lucas H. V.; Postema, Klaas

*Published in:*  
Maturitas

*DOI:*  
[10.1016/j.maturitas.2016.09.001](https://doi.org/10.1016/j.maturitas.2016.09.001)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Final author's version (accepted by publisher, after peer review)

*Publication date:*  
2016

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Ansuategui Echeita, J., Hijmans, J. M., Smits, S., Van der Woude, L. H. V., & Postema, K. (2016). Age-related differences in women's foot shape. *Maturitas*, 94, 64-69.  
<https://doi.org/10.1016/j.maturitas.2016.09.001>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Age-related differences in women's foot shape

Jone Ansuategui Echeita,<sup>a</sup> Juha M. Hijmans,<sup>b</sup> Sharon Smits,<sup>b</sup> Lucas H.V. Van der Woude,<sup>a, b</sup>  
and Klaas Postema<sup>b</sup>

<sup>a</sup> University of Groningen, University Medical Center Groningen, Center for Human Movement Sciences, Groningen, the Netherlands. Mailing address: PO Box 196, 9700 AD Groningen, The Netherlands.

<sup>b</sup> University of Groningen, University Medical Center Groningen, Department of Rehabilitation Medicine, Groningen, the Netherlands. Mailing address: PO Box 30001, 9700 RB Groningen, The Netherlands.

## **Corresponding author:**

Juha M. Hijmans, University of Groningen, University Medical Center Groningen, Department of Rehabilitation Medicine, Groningen, the Netherlands. Mailing address: PO Box 30001, 9700 RB Groningen, The Netherlands. E-mail: [j.m.hijmans@umcg.nl](mailto:j.m.hijmans@umcg.nl)

## **ABSTRACT**

**Purpose:** Describe age-related differences in women's foot shape using a wide range of measurements and ages.

**Study design:** Cross-sectional, observational study.

**Main outcome measurements:** Six foot-shape measurements of each foot: foot lengths, ball widths, ball circumferences, low instep circumferences, high instep circumferences, and heel instep circumference.

**Results:** 168 women from 20 to over 80 years of age, divided into seven age categories, were included. Older women had significantly greater foot-shape measurements, even after adjusting for Body Mass Index. Ball widths increased 3.1-4.0mm per decade, ball circumferences 5.6-7.4mm per decade, high instep circumferences 0.4-4.8mm per decade, and heel instep circumferences 1.8-1.9mm per decade. Ball widths, ball circumferences, and left high instep circumference plateaued in the 70-75 years-of-age category, and decreased in the oldest age category. For low instep circumference, age did not prevail significantly over Body Mass Index. Foot length was not associated with age.

**Conclusion:** This study described women's progressive foot-shape changes with age. The findings provide a better understanding of foot-shape changes, mainly found in the forefoot. It demonstrates that for a good fit, shoe design for older adults and for younger adults should differ.

## **INDEX TERMS**

Foot, Foot pain, Ageing, Female, Ill-fitting

## **ABBREVIATIONS**

FL, foot length; BW, ball width; BC, ball circumference; LI, low instep circumference; HI, high instep circumference; HIC, heel instep circumference.

## 1. INTRODUCTION

Foot pain is present in 20-30% of older adults, displaying a higher prevalence in females than in males [1-4], and in 10-14% of young people aged between 12 and 19 years [5, 6]. Some structural conditions are related to foot pain. Such is the case for plantar fasciitis, hallux valgus, toe deformities, metatarsalgia, calluses and corns, bunions, and ingrowing toenails [1, 7, 8]. These structural conditions may result from friction, repetitive stress in forefoot or heel, toe adaptations to shoe shape, pressure spots on toes, and/or changes in plantar pressure distribution due to wearing an ill-fitting shoe [9-12].

Evidence has shown that individuals with foot pain and foot-morphology problems wore inadequate shoes, and that these were mainly women. The main reason demonstrated for this situation was the use of too small and narrow footwear in relation to foot size [9, 12-14]. In the case of women, the use of high heels was an additional source of foot pain, even later in life [2]. Because off-the-shelf shoes may be based on an “average” foot shape for adult people, the possible effects of age on foot shape may not be taken into consideration in the design of shoes. As a result, incorrect shoe designs may be used, which, in turn, lead to a poor fit and consequent foot pain.

A few studies have investigated the potential age-related changes in foot shape. One study analyzed the differences in foot structure and function between young adults and older adults [15]. In so doing, they demonstrated that older feet are flatter than younger feet. This finding was also supported by another study, one based on the assumption that a better understanding of foot morphology would improve shoe fit [16]. Through their anthropometric study of the foot, they showed that foot circumference was markedly larger in the older group. These two studies have shown differences in foot shape between young and older adults, or young adults, adults and older adults; presenting a limited evidence on the progressive evolution of feet. Another study examined changes, as a result of ageing, in the

size and shape of the foot [17]. Their findings indicated that older Japanese individuals had wider feet than younger groups. Nevertheless, that study only compares a limited number of measurements among age categories. As a result, an overall description of the genuine process of ageing in feet is lacking.

In response to the limitations found in the literature, the aim of this study was to describe women's foot-shape evolution using a wide range of foot measurements and age categories. In particular, our hypothesis is that a woman's foot shape is continuously changing throughout all stages of adulthood. To test this, several foot-shape measurements were taken from different age categories, with gaps between the age groups in order to highlight the changes among contiguous groups.

## **2. METHODS**

### **2.1. Study design and setting**

This is a cross-sectional, observational study that took place between October 2013 and March 2014 at the University Medical Center Groningen (UMCG), the Netherlands. Participants were recruited through advertising material in supermarkets, public places, homes for older people, and local newspapers. The measurement process was conducted once and lasted 45 minutes. Participants were allowed to rest between measurements, if they needed to. The study is part of a larger project, "The effect of age on foot structure, foot complaints, plantar pressure, and center of pressure in adult women," and has the approval of the Medical Ethical Committee of the UMCG (Number: 2013-225). The project was conducted according to the principles of the Declaration of Helsinki (October 2000) and in accordance with the Medical Research Involving Human Subjects Act (WMO).

## **2.2. Participants**

In the current study, women from the age of 20 until over 80, from the north of the Netherlands, took part. They were divided into categories with gaps of five years between them, so that differences among the groups could be highlighted. Subjects were included if they: (1) were Caucasian women, (2) fitted into one of the age categories, and (3) presented a self-reported ability to walk at least ten meters without any walking aid. Contrarily, participants were excluded if they: (1) reported medical conditions that had a major influence on gait (Parkinson's disease or stroke); (2) had undergone a lower limb amputation; and/or (3) currently used orthopedic footwear (the use of insoles was accepted, however). All participants signed the informed consent form.

## **2.3. Measurements**

### *2.3.1. Procedure*

Before the actual assessment, the participants met with the measurer. In these meetings the eligibility of the participants was determined, the measuring procedures were explained, signed informed consents were collected, and participants were given the opportunity to ask questions. Throughout the whole measuring process, participants were barefoot and instructed to maintain a standing position. First, height and weight were measured using a fixed tape on the wall and an analogue weight scale. Then, trained testers took foot-shape measurements as recommended by the orthopedic shoe technicians in the Orthopedic Instrument Manufacturer (OIM) protocol. According to this protocol, measurements were manually taken using a tape, a sliding caliper, and the Brannock device (The Brannock Device®, Liverpool, NY, USA). The tape in the Brannock Device was replaced with a millimeter scale for this study. Both the sliding caliper and tape measure were placed close to the foot over the points marked for foot-shape measurements; under no circumstances were they ever stretched.

### 2.3.2. Foot marking and measurement

*(Insert Table 1 about here)*

Before proceeding to take the measurements, predefined locations were marked with a pen on the skin of the foot. These markers were at the level of 1<sup>st</sup> metatarsal-phalangeal (MTP-I) and 5<sup>th</sup> metatarsal-phalangeal (MTP-V) joints, proximal base of the heads of 1<sup>st</sup> metatarsal (MT-I) and 5<sup>th</sup> metatarsal (MT-V) bones, navicular bone, base of MT-V bone, front of the subtalar joint (bending point between foot and lower leg), and posterior end point of the calcaneus bone. Once the markers were set, the foot-shape measurements were taken (Table 1). One tester took all the measurements for the same subject, beginning with the right foot and followed by the left.

A graphic description of foot markers and foot-shape measurements is shown in Figure 1.

*(Insert Figure 1 about here)*

**Figure 1.** Foot markers and foot-shape measurements.

A, antero-lateral side of the foot; B, plantar side of the foot.

Marks (+), indicate markers placed prior to measurements.

FL, foot length; BW, ball width; BC, ball circumference; LI, low instep circumference; HI, high instep circumference; HIC, heel-instep circumference.

## 2.4. Power calculation

On the basis of the data from a pilot study with four different age categories, a pooled Standard Deviation (SD) of 1.53, giving an effect size of 0.296, was assumed. With  $\alpha$  defined as 0.05, power as 0.8, and using seven age categories, a sample size of 168 participants needed for multiple regression analysis was calculated.



## **2.5. Statistical analysis**

Analyses were performed using SPSS software version 22.0 (IBM Corp., NY, USA). For details on the participant's physical (height, weight, and Body Mass Index [BMI]) and foot-shape characteristics, a descriptive statistical analysis was undertaken. To determine the predictive significance of age on foot-shape differences, a series of simple as well as multiple linear regression models were analyzed. Foot-shape measurements (FL, BW, BC, LI, HI, and HIC) were the outcome variables, and the age of the women was the predictor. In the multiple linear regression analyses, the squared term of age was added as a predictor to ascertain a possible plateau effect. Furthermore, the participant's BMI association with age and foot shape was examined using the Spearman's rho correlation in order to establish a possible confounding effect. All variables were included as continuous variables. The predictors and the confounder were entered stepwise forward into the final multiple linear regression model, and included if  $p < 0.05$ . All statistical analyses were conducted at 5% significance and 95% confidence intervals, and adjusted coefficients of determination were reported.

## **3. RESULTS**

### **3.1. Description of study sample**

In the current study, 168 women participated; full data for all of them was available. Women were divided into seven categories, depending on their age at the time of the measurement, with the same number of participants in each: 20-25, 30-35, 40-45, 50-55, 60-65, 70-75, and >80. Participants height ranged from 145 to 189 cm and weight from 44.2 to 126.7 kg; BMI Mean (SD) was 25.67 (5.20), with almost half of the participants showing

overweight (47%, BMI  $\geq$  25 kg/m<sup>2</sup>). A description of foot-shape measurements per age category is displayed in Table 2.

*(Insert Table 2 about here)*

### **3.2. Univariate linear regression analyses**

The results from the univariate linear regression analyses for foot-shape measurements indicate that older age is related to larger foot-shape measurements. Age was a significant positive predictor for almost all foot-shape measurements in both feet: ball widths, ball circumferences, low instep circumferences, high instep circumferences, and heel instep circumferences (Table 3). Age was also a positive but non-significant predictor for foot length measurements. Age explained from 2.3% to 15.9% of foot-shape variance. In general, being older was a predictor of larger foot-shape measurements.

*(Insert Table 3 about here)*

### **3.3. Multivariate linear regression analyses**

Age and five foot-shape measurements (BW, BC, LI, HIs, and HIC) showed significant correlations with BMI; therefore, it was included in the final multivariate linear regression models. The results of foot-shape measurement changes are depicted in Table 4. Age remained a statistically significant positive predictor for ball widths, ball circumferences, left high instep circumference, and heel circumferences, even when it was adjusted for BMI. The coefficients of the squared term of age indicated a plateau effect in ball widths, ball circumferences, and left high instep circumference. The models explained between 15% and 32.8% of the measurement's variation. Overall, the regression coefficients showed an increase in the measurements over time in most of the foot-shape measurements, although they plateaued for the majority of the measurements.

*(Insert Table 4 about here)*

#### **4. DISCUSSION**

The purpose of the current study was to describe the ageing process on women's foot shape. The results showed that older women had a significant increase in most foot-shape measurements compared to younger women. This relationship remained after adjusting for BMI. Ball width increased between 3.1 and 4 mm per decade, ball circumference between 5.6 and 7.4 mm per decade, high instep circumferences between 0.4 and 4.8 mm per decade, and heel instep circumferences between 1.8 and 1.9 mm per decade. Nevertheless, ball width, ball circumference, and left high instep circumference plateaued in the 70-75 years age category, and even decreased in the oldest age category. Remarkably, for low instep circumferences, age did not prevail significantly over BMI. Foot length was not associated with age. Women revealed a continuous widening of mainly the forefoot over the years.

In agreement with previous research, the current study has proven that foot shape differs from young adults, to adults, and to older adults [15, 16]. Our study also showed the proportion and location of the changes in foot shape with age. The changes in foot morphology can be explained by the changes in body composition due to the aging process. Muscle strength peaks at 20-30 years of age; from this moment on minor declines occur, and at around 60 years of age a major decline of 1-1.5% per year follows. The underlying reason for this decline in strength is a decrease in the size of muscle mass [18]. In addition, tendons in advancing age change their tensile properties, becoming more compliant [19]. Overall, because of the change in the structure of muscle and tendons with age, foot morphology maintained by these soft tissues is modified, and a wider foot can therefore develop. Moreover, individuals at an older age are more likely to be obese ( $BMI \geq 30 \text{ kg/m}^2$ ) as body

weight and BMI peak at 50-59 years of age, and these higher levels of BMI are maintained or decline slightly in the ensuing years [20]. Additionally, overweight and/or obesity have been found to be associated with flatter feet due to the increase in adiposity in the midfoot and altered plantar fat pad [21, 22]. This is consistent with our results, where differences in BMI among participants were significant in all the models, where included. Moreover, BMI showed a major influence in both right and left low-instep circumferences and right high-instep circumference measurements, which could be explained by the increase in adiposity in the midfoot. In addition, flatter wider feet have been also associated with older individuals, when compared to younger adults [15, 17]. Therefore, we suspect that the association of age with BMI could be the underlying reason for these changes.

A singular finding was the significant plateau effect in the 70-75-year age category, and regression in the oldest age category for some of the measurements. Considering that our sample was restricted to women without conditions that have a major influence on gait or who did not use orthopedic footwear, the oldest age category might have been comprised essentially by relatively healthy older women. This, in turn, might explain the decrease in ball width, ball circumference, and left high-instep circumference measurements, given that this age category might have presented higher rates of relatively healthy participants than the other categories.

The difference found between right and left high-instep circumferences is remarkable. The asymmetry within an individual's feet has been described in literature, and, typically, the left foot tends to be larger than the right foot in right-handers [23]. We believe foot laterality might be a viable additional explanation for high-instep discrepancy; however, since laterality was not measured, this fact could not be demonstrated.

To our knowledge this is the first study describing women's foot shape for a wide range of ages. Major strengths of the study are the equal number of participants per age category, which limits bias toward any of the groups measured. Moreover, the groups were measured in a 5-year age bracket per decade for a more defined difference in foot-shape measurements between groups.

There are some limitations that need to be considered. One limitation would be that various trained measurers were involved; thus, we cannot discard the possibility of differences between them. One other limitation would be the reliability of measurement points. Despite anatomical foot-bone projections for easier localization of the measurement point, it might be difficult to repeat the exact same points. One additional limitation would be the absence of repetition in the measurements. Each measurement was taken only once; therefore, there was no average, and this might bias the outcome. One final limitation would be the selection of the participants involved. Since all the participants were healthy volunteers recruited through advertising material, it might be possible that participants in the older age categories were rather unrepresentative for their age cohort.

We consider it important to emphasize that age, the squared term of age, and BMI, although they are significant predictors for most of the measurements, only explain between 15% and 32.8% of foot-measurement variance. As a result, future research should aim to analyze a wider number of factors associated with foot-shape change. Possible comorbidities as diabetes, structural conditions as hallux valgus, or pregnancy effects, could be some of the factors to take into consideration.

## **5. CONCLUSION**

Growing numbers of older people have foot pain, with the majority of them being women. The need for individuals to be able to independently carry out the activities of daily living is a priority; however, this can be inhibited due to foot pain and ill-fitting shoes. The current study not only shows that the foot shape of women becomes wider with age but also how, since ball width, ball circumference, high instep circumference, and heel instep circumference were found to be larger in older age categories. Furthermore, ball widths, ball circumferences, and left high-instep circumference plateaued at the 70-75-year age category and decreased in the oldest age category. Finally, foot length was not related to age. The findings of the current study demonstrate the continuous change in foot shape with age; therefore, for a good fit, shoe design should be different for older adults as compared to younger ones.

## **ACKNOWLEDGEMENTS**

The authors wish to express their gratitude to Prof. P.U. Dijkstra for kindly providing statistical assistance; to Mr. J.N. Wijnholds, orthopedic shoe technician, for his recommendations and expertise concerning the measurements; and to the students who collaborated in this study.

## REFERENCES

1. Benvenuti F, Ferrucci L, Guralnik JM, Gangemi S, Baroni A. Foot pain and disability in older persons: an epidemiologic survey. *J Am Geriatr Soc* 1995; 43(5): 479–84.
2. Dufour AB, Broe KE, Kivell E, Walker AH, Nguyen UD, Gagnon DR, Hillstrom HJ, Hannan MT. Foot Pain: Is Current or Past Shoewear a Factor? The Framingham Foot Study. *Arthritis Rheum* 2009; 61(10), 1352–1358. doi:10.1002/art.24733
3. Hill CL, Gill TK, Menz HB, Taylor AW. Prevalence and correlates of foot pain in a population-based study: the North West Adelaide health study. *J Foot Ankle Res* 2008; 1, 2. doi:10.1186/1757-1146-1-2
4. Thomas MJ, Roddy E, Zhang W, Menz HB, Hannan MT, Peat GM. The population prevalence of foot and ankle pain in middle and old age: a systematic review. *Pain* 2011; 152(12), 2870–80. doi:10.1016/j.pain.2011.09.019
5. Rathleff MS, Roos EM, Olesen JL, Rasmussen S. *BMC Pediatrics* 2013; 13, 191. doi:10.1186/1471-2431-13-191
6. Spahn G, Schiele R, Hell AK, Klinger HM, Jung R, Langlotz A. The prevalence of pain and deformities in the feet of adolescents. Results of a cross-sectional study. *Z Orthop Ihre Grenzgeb* 2004; 142(4): 389–96. doi:10.1055/s-2004-822844
7. Chaiwanichsiri D, Janchai S, Tantisiriwat N. Foot disorders and falls in older persons. *Gerontology* 2009; 55: 296–302. doi:10.1159/000181149
8. Garrow AP, Silman AJ, Macfarlane GJ. The Cheshire Foot Pain and Disability Survey: a population survey assessing prevalence and associations. *Pain* 2004; 110(1-2): 378–84. doi:10.1016/j.pain.2004.04.019

9. Akhtar S, Choudry Q, Kumar R. Incorrectly fitting footwear and associated foot problems. *J Bone Joint Surg Br* 2008; 90-B (SUPP III): 495–496.
10. Edelstein JE. Foot care for the aging. *Phys Ther* 1988; 68: 1882–1886.
11. Goud A, Khurana B, Chiodo C, Weissman BN. Women’s musculoskeletal foot conditions exacerbated by shoe wear: an imaging perspective. *Am J Orthop* 2011; 40(4): 183–91.
12. Menz HB, Morris ME. Footwear characteristics and foot problems in older people. *Gerontology* 2005; 51(5): 346–51. doi:10.1159/000086373
13. Burns SL, Leese GP, McMurdo MET. Older people and ill-fitting shoes. *Postgrad Med J* 2002; 78: 344–346. doi:10.1136/pmj.78.920.344
14. Frey C, Thompson F, Smith J, Sanders M, Horstman H. American Orthopaedic Foot and Ankle Society Women’s Shoe Survey. *Foot Ankle Int* 1993; 14(2): 78–81. doi:10.1177/107110079301400204
15. Scott G, Menz HB, Newcombe L. Age-related differences in foot structure and function. *Gait Posture* 2007; 26(1): 68–75. doi:10.1016/j.gaitpost.2006.07.009
16. Tomassoni D, Traini E, Amenta F. Gender and age related differences in foot morphology. *Maturitas* 2014; 79(4): 421–427. doi:10.1016/j.maturitas.2014.07.019
17. Kouchi M. Foot Dimensions and Foot Shape: Differences Due to Growth, Generation and Ethnic Origin. *Anthropological Science* 1998; 106, 161–188. doi:10.1537/ase.106.Supplement\_161
18. Vandervoort AA. Aging of the human neuromuscular system. *Muscle Nerve* 2002; 25(1): 17–25. doi:10.1002/mus.1215



19. Narici MV, Maganaris CN. Adaptability of older people human muscles and tendons to increased loading. *J Anat* 2006; 208(4): 433–443. doi:10.1111/j.1469-7580.2006.00548.x
20. Villareal DT, Apovian CM, Kushner RF, Klein S. Obesity in older adults: Technical review and position statement of the American Society for Nutrition and NAASO, The Obesity Society. *Am J Clin Nutr* 2005; 82(5): 923–934. doi:10.1038/oby.2005.228
21. Hills AP, Hennig EM, Byrne NM, Steele JR. The biomechanics of adiposity-structural and functional limitations of obesity and implications for movement. *Obes Rev* 2002; 3(1): 35–43. doi:10.1046/j.1467-789X.2002.00054.x
22. Wearing SC, Grigg NL, Lau HC, Smeathers JE. Footprint-based estimates of arch structure are confounded by body composition in adults. *J Orthop Res* 2012; 30(8): 1351–1354. doi:10.1002/jor.22058
23. Peters M. Footedness: asymmetries in foot preference and skill and neuropsychological assessment of foot movement. *Psychol Bull* 1988; 103(2): 179–192. doi:10.1037/0033-2909.103.2.179

## TABLES

**Table 1.** Description of localization of foot-shape measurements of participants.

| Measurement                          | Location  | Device                       |
|--------------------------------------|---|------------------------------|
| <b>FL</b> Foot Length                | Distance from the horizontal line situated at the top of the first toe to the back end point of the calcaneal bone in a straight line | Brannock device              |
| <b>BW</b> Ball Width                 | Space between MTP-I and MTP-V joints  | Sliding caliper              |
| <b>BC</b> Ball Circumference         | Circumference at the level of MTP-I and MTP-V joints  |                              |
| <b>LI</b> Low-Instep Circumference   | Circumference at a proximal level of the heads of MT-I and MT-V bones   | Tape measure around the feet |
| <b>HI</b> High-Instep Circumference  | Circumference at the level of navicular and base of MT-V bones  |                              |
| <b>HIC</b> Heel Instep Circumference | Circumference at the front of the subtalar joint and the posterior end-point of calcaneus bone  |                              |

MTP-I, 1<sup>st</sup> metatarsal-phalanx; MTP-5, 5<sup>th</sup> metatarsal-phalanx; MT-I, 1<sup>st</sup> metatarsal; MT-V, 5<sup>th</sup> metatarsal.

**Table 2.** Description of foot-shape measurements (mm) of participants per age category; mean  $\pm$  SD are shown.

|            |          | <b>TOTAL</b>     | <b>20-25 YEARS</b> | <b>30-35 YEARS</b> | <b>40-45 YEARS</b> | <b>50-55 YEARS</b> | <b>60-65 YEARS</b> | <b>70-75 YEARS</b> | <b>80+ YEARS</b> |
|------------|----------|------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|------------------|
|            |          | (N = 168)        | (N = 24)           | (N = 24)           | (N = 24)           | (N = 24)           | (N = 24)           | (N = 24)           | (N = 24)         |
| <b>FL</b>  | <b>R</b> | 249.0 $\pm$ 11.2 | 244.3 $\pm$ 11.8   | 249.6 $\pm$ 9.8    | 248.4 $\pm$ 9.7    | 249.2 $\pm$ 12.5   | 250.7 $\pm$ 9.4    | 251.2 $\pm$ 11.8   | 249.3 $\pm$ 12.8 |
|            | <b>L</b> | 248.8 $\pm$ 11.5 | 244.0 $\pm$ 11.5   | 249.7 $\pm$ 10.7   | 248.0 $\pm$ 9.5    | 249.2 $\pm$ 13.1   | 250.9 $\pm$ 10.4   | 250.9 $\pm$ 11.8   | 249.0 $\pm$ 13.4 |
| <b>BW</b>  | <b>R</b> | 97.4 $\pm$ 5.8   | 94.1 $\pm$ 5.6     | 95.1 $\pm$ 4.1     | 97.8 $\pm$ 5.8     | 98.4 $\pm$ 5.2     | 98.4 $\pm$ 3.9     | 101.3 $\pm$ 6.7    | 96.7 $\pm$ 6.5   |
|            | <b>L</b> | 96.6 $\pm$ 5.6   | 93.4 $\pm$ 5.5     | 95.0 $\pm$ 4.4     | 96.9 $\pm$ 4.8     | 97.3 $\pm$ 6.2     | 97.3 $\pm$ 4.6     | 99.8 $\pm$ 5.7     | 96.7 $\pm$ 6.2   |
| <b>BC</b>  | <b>R</b> | 239.9 $\pm$ 12.2 | 233.1 $\pm$ 12.1   | 234.6 $\pm$ 8.4    | 240.1 $\pm$ 12.4   | 241.3 $\pm$ 11.9   | 244.5 $\pm$ 8.9    | 246.6 $\pm$ 14.2   | 239.2 $\pm$ 11.7 |
|            | <b>L</b> | 239.2 $\pm$ 11.9 | 232.9 $\pm$ 13.2   | 235.4 $\pm$ 8.8    | 239.8 $\pm$ 10.9   | 240.8 $\pm$ 13.7   | 241.5 $\pm$ 9.3    | 245.0 $\pm$ 12.3   | 239.3 $\pm$ 11.5 |
| <b>LI</b>  | <b>R</b> | 230.5 $\pm$ 10.8 | 227.0 $\pm$ 11.3   | 226.0 $\pm$ 8.5    | 230.7 $\pm$ 10.5   | 230.5 $\pm$ 11.6   | 232.8 $\pm$ 8.7    | 235.1 $\pm$ 12.3   | 231.6 $\pm$ 11.0 |
|            | <b>L</b> | 230.6 $\pm$ 10.9 | 227.4 $\pm$ 11.4   | 227.3 $\pm$ 9.7    | 231.3 $\pm$ 10.2   | 230.9 $\pm$ 12.6   | 231.3 $\pm$ 9.5    | 235.1 $\pm$ 10.9   | 231.1 $\pm$ 11.2 |
| <b>HI</b>  | <b>R</b> | 246.4 $\pm$ 11.2 | 241.3 $\pm$ 11.0   | 241.7 $\pm$ 12.8   | 247.8 $\pm$ 11.9   | 247.0 $\pm$ 8.6    | 249.2 $\pm$ 9.8    | 248.5 $\pm$ 10.9   | 249.5 $\pm$ 11.2 |
|            | <b>L</b> | 245.9 $\pm$ 12   | 239.6 $\pm$ 11.2   | 242.3 $\pm$ 12.2   | 248.1 $\pm$ 12.0   | 247.1 $\pm$ 9.8    | 248.1 $\pm$ 11.6   | 249.3 $\pm$ 12.7   | 246.9 $\pm$ 12.7 |
| <b>HIC</b> | <b>R</b> | 319.0 $\pm$ 16.1 | 308.0 $\pm$ 14.9   | 313.3 $\pm$ 15.4   | 318.8 $\pm$ 14.6   | 316.7 $\pm$ 10.3   | 323.9 $\pm$ 12.9   | 325.2 $\pm$ 15.8   | 327.2 $\pm$ 19.6 |
|            | <b>L</b> | 318.7 $\pm$ 15.7 | 307.6 $\pm$ 14.3   | 311.6 $\pm$ 13.7   | 318.3 $\pm$ 14.1   | 317.7 $\pm$ 10.0   | 323.6 $\pm$ 12.7   | 326.2 $\pm$ 17.4   | 326.1 $\pm$ 17.9 |

FL, foot length; BW, ball width; BC, ball circumference; LI, low-instep circumference; HI, high-instep circumference; HIC, heel-instep circumference.

R, right-foot measurements; L, left-foot measurements.

**Table 3.** Univariate linear regression of age for foot shape characteristics.

|            |          | <b>B</b> | <b>SE (B)</b> | <b>95% CI</b> |           | <b>Sig.</b> | <b>Adjusted R<sup>2</sup></b> |
|------------|----------|----------|---------------|---------------|-----------|-------------|-------------------------------|
|            |          |          |               | <b>LL</b>     | <b>UL</b> |             |                               |
| <b>FL</b>  | <b>R</b> | 0.07     | 0.04          | -0.01         | 0.15      | 0.094       | 0.011                         |
|            | <b>L</b> | 0.07     | 0.04          | -0.02         | 0.16      | 0.107       | 0.010                         |
| <b>BW</b>  | <b>R</b> | 0.07     | 0.02          | 0.03          | 0.11      | 0.001       | 0.054                         |
|            | <b>L</b> | 0.07     | 0.02          | 0.03          | 0.11      | 0.002       | 0.052                         |
| <b>BC</b>  | <b>R</b> | 0.16     | 0.05          | 0.07          | 0.25      | 0.001       | 0.065                         |
|            | <b>L</b> | 0.14     | 0.04          | 0.05          | 0.22      | 0.002       | 0.048                         |
| <b>LI</b>  | <b>R</b> | 0.12     | 0.04          | 0.04          | 0.20      | 0.004       | 0.043                         |
|            | <b>L</b> | 0.09     | 0.04          | 0.01          | 0.17      | 0.028       | 0.023                         |
| <b>HI</b>  | <b>R</b> | 0.14     | 0.04          | 0.06          | 0.22      | 0.001       | 0.056                         |
|            | <b>L</b> | 0.12     | 0.05          | 0.04          | 0.21      | 0.007       | 0.038                         |
| <b>HIC</b> | <b>R</b> | 0.30     | 0.06          | 0.19          | 0.41      | <0.001      | 0.139                         |
|            | <b>L</b> | 0.31     | 0.06          | 0.21          | 0.42      | <0.001      | 0.159                         |

FL, foot length; BW, ball width; BC, ball circumference; LI, low-instep circumference;

HI, high-instep circumference; HIC, heel-instep circumference.

R, right-foot measurements; L, left-foot measurements.

B, unstandardized regression coefficient; SE (B), standard error for the unstandardized regression coefficient; 95% CI, 95% confidence interval for unstandardized regression coefficient; LL, lower limit; UL, upper limit; Sig., significance.

**Table 4.** Final multivariate linear regression analyses of age, squared term of age (age<sup>2</sup>), and BMI for foot-shape measurements.

|           |          | B                | SE (B) | 95% CI |        | Sig.   | Adjusted R <sup>2</sup> |       |
|-----------|----------|------------------|--------|--------|--------|--------|-------------------------|-------|
|           |          |                  |        | LL     | UL     |        |                         |       |
| <b>BW</b> | <b>R</b> | Constant         | 78.99  | 3.23   | 72.61  | 85.37  | <0.001                  | 0.174 |
|           |          | Age              | 0.40   | 0.12   | 0.16   | 0.64   | 0.001                   |       |
|           |          | Age <sup>2</sup> | -0.003 | 0.001  | -0.005 | -0.001 | 0.003                   |       |
|           |          | BMI              | 0.31   | 0.09   | 0.14   | 0.48   | <0.001                  |       |
| <b>L</b>  | <b>R</b> | Constant         | 80.69  | 3.17   | 74.44  | 86.94  | <0.001                  | 0.150 |
|           |          | Age              | 0.31   | 0.12   | 0.08   | 0.54   | 0.010                   |       |
|           |          | Age <sup>2</sup> | -0.002 | 0.001  | -0.005 | 0.000  | 0.021                   |       |
|           |          | BMI              | 0.30   | 0.08   | 0.14   | 0.47   | <0.001                  |       |
| <b>BC</b> | <b>R</b> | Constant         | 197.45 | 6.48   | 184.65 | 210.25 | <0.001                  | 0.242 |
|           |          | Age              | 0.74   | 0.24   | 0.26   | 1.21   | 0.003                   |       |
|           |          | Age <sup>2</sup> | -0.006 | 0.002  | -0.010 | -0.002 | 0.006                   |       |
|           |          | BMI              | 0.91   | 0.17   | 0.57   | 1.24   | <0.001                  |       |
| <b>L</b>  | <b>R</b> | Constant         | 203.62 | 6.57   | 190.64 | 216.59 | <0.001                  | 0.184 |
|           |          | Age              | 0.56   | 0.24   | 0.08   | 1.05   | 0.023                   |       |
|           |          | Age <sup>2</sup> | -0.005 | 0.002  | -0.009 | 0.000  | 0.037                   |       |
|           |          | BMI              | 0.81   | 0.17   | 0.47   | 1.15   | <0.001                  |       |

|            |          |                  |        |       |        |        |        |       |
|------------|----------|------------------|--------|-------|--------|--------|--------|-------|
| <b>LI</b>  | <b>R</b> | Constant         | 202.08 | 3.68  | 194.83 | 209.34 | <0.001 |       |
|            |          | Age              | 0.02   | 0.04  | -0.05  | 0.10   | 0.559  | 0.267 |
|            |          | BMI              | 1.06   | 0.15  | 0.77   | 1.35   | <0.001 |       |
| <b>L</b>   |          | Constant         | 204.01 | 3.79  | 196.54 | 211.49 | <0.001 |       |
|            |          | Age              | -0.003 | 0.04  | -0.08  | 0.07   | 0.934  | 0.235 |
|            |          | BMI              | 1.04   | 0.15  | 0.74   | 1.34   | <0.001 |       |
| <b>HI</b>  | <b>R</b> | Constant         | 216.71 | 3.80  | 209.21 | 224.21 | <0.001 |       |
|            |          | Age              | 0.04   | 0.04  | -0.04  | 0.12   | 0.311  | 0.270 |
|            |          | BMI              | 1.08   | 0.15  | 0.77   | 1.38   | <0.001 |       |
| <b>L</b>   |          | Constant         | 205.97 | 6.31  | 193.50 | 218.44 | <0.001 |       |
|            |          | Age              | 0.48   | 0.24  | 0.12   | 0.94   | 0.045  | 0.263 |
|            |          | Age <sup>2</sup> | -0.004 | 0.002 | -0.008 | 0.000  | 0.050  |       |
|            |          | BMI              | 1.10   | 0.17  | 0.78   | 1.43   | <0.001 |       |
| <b>HIC</b> | <b>R</b> | Constant         | 274.12 | 5.29  | 263.67 | 284.56 | <0.001 |       |
|            |          | Age              | 0.18   | 0.05  | 0.07   | 0.28   | 0.002  | 0.312 |
|            |          | BMI              | 1.39   | 0.21  | 0.97   | 1.81   | <0.001 |       |
| <b>L</b>   |          | Constant         | 274.09 | 5.11  | 264.00 | 284.18 | <0.001 |       |
|            |          | Age              | 0.19   | 0.05  | 0.09   | 0.30   | <0.001 | 0.328 |
|            |          | BMI              | 1.34   | 0.21  | 0.94   | 1.75   | <0.001 |       |

BW, ball width; BC, ball circumference; LI, low-instep circumference; HI, high-instep circumference; HIC, heel-instep circumference.

R, right-foot measurements; L, left-foot measurements.

B, unstandardized regression coefficient; SE (B), standard error for the unstandardized regression coefficient; 95% CI, 95% confidence interval for unstandardized regression coefficient; LL, lower limit; UL, upper limit; Sig., significance.