

University of Groningen

## Ultra diffuse galaxies in the Hydra I cluster from the LEWISProject

Forbes, Duncan A.; Gannon, Jonah; Iodice, Enrichetta; Hilker, Michael; Doll, Goran; Buttitta, Chiara; Marca, Antonio La; Arnaboldi, Magda; Cantiello, Michele; D'ago, G.

*Published in:*  
Monthly Notices of the Royal Astronomical Society: Letters

*DOI:*  
[10.1093/mnrasl/slad101](https://doi.org/10.1093/mnrasl/slad101)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2023

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Forbes, D. A., Gannon, J., Iodice, E., Hilker, M., Doll, G., Buttitta, C., Marca, A. L., Arnaboldi, M., Cantiello, M., D'ago, G., Falcon Barroso, J., Greggio, L., Gullieuszik, M., Hartke, J., Mieske, S., Mirabile, M., Rampazzo, R., Rejkuba, M., Spavone, M., ... Capasso, G. (2023). Ultra diffuse galaxies in the Hydra I cluster from the LEWISProject: Phase-Space distribution and globular cluster richness. *Monthly Notices of the Royal Astronomical Society: Letters*, 525(1), L93-L97. <https://doi.org/10.1093/mnrasl/slad101>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Ultra diffuse galaxies in the Hydra I cluster from the LEWISProject: Phase-Space distribution and globular cluster richness

Duncan A. Forbes,<sup>1★</sup> Jonah Gannon,<sup>1</sup> Enrichetta Iodice,<sup>2</sup> Michael Hilker,<sup>3</sup> Goran Doll,<sup>2,4</sup> Chiara Buttitta<sup>1b</sup>,<sup>2</sup> Antonio La Marca,<sup>5,6</sup> Magda Arnaboldi,<sup>3</sup> Michele Cantiello<sup>1b</sup>,<sup>7</sup> G. D’Ago,<sup>8</sup> Jesus Falcon Barroso,<sup>9,10,11</sup> Laura Greggio,<sup>12</sup> Marco Gullieuszik,<sup>12</sup> Johanna Hartke,<sup>13,14</sup> Steffen Mieske,<sup>15</sup> Marco Mirabile,<sup>7,16</sup> Roberto Rampazzo,<sup>12</sup> Marina Rejkuba,<sup>3</sup> Marilena Spavone<sup>1b</sup>,<sup>2</sup> Chiara Spiniello<sup>1b</sup><sup>17</sup> and Giulio Capasso<sup>2</sup>

<sup>1</sup>Centre for Astrophysics & Supercomputing, Swinburne University, Hawthorn, VIC 3122, Australia

<sup>2</sup>INAF - Astronomical Observatory of Capodimonte, Salita Moiarillo 16, I-80131 Naples, Italy

<sup>3</sup>European Southern Observatory, Karl-Schwarzschild-Strasse 2, D-85748 Garching, Germany

<sup>4</sup>University of Naples ‘Federico II’, C.U. Monte Sant’Angelo, Via Cinthia, I-80126 Naples, Italy

<sup>5</sup>SRON Netherlands Institute for Space Research, Landleven 12, NL-9747 AD Groningen, The Netherlands

<sup>6</sup>Kapteyn Astronomical Institute, University of Groningen, Postbus 800, NL-9700 AV Groningen, The Netherlands

<sup>7</sup>INAF - Astronomical Observatory of Abruzzo, Via Maggini, I-64100 Teramo, Italy

<sup>8</sup>Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

<sup>9</sup>INAF - Astronomical Observatory of Padova, Vicolo dell’Osservatorio 5, I-35122 Padova, Italy

<sup>10</sup>Instituto de Astrofísica de Canarias, Vía Láctea s/n, E-38205 La Laguna, Tenerife, Spain

<sup>11</sup>Departamento de Astrofísica, Universidad de La Laguna, E-38200 La Laguna, Tenerife, Spain

<sup>12</sup>Osservatorio Astronomico di Padova, Via dell’Osservatorio 8, I-36012 Asiago (VI), Italy

<sup>13</sup>Finnish Centre for Astronomy with ESO (FINCA), University of Turku, FI-20014, Finland

<sup>14</sup>Tuorla Observatory, Department of Physics and Astronomy, University of Turku, FI-20014, Finland

<sup>15</sup>European Southern Observatory, Alonso de Cordova 3107, Santiago 7630355, Chile

<sup>16</sup>Gran Sasso Science Institute, L’Aquila, Italy

<sup>17</sup>Sub-department of Astrophysics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, United Kingdom

Accepted 2023 July 19. Received 2023 July 18; in original form 2023 June 14

## ABSTRACT

Although ultra diffuse galaxies (UDGs) are found in large numbers in clusters of galaxies, the role of the cluster environment in shaping their low surface brightness and large sizes is still uncertain. Here, we examine a sample of UDGs in the Hydra I cluster ( $D = 51$  Mpc) with new radial velocities obtained as part of the LEWIS (Looking into the faintest with MUSE) project using VLT/MUSE data. Using a phase-space, or infall diagnostic, diagram we compare the UDGs to other known galaxies in the Hydra I cluster and to UDGs in other clusters. The UDGs, along with the bulk of regular Hydra I galaxies, have low relative velocities and are located near the cluster core, and thus consistent with very early infall into the cluster. Combining with literature data, we do not find the expected trend of GC-rich UDGs associated with earlier infall times. This result suggests that quenching mechanisms other than cluster infall should be further considered, e.g. quenching by strong feedback or in cosmic sheets and filaments. Tidal stripping of GCs in the cluster environment also warrants further modelling.

**Key words:** galaxies: haloes – galaxies: photometry – galaxies: star clusters: general – galaxies: structure.

## 1 INTRODUCTION

The possible formation pathways of ultra diffuse galaxies (UDGs) have been a subject of an ongoing vigorous debate since 2015, when a population of these extremely diffuse galaxies was identified in the Coma cluster using the Dragonfly Telephoto Array (van Dokkum et al. 2015). Existing in all environments, they are most common in clusters with several hundred found in the Coma cluster (Yagi et al. 2016; Alabi et al. 2018). This significant contribution to our

‘census of galaxies’ has prompted numerous simulation studies and accompanying predictions (see Sales et al. 2020 and references therein). These simulations can be broadly placed in two categories: internal processes (e.g. episodic supernova feedback) or external (e.g. tidal effects in a dense environment). Some combination of both processes may be operating along with past galaxy infall (and subsequent quenching) into clusters.

UDGs have low surface brightnesses (LSBs; they are defined to have central values in the  $G$  band of  $\mu > 24$  mag arcsec<sup>-2</sup>) so that spectroscopic studies of them push even 8–10 m class telescopes, with efficient LSB instruments, such as KWCI on Keck or MUSE on VLT, to their limits. While strictly speaking dwarf galaxies with  $M_* <$

\* E-mail: dforbes@swin.edu.au

$10^9 M_{\odot}$ , UDGs are unlike classical dwarfs as they have extreme sizes with effective radii  $R_e > 1.5$  kpc (i.e. comparable to the disc of the Milky Way with  $R_e \sim 3.5$  kpc). They also reveal another unexplained feature, with some hosting up to ten times more globular clusters (GCs) than classical dwarf galaxies of the same luminosity (Forbes et al. 2020). Their very existence in clusters and their generally old stellar populations suggests that some may be protected within an overly massive dark matter halo. The latter is supported by the correlation between GC numbers and host galaxy halo mass for normal galaxies (e.g. Burkert & Forbes 2020).

In the standard picture of dwarf galaxy evolution (Mistani et al. 2016), dwarfs that fell into clusters at early times will have experienced intense star formation (SF), prior to, or at the start of, infall (which is also expected to give rise to a high fraction of stars in bound star clusters). This is followed by quenching of any further SF as the infall proceeds. Both of these effects would lead to a high fraction of GCs relative to their field stars (Mistani et al. 2016; Ramos-Almendares et al. 2020). Indeed trends of GC richness and  $[\alpha/\text{Fe}]$  ratios with clustercentric radius provide some observational support for this interpretation (Peng et al. 2008; Liu et al. 2016; Romero-Gómez et al. 2023). This early-infall, or biasing, has been invoked for UDGs by Carleton et al. (2021) who include cluster tidal effects within the IllustrisTNG simulation and simplified GC formation physics. Similar to classical dwarfs, they predict that early-infall UDGs should be rich in GCs. Based on a semi-empirical model, Trujillo-Gomez, Kruijssen & Reina-Campos (2022) also predict that galaxies near the cluster core form more GCs.

Using phase-space, or infall diagnostic, diagrams of the type proposed by Rhee et al. (2017) one can investigate whether GC richness depends on UDG cluster infall time. No trend between GC richness and very early infall times might suggest that GC formation and quenching occurred before cluster infall. While low mass galaxies typically quench at late times, there is a considerable range in quenching times with some low mass galaxies quenching at  $z \sim 2$  or 10.5 Gyr ago (Moster, Naab & White 2020). Quenching at early times via stellar feedback (Stinson et al. 2013) may be one possibility. This early quenching applied to UDGs has been described by Danieli et al. (2022). Another possibility may be quenching via the interaction with cosmic sheets or filaments (Pasha et al. 2023). A first attempt at this sort of infall analysis applied to GCs was presented in Gannon et al. (2022) for several UDGs in the Coma and Perseus clusters. No clear signal was found but the sample was small with just over a dozen UDGs and with a bias towards GC-rich UDGs.

In this letter, we examine the infall diagnostic diagram for a new sample of UDGs in the Hydra I cluster (A1060;  $D = 51 \pm 4$  Mpc). The Hydra I cluster appears to be fairly dynamically relaxed (Ventimiglia, Arnaboldi & Gerhard 2011) but also reveals hints of substructures (Lima-Dias et al. 2021; La Marca et al. 2022a), an infalling group of galaxies (Arnaboldi et al. 2012), and evidence for ram pressure stripping (Iodice et al. 2021; Wang et al. 2021). The observed UDGs are located near the cluster core and the northern subgroup, with all lying within the 0.3 virial radii ( $R_{200}$ ) of the Hydra I cluster centre. Each was observed using MUSE on the VLT as part of the ongoing LEWIS (Looking into the faintEst With MUSE) project. Details of the project, including galaxy radial velocities, positions, GC counts etc, are given in Paper I by Iodice et al. (2023, submitted). The GC counts are based on deep, optical multi-filter imaging with the VST as part of the VEGAS project (Iodice et al. 2020) and will be updated after the full analysis of the MUSE data. Here, we explore the distribution of UDGs in phase space and investigate whether


they reveal any trend in this space with their GC richness. We also include similar data for UDGs in other nearby clusters. For the Hydra I cluster we adopt the same parameters as used by La Marca et al. (2022b), i.e.  $cz = 3683 \pm 46 \text{ km s}^{-1}$ ,  $\sigma = 724 \pm 31 \text{ km s}^{-1}$ , and virial parameters  $R_{200} = 1.6 \text{ Mpc}$ ,  $M_{200} = 2 \times 10^{14} h^{-1} M_{\odot}$  and take its centre as NGC 3311 (RA = 159.17842, Dec. =  $-27.528339$ ). These values are similar to those found by Lima-Dias et al. (2021) who recently studied Hydra I galaxies out to the virial radius.

## 2 INFALL DIAGNOSTIC DIAGRAM FOR HYDRA I CLUSTER GALAXIES

Rhee et al. (2017) carried out cosmological simulations of several galaxy clusters and examined the resulting distribution of galaxies in phase-space (i.e. velocity of the galaxy relative to the mean cluster velocity normalized by the cluster velocity dispersion versus the galaxy's clustercentric radius normalized by the cluster virial radius). Based on the infall time of galaxies, they divided this diagram into several infall zones, ranging from those that fell into the cluster at very early times, to those that are yet to fall in. Thus the location of galaxies in this diagram provides an 'infall diagnostic' which is statistical in nature and additional scatter is introduced when using 2D projected radii (as is the case for observational data). For example, the 'very early infall' (or ancient infaller) zone in the simulation is occupied by a slight majority (52 percent) of galaxies that have resided in the cluster for more than 6.45 Gyr. Projection effects mean that the true clustercentric radius for some galaxies is larger in 3D than observed in 2D. For most galaxies this effect should be less a factor of two from the projected one.

In Fig. 1, we show such an infall diagnostic diagram for all galaxies in the Hydra I cluster out to half the virial radius. This includes giant and dwarf galaxies from the study of Christlein & Zabludoff (2003) plus the addition of UDGs and three LSB galaxies that have UDG-like sizes but are slightly brighter from Iodice et al. (2023, submitted). We find that the bulk of the non-UDG Hydra I galaxies are located within the 'very early infall' zone. The simulation of Rhee et al. (2017) predicts that just over half of these would have been part of the cluster for at least 6.45 Gyr. There are also galaxies located in later infall zones and three galaxies that may lie outside of the cluster with large relative velocities – these could be backplash galaxies (having passed through the cluster) or simply galaxies that are yet to fall into the cluster.

If we examine giant and classical dwarf galaxies separately (divided at  $M_R = -18$  or  $m_R = 15.5$ ) there is no clear difference between them in terms of their infall properties. Compared to the UDGs they appear to scatter to higher relative velocities on average. A more quantitative measure of the differences in their infall properties can be obtained from the product of their relative velocity from the cluster mean and their radial position:  $\Delta V/\sigma \times R/R_{200}$ . Restricting to  $0.3R/R_{200}$ , as probed by the imaging, we find mean values (and error on the mean) of  $\Delta V/\sigma \times R/R_{200} = 0.83 (\pm 0.07) \times 0.15 (\pm 0.01) = 0.12 (\pm 0.02)$  for giant galaxies and  $0.88 (\pm 0.06) \times 0.16 (\pm 0.01) = 0.14 (\pm 0.02)$  for classical dwarfs. For the UDGs the mean value is  $\Delta V/\sigma \times R/R_{200} = 0.80 (\pm 0.17) \times 0.16 (\pm 0.02) = 0.13 (\pm 0.04)$ . This indicates that UDGs are similarly concentrated in phase-space to the other cluster galaxies. Also, while UDGs have a similar distribution in clustercentric radius, their velocities are closer to the cluster mean than either giants or classical dwarfs. We note that Lima-Dias et al. (2021) also found passive early-type galaxies to be concentrated in the cluster core. The LSB galaxies in Fig. 1 are found in a range of infall zones, from early to late infall.



**Figure 1.** Infall diagnostic diagram for non-UDG (giants, dwarfs, and LSB galaxies) and UDGs in the Hydra I cluster. The diagram shows the relative line-of-sight velocity of each galaxy normalized by the cluster velocity dispersion against the projected radius normalized by the virial radius. Regions of the diagram are shaded according to their infall times from the cosmological simulations of Rhee et al. (2017) as indicated in the legend. The plot shows that most UDGs and non-UDG galaxies of the Hydra I cluster lie within the very early infall zone – the simulations of Rhee et al. (2017) indicate that around half of the galaxies in this zone were part of the cluster at least 6.45 Gyr ago.

As might be expected from their inner cluster position, our UDGs were among the earliest inhabitants of the cluster, infalling at least 6.45 Gyr ago according to simulations of Rhee et al. (2017). They would be expected to have SF histories that indicate early quenching. A preliminary analysis by Iodice et al. (2023, submitted) for one UDG (UDG11) indicates an old age of  $\sim 10$  Gyr, suggestive of early quenching. Future analysis will also include the  $[\alpha/\text{Fe}]$  ratios which appears to be a sensitive indicators of SF histories for low mass galaxies (see Ferre-Mateu et al. 2023, in prep. for results on UDGs in other clusters and Romero-Gómez et al. 2023, for dwarf galaxies in the Fornax cluster). We note that the study of Lima-Dias et al. (2021) found 88 per cent of Hydra I galaxies (with  $\log M_* > 8.5$ ) to be quenched, i.e. no sign of ongoing SF.


### 3 INFALL DIAGNOSTIC DIAGRAM FOR UDGs IN SEVERAL CLUSTERS

In Fig. 2, we show the UDGs from the Hydra I cluster along with those from the literature and coded by GC richness. Total GC counts for the Hydra I UDGs are determined in (Iodice et al. 2020; La Marca et al. 2022b) and listed again in Paper I (Iodice et al. 2023, in prep.). Literature data comes from Gannon et al. (2022) and the recent work of Toloba et al. (2023). The GC counts are almost exclusively based on imaging (i.e. lacking radial velocities) and we follow Gannon et al. (2022) assigning a somewhat arbitrary separation between rich and poor GC systems at 20 GCs. This corresponds to a halo mass of  $10^{11} M_\odot$  using the scaling relation of Burkert & Forbes (2020). Below 20 GCs the scaling relation is less predictive of halo mass due to increased scatter. By this definition, all of the UDGs in the Hydra

I cluster are GC-poor (ranging from no GCs for several UDGs to 15 GCs for UDG3) and this is unlikely to change significantly when the full set of MUSE spectroscopic data is available. Given the relatively small stellar mass range of the Hydra I UDGs, a fixed GC number corresponds closely to a GC system total mass per host galaxy stellar mass. If we assume the same average mass for a GC of  $10^5 M_\odot$ , this ratio is  $< 1.2$  per cent for all of the observed Hydra I UDGs. While some Coma cluster UDGs also have a ratio  $< 1.2$  per cent the majority have much higher ratios, with up to  $\sim 10$  per cent of the galaxy stellar mass in their GC system, see fig. 4 of Forbes et al. (2020).

Before interpreting Fig. 2 there are various caveats and selection effects that should be born in mind. First, we note that some of the literature UDGs lack firm GC counts and their rich/poor status is on the basis of a visual estimate only (Gannon et al. 2022). Secondly, the literature sample is subject to sample selection effects. The Coma cluster sample of UDGs comes from studies that have focused on GC-rich galaxies or they have focused on a narrow range in clustercentric radius (i.e. around  $0.12 R/R_{200}$  in the Coma cluster). Observations of the Perseus cluster UDGs have so far avoided the cluster inner regions. The Virgo UDG sample is relatively small and mostly GC-poor. In terms of a selection bias, the Hydra I UDGs are the closest to being a representative sample of UDGs in the cluster, however only the inner  $0.3 R/R_{200}$  was imaged in Iodice et al. (2020). Thus, we may be missing the late infalling UDGs. We note that La Marca et al. (2022b) estimated a total UDG population out to the virial radius of  $48 \pm 10$  and so many outer region UDGs, which may be late infallers, remain to be studied.

The UDG infall diagram does *not* clearly show GC-rich UDGs to be located in earlier infall zones as might be expected in the standard



**Figure 2.** Infall diagnostic diagram for only UDGs in the Hydra I, Coma, Virgo, and Perseus clusters. Regions of the diagram are shaded according to their infall times from the cosmological simulations of Rhee et al. (2017). As per the legend, UDGs in different clusters are denoted by different symbols. Symbols are outlined in red (if GC-rich) or blue (if GC-poor), and without an outline if the GC properties are unknown. See main text for discussion of selection effects in the UDG samples. GC-rich UDGs are *not* predominately found in the very early infall region, indeed the data suggest that very early infall UDGs tend to be GC-poor.

picture of dwarf galaxy quenching due to infall which leads to richer GC systems (as described in the Introduction). Indeed, the opposite trend may be present, such that in the very early infall region there are 13 GC-poor UDGs and five GC-rich ones, whereas outside of this region (but within  $0.5 R/R_{200}$ ) there are only six GC-poor and six GC-rich UDGs. Again, we caution that selection and projection effects make conclusions tentative.

#### 4 DISCUSSION

Alabi et al. (2018) used the phase-space diagram to investigate the infall epoch of UDGs, classical dwarfs, and other galaxies in the Coma cluster (a massive dynamically relaxed cluster). Similar to the Hydra I cluster, they saw little difference between classical dwarfs and the giant galaxies. For the UDGs, they identified both early and late infallers. A similar situation might be present for Hydra I UDGs if outer region UDGs were probed. Alabi et al. (2018) did not include GC-richness in their study.

Given the lack of a clear signal for ‘infall bias’ in the GC richness of UDGs alternatives should be further investigated. As noted in the Introduction, quenching at very early times prior to cluster infall should be considered. For such UDGs, we would expect very old ages, low metallicities (similar to the metal-poor subpopulation of GCs) and high alpha overabundances (indicative of rapid SF). A high fraction of mass in GCs relative to field stars might also be expected. A UDG in the NGC 5846 group, (NGC5846\_UDG1) discovered in VEGAS imaging (Forbes et al. 2019), may be an example of such a failed galaxy having a remarkable 13 per cent of its current stellar mass in the form of GCs (Danieli et al. 2022). As noted above, the

observed Hydra I UDGs (from the inner cluster regions) all have less than 1.2 per cent of their stellar mass in GCs.

Another possibility is that the Hydra I UDGs are GC-poor because they have been tidally stripped from their host galaxy. This tidal stripping would have to remove most of the dark matter halo before any GCs, since the dark matter is more radially extended than GC systems. Continued stripping would be expected to remove GCs and stars in roughly equal proportions since the radial extent of GC systems for UDGs closely follows that of the galaxy stars. As well as operating in clusters, tidal stripping of UDGs may occur in galaxy groups. We note that UDGs in the field do tend to be GC-poor (Jones et al. 2023), however this is unlikely to be due to tidal effects and rather some internal process.

The Hydra I UDGs are generally well-fit by a single Sersic profile however a few show hints of asymmetries that might point to a tidal interaction (Iodice et al. 2020; La Marca et al. 2022a). For the one UDG examined in detail by Iodice et al. (2023, submitted) there is some evidence for an isophotal twist in the MUSE data. This might indicate tidal interaction (or a triaxial potential). Furthermore, a Hydra I UDG first identified by Misgeld, Mieske & Hilker (2008) reveals a clear S-shape indicative of ongoing tidal interaction (Koch et al. 2012). In the case of Coma cluster UDGs, Mowla et al. (2017) looked specifically for signs of tidal features via position angle twists in a stacked sample, finding no evidence for such twists.

Sales et al. (2020) have simulated UDGs in clusters of similar mass to Hydra I using Illustris-TNG100. They identify two types of UDGs in clusters, i.e. Tidal-UDGs and Born-UDGs (see also Jiang et al. 2019). The Tidal-UDGs were originally massive galaxies (up to  $10^{10} M_{\odot}$ ) that have been tidally stripped of stars and puffed-up by

the cluster. Born-UDGs were formed as UDGs outside of the cluster and more recently entered the cluster. Thus Tidal-UDGs dominate the inner  $\sim 0.5R/R_{200}$  since they were accreted at early times, while Born-UDGs dominate the outer regions with some only recently falling into the cluster. We remind the reader that we only probe out to  $0.3R/R_{200}$  in Hydra I. The Sales et al. (2020) model would also predict on average higher metallicities, older ages, and lower internal velocity dispersions, for a given stellar mass, for their Tidal-UDG compared to the Born-UDGs. These stellar population, kinematic, GC colours, and dark matter content predicted for Tidal-UDGs can be tested when the full LEWIS data set is available.

## 5 CONCLUSIONS

As part of the LEWIS project (Iodice et al. 2023, submitted), we obtained new VLT/MUSE observations of the radial velocities of UDGs in the Hydra I cluster (at  $D = 51$  Mpc). Here, we examine the location of Hydra I UDGs in infall phase-space diagrams based on simulations of cluster galaxies. We find all of the observed UDGs (and three LSB galaxies) to be associated with the cluster. From comparison with the Rhee et al. (2017) simulations, we conclude that most giants, classical dwarfs, and UDGs fell into the Hydra I cluster long ago, with UDGs being among the earliest infallers. Projection effects in observations and the statistical nature of the infall diagnostic diagram limit our ability to determine the true fraction of ancient infallers. Nevertheless, we might expect UDGs in the Hydra I cluster to reveal old stellar populations consistent with early quenching.

We also compare Hydra I UDGs with their counterparts in the Coma, Perseus, and Virgo clusters in terms of their GC richness. If very early infall into a cluster is associated with enhanced GC richness (as has been suggested for classical dwarf galaxies) then such a trend is expected. The data from these clusters do *not* show a clear trend of GC richness with earlier infall times, indeed the data suggest the opposite trend. If verified by larger and more complete samples, then UDGs may be quenched by a different mechanism than that thought to operate on classical dwarf galaxies. As more data for UDGs is acquired, trends, or the lack of, may become more apparent in an infall diagnostic diagram. A future analysis of SF histories will give an indication of when quenching occurred for the Hydra I UDGs. Once the full data set of the LEWIS project is available, we will be able to test other mechanisms, such as pre-infall quenching and/or tidal stripping, and their possible role in shaping UDGs and their GC systems.

## ACKNOWLEDGEMENTS

We wish to thank the anonymous referee for their comments. We thank A. Romanowsky, L. Buzzo, L. Haacke, and O. Gerhard for useful suggestions. This work is based on visitor mode observations collected at the European Southern Observatory (ESO) La Silla Paranal Observatory and collected at the European Southern Observatory under ESO programmes 099.B-0560(A) and 108.222P. INAF authors acknowledge financial support for the VST project (PI: P. Schipani). DAF thanks the ARC for support via DP220101863. Parts of this research were supported by the Australian Research Council Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D), through project number CE170100013. MC acknowledges support from the INAF-EDGE program (PI Leslie K. Hunt). JF-B acknowledges support through the RAVET project by the grant PID2019-107427GB-C32 from the Spanish Ministry

of Science, Innovation and Universities (MCIU), and through the IAC project TRACES which is partially supported through the state budget and the regional budget of the Consejería de Economía, Industria, Comercio y Conocimiento of the Canary Islands Autonomous Community.

## DATA AVAILABILITY

Raw data is available from the ESO archive.

## REFERENCES

- Alabi A. et al., 2018, *MNRAS*, 479, 3308  
 Arnaboldi M., Ventimiglia G., Iodice E., Gerhard O., Coccato L., 2012, *A&A*, 545, A37  
 Burkert A., Forbes D. A., 2020, *AJ*, 159, 56  
 Carleton T., Guo Y., Munshi F., Tremmel M., Wright A., 2021, *MNRAS*, 502, 398  
 Christlein D., Zabludoff A. I., 2003, *ApJ*, 591, 764  
 Danieli S. et al., 2022, *ApJ*, 927, L28  
 Forbes D. A., Gannon J., Couch W. J., Iodice E., Spavone M., Cantiello M., Napolitano N., Schipani P., 2019, *A&A*, 626, A66  
 Forbes D. A., Alabi A., Romanowsky A. J., Brodie J. P., Arimoto N., 2020, *MNRAS*, 492, 4874  
 Gannon J. S. et al., 2022, *MNRAS*, 510, 946  
 Iodice E. et al., 2020, *A&A*, 642, A48  
 Iodice E. et al., 2021, *A&A*, 652, L11  
 Jiang F., Dekel A., Freundlich J., Romanowsky A. J., Dutton A. A., Macciò A. V., Di Cintio A., 2019, *MNRAS*, 487, 5272  
 Jones M. G. et al., 2023, *ApJ*, 942, L5  
 Koch A., Burkert A., Rich R. M., Collins M. L. M., Black C. S., Hilker M., Benson A. J., 2012, *ApJ*, 755, L13  
 La Marca A. et al., 2022a, *A&A*, 659, A92  
 La Marca A. et al., 2022b, *A&A*, 665, A105  
 Lima-Dias C. et al., 2021, *MNRAS*, 500, 1323  
 Liu Y. et al., 2016, *ApJ*, 818, 179  
 Mispeld I., Mieske S., Hilker M., 2008, *A&A*, 486, 697  
 Mistani P. A. et al., 2016, *MNRAS*, 455, 2323  
 Moster B. P., Naab T., White S. D. M., 2020, *MNRAS*, 499, 4748  
 Mowla L., van Dokkum P., Merritt A., Abraham R., Yagi M., Koda J., 2017, *ApJ*, 851, 27  
 Pasha I., Mandelker N., van den Bosch F. C., Springel V., van de Voort F., 2023, *MNRAS*, 520, 2692  
 Peng E. W. et al., 2008, *ApJ*, 681, 197  
 Ramos-Almendares F., Sales L. V., Abadi M. G., Doppel J. E., Muriel H., Peng E. W., 2020, *MNRAS*, 493, 5357  
 Rhee J., Smith R., Choi H., Yi S. K., Jaffé Y., Candlish G., Sánchez-Jánsen R., 2017, *ApJ*, 843, 128  
 Romero-Gómez J. et al., 2023, *MNRAS*, 522, 130  
 Sales L. V., Navarro J. F., Peñafiel L., Peng E. W., Lim S., Hernquist L., 2020, *MNRAS*, 494, 1848  
 Stinson G. S., Brook C., Macciò A. V., Wadsley J., Quinn T. R., Couchman H. M. P., 2013, *MNRAS*, 428, 129  
 Toloba E. et al., 2023, *ApJ*, 951, 19  
 Trujillo-Gomez S., Kruijssen J. M. D., Reina-Campos M., 2022, *MNRAS*, 510, 3356  
 van Dokkum P. G., Abraham R., Merritt A., Zhang J., Geha M., Conroy C., 2015, *ApJ*, 798, L45  
 Ventimiglia G., Arnaboldi M., Gerhard O., 2011, *A&A*, 528, A24  
 Wang J. et al., 2021, *ApJ*, 915, 70  
 Yagi M., Koda J., Komiyama Y., Yamao H., 2016, *ApJS*, 225, 11

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.