

University of Groningen

Use of real-world evidence in pharmacoeconomic analysis

Huang, Yunyu

DOI:
[10.33612/diss.95669767](https://doi.org/10.33612/diss.95669767)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Huang, Y. (2019). *Use of real-world evidence in pharmacoeconomic analysis: illustrations in The Netherlands and China*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen. <https://doi.org/10.33612/diss.95669767>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

2

Economic evaluations of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers in type 2 diabetic nephropathy: a systematic review

Yunyu Huang*^{1,2,3}

Qiyun Zhou¹

Flora M. Haaijer-Ruskamp²

Maarten J. Postma¹

¹Department of Pharmacy, Unit of Pharmaco Epidemiology & Pharmaco Economics, University of Groningen, Groningen, The Netherlands

²Department of Clinical Pharmacology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

³School of Public Health, Fudan University, Shanghai, China

*Corresponding author

ABSTRACT

Background:

Structured comparison of pharmacoeconomic analyses for ACEIs and ARBs in patients with type 2 diabetic nephropathy is still lacking. This review aims to systematically review the cost-effectiveness of both ACEIs and ARBs in type 2 diabetic patients with nephropathy.

Methods:

A systematic literature search was performed in MEDLINE and EMBASE for the period from November 1, 1999 to Oct 31, 2011. Two reviewers independently assessed the quality of the articles included and extracted data. All cost-effectiveness results were converted to 2011 Euros.

Results:

Up to October 2011, 434 articles were identified. After full-text checking and quality assessment, 30 articles were finally included in this review involving 39 study settings. All 6 ACEIs studies were literature-based evaluations which synthesized data from different sources. Other 33 studies were directed at ARBs and were designed based on specific trials. The Markov model was the most common decision analytic method used in the evaluations. From the cost-effectiveness results, 37 out of 39 studies indicated either ACEIs or ARBs were cost-saving comparing with placebo/conventional treatment, such as amlodipine. A lack of evidence was assessed for valid direct comparison of cost-effectiveness between ACEIs and ARBs.

Conclusion:

There is a lack of direct comparisons of ACEIs and ARBs in existing economic evaluations. Considering the current evidence, both ACEIs and ARBs are likely cost-saving comparing with conventional therapy, excluding such RAAS inhibitors.

BACKGROUND

Approximately one fourth to one third of patients with diabetes mellitus develop renal manifestations [1-4]. Clinical stages of diabetic nephropathy are generally categorized into stages based on the values of urinary albumin excretion: microalbuminuria (MiA) and macroalbuminuria (MaA) [5]. The prevalence of MiA and MaA in type 2 diabetes is as high as 37–40% in western countries and 57.4–59.8% in Asian countries [6-8]. 20–40% of type 2 diabetic patients with MiA progress to overt nephropathy, and by 20 years after onset of overt nephropathy, about 20% will have progressed to end-stage renal diseases (ESRD) [9]. Because of the large prevalence, diabetes has become the most common single cause of ESRD in the U.S. and Europe [10,11]. As therapies and interventions for coronary artery disease continue to improve, more patients with type 2 diabetes may be expected to survive long enough to develop renal failure.

In developed countries, ESRD is a major cost driver for health-care systems, with annual growth of dialysis programs ranging between 6% and 12% over the past two decades and continuing to grow, particularly in developing countries [12]. Although there are no definitive cure solutions, there is good evidence that adequate treatment can delay or prevent the progress of diabetic nephropathy including strict control of glycaemia, early treatment of hypertension, dietary protein restriction and lipid-lowering therapy [13]. Targeting renin–angiotensin–aldosterone system (RAAS) is the most effective way to delay renal disease progression. Treatment guidelines therefore recommended angiotensin-converting enzyme inhibitors (ACEIs) and angiotensin II receptor blockers (ARBs) as the first-choice agents for treating nephropathy in diabetic patients [14].

Both ACEIs and ARBs target the RAAS and have proven their renal protective effects in diabetic patients in various clinical trials. One disadvantage of ACEIs [15-17] in comparison with ARBs is the higher risk of dry cough while significant differences in effectiveness between these two drug classes have not been shown convincingly although ARBs have been more thoroughly investigated in controlled settings in the recent decade providing relatively high levels of evidence. Often clinical practice guidelines recommend both ACEIs and ARBs in diabetic patients with or even without (micro)albuminuria [18].

Pharmacoeconomic evaluations of ACEIs and ARBs have been widely applied based on clinical trials' results. The pharmacoeconomic results of ARBs have been reviewed previously [19-26]. ARBs were suggested to be cost saving in type 2 diabetic patients with nephropathy versus conventional therapy, largely due to the high costs of treatment

of ESRD. However, a systematic review of cost-effectiveness results of ACEIs in type 2 diabetic patients with renal disease is still lacking. In addition, the need of a structured pharmacoeconomic comparison of the ACEIs with ARBs is pointed out by some researchers [21,26].

The aim of this study is to address the similarities and differences in cost-effectiveness analyses for both ACEIs and ARBs in type 2 diabetic patients with nephropathy. In particular, three objectives are addressed: 1) to summarize the cost-effectiveness of ACEIs; 2) to update the cost-effectiveness of ARBs; 3) to compare the characteristics of different economic evaluations and analyze potential differences and similarities in the cost-effectiveness between the two drug classes reviewed.

METHODS

Literature search strategy

A systematic literature search was performed in MEDLINE and EMBASE for the period November 1, 1999 to Oct 31, 2011. The key words (MeSH headings in MEDLINE, EMtree terms in EMBASE and other text terms) included were (**Table 1**):

- Indicating target drugs, the variations in and abbreviations of ACEIs and ARBs were searched, such as 'angiotensin receptor antagonists', 'renin angiotensin aldosterone system inhibitors', and specific drug names of different ACEIs or ARBs, including 10 specific ACEIs (such as captopril, enalapril, etc.) and 8 ARBs (such as losartan, irbesartan, etc.).
- Indicating diabetic nephropathy, key words were limited to 'type 2 diabetes' and its variations. Variations of nephropathy were combined with diabetes, such as 'diabetic renal diseases' or 'diabetic kidney diseases'.
- Indicating economic evaluations, various key words relating to different evaluation types, pharmacoeconomics, cost of drugs and cost analysis were searched, including 'cost-effectiveness analysis' (CEA), 'cost-utility analysis' (CUA), 'cost-benefit analysis' (CBA), and 'cost savings', etc.

The references of identified articles were manually screened for relevant economic evaluations not identified in the above-mentioned searches (snowballing).

Table 1. Search terms for systematic review

Search terms	MEDLINE	EMBASE
Drug	<p>Mesh: Angiotensin-Converting Enzyme Inhibitors; Angiotensin Receptor Antagonists;</p> <p>TIAB (Title and Abstract): ACEIs; ARBs; ACEI; ARB; renin angiotensin system inhibitor*^a; renin angiotensin aldosterone system inhibitor*; ACE inhibitor*; RAS inhibitor*; RAAS inhibitor*; angiotensin converting enzyme inhibitor*; renin angiotensin system inhibitor*; angiotensin receptor blocker*; Losartan; Candesartan; Valsartan; Irbesartan; Telmisartan; Eprosartan; Olmesartan; Azilsartan; Benazepril; Captopril; Enalapril; Fosinopril; Lisinopril; Moexipril; Perindopril; Quinapril; Ramipril; Trandolapril</p>	<p>EMtree: dipeptidyl carboxypeptidase inhibito; angiotensin receptor antagonist;</p> <p>ab,ti (Abstract and Title): angiotensin receptor blocker; angiotensin receptor blockers; arb; arbs; ace inhibitor; ace inhibitors; angiotensin converting enzyme inhibitor; angiotensin converting enzyme inhibitors; angiotensin converting enzyme (ace) inhibitor; angiotensin converting enzyme (ace) inhibitors; acei; aceis; renin angiotensin system inhibitor; renin angiotensin system inhibitors; renin angiotensin system (ras) inhibitor; renin angiotensin system (ras) inhibitors; ras inhibitor; ras inhibitors; renin angiotensin aldosterone system inhibitor; renin angiotensin aldosterone system inhibitors; raas inhibitor; raas inhibitors; losartan; candesartan; valsartan; irbesartan; telmisartan; eprosartan; olmesartan; azilsartan; benazepril; captopril; enalapril; fosinopril; lisinopril; moexipril; perindopril; quinapril; ramipril; trandolapril;</p>
Diabetic Nephropathy (DN)	<p>Mesh: Diabetes Mellitus, Type 2; Diabetic Nephropathies; Kidney Failure, Chronic;</p> <p>TIAB: diabetic nephropathy*; diabetic renal disease*; diabetic kidney disease*;</p>	<p>EMtree: non insulin dependent diabetes mellitus; diabetic nephropathy;</p> <p>ab,ti: diabetic nephropathy; diabetic nephropathies; diabetic renal diseases; diabetic renal disease; diabetic kidney diseases; diabetic kidney disease</p>
Economic Evaluation (EE)	<p>Mesh: Economics, Pharmaceutical; Costs and Cost Analysis; Drug Costs; Cost Savings; Cost of Illness; Cost-Benefit Analysis;</p> <p>TIAB: cost effect*; cost utility; cost benefit*; economic evaluation*; cost analys*</p>	<p>EMtree: pharmacoeconomics; economic evaluation; drug cost; cost control; cost of illness; cost benefit analysis; cost effectiveness analysis;</p> <p>ab,ti: cost effectiveness; cost utility; cost benefit; economic evaluation; economic evaluations; cost analys;</p>

Table 1. Continued

Search terms	MEDLINE	EMBASE
Search Strategy	(" Drug Term 1 " [Mesh] OR " Drug Term 2 " [TIAB] ...) AND (" DN Term 1 " [Mesh] OR " DN Term 2 " [TIAB] ...) AND (" EE Term 1 " [Mesh] OR " EE Term 2 " [TIAB] ...)	(' Drug Term 1 ' /exp OR ' Drug Term 2 ' :ab,ti ...) AND (' DN Term 1 ' /exp OR ' DN Term 2 ' :ab,ti ...) AND (' EE Term 1 ' /exp OR ' EE Term 2 ' :ab,ti ...) NOT [medline]/lim ^b

a: An asterisk (*) following the word is the wildcard character, which means to search in MEDLINE for all terms that begin with a word; b: To exclude articles that can be found in MEDLINE.

Study selection

Inclusion criteria for the review were as follows (following the PICOS-design):

- **Population:** patients in studies had to have type 2 diabetes with symptoms of renal diseases;
- **Interventions and Comparators:** studies must examine an ACEI- or ARB-based treatment regimen for the progression of diabetic nephropathy compared with regimens that did not include these medications, or if available, compare ACEIs with ARBs directly;
- **Outcomes:** clinical outcomes should be relevant to renal disease symptoms, including overt diabetic nephropathy, ESRD (kidney transplantation or dialysis), all-cause mortality, etc.; and
- **Study design:** studies had to be original economic evaluations.

Other criteria concerned that studies had to have been published as full-length articles and were peer-reviewed for English-language journals.

Study selection was performed in three rounds. First, titles and abstracts of searched articles were scanned and checked. In the second round, the full-texts of included articles were read carefully and quality was assessed in the last round. Two authors independently assessed the quality of the articles included and extracted the data. Differences were resolved by consensus.

Quality assessment

Quality assessment was conducted at the 'study' level, i.e. each study was analyzed one by one. A checklist for critical appraisal of economic evaluations [27] was used to evaluate the study quality. The checklist comprises 12 criteria assessing the study design, outcomes

and costs and the extrapolation of the results of an economic evaluation. An additional file shows this checklist in more detail (see Additional file 1).

The criterion 'applicable to local population' was not included in the assessment as we didn't feel this was relevant for the current study; i.e. 11 criteria were considered in the end. In case studies showing cost savings, the absence of an explicit incremental cost-effectiveness ratio (ICER) was classified as adequate, since in that case no incremental ratio is necessary or meaningful.

Studies were subsequently included in the full review if: 1) the outcomes and costs have been assessed as being credibly, 2) at least 6 of the 11 quality criteria were rated as adequate or good; and 3) not more than three quality criteria were assessed as being inadequate.

Data extraction

Data extraction was based on the 11 criteria included in the quality assessment checklist which concerned: 1) basic information of study design; 2) data on outcomes and costs; and 3) results and conclusions. We grouped articles into two groups, reflecting ACEIs and ARBs. The latter group was subdivided into three subgroups in line with the three mostly analyzed ARBs, irbesartan, losartan and valsartan.

To make the results comparable across the studies, cost-saving or ICER results were standardized to 2011 price levels, by applying the appropriate annual deflators for each country, based on the statistics from the World Bank [28]. Since the deflator data for Taiwan was not available from the World Bank, cost data of this region was not standardized. The original cost-saving result was showed as reference.

All the currencies were converted to 2011 Euros, based on the Euro rate as of June 30th, 2011 [29].

The results of selected studies were classified in 5 categories: 1) cost-saving: net life years or QALYs gained in conjunction with \geq €1,000 saved per patient as compared with the comparison intervention; 2) almost cost-neutral: net life years or QALYs gained, with $<$ €1,000 saved per patient; 3) very cost-effective: $0 < \text{ICER} \leq \text{€}20,000$; 4) cost-effective: $\text{€}20,000 < \text{ICERs} \leq \text{€}40,000$; 5) not cost-effective: $\text{ICERs} > \text{€}40,000$. The classification was based on both literature and suggestions in identified studies in this review [30,31].

RESULTS

Up to October 2011, 434 articles (141 articles from PubMed and 293 articles from EMBASE) were identified. After full-text checking, 32 articles were included into the quality assessment. After quality assessment, 30 articles were finally included in this review (Figure 1). One of the excluded articles had 4 criteria assessed as inadequate and only 4 criteria assessed as good. The other one merely got 5 criteria rated as adequate among the 11 criteria considered.

Among these 30 selected articles, in one article on losartan for an Asian population [32] only the data from Hong Kong were considered as the cost data from other Asian countries or regions assessed seemed not to be of adequate quality. Finally, 39 studies in different countries or regions contained in these 30 articles were included in the analysis.




Figure 1. Flow chart summarizing systematic study selection process.

Summary of selected studies

Table 2 summarizes the basic features of studies included. All six ACEIs studies [33-38] were literature-based evaluations which synthesized data from different sources. All ARBs studies [32,39-62] were designed based on specific trials. The Markov model was the most common decision analytic method used in these evaluations. From the cost-effectiveness results, 37 out of 39 studies indicated both ACEIs and ARBs were cost-saving comparing with placebo/conventional treatment or amlodipine. In the absence of clear cost savings, cost neutrality of very favorable cost-effectiveness was achieved minimally. No studies were identified with a direct cost-effectiveness comparison between ACEIs and ARBs.

Table 2. Summary of selected studies (number of study)

		ACEIs (Total 6)	ARBs (Total 33)	ARBs Losartan (Total 14)	ARBs Irbesartan (Total 18)	ARBs Valsartan (Total 1)
Data source	Trial based	0	33	14	18	1
	Literature based	6	0	0	0	0
Intervention and control group	Comparing with placebo / conventional therapy	2	22	14	8	0
	Comparing with other drugs	0	12	0	11	1
	Comparing different strategies	4	10	0	10	0
Decision Model	Markov model	6	20	1	18	1
	Weibull model	0	3	3	0	0
	Regression method	0	10	10	0	0
Perspective	Third party payer	4	33	14	18	1
	Societal	2	0	0	0	0
CE results	Cost-saving	5	32	13	18	1
	Cost-neutral	0	1	1	0	0
	Very cost- effective	1	0	0	0	0
	Cost-effective	0	0	0	0	0
	Not cost- effective	0	0	0	0	0

Design of selected studies

Key features of the design of the selected studies were summarized in **Table 3**. Six studies of ACEIs [33-38] were diverse in data sources, intervention and control groups. The 33 studies on ARBs showed much more consistency within each ARB drug class (losartan, irbesartan and valsartan) regarding control and data sources concerning the various clinical trials done in ARBs.

ACEIs

Six studies [33-38] evaluated the cost-effectiveness of ACEIs, all using a Markov model as the method for decision modeling. The transition probabilities in these Markov models, i.e. the sources and sizes of effectiveness data in these studies, were diverse. All six studies obtained their effectiveness data from more than one RCT [63-67] or from meta-analyses [37,38]. Only one of the studies [34] included a specific ACEI, enalapril, to compare with placebo, while the other five studies treated ACEIs as a group or drug class. ARBs were also included in the analytic model as a substitute for ACEIs when patients got cough side-effect in the two articles written by Adarkwah et al. [37,38].

ARBs

The 33 studies (included in 24 articles [32,39-61]) targeting ARBs have major similarities in study design. Fourteen evaluations for losartan [32,39-47] were based on The Reduction of Endpoints in Non-insulin Dependent Diabetes Mellitus with the Angiotensin II Antagonist Losartan (RENAAL) trial [62]. Eighteen evaluations of irbesartan [48-60] used data from the Irbesartan in Diabetic Nephropathy Trial (IDNT) [68] to assess the cost-effectiveness for patients with type 2 diabetes and overt nephropathy before 2004. Later the Irbesartan in Reduction of Microalbuminuria-2 (IRMA-2) [69] trial was added into the model to expand the progress of diabetic renal development from nephropathy back to the onset of MiA. The only study for valsartan was based on the MicroAlbuminuria Reduction With VALsartan (MARVAL) study [70].

All 14 losartan studies can be subdivided into two groups based on different time horizon. Eleven studies [32,39-42,46,47] were within-trial analyses, while the other three [43-45] extrapolated to beyond-trial time-horizon analyses. Ten within-trial analyses [32,39-42,47] used a straightforward method to calculate the effectiveness and cost. In this method, the patient-days spent in the stage of ESRD were estimated by subtracting the area under curve (AUC) of the Kaplan-Meier survival curve for time to the minimum of ESRD or all-cause death for both groups in the trial. The costs of ESRD were calculated by multiplying ESRD

days and daily cost of ESRD. Only one within-trial study [46] performed a Markov model as the analytic method to evaluate the cost-effectiveness. Three beyond-trial studies [43-45] used a Weibull model to prolong the time horizon to lifetime. Cumulative incidence of ESRD and life expectancy were assessed as the effectiveness measurements.

Irbesartan for overt nephropathy was compared with conventional treatment and amlodipine in five studies [48-51]. These five studies were based on the IDNT trial and a Markov model with five stages (from ' overt nephropathy ' via ' double of serum creatinine ', ' ESRD + dialysis ' and ' ESRD + transplant ' to ' death ') was developed to evaluate life expectancy and lifetime cost. In particular, Palmer et al. combined the IRMA-2 trial with the IDNT trial and applied a seven-stage Markov model, extrapolating the Markov model with a previous MiA state [48,49,51-60]. ' Early irbesartan ' (standard antihypertensive therapy plus irbesartan at the onset of MiA) was then compared with conventional therapy and ' late irbesartan ' or ' late amlodipine ' (standard antihypertensive therapy plus administration of irbesartan/amlodipine once the patients reach the advanced diabetic nephropathy stage).

Cost-effectiveness of Valsartan [61] was evaluated in one study using amlodipine as the control. A Markov model with seven stages was designed and QALYs were calculated as the effectiveness results.

Table 3. Study design of economic evaluations on ACEIs and ARBs

Study, country/region	Source of effectiveness data	Intervention group	Control group	Decision model Type	Time horizon (years)	Evaluation type
ACEIs						
Golan et al. 1999 US ³²	UERNN, LEAPP and EADN trial	‘‘Treat all’’ strategy ^a	(1) Screen for MIA ^b ; (2) Screen for gross proteinuria ^c .	Markov model with 5 states	10	CEA & CUA (Life-years & QALYs)
Sakthong et al. 2001 Thailand ³³	LEAN trial and the opinion of nephrologists	Enalapril at the dose of 10 mg/day	Placebo	Markov model with 4 stages	25	CEA (Life-years)
Rosen et al. 2005 US ³⁴	UERNN, EADN, LEAN, H-MH studies and HOPE trial	Medicare first-dollar coverage of ACEIs	Year 2005’s Medicare practice	Markov model adding a cardiovascular events component.	Lifetime	CEA & CUA (Life-years & QALYs)
Campbell et al. 2007 US ³⁵	UERNN, EADN, H-MH studies and IRMA-2 trial	ACEI therapy in normoalbuminuric, microalbuminuric, and macroalbuminuric patients	No ACEI initiation in patients	Markov model	8	CEA (CVD event avoided, life saved, dialysis prevented, composite endpoint avoided)
Adarkwah et al. 2010 Germany ³⁶	EADN and two meta-analyses	‘‘Treat all’’ strategy ^a	(1) Screen for MIA ^b ; (2) Screen for MaA ^c ; (3) no-screening and no-treatment alternative.	Markov model with 5 states	50	CUA (QALY)
Adarkwah et al, 2011 Netherlands ³⁷	EADN and two meta-analyses	‘‘Treat all’’ strategy ^a	(1) Screen for MIA ^b ; (2) Screen for MaA ^c .	Markov model with 5 states	50	CUA (QALY)

Table 3. Continued

Study, country/region	Source of effectiveness data	Intervention group	Control group	Decision model Type	Time horizon (years)	Evaluation type
ARBs						
Losartan						
Herman et al. 2003 US ⁴³	RENAAL trial	Losartan	Placebo ^d	A regression-based method	3.5 / 4	CEA (Number of ESRD days)
Souchet et al. 2003 France ⁴⁴	RENAAL trial	Losartan (initial daily dosing of losartan was 50 mg, with the possibility of titration to 100 mg/day)	Placebo ^d	A regression-based method	3.5 / 4	CEA (Number of ESRD days)
Burgess et al. 2004 Canada ⁴⁵	RENAAL trial	Losartan	Placebo ^d	A regression-based method	3.5 / 4	CEA (Number of ESRD days)
Szucs et al. 2004 Switzerland ⁴⁶	RENAAL trial	Losartan (initial daily dosing of losartan was 50 mg, with the possibility of titration to 100 mg/day)	Placebo ^d	A regression-based method	3.5 / 4	CEA (Number of ESRD days)
Seng et al. 2005 Hong Kong ⁴⁷ (only data of Hong Kong were included)	RENAAL trial	Losartan	Placebo ^d	A regression-based method	3.5	CEA (Number of ESRD days)

Table 3. Continued

Study, country/region	Source of effectiveness data	Intervention group	Control group	Decision model Type	Time horizon (years)	Evaluation type
Arredondo et al. 2005 Mexico ⁴⁸	RENAAL trial	Losartan	Placebo ^d	A variation of the cumulative incidence competing risk method / Weibull model	25 (life time)	CEA (Cumulative incidence of ESRD, life expectancy)
Vora et al. 2005 UK ⁴⁹	RENAAL trial	Losartan (50-100 mg QD)	Conventional antihypertensive treatment ^d (excluding ACEIs or angiotensin II antagonists)	Weibull model	life time	CEA (Cumulative incidence of ESRD, life expectancy)
Carides et al. 2006 US ⁵⁰	RENAAL trial	Losartan	Placebo ^d	A cumulative incidence competing risk method / Weibull model	25 (life time)	CEA (Cumulative incidence of ESRD, life expectancy)
Stafylas et al. 2007 Greece ⁵¹	RENAAL trial	Losartan (50-100 mg QD)	Placebo ^d	Markov model with 6 states	3.5 / 4	CEA (Number of ESRD days)
de Portu et al. 2011 Italy, France, Germany, Switzerland, US ⁵²	RENAAL trial	Losartan	Standard care ^d	Standard methods by comparing the economic outcomes deriving from additional losartan to standard care vs standard care alone	3.4	CEA (Number of ESRD days)
Irbesartan						
Rodby RA et al. 2003 US ⁵³	IDNT trial	Irbesartan titrated from 75 to 300 mg/day	(1) 'Control'; (2) Amlodipine titrated from 2.5 to 10 mg/day.	Markov model with 5 stages	25	CEA (Life expectancy)

Table 3. Continued

Study, country/region	Source of effectiveness data	Intervention group	Control group	Decision model Type	Time horizon (years)	Evaluation type
Palmer A] et al. 2003 Belgium, France ⁵⁴	IDNT trial	Irbesartan titrated from 75 to 300 mg/day	(1) 'Control' ^g ; (2) Amlodipine titrated from 2.5 to 10 mg/day.	Markov model with 5 stages	25	CEA (Life expectancy)
Coyle D et al. 2004 Canada ⁵⁵	IDNT trial	Irbesartan	(1) Amlodipine; (2) Standard care ^d .	Markov model with 5 stages	25	CEA (Life expectancy)
Palmer A] et al. 2004 UK ⁵⁶	IDNT trial	Irbesartan 300mg per day	(1) 'Control' ^g ; (2) Amlodipine 10mg per day.	Markov model with 5 stages	25	CEA (Life expectancy)
Palmer A] et al. 2004 US ⁵⁷	IRMA-2 study and IDNT	'Early irbesartan' ^e	(1) 'Control' ^g ; (2) 'Late irbesartan' ^f	Markov model with 7 stages	25	CEA (Years free of ESRD, cumulative incidence ESRD, life expectancy)
Palmer A] et al. 2005 Spain ⁵⁸	IRMA-2 study and IDNT	'Early irbesartan' ^e	Standard antihypertensive medications ^d	Markov model with 7 stages	25	CEA (Years free of ESRD, cumulative incidence ESRD, life expectancy)
Palmer A] et al. 2006 Switzerland ⁵⁹	IRMA-2 study and IDNT	'Early irbesartan' ^e	Conventional antihypertensive treatment ^d initiated when patients had developed MiA.	Markov model with 7 stages	25	CEA (Years free of ESRD, cumulative incidence of ESRD, life expectancy)

Table 3. Continued

Study, country/region	Source of effectiveness data	Intervention group	Control group	Decision model Type	Time horizon (years)	Evaluation type
Palmer A] et al. 2006 France ⁶⁰	IRMA-2 study and IDNT	'Early irbesartan' ^e	(1) 'Control' ^h ; (2) 'Late irbesartan' ^r	Markov model with 7 stages	25	CEA & CUA (Years free of ESRD, life expectancy, QALY)
Palmer A] et al. 2007 Hungary ⁶¹	IRMA-2 study and IDNT	'Early irbesartan' ^e	'Placebo' ^h ; standard antihypertensive medications initiated when patients developed MiA.	Markov model with 7 stages	25	CEA (Years free of ESRD, cumulative incidence ESRD, life expectancy)
Palmer A] et al. 2007 UK ⁶²	IRMA-2 study and IDNT trial	'Early irbesartan' ^e	(1) 'Control' ^h ; (2) 'Late irbesartan' ^r	Markov model with 7 stages	25	CEA (Years free of ESRD, cumulative Incidence of ESRD, life expectancy)
Coyle D et al. 2007 Canada ⁶³	IRMA-2 study and IDNT	'Early irbesartan' ^e	(1) 'Late irbesartan' ^r ; (2) 'Conventional' ^d	Markov model with 7 stages	25	CEA (Life expectancy)
Yang W.C. et al. 2007 Taiwan ⁶⁴	IRMA-2 study and IDNT	'Early irbesartan' ^e	(1) 'Standard' ^h ; (2) 'Late irbesartan' ^r ; (3) 'Late amlodipine' ^g	Markov model with 7 stages	25	CEA (Life expectancy, number of years free of ESRD, cumulative incidence of ESRD)

Table 3. Continued

Study, country/region	Source of effectiveness data	Intervention group	Control group	Decision model Type	Time horizon (years)	Evaluation type
Annemans et al. 2008 China, Taiwan, Malaysia, Thailand, South Korea ⁶⁵	IRMA-2 study and IDNT trial	'Early irbesartan' ^b	(1) 'Standard' ^a ; (2) 'Late irbesartan' ^c ; (3) 'Late amlodipine' ^d	Markov model with 7 stages	25	CEA (Cumulative incidence of ESRD, number of days in dialysis, number of years free of ESRD, life expectancy)
Valsartan						
Smith DG et al. 2004 US ⁶⁶	MARVAL study	Valsartan	Amlodipine	Markov model with 7 stages	8	CUA (Quality-adjusted survival)

a: no screening was performed at all and patients started on ACEI therapy at the time of diagnosing type 2 diabetes

b: patients were screened for MIA once a year and ACEI treatment was started if the test result is positive

c: patients were screened for MaA once a year and ACEI treatment was started if the test result is positive

d: standard antihypertensive therapy alone, excluding the use of ACEIs, ARBs

e: standard antihypertensive therapy plus administration of irbesartan 300 mg/d at the onset of MiA

f: standard antihypertensive therapy plus administration of irbesartan 300 mg/d once the patients reach the advanced diabetic nephropathy stage

g: standard antihypertensive therapy plus administration of amlodipine titrated from 5 to 10 mg/d once the patients reach the advanced diabetic nephropathy stage

UERNN = Use of enalapril to attenuate decline in renal function in normotensive, normoalbuminuric patients with type 2 diabetes mellitus; **LEAPP** = Long-term stabilizing effect of angiotensin-converting enzyme inhibition on plasma creatinine and on proteinuria in normotensive type II diabetic patients; **EADN** = The effect of angiotensin-converting-enzyme inhibition on diabetic nephropathy; **LEAN** = Long-term renoprotective effect of angiotensin-converting enzyme inhibition in non-insulin-dependent diabetes mellitus; **H-MH** = Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy; **HOPE** = The Heart Outcomes Prevention Evaluation; **RENAAL** = The reduction of endpoints in non-insulin dependent diabetes mellitus with the angiotensin II antagonist losartan; **IDNT** = The irbesartan in diabetic nephropathy trial; **IMRA-2** = The irbesartan in reduction of microalbuminuria-2; **MARVAL** = The microalbuminuria reduction with valsartan.

CEA = cost-effectiveness analysis; **CUA** = cost-utility analysis.

Cost-effectiveness results

The key features and main results of all included evaluations are summarized in **Table 4**.

ACEIs

Of the six ACEIs' studies, two [33,35] adopted a societal perspective. This contained additional cost analyses including productivity gains and losses, caregiver time costs. The other four [34,36-38] took the third party payer/health care perspective including only direct costs of nephropathy, ACEIs or other related treatment such as those for cardiovascular disease (CVD). All studies except one [33] favored ACEIs due to the cost-saving results. The exception was the evaluation from Golan et al. [33], showing that compared to 'screen for MiA' (patients were screened for MiA once a year and ACEI treatment was started if the test result is positive), the 'treat all' strategy with ACEIs (no screening was performed at all and patients started on ACEI therapy at the time of diagnosing type 2 diabetes) raised the costs by \$300, but the results still supported 'treat all' strategy as very cost-effective. – It should be noted that these positive results were based on the comparison between ACEIs and no blood pressure (BP) control treatment but not other BP control interventions.

ARBs

Based on the RENAAL trial, all the results over 3.5 years indicated losartan was cost-saving or cost-neutral (Hong Kong) [32] comparing to placebo/conventional therapy. The cost savings per patients ranged from € 2,079 in Greece [46] to € 4,641 in France [47]. When the time horizon was prolonged to lifetime or 25 years, beyond-trial studies showed that the net cost savings by adding losartan to conventional therapy were € 9,182 in UK [44], € 1,861 in Mexico [43] and € 22,757 in U.S [45].

For irbesartan, results consistently showed cost-savings comparing with conventional therapy or amlodipine, even when already started at the onset of MiA. Such early start of irbesartan would economically be even more attractive as compared with late irbesartan starting at overt nephropathy. The five studies [48-51] based on the IDNT trial demonstrated that irbesartan for overt nephropathy could prolong life expectancy with 0.43 years (Canada) [50] to 0.74 years (U.S.) [48] and save € 7,075 (U.K.) [51] to € 19,132 (France) [49] per patient comparing with control over 25 years. When the MiA stage was introduced into the model, early irbesartan remained cost-saving at € 2,564 in Hungary [56] to € 57,871 in Canada [58] compared with control, being more cost-saving than late irbesartan.

The only study for valsartan [61] also supported the using of ARBs in patients with type 2 diabetes and MiA because of saving QALYs and costs. Over 8 years, valsartan treatment had 0.555 discounted QALYs advantage over amlodipine with savings at € 30,424 compared to amlodipine.

Table 4. Main results of economic evaluations on ACEIs and ARBs

Study, country/ region	Discount rate (per annum)	Perspective	Cost categories	Discounted life expectancy /QALY	Incremental cost per patients [year of value]	Incremental cost per patient (standardized to 2011 Euro)	CE with Interventions
	Effects (%)	Costs (%)					
ACEIs							
Golan et al. 1999 US ³²	3	3	Societal The cost of ESRD (dialysis & transplant), ACEIs and screening	15.63 years /11.82 QALYs with 'treat all', 15.59 years /11.78 QALYs with 'screen for MiA', 15.39 years /11.59 QALYs with 'screen for gross proteinuria'	'Treat all' vs 'screen for MiA': \$300 ^a	'Treat all' vs 'screen for MiA': €299	Very cost-effective ['Treat all' vs. 'screen for MiA': €8,062/ QALY]
Sakthong et al. 2001 Thailand ³³	8	8	Not mentioned The cost of ESRD (haemodialysis) and ACEI	9.04 years with enalapril, 7.54 years with control	-\$1,198 [1999]	-€1,269	Cost saving [Enalapril]
Rosen t al. 2005 US ³⁴	3	3	Medicare and societal (1) Medicare perspective: direct medical costs and future health care costs. (2) Societal perspective: additional analyses included productivity gains and losses, caregiver time costs	10.55 years /8.36 QALYs with Medicare first- dollar coverage of ACEIs, 10.30 years /8.13 QALYs with at the time practice costs.	-\$1,606 [2003]	-€1,453	Cost saving [Medicare first- dollar coverage of ACEIs]

Table 4. Continued

Study, country/region	Discount rate (per annum) Effects (%) Costs (%)	Perspective	Cost categories	Discounted life expectancy /QALY	Incremental cost per patients [year of value]	Incremental cost per patient (standardized to 2011 Euro)	CE with Interventions
Campbell et al. 2007 US ³⁵	3 3 3	Health payer	Direct medical costs of nephropathy, CVD, and ACEIs		-\$772 for normoalbuminuria on diagnosis, -\$7,098 for MiA on diagnosis, \$7,987 for MaA on diagnosis [2005]	-\$658 for normoalbuminuria on diagnosis, -\$6,048 for MiA on diagnosis, €6,806 for MaA on diagnosis	Cost-neutral [ACEIs used on normoalbuminuria] Cost saving [ACEIs on MiA]
Adarkwah et al. 2010 Germany ³⁶	3 3 3	the German statutory health insurance	The cost of ESRD (dialysis & transplant), ACEIs, ARBs and screening	15.21 QALYs with 'treat all', 15.14 QALYs with 'screen for MiA', 14.83 QALYs with 'screen for MaA', 14.46 QALYs with 'placebo'	'Treat all' vs. 'placebo': -€16,024 [2006]	'Treat all' vs. 'placebo': -€16,841 [2006]	Cost saving [Treat all using ACEIs]
Adarkwah et al. 2011 Netherlands ³⁷	1.5 4 4	Health care	The cost of ESRD (dialysis & transplant), ACEIs, ARBs and screening	19.63 QALYs with 'treat all', 19.54 QALYs with 'screen for MiA', 19.15 with 'screen for MaA'	'Treat all' vs. 'screen for MiA': -€2,719, 'treat all' vs. 'screen for MaA': -€12,356 [2010]	'Treat all' vs. 'screen for MiA': -€2,749, 'treat all' vs. 'screen for MaA': -€12,492	Cost saving [Treat all using ACEIs]

Table 4. Continued

Study, country/region	Discount rate (per annum)	Perspective	Cost categories	Discounted life expectancy /QALY	Incremental cost per patients [year of value]	Incremental cost per patient (standardized to 2011 Euro)	CE with Interventions
	Effects (%)						
ARBS							
Losartan							
Herman WH et al, 2003 US ⁴³	none	3	Health care system	The cost of ESRD (hemodialysis) and losartan therapy	Over 3.5 years: -\$3,522 [2001]	Over 3.5 years: -€3,306	Cost saving [losartan]
Souchet T et al, 2003 France ⁴⁴	none	8.1% ^b	French healthcare system	The cost of ESRD (dialysis) and losartan therapy	Over 3.5 years: -€3,863 [2002]	Over 3.5 years: -€4,522	Cost saving [losartan]
Burgess ED et al, 2004 Canada ⁴⁵	none	none	Health care system	The cost of ESRD (dialysis & transplant) and losartan therapy	Over 3.5 years: -\$3,675 ^a	Over 3.5 years: -€3,368	Cost saving [losartan]
Szucs TD et al, 2004 Switzerland ⁴⁶	none	none	Swiss healthcare payer	The cost of ESRD (dialysis & transplant) and losartan therapy (only the insurance-paid part)	Over 3.5 years: -CHF4,084 ^a	Over 3.5 years: -€3,660	Cost saving [losartan]

Table 4. Continued

Study, country/region	Discount rate (per annum)	Perspective	Cost categories	Discounted life expectancy /QALY	Incremental cost per patients [year of value]	Incremental cost per patient (standardized to 2011 Euro)	CE with Interventions
Effects (%)	Costs (%)						
Seng WK et al, 2005	3	Health care system	The cost of ESRD (dialysis) and losartan therapy		-\$515 [2004]	-€413	Cost-neutral [losartan]
Hong Kong ⁴⁷ (only data of Hong Kong were included)							
Arredondo A et al, 2005	3	Healthcare system	The cost of ESRD (dialysis), diabetes and losartan therapy	0.697 life years gained for losartan	-\$24,073 [2004]	-€1,861	Cost saving [losartan]
Mexico ⁴⁸							
Vora] et al, 2005	3.5	The UK National Health Service (NHS)	The cost of ESRD (dialysis) and losartan therapy	7.82 life years with losartan, 7.38 life years with placebo (0.44 life years gained for losartan)	£6,622 [2004]	-€9,182	Cost saving [losartan]
UK ⁴⁹							
Carides GW et al, 2006	3	Healthcare system	The cost of ESRD (dialysis), diabetes and losartan therapy	0.697 life years gained for losartan	-\$24,632 [2002]	-€22,757	Cost saving [losartan]
US ⁵⁰							
Stafylas PC et al, 2007	3	The Greek social insurance system	The cost of ESRD (dialysis & transplant) and 75% of drug treatment costs		Over 3.5 years: -€1,665,43 [2003]	Over 3.5 years: -€2,079	Cost saving [losartan]
Greece ⁵¹							

Table 4. Continued

Study, country/region	Discount rate (per annum) Effects (%)	Perspective	Cost categories	Discounted life expectancy /QALY	Incremental cost per patients [year of value]	Incremental cost per patient (standardized to 2011 Euro)	CE with Interventions
de Portu Set al, 2011 Italy ⁵²	3 3	National Healthcare Service	The cost of ESRD (hemodialysis) and losartan therapy		-€3,602.98 [2009]	-€3,664	Cost saving [losartan]
de Portu Set al, 2011 France ⁵²	3 3	Health Insurance	The cost of ESRD (hemodialysis) and losartan therapy		-€4,531.35 [2009]	-€4,641	Cost saving [losartan]
de Portu Set al, 2011 Germany ⁵²	3 3	Health Insurance	The cost of ESRD (hemodialysis) and losartan therapy		-€3,019.66 [2009]	-€3,062	Cost saving [losartan]
de Portu Set al, 2011 Switzerland ⁵²	3 3	Medical Insurance	The cost of ESRD (hemodialysis) and losartan therapy		-€3,949.50 [2009]	-€3,977	Cost saving [losartan]
de Portu Set al, 2011 US ⁵²	3 3	Centers for Medicare & Medicaid Services	The cost of ESRD (hemodialysis) and losartan therapy		-€3,855.50 [2009]	-€4,007	Cost saving [losartan]
Irbesartan							
Rodby RA et al, 2003 US ⁵³	3 3	Health care system	The cost of ESRD (dialysis & transplant), hospitalizations, irbesartan & concomitant antihypertensive drugs	8.225 years with irbesartan, 7.484 years with control (0.741 years gained for irbesartan)	-\$15,607 [2000]	-€14,987	Cost saving [irbesartan]

Table 4. Continued

Study, country/region	Discount rate (per annum)	Perspective	Cost categories	Discounted life expectancy /QALY	Incremental cost per patients [year of value]	Incremental cost per patient (standardized to 2011 Euro)	CE with Interventions
	Effects (%)						
Palmer AJ et al, 2003 Belgium ⁵⁴	3	Institut National d'Assurance de Maladie et Invalidite' (INAMI)	The cost of ESRD (dialysis & transplant) and irbesartan & concomitant antihypertensive drugs	8.57 years with irbesartan, 7.95 years with control (0.62 years gained for irbesartan)	-€11,885 [2002]	-€14,231	Cost saving [irbesartan]
Palmer AJ et al, 2003 France ⁵⁴	3	Social security	The cost of ESRD (dialysis & transplant) and irbesartan & concomitant antihypertensive drugs	8.58 years with irbesartan, 7.97 years with control (0.61 years gained for irbesartan)	-€16,345 [2002]	-€19,132	Cost saving [irbesartan]
Coyle D et al, 2004 Canada ⁵⁵	5	Third party payer	The cost of ESRD (dialysis & transplant), irbesartan & concomitant antihypertensive drugs and other medical costs	6.80 years with irbesartan, 6.37 years with control (0.43 years gained for irbesartan)	-CAD12,564 [2001]	-€11,457	Cost saving [irbesartan]
Palmer AJ et al, 2004 UK ⁵⁶	1.5	National Health Service (NHS) payer	The cost of ESRD (dialysis & transplant) and irbesartan & concomitant antihypertensive drugs	0.58 years gained for irbesartan vs control	-£4,978 ^a	-€7,075	Cost saving [irbesartan]

Table 4. Continued

Study, country/region	Discount rate (per annum) Effects (%)	Costs (%)	Perspective	Cost categories	Discounted life expectancy /QALY	Incremental cost per patients [year of value]	Incremental cost per patient (standardized to 2011 Euro)	CE with Interventions
Palmer AJ et al, 2004 US ⁵⁷	3	3	Third party reimbursement	The cost of ESRD (dialysis & transplant) and irbesartan	11.46 years with 'early irbesartan', 10.54 years with 'late irbesartan', 10.50 years with control (0.96 years gained for irbesartan vs control)	Early irbesartan vs. control: -\$11,922, late irbesartan vs. control: -\$3,252 [2000]	Early irbesartan vs. control: -€11,448, late irbesartan vs. control: -€3,123	Cost saving [early irbesartan]
Palmer AJ et al, 2005 Spain ⁵⁸	3	3	Third party payer	The cost of ESRD (dialysis & transplant) and irbesartan	12.37 years with 'early irbesartan', 11.53 years with control (0.84 years gained for irbesartan)	-€11,082 ^a	-€12,971	Cost saving [early irbesartan]
Palmer AJ et al, 2006 Switzerland ⁵⁹	5	5	Third party Swiss health insurance payer	The cost of ESRD (dialysis & transplant) and irbesartan	10.37 years with 'early irbesartan', 9.80 years with control (0.57 years gained for irbesartan)	-CHF21,487 [2003]	-€19,257	Cost saving [early irbesartan]
Palmer AJ et al, 2006 France ⁶⁰	3	3	Third party French social security insurance payer	The cost of ESRD (dialysis & transplant) and irbesartan	12.17 years /10.55 QALYs with 'early irbesartan', 11.27 years /9.58 QALYs with 'late irbesartan', 11.23 years /9.52 QALYs with control (0.94 years /1.03 QALYs gained for irbesartan vs control)	'Early irbesartan' vs. control: -€22,314, 'late irbesartan vs. control': -€6,619 [2002]	'Early irbesartan' vs. control: -€26,119, 'late irbesartan' vs. control: -€7,748	Cost saving [early irbesartan]
Palmer AJ et al, 2007 Hungary ⁶¹	5	5	Third-party Hungarian health insurance payer	The cost of ESRD (dialysis & transplant) and irbesartan	8.16 years with 'early irbesartan', 7.62 years with control (0.54 years gained for irbesartan)	-HUF519,993 [2002]	-€2,564	Cost saving [early irbesartan]

Table 4. Continued

Study, country/region	Discount rate (per annum) Effects (%)	Discount rate (%)	Perspective	Cost categories	Discounted life expectancy /QALY	Incremental cost per patients [year of value]	Incremental cost per patient (standardized to 2011 Euro)	CE with Interventions
Palmer AJ, 2007 UK ⁶²	3.5	3.5	Third party UK National Health Service (NHS) payer	The cost of ESRD (dialysis & transplant) and irbesartan	11.00 years with 'early irbesartan', 10.20 years with 'late irbesartan', 10.18 years with control (0.82 years gained for irbesartan vs control)	'Early irbesartan' vs. control: -£3,801, 'late irbesartan' vs. control:-£1,491 [2002]	'Early irbesartan' vs. control: -€5,532, 'late irbesartan' vs. control: -€2,170	Cost saving [early irbesartan]
Coyle D et al, 2007 Canada ⁶³	5	5	Canadian health and social care system	All direct costs, including the costs of health, social services, long-term care.	11.52 years with 'early irbesartan', 11.06 years with 'late irbesartan', 10.90 years with control (0.62 years gained for irbesartan vs control)	'Early irbesartan' vs. control: -CAD68,400, 'late irbesartan' vs. control: -CAD14,300 [2006]	'Early irbesartan' vs. control: -€57,871, 'late irbesartan' vs. control: -€12,099	Cost saving [early irbesartan]
Yang W.C. et al, 2007 Taiwan ⁶⁴	3	3	Third-party payer in Taiwan (Taiwan National Health Insurance Program)	The cost of ESRD (dialysis & transplant) and irbesartan	12.003 years with 'early irbesartan', 11.332 years with 'late irbesartan', 11.223 years with control (0.780 years gained for irbesartan vs control)	'Early irbesartan' vs. control: -\$7,603, 'late irbesartan' vs. control: -\$3,233 [2004]		Cost saving [early irbesartan]
Annemans L et al, 2008 China, Taiwan, Malaysia, Thailand, South Korea ⁶⁵	5	5	Third party payer	The cost of ESRD (dialysis & transplant) and irbesartan	'Early irbesartan' strategy had the longest life expectancy (no detail data)	The least expensive strategy: 'early irbesartan' (no detail data)		Cost saving [early irbesartan]

Table 4. Continued

Study, country/region	Discount rate (per annum) Effects (%)	Discount rate (%)	Perspective	Cost categories	Discounted life expectancy /QALY	Incremental cost per patients [year of value]	Incremental cost per patient (standardized to 2011 Euro)	CE with Interventions
Valsartan								
Smith DG et al, 2004 US ⁶⁶	3	3	Third-party payer	Medical care costs including costs of study drugs, routine health care services, and aggregate estimates of medical care associated with the various health states.	6,390 QALYs with valsartan, 5,835 QALYs with amlodipine (0.555 QALYs gained for valsartan)	-\$32,412 [2001]	-€30,424	Cost saving [valsartan]

a: In which year the value of money standardized was not clear. It was assumed to be one year before the publication.

b: The total discount rate within time horizon, not annually.

DISCUSSION

To our knowledge, this is the first review that summarizes all information on the cost-effectiveness of both ACEIs and ARBs. Our systematic review confirms earlier results evidencing the cost-saving potentials of ARBs for type 2 diabetic patients with nephropathy compared with conventional therapy excluding a RAAS inhibitor. Also, our review shows that such potentials might even stronger exist in early treatments prior to the stage of nephropathy, for example, in the MiA-stage. In addition, we found similar cost-saving results for ACEIs due to avoidance of ESRD in combination with prolonging life expectancy. Differences in cost effectiveness of ACEI versus ARB could not be solidly established because of differences in model design, time horizon and country setting among all included studies and lack of head-to-head comparisons in economic evaluations. Yet, cost-saving potentials were unequivocally assessed for both drug groups.

Cost-effectiveness of ACEIs

The number of articles concerning ACEIs was limited compared with the number for ARBs. The reason for not basing studies on single clinical trials may be related to the chronology of ACEIs being available on the market, i.e. the 1980s, before the ARBs. The relevance of CVD in diabetes became only clear in 1990s when the benefits of RAAS inhibitors started also to become clear for diabetic patients. From the six articles included in this review, ACEIs were cost-saving in articles published after 2000s [34-38] and not cost saving (but very cost-effective) in the only one article before 2000s [33]. This may be explained by the patent protection of ACEIs which became generic in the late of 1990s.

Three articles [33,37,38] combined screening for MiA or MaA as the start time point of ACEIs treatment in their analyses. Previous studies of screening for albuminuria with subsequent ACEIs treatment on cardiovascular and renal diseases also support the conclusions on favorable cost-effectiveness and early treatments. Atthobari et al. [71] found that the estimated cost-effectiveness of screening for albuminuria with ACEIs treatment was approximately € 16,700/LYG (2006 value) for subjects with a urinary albumin excretion >15 mg/d compared with no screening when adopting the Dutch health care perspective. This was in accordance with the analyses from Boersma C et al. [72] suggesting the potentially favorable cost-effectiveness of population-based screening for MiA compared with other alternatives. Notably, however the latter two articles were for prevention of cardiovascular and renal events in the general population, not particularly for diabetic patients.

Cost-effectiveness of ARBs

The pharmacoeconomic results of ARBs for renal disease in patients with type 2 diabetes were reviewed previously. Ravera et al. [22] and Boersma et al. [21] reviewed the economic evaluations for ARBs and concluded that evaluations derived from RENAAL, IDNT, IRMA-2 and MARVAL all suggested ARBs to be cost saving compared with conventional therapy in type 2 diabetes patients with nephropathy. Postma & de Zeeuw [26] reviewed the economic benefits of preventing ESRD in patients with type 2 diabetes. They divided the RAAS drug treatment into early and late interventions and concluded that early intervention strategies appear more effective in reducing the risk and the pharmacoeconomic profiles of early intervention clearly outweigh those of late intervention.

From our literature search, there were various economic evaluations on the ARBs losartan, irbesartan and valsartan. There were little differences between studies in each subgroups of ARBs concerning the analysis model, time horizon and measurement of costs and benefits. Although the results varied in different studies and countries, all conclusions supported ARBs as a cost-saving choice.

Differences in economic evaluations of ACEIs and ARBs

The trials referred to in the studies included in this review had different patient characteristics and treatment strategies. Patients enrolled in ACEIs trials were mainly normotensive, while patients enrolled in ARBs trials were mainly hypertensive. Trails with ACEIs had no equal BP control in placebo groups, whereas trials with ARBs had active BP control in placebo groups. Differences in time horizons used for ACEIs and ARBs present another reason hindering comparison of cost effectiveness between these two drug classes.

Referring to the analytic models, the transition probabilities between two states in the Markov model adopted in these ACEIs studies were from different trials, which may weaken the internal validity of the simulation model used and effectiveness results generated. The analytic models used for ARBs were relatively consistent in their strong alignment to the clinical trials available. Similar methods were adjusted to different country settings. This enhanced similarity in cost-effectiveness results of the same ARB drug in different countries. One might argue that the majority of economic evaluations for losartan were cost analyses with existing trial-based effectiveness as the building block.

Differences in evaluation results of ACEIs and ARBs

Previous reviews [18-26] of ACEIs and ARBs didn't summarize the differences between ACEIs and ARBs in the absence of direct comparisons between ARBs and ACE inhibitors

in terms of cost-effectiveness. In this review, also no valid comparison between ACEIs and ARBs is possible regarding cost-effectiveness.

In the lifetime treatment for diabetic nephropathy, cost of dialysis when patients develop to ESRD plays an important role in the burden of disease. Comparing to the cost of ESRD, the cost of drugs comprise a relatively low proportion in the total disease expenditure. As ACEIs and ARBs both can delay the deterioration of kidney function to save huge cost due to treatment, results of the economic evaluations included in this review are all pointing into the same direction that these two drug classes are cost-saving or very cost-effective. Furthermore, most ARBs now are available in generic forms and thus cheaper than when these evaluations were performed, which makes ARBs and ACEIs more similar in both effectiveness and cost. Therefore, similar cost-effectiveness result between ACEIs and ARBs can be hypothesized and results in this review strengthen the relevance of the choice made in guidelines [14,18] of recommending ACEIs or ARBs as both presenting cost-effective choices for patients with diabetic nephropathy.

Limitations

In our review, although the standardized results showed an overview of the cost-effectiveness results of ACEIs and ARBs, to calculate a synthesized economic evaluation result of ACEIs and ARBs using the cost-effectiveness results in different economic evaluations could not validly be done, given all the aforementioned differences. This is mainly due to two limitations. Firstly, the baseline characteristics of the populations varied in the studies included. Secondly, the effectiveness outcomes varied in different studies.

Various selected studies in this review were strongly based on clinical trial settings. Trials are the gold standard for internal validity, but the problem is the lack of external validity [73]. The challenges and the need to include the real-world evidence in economic evaluations have been pointed out by pharmaco-economic researchers [74]. In the mentioned cost-effectiveness analysis of screening for MiA by Boersma C et al. [72], they used population-based observational data, rather than efficacy data from clinical trials. The obvious problem in these settings is how to adjust for potential confounders and this requires careful consideration. For example, the extent in which data cover the population actually using the drugs, the adverse drug events and the drug use pattern all influence the results of effectiveness analysis. Findings from drug utilization studies relevant to aspects involving (non-)adherence or safety issues should be used in future analyses of drugs' (cost-)effectiveness. In our efforts to extract some safety information from our

current included studies, only two articles [37,38] mention a higher risk of dry cough associated with ACE inhibitors and discuss whether this side effect would influence the cost-effectiveness of ACEIs. This systematic review illustrates the lack of inclusion of observational data in the pharmacoeconomic evaluations so far performed.

CONCLUSION

Considering the current evidence, both ACEIs and ARBs are cost-saving compared with conventional therapy excluding a RAAS inhibitors. There is a lack of evidence in direct comparison of these two drug classes in consistent economic evaluations. Because of the limited external validity in using RCT data and the simulation results derived from trial-based analytical models, observational data should be used to confirm these trial-based cost-effectiveness analyses' results.

COMPETING INTERESTS

YH works as a PhD student in University Medical Center Groningen (UMCG). Funding was received from University of Groningen. The results of this paper have not been presented or published elsewhere, in whole or in part.

AUTHORS' CONTRIBUTIONS

YH and QZ performed the literature search, data collection and analysis and wrote the draft. YH, FH and MP contributed to the study design, and reviewing the manuscript. All authors contributed to the conception and design, and read and approved the final manuscript.

ACKNOWLEDGEMENTS

The authors are grateful to Professor Dr. Dick de Zeeuw (UMCG) for supporting the study design and Dr. Petra Denig (UMCG) for commenting on an earlier version of this manuscript.

REFERENCES

1. Ritz E, Orth SR: Nephropathy in patients with type 2 diabetes mellitus. *N Engl J Med* 1999, 341(15):1127-1133.
2. New JP, Middleton RJ, Klebe B, Farmer CK, de Lusignan S, Stevens PE, O'Donoghue DJ: Assessing the prevalence, monitoring and management of chronic kidney disease in patients with diabetes compared with those without diabetes in general practice. *Diabet Med* 2007, 24(4):364-369.
3. Koro CE, Lee BH, Bowlin SJ: Antidiabetic medication use and prevalence of chronic kidney disease among patients with type 2 diabetes mellitus in the United States. *Clin Ther* 2009, 31(11):2608-2617.
4. Bakris GL: Recognition, pathogenesis, and treatment of different stages of nephropathy in patients with type 2 diabetes mellitus. *Mayo Clin Proc* 2011, 86(5):444-456.
5. Mogensen CE, Christensen CK, Vittinghus E: The stages in diabetic renal disease. With emphasis on the stage of incipient diabetic nephropathy. *Diabetes* 1983, 32(Suppl 2):64-78.
6. Scheffel RS, Bortolanza D, Weber CS, Costa LA, Canani LH, Santos KG, Crispim D, Roisenberg I, Lisbôa HR, Tres GS, Tschiedel B, Gross JL: Prevalence of micro and macroangiopathic chronic complications and their risk factors in the care of outpatients with type 2 diabetes mellitus. *Rev Assoc Med Bras* 2004, 50(3):263-267.
7. Ubink-Veltmaat LJ, Bilo HJ, Meyboom-de Jong B: Microalbuminuria in patients with type 2 diabetes mellitus in general practice. *Ned Tijdschr Geneesk* 2004, 148(41):2026-2030.
8. Wu AY, Kong NC, de Leon FA, Pan CY, Tai TY, Yeung VT, Yoo SJ, Rouillon A, Weir MR: An alarmingly high prevalence of diabetic nephropathy in Asian type 2 diabetic patients: the MicroAlbuminuria Prevalence (MAP) Study. *Diabetologia* 2005, 48(1):17-26.
9. Palmer AJ, Valentine WJ, Chen R, Mehin N, Gabriel S, Bregman B, Rodby RA: A health economic analysis of screening and optimal treatment of nephropathy in patients with type 2 diabetes and hypertension in the USA. *Nephrol Dial Transplant* 2008, 23(4):1216-1223.
10. Lea JP, Nicholas SB: Diabetes mellitus and hypertension: Key risk factors for kidney disease. *J Natl Med Assoc* 2002, 94(Suppl 8):s7-s15.
11. U.S. Renal Data System: USRDS 2011 Annual Data Report: Atlas of end-stage renal disease in the United States. Bethesda, MD, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases 2011.
12. Couser WG, Remuzzi G, Mendis S, Tonelli M: The contribution of chronic kidney disease to the global burden of major noncommunicable diseases. *Kidney Int* 2011, 80(12):1258-1270.
13. Rippin JD, Barnett AH, Bain SC: Cost-effective strategies in the prevention of diabetic nephropathy. *Pharmacoeconomics* 2004, 22(1):9-28.
14. American Diabetes Association: Standards of medical care in diabetes. *Diabetes Care* 2005, 28(Suppl 1):s4-s36.

15. Matchar DB, McCrory DC, Orlando LA, Patel MR, Patel UD, Patwardhan MB, Powers B, Samsa GP, Gray RN: Systematic review: comparative effectiveness of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers for treating essential hypertension. *Ann Intern Med* 2008, 148(1):16-29.
16. Pylypchuk GB: ACE-inhibitor versus angiotensin II blocker induced cough and angioedema. *Ann Pharmacother* 1998, 32(10):1060-1066.
17. Fuchs SA, Meyboom RH, van Puijenbroek EP, Guchelaar HJ: Use of angiotensin receptor antagonists in patients with ACE inhibitor induced angioedema. *Pharm World Sci* 2004, 26(4):191-192.
18. Chadban S, Howell M, Twigg S, Thomas M, Jerums G, Cass A, Campbell D, Nicholls K, Tong A, Mangos G, Stack A, MacIsaac RJ, Girgis S, Colagiuri R, Colagiuri S, Craig J; CARI: The CARI guidelines. Cost-effectiveness and socioeconomic implications of prevention and management of chronic kidney disease in type 2 diabetes. *Nephrology (Carlton)* 2010, 15(Suppl 1):s195-s203.
19. Carides GW: Losartan: A pharmaco-economic review. *JME* 2007, 10(4):573-585.
20. Bramlage P, Schindler C: Differences in pharmacology and their translation into differences in clinical efficacy--a comparison of the renin angiotensin blocking agents irbesartan and losartan. *Expert Opin Pharmacother* 2010, 11(4):521-535.
21. Boersma C, Atthobari J, Gansevoort RT, de Jong-Van den Berg LT, de Jong PE, de Zeeuw D, Annemans LJ, Postma MJ: Pharmacoeconomics of angiotensin II antagonists in type 2 diabetic patients with nephropathy: implications for decision making. *Pharmacoeconomics* 2006, 24(6):523-535.
22. Ravera M, Re M, Vettoretti S: Economic evaluation of angiotensin receptor blockers in type 2 diabetes, hypertension, and nephropathy. *J Am Soc Nephrol* 2006, 17(4 Suppl 2):s44-s48.
23. Palmer AJ, Tucker DM, Valentine WJ, Roze S, Gabriel S, Cordonnier DJ: Cost-effectiveness of irbesartan in diabetic nephropathy: a systematic review of published studies. *Nephrol Dial Transplant* 2005, 20(6):1103-1109.
24. Palmer AJ, Rodby RA: Health economics studies assessing irbesartan use in patients with hypertension, type 2 diabetes, and microalbuminuria. *Kidney Int Suppl* 2004, (92):s118-s120.
25. Postma MJ, Kruidhof H, de Jong-van den Berg LT, de Zeeuw D: Pharmacoeconomic aspects of losartan treatment to delay progression of renal disease in patients with Type 2 diabetes. *Expert Opin Pharmacother* 2003, 4(9):1543-1550.
26. Postma MJ, de Zeeuw D: The economic benefits of preventing end-stage renal disease in patients with type 2 diabetes mellitus. *Nephrol Dial Transplant* 2009, 24(10):2975-2983.
27. Drummond MF, Sculpher MJ, Torrance GW, O'Brien BJ, Stoddart GL: *Methods for the Economic Evaluation of Health Care Programmes*. Oxford University Press; 2005.
28. Inflation, GDP deflator (annual %) [<http://data.worldbank.org/indicator/NY.GDP.DEFL.KD.ZG>]
29. Currency and foreign exchange [<http://www.xe.com>]

30. Grosse SD: Assessing cost-effectiveness in health care: history of the \$50,000 per QALY threshold. *Value Health* 2008, 8(2):165-178.
31. Li R, Zhang P, Barker LE, Chowdhury FM, Zhang X: Cost-effectiveness of interventions to prevent and control diabetes mellitus: a systematic review. *Diabetes Care* 2010, 33(8):1872-1894.
32. Golan L, Birkmeyer JD, Welch HG: The cost-effectiveness of treating all patients with type 2 diabetes with angiotensin-converting enzyme inhibitors. *Ann Intern Med*. 1999, 131(9):660-667.
33. Sakthong P, Tangphao O, Eiam-Ong S, Kamolratanakul P, Supakankunti S, Himathongkam T, Yathavong K: Cost-effectiveness of using angiotensin-converting enzyme inhibitors to slow nephropathy in normotensive patients with diabetes type II and microalbuminuria. *Nephrology* 2001, 6(2):71-77.
34. Rosen AB, Hamel MB, Weinstein MC, Cutler DM, Fendrick AM, Vijan S: Cost-effectiveness of full medicare coverage of angiotensin-converting enzyme inhibitors for beneficiaries with diabetes. *Ann Intern Med* 2005, 143(2):89-99.
35. Campbell HM, Boardman KD, Dodd MA, Raisch DW: Pharmacoeconomic analysis of angiotensin-converting enzyme inhibitors in type 2 diabetes: a Markov model. *Ann Pharmacother* 2007, 41(7):1101-1110.
36. Adarkwah CC, Gandjour A: Cost-effectiveness of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers in newly diagnosed type 2 diabetes in Germany. *Int J Technol Assess Health Care* 2010, 26(1):62-70.
37. Adarkwah CC, Gandjour A, Akkerman M, Evers SM: Cost-effectiveness of angiotensin-converting enzyme inhibitors for the prevention of diabetic nephropathy in The Netherlands--a Markov model. *PLoS One* 2011, 6(10):e26139.
38. Lewis EJ, Hunsicker LG, Bain RP, Rohde RD: The effect of angiotensin-converting-enzyme inhibition on diabetic nephropathy. The Collaborative Study Group. *N Engl J Med* 1993, 329(20):1456-1462.
39. Ravid M, Savin H, Jutrin I, Bental T, Katz B, Lishner M: Long-term stabilizing effect of angiotensin-converting enzyme inhibition on plasma creatinine and on proteinuria in normotensive type II diabetic patients. *Ann Intern Med* 1993, 118(8):577-581.
40. Ravid M, Brosh D, Levi Z, Bar-Dayyan Y, Ravid D, Rachmani R: Use of enalapril to attenuate decline in renal function in normotensive, normoalbuminuric patients with type 2 diabetes mellitus. A randomized, controlled trial. *Ann Intern Med* 1998, 128(12 Pt 1):982-988.
41. Ravid M, Lang R, Rachmani R, Lishner M: Long-term renoprotective effect of angiotensin-converting enzyme inhibition in non-insulin-dependent diabetes mellitus. A 7-year follow-up study. *Arch Intern Med* 1996, 156(3):286-289.
42. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. Heart Outcomes Prevention Evaluation Study Investigators. *Lancet* 2000, 355(9200):253-259.

43. Herman WH, Shahinfar S, Carides GW, Dasbach EJ, Gerth WC, Alexander CM, Cook JR, Keane WF, Brenner BM: Losartan reduces the costs associated with diabetic end-stage renal disease: the RENAAL study economic evaluation. *Diabetes Care* 2003, 26(3):683-687.
44. Souchet T, Durand Zaleski I, Hannedouche T, Rodier M, Gaugris S, Passa P; RENAAL study: An economic evaluation of Losartan therapy in type 2 diabetic patients with nephropathy: an analysis of the RENAAL study adapted to France. *Diabetes Metab* 2003, 29(1):29-35.
45. Burgess ED, Carides GW, Gerth WC, Marentette MA, Chabot I; Canadian Hypertension Society: Losartan reduces the costs associated with nephropathy and end-stage renal disease from type 2 diabetes: Economic evaluation of the RENAAL study from a Canadian perspective. *Can J Cardiol* 2004, 20(6):613-618.
46. Szucs TD, Sandoz MS, Keusch GW: The cost-effectiveness of losartan in type 2 diabetics with nephropathy in Switzerland--an analysis of the RENAAL study. *Swiss Med Wkly* 2004, 134(31-32):440-447.
47. Seng WK, Hwang SJ, Han DC, Teong CC, Chan J, Burke TA, Carides GW, Choi YJ: Losartan reduces the costs of diabetic end-stage renal disease: an Asian perspective. *Nephrology (Carlton)* 2005, 10(5):520-524.
48. Arredondo A, Burke TA, Carides GW, Lemus E, Querol J: The impact of losartan on the lifetime incidence of ESRD and costs in Mexico. *Rev Invest Clin* 2005, 57(3):399-405.
49. Vora J, Carides G, Robinson P: Effects of Losartan-based therapy on the incidence of end-stage renal disease and associated costs in type 2 diabetes mellitus: A retrospective cost-effectiveness analysis in the United Kingdom. *Curr Ther Res Clin E* 2005, 66(6):475-485.
50. Carides GW, Shahinfar S, Dasbach EJ, Keane WF, Gerth WC, Alexander CM, Herman WH, Brenner BM; RENAAL Investigators: The impact of losartan on the lifetime incidence of end-stage renal disease and costs in patients with type 2 diabetes and nephropathy. *Pharmacoeconomics* 2006, 24(6):549-558.
51. Stafylas PC, Sarafidis PA, Lasaridis AN, Tsakni E, Niakas DA, Dombros NV, Grekas DM, Bakris GL: Cost-effectiveness of losartan in diabetic nephropathy: a Greek perspective. *J Nephrol* 2007, 20(6):703-715.
52. de Portu S, Citarella A, Cammarota S, Menditto E, Mantovani LG: Pharmaco-economic consequences of losartan therapy in patients undergoing diabetic end stage renal disease in EU and USA. *Clin Exp Hypertens* 2011, 33(3):174-178.
53. Rodby RA, Chiou CF, Borenstein J, Smitten A, Sengupta N, Palmer AJ, Roze S, Annemans L, Simon TA, Chen RS, Lewis EJ; Collaborative Study Group: The cost-effectiveness of irbesartan in the treatment of hypertensive patients with type 2 diabetic nephropathy. *Clin Ther* 2003, 25(7):2102-2119.

54. Palmer AJ, Annemans L, Roze S, Lamotte M, Rodby RA, Cordonnier DJ: An economic evaluation of irbesartan in the treatment of patients with type 2 diabetes, hypertension and nephropathy: cost-effectiveness of Irbesartan in Diabetic Nephropathy Trial (IDNT) in the Belgian and French settings. *Nephrol Dial Transplant* 2003, 18(10):2059-2066.
55. Coyle D, Rodby RA: Economic evaluation of the use of irbesartan and amlodipine in the treatment of diabetic nephropathy in patients with hypertension in Canada. *Can J Cardiol* 2004, 20(1):71-79.
56. Palmer AJ, Annemans L, Roze S, Lamotte M, Rodby RA, Bilous RW: An economic evaluation of the Irbesartan in Diabetic Nephropathy Trial (IDNT) in a UK setting. *J Hum Hypertens* 2004, 18(10):733-738.
57. Palmer AJ, Annemans L, Roze S, Lamotte M, Lapuerta P, Chen R, Gabriel S, Carita P, Rodby RA, de Zeeuw D, Parving HH: Cost-effectiveness of early irbesartan treatment versus control (standard antihypertensive medications excluding ACE inhibitors, other angiotensin-2 receptor antagonists, and dihydropyridine calcium channel blockers) or late irbesartan treatment in patients with type 2 diabetes, hypertension, and renal disease. *Diabetes Care* 2004, 27(8):1897-1903.
58. Palmer AJ, Annemans L, Roze S, Lapuerta P, Chen R, Gabriel S, Carita P, Rodby RA, de Zeeuw D, Parving HH, De Alvaro F: Irbesartan is projected to be cost and life saving in a Spanish setting for treatment of patients with type 2 diabetes, hypertension, and microalbuminuria. *Kidney Int Suppl* 2005, (93):s52-s54.
59. Palmer AJ, Roze S, Valentine WJ, Ray JA, Frei A, Burnier M, Hess B, Spinass GA, Brändle M: Health economic implications of irbesartan plus conventional antihypertensive medications versus conventional blood pressure control alone in patients with type 2 diabetes, hypertension, and renal disease in Switzerland. *Swiss Med Wkly* 2006, 136(21-22):346-352.
60. Palmer AJ, Valentine WJ, Tucker DM, Ray JA, Roze S, Annemans L, Lapuerta P, Chen R, Gabriel S, Carita P, Rodby RA, de Zeeuw D, Parving HH, Laville M: A French cost-consequence analysis of the renoprotective benefits of irbesartan in patients with type 2 diabetes and hypertension. *Curr Med Res Opin* 2006, 22(11):2095-2100.
61. Palmer AJ, Valentine WJ, Ray JA, Roze S, Muszbek N: Health economic implications of irbesartan treatment versus standard blood pressure control in patients with type 2 diabetes, hypertension and renal disease: a Hungarian analysis. *Eur J Health Econ* 2007, 8(2):161-168.
62. Palmer AJ, Valentine WJ, Ray JA: Irbesartan treatment of patients with type 2 diabetes, hypertension and renal disease: a UK health economics analysis. *Int J Clin Pract* 2007, 61(10):1626-1633.
63. Coyle D, Rodby R, Soroka S, Levin A, Muirhead N, de Cotret PR, Chen R, Palmer A: Cost-effectiveness of irbesartan 300 mg given early versus late in patients with hypertension and a history of type 2 diabetes and renal disease: a Canadian perspective. *Clin Ther* 2007, 29(7):1508-1523.
64. Yang WC, Hwang SJ, Annemans L, Ray JA, Roze S, Valentine WJ, Palmer AJ: Irbesartan and amlodipine in the treatment of patients with microalbuminuria, hypertension and type 2 diabetes in Taiwan: a modelling projection over 25 years. *JME* 2007, 10(4):539-552.

65. Annemans L, Demarteau N, Hu S, Lee TJ, Morad Z, Supaporn T, Yang WC, Palmer AJ: An Asian regional analysis of cost-effectiveness of early irbesartan treatment versus conventional antihypertensive, late amlodipine, and late irbesartan treatments in patients with type 2 diabetes, hypertension, and nephropathy. *Value Health* 2008, 11(3):354-364.
66. Smith DG, Nguyen AB, Peak CN, Frech FH: Markov modeling analysis of health and economic outcomes of therapy with valsartan versus amlodipine in patients with type 2 diabetes and microalbuminuria. *J Manag Care Pharm* 2004, 10(1):26-32.
67. Brenner BM, Cooper ME, de Zeeuw D, Keane WF, Mitch WE, Parving HH, Remuzzi G, Snapinn SM, Zhang Z, Shahinfar S; RENAAL Study Investigators: Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001, 345(12):861-869.
68. Lewis EJ, Hunsicker LG, Clarke WR, Berl T, Pohl MA, Lewis JB, Ritz E, Atkins RC, Rohde R, Raz I; Collaborative Study Group: Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001, 345(12):851-860.
69. Parving HH, Lehnert H, Bröchner-Mortensen J, Gomis R, Andersen S, Arner P; Irbesartan in Patients with Type 2 Diabetes and Microalbuminuria Study Group: The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. *N Engl J Med* 2001, 345(12):870-878.
70. Viberti G, Wheeldon NM; MicroAlbuminuria Reduction With VALsartan (MARVAL) Study Investigators: Microalbuminuria reduction with valsartan in patients with type 2 diabetes mellitus: a blood pressure-independent effect. *Circulation* 2002, 106(6):672-678.
71. Atthobari J, Asselbergs FW, Boersma C, de Vries R, Hillege HL, van Gilst WH, Gansevoort RT, de Jong PE, de Jong-van den Berg LT, Postma MJ; PREVEND IT Study Group: Cost-effectiveness of screening for albuminuria with subsequent fosinopril treatment to prevent cardiovascular events: A pharmaco-economic analysis linked to the prevention of renal and vascular endstage disease (PREVEND) study and the prevention of renal and vascular endstage disease intervention trial (PREVEND IT). *Clin Ther* 2006, 28(3):432-444.
72. Boersma C, Gansevoort RT, Pechlivanoglou P, Visser ST, van Toly FF, de Jong-van den Berg LT, de Jong PE, Postma MJ; Prevention of Renal and Vascular End Stage Disease Study Group: Screen-and-treat strategies for albuminuria to prevent cardiovascular and renal disease: cost-effectiveness of nationwide and targeted interventions based on analysis of cohort data from the Netherlands. *Clin Ther* 2010, 32(6):1103-1121.
73. Drummond MF: Experimental versus observational data in the economic evaluation of pharmaceuticals. *Med Decis Making* 1998, 18(Suppl 2):s12-s18.
74. Baltussen R, Leidl R, Ament A: Real world designs in economic evaluation. Bridging the gap between clinical research and policy-making. *Pharmacoeconomics* 1999, 16(5 Pt 1):449-458.

