

## University of Groningen

### Nitric oxide donor molsidomine favors features of atherosclerotic plaque stability and reduces myocardial infarction in mice

Roth, Lynn; Van der Donckt, Carole; Veseli, Besa Emini; Van Dam, Debby; De Deyn, Peter P.; Martinet, Wim; Herman, Arnold G.; De Meyer, Guido R. Y.

*Published in:*  
Vascular pharmacology

*DOI:*  
[10.1016/j.vph.2019.05.001](https://doi.org/10.1016/j.vph.2019.05.001)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2019

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Roth, L., Van der Donckt, C., Veseli, B. E., Van Dam, D., De Deyn, P. P., Martinet, W., Herman, A. G., & De Meyer, G. R. Y. (2019). Nitric oxide donor molsidomine favors features of atherosclerotic plaque stability and reduces myocardial infarction in mice. *Vascular pharmacology*, 118-119, Article 106561. <https://doi.org/10.1016/j.vph.2019.05.001>

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*



## Nitric oxide donor molsidomine favors features of atherosclerotic plaque stability and reduces myocardial infarction in mice



Lynn Roth<sup>a,1</sup>, Carole Van der Donckt<sup>a,1</sup>, Besa Emini Veseli<sup>a</sup>, Debby Van Dam<sup>b,c</sup>, Peter P. De Deyn<sup>b,c,d</sup>, Wim Martinet<sup>a</sup>, Arnold G. Herman<sup>a</sup>, Guido R.Y. De Meyer<sup>a,\*</sup>

<sup>a</sup> Laboratory of Physiopharmacology, University of Antwerp, Antwerp, Belgium

<sup>b</sup> Laboratory of Neurochemistry and Behaviour, Institute Born-Bunge, University of Antwerp, Antwerp, Belgium

<sup>c</sup> Department of Neurology and Alzheimer Research Center, University of Groningen and University Medical Center Groningen, Hanzeplein 1, 9713, GZ, Groningen, The Netherlands

<sup>d</sup> Department of Neurology, Memory Clinic of Hospital Network Antwerp (ZNA) Middelheim and Hoge Beuken, Lindendreef 1, 2020 Antwerp, Belgium

### ABSTRACT

Nitric oxide (NO) donors are commonly used for the prevention and treatment of ischemic heart disease. Besides their effects on the heart, NO donors may also prevent hypoxic brain damage and exert beneficial effects on atherosclerosis by favoring features of plaque stability. We recently described that *apolipoprotein E* (*ApoE*) deficient mice with a mutation in the *fibrillin-1* (*Fbn1*) gene (*ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup>) develop accelerated atherosclerosis, plaque rupture, myocardial infarction, cerebral hypoxia and sudden death. In the present study, we evaluated the effects of chronic treatment with the NO donor molsidomine on atherosclerotic plaque stability, cardiac function, neurological symptoms and survival in the *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> mouse model. Female *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> mice were fed a Western diet (WD). After 8 weeks of WD, the mice were divided into two groups receiving either molsidomine via the drinking water (1 mg/kg/day; n = 34) or tap water (control; n = 36) until 25 weeks of WD. Survival tended to increase after molsidomine treatment (68% vs. 58% in controls). Importantly, atherosclerotic plaques of molsidomine-treated mice had a thicker fibrous cap (11.1 ± 1.2 vs. 8.1 ± 0.7 μm) and showed an increased occurrence of plaque macrocalcifications (30% vs. 0%), indicative of a more stable phenotype. Molsidomine also improved cardiac function, as fractional shortening was increased (40 ± 2% vs. 27 ± 2%) combined with a decreased end diastolic (3.1 ± 0.2 vs. 3.9 ± 0.2 mm) and end systolic diameter (1.9 ± 0.1 vs. 2.9 ± 0.2 mm). Furthermore, perivascular fibrosis (23 ± 2 vs. 30 ± 2%) and the occurrence of myocardial infarctions (12% vs. 36%) was significantly reduced. Track width, a measure of the animal's hind limb base of support and representative of hypoxic brain damage, was also normalized as a result of molsidomine treatment (2.54 ± 0.04 vs. 2.91 ± 0.09 cm in controls). These findings demonstrate that the NO donor molsidomine improves cardiac function, reduces neurological symptoms and enhances atherosclerotic plaque stability.

### 1. Introduction

Nitric oxide (NO) is involved in a wide range of physiological processes in the cardiovascular system [1]. In blood vessels, NO is an important determinant of vascular tone, regulating blood pressure and vascular compliance. Moreover, NO maintains vascular integrity by inhibiting smooth muscle cell (SMC) proliferation, platelet aggregation and leukocyte adhesion to the endothelium [1–3]. In the heart, NO plays a major role in determining the resting vascular tone of coronary resistance vessels [4,5] and is produced in cardiac SMC, where it regulates myocardial contractility [2,5]. Physical or biological injury to the endothelium impairs the production and/or bioavailability of NO, which may play a role in hypertension, reperfusion injury, atherosclerosis and myocardial depression [1,6,7].

In line with these findings, patients suffering from ischemic cardiac failure and atherosclerosis might benefit from chronic treatment with

an exogenous NO donor. The NO donor molsidomine has the advantage that it releases NO non-enzymatically, avoiding tolerance. Our group and others have previously shown that molsidomine, which exerts its effects through the metabolite 3-morpholinolysidonimine (SIN-1), favoured cardiac function [8–12], decreased platelet aggregation [11,13,14] and signs of oxidative stress and increased features of plaque stability [15] in humans, as well as in animal models. Moreover, patients treated with molsidomine for over a year had significantly lower plasma levels of soluble adhesion molecule ICAM-1, which may slow down the progression of atherosclerosis [16]. However, the beneficial effects of molsidomine are dose-dependent, since high amounts can generate peroxynitrite, causing enhanced oxidative stress, thereby aggravating atherosclerosis [17].

We recently described a novel atherosclerosis model, apolipoprotein E-deficient mice with a mutation in the fibrillin-1 gene (*ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup>), characterized by accelerated atherosclerosis, plaque

\* Corresponding author at: Laboratory of Physiopharmacology, University of Antwerp, Universiteitsplein 1, B-2610 Antwerp, Belgium.

E-mail address: [guido.demeyer@uantwerpen.be](mailto:guido.demeyer@uantwerpen.be) (G.R.Y. De Meyer).

<sup>1</sup> Equal contribution.

rupture, myocardial infarction, cerebral hypoxia and sudden death [18]. In these mice, the elastic fibers in the vessel wall are fragmented, resulting in enhanced arterial stiffness and thereby mimicking vascular aging [19]. As a result, *ApoE<sup>-/-</sup>Fbn1<sup>C1039G+/-</sup>* mice develop large and highly unstable plaques with a phenotype similar to human plaques. Moreover, these mice develop coronary artery plaques and left ventricular hypertrophy, resulting in decreased cardiac function and myocardial infarction. In the present study, we evaluated the effects of chronic molsidomine treatment in *ApoE<sup>-/-</sup>Fbn1<sup>C1039G+/-</sup>* mice on atherosclerotic plaque stability, cardiac function, neurological symptoms and sudden death.

## 2. Methods

### 2.1. Mice

Female *ApoE<sup>-/-</sup>Fbn1<sup>C1039G+/-</sup>* mice (n = 70, 6 weeks of age) were fed a Western-type diet (WD; TD88137, Harlan Teklad, Madison, WI) for up to 25 weeks. After 8 weeks of WD feeding, animals were divided into two groups receiving either molsidomine via the drinking water (1 mg/kg body weight/day; n = 34) or tap water (control; n = 36) until 25 weeks of WD. The animals were housed in a temperature-controlled room with a 12-hour light/dark cycle with free access to water and food. Mice were inspected daily for occurrence of sudden death. At the end of the experiment, mice were anesthetized with sodium pentobarbital (75 mg/kg, i.p.) and blood samples were taken from the retro-orbital plexus for plasma cholesterol analysis. Subsequently, animals were euthanized with an overdose of sodium pentobarbital (250 mg/kg, i.p.). Total plasma cholesterol was analysed by means of a commercial kit (Randox, Crumlin, UK). All experiments were approved by the ethical committee of the University of Antwerp and were performed according to the guidelines from Directive 2010/63/EU of the European Parliament on the protection of animals used for scientific purposes.

### 2.2. Echocardiography

Transthoracic echocardiograms (ECGs) of *ApoE<sup>-/-</sup>Fbn1<sup>C1039G+/-</sup>* mice were performed at the start (baseline, 8 weeks on WD) and after 12 weeks of molsidomine treatment. ECGs were performed on anesthetized mice (isoflurane, IsoFlo, Abbott) using a Toshiba diagnostic ultrasound system (SSA-700A), equipped with a 15 MHz transducer.

### 2.3. Blood pressure

Peripheral blood pressure was measured with a tail cuff (Coda™ High throughput, Kent Scientific, Torrington, CT). Mice were trained for four weeks before the baseline measurement and every week thereafter, until 12 weeks of molsidomine treatment, when a final measurement was performed.

### 2.4. Motor coordination

Track width was analysed after 12 weeks of treatment as described previously [20]. Briefly, ink was applied to the animals' hind paws and the mice were required to walk on a strip of paper towards a dark goal box. The median value of a minimum of 10 measurements per mouse was used.

### 2.5. Histological evaluation

After euthanasia or sudden death, mice were dissected and aorta, heart, spleen and kidneys were weighed (dry weight). Heart and arteries were post-fixed in 4% formaldehyde (pH 7.4) for 24 h, processed and embedded in paraffin. Histological analyses were performed on serial cross sections (5 µm) of the brachiocephalic artery and proximal ascending aorta. The percentage stenosis and the size of atherosclerotic

plaques and necrotic cores was measured on hematoxylin and eosin (H & E) stained sections. The necrotic cores were defined as acellular/anuclear areas (negative for hematoxylin) with a 3000 µm<sup>2</sup> threshold implemented to avoid counting of very small H&E negative areas that likely do not represent areas of necrosis. The presence of macrocalcifications, identified as basophilic (dark purple) crystalline structures, was determined on H&E stained sections. The presence of 3-nitrotyrosine in plaques of the proximal ascending aorta was identified by immunohistochemical staining with an anti-nitrotyrosine antibody (sc-55,256, Santa cruz). Cellular composition of the atherosclerotic plaques was analysed by immunohistochemistry using anti-mannose receptor (macrophages, ab64693, Abcam) and anti-α-smooth muscle cell actin (F3777, Sigma-Aldrich). Fibrous cap thickness was determined as the median value of 10 measurements per atherosclerotic plaque on α-SMC actin (F3777, Sigma-Aldrich) stained sections. Collagen content was determined on Sirius red stained sections with light microscopy (total collagen) and polarized light (type I collagen). Fibrous cap thickness was determined as the median value of 10 measurements per atherosclerotic plaque on α-SMC actin (F3777, Sigma-Aldrich) stained sections. The ICAM-1 expression of the endothelial cells was measured via immunohistochemical staining with an anti-vWF (PC054, The Binding Site) and anti-ICAM-1 antibody (550287, BD Biosciences). The length of CD31 positive cells along the luminal border was quantified in a complete cross-section of the proximal ascending aorta and ICAM-1 positivity was expressed as percentage of CD31 positivity. Intraplaque neovascularization and hemorrhages in the left common carotid artery were examined on slides that were stained with anti-vWF (PC054, The Binding Site) and anti-TER119 (550,565, BD Biosciences), respectively. The occurrence of myocardial infarctions, coronary plaques and perivascular fibrosis, measured as the perivascular collagen area divided by the luminal area (PVCA/LA) of 10 coronary arteries per mouse, was analysed on Masson's trichrome stained transversal sections (cut from the middle of the heart to the apex). The myocardial infarctions were defined as large fibrotic areas with infiltration of inflammatory cells. If plaques were present in the coronary arteries, the plaque size and percentage of stenosis were measured on Masson's trichrome stained sections. All images were acquired with the Universal Grab 6.1. (IDL) software (Exelis, Boulder, CO) using an Olympus BX40 microscope (Tokyo, Japan). Images were quantified using Image J software (NIH).

### 2.6. Statistical analyses

Normally distributed data are expressed as mean ± SEM and non-normally distributed variables are represented as median [min-max]. Statistical analyses were performed using SPSS software (version 24, SPSS Inc., Chicago). Statistical tests are specified in the figure and table legends. Histological data of the proximal ascending aorta and left common carotid artery only include mice that survived the experiment. Data on MI and coronary arteries include all mice. Differences were considered significant at P < 0.05.

## 3. Results

Body weight and total plasma cholesterol was similar in treated and control mice (Table 1). Survival improved slightly, albeit not significantly, after molsidomine treatment (68% vs. 58% in controls, Table 1). A marked decrease in kidney and spleen weight was observed in the molsidomine-treated group, whereas lung weight remained equal between both groups (Table 1).

### 3.1. Molsidomine improves atherosclerotic plaque stability

Molsidomine did not affect plaque size in the proximal ascending aorta nor the degree of stenosis in *ApoE<sup>-/-</sup>Fbn1<sup>C1039G+/-</sup>* mice (Table 2, Fig. S1). Furthermore, the NO donor had no effect on the size of the necrotic core, the amount of macrophages, smooth muscle cells (SMCs),

**Table 1**  
Characteristics of control and molsidomine-treated *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> mice.

	Control	Molsidomine
Survival (%)	21/36 (58%)	23/34 (68%)
Body weight (g)	23 ± 1	23 ± 1
Plasma cholesterol (mg/dl)	616 ± 57	621 ± 51
Organ weights (mg)		
Spleen	174 ± 15	107 ± 9**
Lungs	229 ± 25	219 ± 18
Kidney	316 ± 23	242 ± 9**

Survival was evaluated using a Pearson's chi-square test. Body weight, plasma cholesterol and organ weights are presented as mean ± SEM (n = 13–18 per group) and analysed by an independent samples *t*-test. \*P < 0.05, \*\*P < 0.01 vs. control mice.

total collagen, type I collagen and endothelial ICAM-1 expression in the plaque (Table 2, Fig. S1). Nonetheless, the fibrous cap was significantly thicker in molsidomine-treated mice as compared to controls (Table 2, Fig. S1). Molsidomine also induced the formation of intraplaque macrocalcifications, as none of the control mice showed any signs of calcified plaque regions as compared to 30% in the molsidomine-treated mice (Fig. 1). The macrocalcifications were always located at the medial side of the plaque. No signs of calcification were observed in the fibrous cap. The calcified plaques showed a significantly higher type I collagen content as compared to non-calcified plaques of molsidomine-treated mice and controls (Table 2). Furthermore, the calcified plaque area was negatively correlated with plaque size in the proximal ascending aorta (Pearson correlation; R square = 0.868, P = 0.021). Importantly, there was no difference in the percentage of 3-nitrotyrosine, indicative of peroxynitrite-related oxidative stress, in plaques of control and molsidomine-treated mice (Fig. S2). The occurrence of intraplaque neovascularization (control: 83%; molsidomine: 75%), the number of microvessels (control: 4 [0–19]; molsidomine: 3 [0–19]) and the occurrence of hemorrhages in the left common carotid artery (control: 17%; molsidomine: 25%) was not affected by molsidomine treatment.

### 3.2. Molsidomine reduces cardiac mass and improves cardiac function and left ventricular hypertrophy

Systolic and diastolic peripheral blood pressure did not differ between the molsidomine and control group (Table 3). Echocardiography of the heart revealed a significantly higher fractional shortening (FS) and lower end diastolic diameter (EDD) and end systolic diameter (ESD) in molsidomine-treated mice compared to controls (Fig. 2A–C). Additionally, molsidomine significantly reduced heart mass as compared to the *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> control mice (Fig. 2D). Cross-

**Table 2**  
Atherosclerotic plaque characteristics.

	Control <sup>a,b</sup>	Molsidomine <sup>a</sup>	Molsidomine non calcified <sup>b</sup>	Molsidomine calcified <sup>b</sup>
Vessel area (× 10 <sup>3</sup> μm <sup>2</sup> )	1379 ± 100	1284 ± 117	1399 ± 133	1015 ± 210
Plaque size (× 10 <sup>3</sup> μm <sup>2</sup> )	798 ± 67	792 ± 73	861 ± 83	630 ± 133
Stenosis (%)	58 ± 3	62 ± 2	62 ± 3	62 ± 4
Necrotic core (%)	8.6 ± 1.1	10.1 ± 1.4	10.9 ± 1.9	8.0 ± 1.5
Macrophages (%)	1.8 ± 0.3	2.7 ± 0.5	2.6 ± 0.6	3.0 ± 0.7
Total collagen (%)	39.8 ± 1.5	39.7 ± 1.2	38.2 ± 1.5	43.2 ± 1.4
Type I collagen (%)	3.4 ± 0.4	3.5 ± 0.4	2.8 ± 0.4	5.0 ± 0.5*, <sup>§§</sup>
Smooth muscle cells (%)	15.5 ± 1.1	14.2 ± 0.9	13.8 ± 1.3	15.3 ± 0.8
Fibrous cap thickness (μm)	8.1 ± 0.7	11.1 ± 1.2*	10.5 ± 1.3	12.4 ± 2.6
ICAM-1 positive ECs (%)	94.4 ± 1.5	94.8 ± 1.4	94.5 ± 1.6	95.5 ± 2.8

Data from proximal ascending aorta, mean ± SEM, ECs = endothelial cells.

<sup>a</sup> n = 20 per group. Independent samples *t*-test.

<sup>b</sup> Control: n = 20, molsidomine non calcified: n = 14 and molsidomine calcified: n = 6. One-way ANOVA, post hoc LSD.

\* P < 0.05 vs. control mice.

<sup>§§</sup> P < 0.01 vs. molsidomine non calcified.

sectional histological evaluation of the heart showed a decrease of one third in the occurrence of myocardial infarctions after molsidomine treatment (Table 3 and Fig. 2E). The infarcted area remained equal in treated and control mice (Table 3). The occurrence of coronary artery plaque and the percentage stenosis did not differ between molsidomine-treated and control mice (Table 3). However, coronary perivascular fibrosis was significantly reduced in the treatment group (Fig. 2F).


### 3.3. Molsidomine improves motor function of *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> mice

Track width measurements were previously validated as a reliable marker for hypoxic brain damage in *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> mice [20]. Molsidomine was able to inhibit the increase in track width normally observed in *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> mice fed a WD. Both the median and maximum track width values were significantly lower after 12 weeks of molsidomine treatment compared with control mice (Fig. 3A–B). Track width was also positively correlated with plaque size in the proximal ascending aorta (Fig. 3C).

## 4. Discussion

The present study confirms the beneficial effects of molsidomine in the treatment of ischemic and left ventricular cardiac failure, as seen in humans [8]. Moreover, molsidomine prevented a loss in motor function and favoured atherosclerotic plaque stability. Molsidomine is a venodilator that reduces the cardiac preload, thereby lowering cardiac output, ventricular work and myocardial oxygen consumption [9]. Furthermore, it evokes a local dilatation of the coronary arteries, reducing coronary artery stenosis and preventing cardiac ischemia [10]. *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> mice are characterized by the development of pronounced atherosclerosis, which is caused by stiffening and inflammation of the arteries, as a result of elastin fragmentation in the vessel wall [18,19]. These mice develop progressive cardiac failure, coronary artery plaque, myocardial infarctions and hypoxic brain damage, similar as seen in humans [18]. Therefore, they could benefit from treatment with an exogenous NO donor. We chose for a chronic treatment with a low dose of molsidomine (1 mg/kg/day). It is very important to select the most optimal dose, because high dosages can generate peroxynitrite, causing enhanced oxidative stress, which may lead to endothelial dysfunction [17]. This was also reported during long-term treatment with other NO donors (nitroglycerin, isosorbide mononitrate) [21]. Therefore, it is important to evaluate 3-nitrotyrosine as a marker of peroxynitrite formation. In the current study, we could not detect a difference between 3-nitrotyrosine in plaques of control and molsidomine-treated mice. Thus, molsidomine at a dose of 1 mg/kg/day does not lead to oxidative stress-related side effects [15].

Importantly, molsidomine significantly improved cardiac function



**Fig. 1.** Presence of plaque macrocalcifications. (A) H&E stained sections of the proximal ascending aorta showing the presence of a macrocalcification (dark purple) as a result of molsidomine treatment (scale bar = 100  $\mu\text{m}$ ). (B) Percentage of mice showing plaque macrocalcifications in the proximal ascending aorta. Pearson's chi-square test; \*\* $P < 0.01$  vs. control mice. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

**Table 3**

Blood pressure, MI and coronary artery plaque characteristics.

	Control	Molsidomine
Blood pressure (mm Hg)		
Systolic BP	112 $\pm$ 7	111 $\pm$ 5
Diastolic BP	79 $\pm$ 7	77 $\pm$ 4
Pulse pressure	35 $\pm$ 1	33 $\pm$ 2
Myocardial infarction (MI)		
Occurrence of MI	12/33 (36%)	4/33 (12%)*
Infarcted area (%)	1.8 $\pm$ 0.4	1.3 $\pm$ 0.5
Coronary plaque		
Occurrence of coronary plaque	15/33 (45%)	14/33 (42%)
Stenosis (%)	58 $\pm$ 9	72 $\pm$ 8

Blood pressure is presented as mean  $\pm$  SEM (n = 13–18 per group) and analysed by Independent samples *t*-test. Occurrence of MI and coronary plaque were evaluated using a Pearson's chi-square test. Infarcted area (in mice with MI) and coronary stenosis (in mice with coronary plaque) are presented as mean  $\pm$  SEM and were analysed by an Independent samples *t*-test.


\*  $P < 0.05$  vs. control mice.

(increased FS) and left ventricular hypertrophy (decreased EDD and ESD) in *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> mice. Concomitant with these observations, a significant decrease in heart weight was observed, as well as a clear decrease in the occurrence of myocardial infarctions. Molsidomine did not significantly affect systolic or diastolic blood pressure, consistent with previous observations in humans [8]. The observed influence of molsidomine treatment on the heart is probably related to its hemodynamic effects, but we cannot exclude that also other pharmacological actions play a role. Further research, in which the response to other NO donors is compared with molsidomine, is necessary to study which mechanism of action contributes to the protective effects of molsidomine on the heart.


Apart from its cardiac effects, molsidomine prevented an increase in track width in *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> mice. We previously reported that measuring track width in this mouse model is a reliable technique to monitor hypoxic brain damage, as high track width values correspond to the presence of more pyknotic neurons in the cerebral cortex [20]. Furthermore, track width correlated with plaque size in the proximal ascending aorta [20], which we confirmed in the current study. As molsidomine treatment resulted in lower track width values, it is likely that it is able to prevent cerebral damage in *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> mice. It is well-known that NO plays an important role in the regulation of cerebral blood flow in normal and pathological conditions [22].

Therefore NO donors have been investigated as a therapeutic strategy for preventing hypoxic brain damage. For instance, treatment with NO donors prevented hypoxia-ischemia induced neurological damage in rats [23,24] and a reduced brain infarct size was observed in several animal models of stroke [25]. The trials involving the use of molsidomine are limited, but a recent observational study showed that after subarachnoid haemorrhage, molsidomine treatment resulted in less vasospasm-related infarctions and a better clinical outcome [26]. These beneficial effects are not only attributed to the improved cerebral blood flow, but also to other cellular and metabolic effects, such as the inhibition of neuronal apoptosis after ischemia [26]. Thus, the current study further supports the finding that molsidomine treatment can prevent hypoxic brain damage, but further clinical trials are needed to confirm these results.

We also investigated whether molsidomine could have beneficial effects on atherosclerosis. We did not observe a reduction in plaque size nor degree of stenosis, confirming our previous observations [15]. Although macrophage, SMC and total collagen content remained unaltered, the NO donor significantly increased fibrous cap thickness, a major determinant of tensile strength of the cap. NO is able to down-regulate p53 and reduce the expression of pro-inflammatory cytokines IL-1 and TNF- $\alpha$ , thereby decreasing apoptosis and inflammation [27]. In this way, SMC survival is favoured, leading to a thicker fibrous cap and a more stable plaque phenotype. Moreover, molsidomine treatment resulted in a significant increase in plaque macrocalcifications located near the medial border. It is known that calcified coronary nodules are a feature of acute coronary syndromes [28,29]. However, the type of calcification observed in the present study does not resemble calcified nodules since it is not eruptive and does not protrude into the lumen. Furthermore, it is described that coronary artery calcification (CAC) is associated with adverse cardiovascular events in humans, although it is not correlated with plaque vulnerability [30–32]. Importantly, statin therapy increases CAC, which is not related with an increased risk of clinical complications [33–36]. This suggests that the increase in CAC by statin therapy is plaque stabilizing rather than harmful. Thus, there is still no consensus on the impact of calcification on plaque vulnerability and the overall prognosis of cardiovascular disease. In this regard, it is important to differentiate between the presence of microcalcifications and macrocalcifications when evaluating the effect on plaque stability. The formation of 'spotty' or microcalcifications is the result of chronic plaque inflammation characterized by the presence of pro-inflammatory cytokines such as IL-6 and TNF- $\alpha$  [37]. Under these



**Fig. 2.** Cardiac parameters of control and molsidomine-treated mice. (A–C) *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> mice treated with molsidomine (n = 9) showed a significantly lower end systolic diameter (ESD) and end diastolic diameter (EDD) as compared to controls (n = 9). Moreover, fractional shortening (FS) was significantly higher in the molsidomine treatment group. Independent samples *t*-test; \*\**P* < 0.01 and \*\*\**P* < 0.001. (D) Molsidomine treatment normalized heart weight of *ApoE*<sup>-/-</sup>*Fbn1*<sup>C1039G+/-</sup> mice (n = 15 in both groups). Independent samples *t*-test; \**P* < 0.05. (E) Histological images of Masson's trichrome stained hearts showing a myocardial infarction (blue staining, arrow) in the right ventricle of the control-treated mouse. Scale bar = 500  $\mu$ m. (F) Masson's trichrome staining of coronary arteries indicated that perivascular fibrosis, measured as perivascular collagen area (PVCA) divided by the lumen area (LA), was significantly reduced in molsidomine-treated mice as compared to controls. 2-way ANOVA; \*\**P* < 0.01. Scale bar = 100  $\mu$ m. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)



**Fig. 3.** Track width of control and molsidomine-treated mice. (A–B) Both the median and maximum track width values were significantly lower in *ApoE*<sup>-/-</sup> *Fbn1*<sup>C1039G +/-</sup> mice treated with molsidomine (n = 10) as compared to controls (n = 7). Independent samples *t*-test; \*\**P* < 0.01. (C) The median track width values correlated with plaque size in the proximal ascending aorta. Pearson correlation; R square = 0.347, *P* = 0.044.

circumstances, remnants of apoptotic SMCs, which remain in the plaque as matrix vesicles, will act as nucleating structures for calcification [38,39]. These microcalcifications are associated with enhanced plaque progression [32] and have the potential to increase the occurrence of plaque rupture via the induction of biomechanical stress [32,40,41]. Considering the latter, the location in the plaque is also of great importance, because only microcalcifications in the fibrous cap negatively affect the mechanical stability, while calcifications found in the necrotic core or at the medial border, as seen in the present study, do not contribute to plaque instability [32,41]. In contrast to microcalcifications, macrocalcifications are considered plaque stabilizing, because they are the result of a healing process that is able to separate the inflammatory plaque content from the vessel lumen [42]. These large calcified areas are formed when the initial inflammation in the plaque is resolved, leading to a tissue repair response and osteogenesis. Therefore, large plaque calcifications have been associated with decreased fibrous cap inflammation [43]. As bone formation is inhibited by chronic inflammation, microcalcifications will not evolve into macrocalcifications, unless the inflammatory status of the plaque decreases [37]. NO has anti-inflammatory properties, which can explain the increase in plaque macrocalcifications [27]. Molsidomine treatment was introduced at a timepoint where advanced atherosclerotic plaques are present in *ApoE*<sup>-/-</sup> *Fbn1*<sup>C1039G +/-</sup> mice, which is the ideal environment for the development of microcalcifications, but the anti-inflammatory effects of the therapy allowed osteogenesis to occur followed by the formation of macrocalcifications. Moreover, we could detect a clear increase in collagen type I in the calcified plaques, which is also a typical hallmark of a healing response after inflammation and further stabilizes the atherosclerotic plaque. All these data indicate that molsidomine treatment is able to favour plaque stability in advanced atherosclerosis.

Interestingly, we also observed a lower spleen and kidney weight, indicative of decreased inflammation. Other research groups have demonstrated that molsidomine was able to reduce kidney weight [12], which was attributed to the anti-inflammatory and local dilating effects of NO. [44] In addition, molsidomine modulates inflammation in various conditions such as muscular atrophy [45,46], wound healing [47] and bacterial inflammation [48].

In conclusion, we demonstrated that administration of the NO donor molsidomine improves heart function, reduces neurological signs and enhances atherosclerotic plaque stability in a mouse model of vulnerable plaque and spontaneous myocardial infarction. The effect of molsidomine treatment on the phenotype of rupture-prone plaques in mice

is new and might offer opportunities to further investigate the use of NO donors in atherosclerotic plaque stabilization and the prevention of cardiac and cerebral ischemia.

#### Sources of funding

This work was financially supported by the Agency for Innovation by Science and Technology (IWT), the Fund for Scientific Research (FWO)-Flanders, the University of Antwerp (BOF), agreement between Institute Born-Bunge and University of Antwerp, the Medical Research Foundation Antwerp, and Neurosearch Antwerp. Besa Emini Veseli is a PhD fellow of the Horizon 2020 program of the European Union - Marie Skłodowska-Curie Actions, Innovative Training Networks (ITN), Call: H2020-MSCA-ITN-2015, NUMBER — 675,527 — MOGLYNET.

#### Declaration of interest

None.

#### Acknowledgements

The authors would like to thank Bronwen Martin for critically reviewing the manuscript and Rita Van den Bossche, Hermine Fret, Anne-Elise Van Hoydonck, Sanne Laurysen, Michiel Knaapen and Anne-Sophie Hervent for their valuable assistance.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.vph.2019.05.001>.

#### References

- [1] J. Loscalzo, G. Welch, Nitric oxide and its role in the cardiovascular system, *Prog. Cardiovasc. Dis.* 38 (1995) 87–104.
- [2] V.W. Liu, P.L. Huang, Cardiovascular roles of nitric oxide: a review of insights from nitric oxide synthase gene disrupted mice, *Cardiovasc. Res.* 77 (2008) 19–29.
- [3] C. Napoli, F. de Nigris, S. Williams-Ignarro, O. Pignalosa, V. Sica, L.J. Ignarro, Nitric oxide and atherosclerosis: an update, *Nitric Oxide* 15 (2006) 265–279.
- [4] M. Kelm, M. Feelisch, R. Spahr, H.M. Piper, E. Noack, J. Schrader, Quantitative and kinetic characterization of nitric oxide and EDRF released from cultured endothelial cells, *Biochem. Biophys. Res. Commun.* 154 (1988) 236–244.
- [5] P.B. Massion, O. Feron, C. Dessy, J.L. Balligand, Nitric oxide and cardiac function: ten years after, and continuing, *Circ. Res.* 93 (2003) 388–398.
- [6] L.J. Ignarro, C. Napoli, J. Loscalzo, Nitric oxide donors and cardiovascular agents modulating the bioactivity of nitric oxide: an overview, *Circ. Res.* 90 (2002) 21–28.

- [7] S. Kawashima, M. Yokoyama, Dysfunction of endothelial nitric oxide synthase and atherosclerosis, *Arterioscler. Thromb. Vasc. Biol.* 24 (2004) 998–1005.
- [8] G.M. Berkenboom, J.C. Sobolski, P.P. Vandermoten, E.E. Stoupe, S.G. Degre, Effects of molsidomine on left ventricular dimensions and cardiac function in patients with chronic heart failure, *Am. Heart J.* 109 (1985) 691–693.
- [9] R.E. Nitz, P.A. Martorana, The activity of molsidomine in experimental models of ischemic cardiac disease, *Am. Heart J.* 109 (1985) 631–636.
- [10] E. Bassenge, U. Pohl, Effect of molsidomine on cardiac preload, coronary artery diameter, and coronary resistance, *Am. Heart J.* 109 (1985) 627–630.
- [11] V.B. Fielder, Effects of molsidomine on coronary artery thrombosis and myocardial ischemia in acute canine experiments, *Eur. J. Pharmacol.* 73 (1981) 85–89.
- [12] L.G. Bongartz, B. Braam, M.C. Verhaar, M.J. Cramer, R. Goldschmeding, C.A. Gaillard, P. Steendijk, P.A. Doevendans, J.A. Joles, The nitric oxide donor molsidomine rescues cardiac function in rats with chronic kidney disease and cardiac dysfunction, *Am. J. Physiol. Heart Circ. Physiol.* 299 (2010) H2037–H2045.
- [13] H. Bult, H.R.L. Fret, A.G. Herman, Interaction between sin-1 and prostacyclin in inhibiting platelet-aggregation, *J. Cardiovasc. Pharmacol.* 14 (1989) S120–S123.
- [14] H. Bult, H.R. Fret, R.M. Van den Bossche, A.G. Herman, Platelet inhibition by endothelium-derived relaxing factor from the rabbit perfused aorta, *Br. J. Pharmacol.* 95 (1988) 1308–1314.
- [15] G.R. De Meyer, M.M. Kockx, M.W. Knaapen, W. Martinet, D.M. De Cleen, H. Bult, A.G. Herman, Nitric oxide donor molsidomine favors features of atherosclerotic plaque stability during cholesterol lowering in rabbits, *J. Cardiovasc. Pharmacol.* 41 (2003) 970–978.
- [16] C. Van Hove, F. Carreer-Bruhwyler, J. Geczy, A.G. Herman, Long-term treatment with the NO-donor molsidomine reduces circulating ICAM-1 levels in patients with stable angina, *Atherosclerosis* 180 (2005) 399–405.
- [17] H. Bult, G.R. De Meyer, A.G. Herman, Influence of chronic treatment with a nitric oxide donor on fatty streak development and reactivity of the rabbit aorta, *Br. J. Pharmacol.* 114 (1995) 1371–1382.
- [18] C. Van der Donck, J.L. Van Herck, D.M. Schrijvers, G. Vanhoutte, M. Verhoye, I. Blockx, A. Van Der Linden, D. Bauters, H.R. Lijnen, J.C. Sluimer, L. Roth, C.E. Van Hove, P. Franssen, M.W. Knaapen, A.S. Hervent, G.W. De Keulenaer, H. Bult, W. Martinet, A.G. Herman, G.R. De Meyer, Elastin fragmentation in atherosclerotic mice leads to intraplaque neovascularization, plaque rupture, myocardial infarction, stroke, and sudden death, *Eur. Heart J.* 36 (2015) 1049–1058.
- [19] J.L. Van Herck, G.R. De Meyer, W. Martinet, C.E. Van Hove, K. Foubert, M.H. Theunis, S. Apers, H. Bult, C.J. Vrints, A.G. Herman, Impaired fibrillin-1 function promotes features of plaque instability in apolipoprotein E-deficient mice, *Circulation* 120 (2009) 2478–2487.
- [20] L. Roth, D. Van Dam, C. Van der Donck, D.M. Schrijvers, K. Lemmens, I. Van Brussel, P.P. De Deyn, W. Martinet, G.R.Y. De Meyer, Impaired gait pattern as a sensitive tool to assess hypoxic brain damage in a novel mouse model of atherosclerotic plaque rupture, *Physiol. Behav.* 139 (2015) 397–402.
- [21] T. Munzel, S. Steven, A. Daiber, Organic nitrates: update on mechanisms underlying vasodilation, tolerance and endothelial dysfunction, *Vasc. Pharmacol.* 63 (2014) 105–113.
- [22] P.S. Garry, M. Ezra, M.J. Rowland, J. Westbrook, K.T. Pattinson, The role of the nitric oxide pathway in brain injury and its treatment—from bench to bedside, *Exp. Neurol.* 263 (2015) 235–243.
- [23] H. Kakizawa, F. Matsui, Y. Tokita, K. Hirano, M. Ida, K. Nakanishi, M. Watanabe, Y. Sato, A. Okumura, S. Kojima, A. Oohira, Neuroprotective effect of nipradilol, an NO donor, on hypoxic-ischemic brain injury of neonatal rats, *Early Hum. Dev.* 83 (2007) 535–540.
- [24] M.S. Wainwright, D. Grundhoefer, S. Sharma, S.M. Black, A nitric oxide donor reduces brain injury and enhances recovery of cerebral blood flow after hypoxia-ischemia in the newborn rat, *Neurosci. Lett.* 415 (2007) 124–129.
- [25] M. Willmot, L. Gray, C. Gibson, S. Murphy, P.M. Bath, A systematic review of nitric oxide donors and L-arginine in experimental stroke; effects on infarct size and cerebral blood flow, *Nitric Oxide* 12 (2005) 141–149.
- [26] A. Ehlert, C. Schmidt, J. Wolfer, G. Manthei, A.H. Jacobs, R. Bruning, W. Heindel, E.B. Ringelstein, W. Stummer, R.M. Pluta, V. Hesselmann, Molsidomine for the prevention of vasospasm-related delayed ischemic neurological deficits and delayed brain infarction and the improvement of clinical outcome after subarachnoid hemorrhage: a single-center clinical observational study, *J. Neurosurg.* 124 (2016) 51–58.
- [27] L. Phillips, A.H. Toledo, F. Lopez-Neblina, R. Anaya-Prado, L.H. Toledo-Pereyra, Nitric oxide mechanism of protection in ischemia and reperfusion injury, *J. Invest. Surg.* 22 (2009) 46–55.
- [28] F. Alfonso, M. Joner, Untangling the diagnosis and clinical implications of calcified coronary nodules, *JACC Cardiovasc. Imaging* 10 (2017) 892–896.
- [29] N. Kobayashi, M. Takano, M. Tsurumi, Y. Shibata, S. Nishigoori, S. Uchiyama, H. Okazaki, A. Shirakabe, Y. Seino, N. Hata, W. Shimizu, Features and outcomes of patients with calcified nodules at culprit lesions of acute coronary syndrome: an optical coherence tomography study, *Cardiology* 139 (2018) 90–100.
- [30] P. Raggi, T.Q. Callister, B. Cool, Z.X. He, N.J. Lippolis, D.J. Russo, A. Zelinger, J.J. Mahmarian, Identification of patients at increased risk of first unheralded acute myocardial infarction by electron-beam computed tomography, *Circulation* 101 (2000) 850–855.
- [31] A.R. Folsom, R.A. Kronmal, R.C. Detrano, D.H. O'Leary, D.E. Bild, D.A. Bluemke, M.J. Budoff, K. Liu, S. Shea, M. Szklo, R.P. Tracy, K.E. Watson, G.L. Burke, Coronary artery calcification compared with carotid intima-media thickness in the prediction of cardiovascular disease incidence: the multi-ethnic study of atherosclerosis (MESA), *Arch. Intern. Med.* 168 (2008) 1333–1339.
- [32] F. Otsuka, K. Sakakura, K. Yahagi, M. Joner, R. Virmani, Has our understanding of calcification in human coronary atherosclerosis progressed? *Arterioscler. Thromb. Vasc. Biol.* 34 (2014) 724–736.
- [33] M. Henein, G. Granasen, U. Wiklund, A. Schermund, A. Guerci, R. Erbel, P. Raggi, S.E. Nissen, Impact of statins on serial coronary calcification during atheroma progression and regression, *J. Am. Coll. Cardiol.* 65 (2015) 1273–1282.
- [34] R. Puri, S.J. Nicholls, M. Shao, Y. Kataoka, K. Uno, S.R. Kapadia, E.M. Tuzcu, S.E. Nissen, Impact of statins on serial coronary calcification during atheroma progression and regression, *J. Am. Coll. Cardiol.* 65 (2015) 1273–1282.
- [35] A. Saremi, G. Bahn, P.D. Reaven, V. Investigators, Progression of vascular calcification is increased with statin use in the veterans affairs diabetes trial (VADT), *Diabetes Care* 35 (2012) 2390–2392.
- [36] I.C. Thomas, N.I. Forbang, M.H. Criqui, The evolving view of coronary artery calcification and cardiovascular disease risk, *Clin. Cardiol.* 41 (2018) 144–150.
- [37] A. Shioi, Y. Ikari, Plaque calcification during atherosclerosis progression and regression, *J. Atheroscler. Thromb.* 25 (2018) 294–303.
- [38] M.M. Kockx, G.R. De Meyer, J. Muhring, W. Jacob, H. Bult, A.G. Herman, Apoptosis and related proteins in different stages of human atherosclerotic plaques, *Circulation* 97 (1998) 2307–2315.
- [39] D. Proudfoot, J.N. Skepper, L. Hegyi, M.R. Bennett, C.M. Shanahan, P.L. Weissberg, Apoptosis regulates human vascular calcification in vitro: evidence for initiation of vascular calcification by apoptotic bodies, *Circ. Res.* 87 (2000) 1055–1062.
- [40] P. Libby, I. Tabas, G. Fredman, E.A. Fisher, Inflammation and its resolution as determinants of acute coronary syndromes, *Circ. Res.* 114 (2014) 1867–1879.
- [41] N. Maldonado, A. Kelly-Arnold, Y. Vengrenyuk, D. Laudier, J.T. Fallon, R. Virmani, L. Cardoso, S. Weinbaum, A mechanistic analysis of the role of microcalcifications in atherosclerotic plaque stability: potential implications for plaque rupture, *Am. J. Physiol. Heart Circ. Physiol.* 303 (2012) H619–H628.
- [42] N.V. Joshi, A. Vesey, D.E. Newby, M.R. Dweck, Will 18F-sodium fluoride PET-CT imaging be the magic bullet for identifying vulnerable coronary atherosclerotic plaques? *Curr. Cardiol. Rep.* 16 (2014) 521.
- [43] W.E. Shaalan, H.W. Cheng, B. Gewertz, J.F. McKinsey, L.B. Schwartz, D. Katz, D.C. Cao, T. Desai, S. Glagov, H.S. Bassiouny, Degree of carotid plaque calcification in relation to symptomatic outcome and plaque inflammation, *J. Vasc. Surg.* 40 (2004) 262–269.
- [44] S.R. Vercauteren, D.K. Ysebaert, K.E. De Greef, E.J. Eyskens, M.E. De Broe, Chronic reduction in renal mass in the rat attenuates ischemia/reperfusion injury and does not impair tubular regeneration, *J. Am. Soc. Nephrol.* 10 (1999) 2551–2561.
- [45] P. Zordan, C. Sciorati, L. Campana, L. Cottone, E. Clementi, P.R. Querini, S. Brunelli, The nitric oxide-donor molsidomine modulates the innate inflammatory response in a mouse model of muscular dystrophy, *Eur. J. Pharmacol.* 715 (2013) 296–303.
- [46] E. Rigamonti, P. Zordan, C. Sciorati, P. Rovere-Querini, S. Brunelli, Macrophage plasticity in skeletal muscle repair, *Biomed. Res. Int.* 2014 (2014) 560629.
- [47] M.B. Witte, T. Kiyama, A. Barbul, Nitric oxide enhances experimental wound healing in diabetes, *Br. J. Surg.* 89 (2002) 1594–1601.
- [48] N.H. Kumins, J. Hunt, R.L. Gamelli, J.P. Filkins, Molsidomine increases endotoxin survival and decreases cytokine production, *Shock* 7 (1997) 200–205.