

University of Groningen

Collaborative partnership between family caregivers and nurses in the care of older hospitalized persons

Hagedoorn, Ellen Ingrid

DOI:
[10.33612/diss.97727618](https://doi.org/10.33612/diss.97727618)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hagedoorn, E. I. (2019). *Collaborative partnership between family caregivers and nurses in the care of older hospitalized persons*. Rijksuniversiteit Groningen. <https://doi.org/10.33612/diss.97727618>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

**Translation and Psychometric Evaluation of the Dutch Families
Importance in Nursing Care – Nurses’ Attitudes Scale based on
the Generalized Partial Credit Model**

E.I. Hagedoorn

W. Paans

T. Jaarsma

J.C. Keers

C.P. van der Schans

ML.A. Luttik

W.P. Krijnen

Abstract

The Families Importance in Nursing Care– Nursing Attitudes (FINC-NA) instrument is used to measure nurses' attitudes towards families. The aim of this study is to evaluate the FINC-NA scale in a population of Dutch nurses and add new psychometric information to existing knowledge. Using a cross sectional design, 1,211 nurses received an online application in 2015. Psychometric properties were based on Polychoric correlations and the Generalized Partial Credit Model. A total of 597 (49%) nurses responded. Results confirmed a four-subscale structure. All response categories were utilized, although some ceiling effects occurred. Most items increase monotonically, and the majority of items discriminate well between different latent trait scores of nurses with some items providing more information than others. This study shows good psychometric properties of the Dutch FINC-NA instrument. New insights into the construct and content of items enable generating a more generic instrument that could be valid across several cultures.

Introduction

Family members have considerable influence on the health and self-care of elderly individuals with chronic diseases.¹⁻³ Therefore, it is important for nurses to involve family members in nursing care to promote continuity of care for the patient^{4,5} and collaborate with them as partners in care.^{6,7} When nurses have a more positive attitude toward involving families in their nursing care, they are more likely to do so.^{8,9} A widely used instrument to measure nurses' attitudes towards involvement of family members is the Families Importance in Nursing Care – Nursing Attitudes.^{10,11}

The FINC-NA instrument was first developed in Sweden and has been employed to assess nurses' attitudes toward families in nursing care^{9,12,13} and to measure the effect of educational types of program interventions on the attitudes of nurses.^{14,15} Since its initial use in 2008 in Sweden,¹⁰ the FINC-NA has also been widely used in other countries (See Table 1), for example Iceland,¹³ Portugal,¹⁶⁻¹⁸ Spain,¹⁹ Taiwan,⁸ Denmark, Norway, Belgium,¹² and Australia.²⁰

To use the FINC-NA instrument in intervention research to measure changes in nurses' attitudes in the Netherlands, the instrument needed to be translated and psychometrically evaluated in Dutch. This study was part of a research program with the focus on family care interventions aiming to improve collaboration between family caregivers and nurses in hospitals, as well as in home health care. The use of a Dutch language FINC-NA was needed to be able to measure changes in nurses' attitudes in follow-up pre- and post-intervention research, and to identify differences in groups.

Psychometric properties are important for assessing the validity and reliability of instruments in nursing research.²¹ Psychometric properties most commonly investigated in nursing research are internal consistencies by Cronbach's alpha to quantify the internal reliability of a scale or subscale and factor analysis to assess the dimensionality of a scale.²² As indicated in Table 1, many studies have reported various psychometric properties of the FINC-NA instrument. The most commonly reported is the internal consistency using the Cronbach's alpha of the FINC-NA scale and subscales (11 out of 15 studies). Five studies reported scale dimensionality of the FINC-NA based on confirmative as well as explorative methods of factor analysis. Two studies reported a score distribution of the items by the standardized response mean which is important for assessing the concept of the

responsiveness of the items. Finally, three studies reported item-total correlations to evaluate the discrimination ability of the items.

The psychometric properties of studies exploring the FINC-NA in Table 1 are all based on classical test theories.^{21,22} Until recently, the Rasch type of models²³ were not used to investigate scale properties. Such models represent constructs as latent traits allowing for a more detailed investigation of item and scale performance compared to the classical test theory and are beneficial for reviewing the psychometric properties of existing ordinal scales.²⁴ Classical Test Theories do not evaluate instruments' psychometric properties of monotonicity, discrimination capacities of items, or item information curves per subscale.²⁵ The Generalized Partial Credit Model (GPCM), an extension of the Rasch model,²⁵ was therefore used to evaluate these missing psychometric properties of the FINC-NA instrument.²⁵ This method is useful for reviewing the psychometric properties of ordinal scales²⁴ and allows for a more rigorous examination of measurement instruments in nursing.²⁶ The GPCM assumes that each nurse has a fixed value on a latent trait continuum. Latent trait values determine the probability of specific nurse responses on categories of items belonging to a subscale.²⁵ Monotonicity refers to the models' assumption that when there is an increase in nurses' positive attitudes towards families, there should also be an increase in the proportion of nurses with a high latent trait score. Items that violate this assumption might not optimally measure the construct to be evaluated. A discrimination parameter is important for examining the degree to which each item discriminates between nurses having different values on the trait.²⁵ Item information curves demonstrate how much information each item contributes to values of the latent trait for each scale. By using the GPCM, more information can be generated on the variability of the scores.²⁷ An application of the model thus facilitates detailed information on the variability of scores and their importance with respect to the latent attitude trait.²⁶

Based on these results, we ascertained a need to further explore the psychometric properties of the FINC-NA. The aim of this study was to determine psychometric properties of the FINC-NA regarding dimensionality, score distribution, internal consistency, monotonicity, discrimination parameters, and item information curves of each subscale using the GPCM.

Table 1. FINC-NA used in studies and reported psychometric properties

Author	Country	Nurses sample	N	Likert scale	Number of Subscales	Cronbach's alphas	Factor loadings	Scale dimensions	Distribution of scores	Item total correlations
Benzein, Johansson, Årestedt, Berg & Saveman (2008)	Sweden	Registered nurses	634	4 point	4	Yes	Yes	Yes	Yes	Yes
Skúladóttir, Konráðsdóttir & Ágústsdóttir (2010)	Iceland	Unknown	140	4 point	4	Yes	No	No	No	No
Saveman, Benzein, Engström & Årestedt (2011)	Sweden	Nurse specialist students (e.g. midwives)	246	5 point	4	Yes	Yes	Yes	Yes	Yes
Sveinbjarnardóttir, Svavarsdóttir & Saveman (2011)	Iceland	Psychiatric nurses	81	4 point	4	No	No	No	No	No
Oliveira et al. 2011	Portugal	Primary healthcare nurses	136	4 point	3	Yes	Yes	No	No	No
Silvia et al. 2015	Portugal	Primary healthcare nurses	871	4 point	3	Yes	No	No	No	No

Table 1. Continued

Author	Country	Nurses sample	N	Likert scale	Number of Subscales	Cronbach's alphas	Factor loadings	Scale dimensions	Distribution of scores	Item total correlations
Blöndal et al. 2014	Iceland	Surgical nurses	181	4 point	4	Yes	No	No	No	No
Angelo et al. 2014	Brazil	Paediatric nurses	50	4 point	3	No	No	No	No	No
Linnarsson, Benzein & Årestedt (2015)	Sweden	Emergency depart. nurses	457	5 point	4	Yes	No	No	No	No
Hsiao & Tsai (2015)	Taiwan	Psychiatric nurses	175	5 point	4	Yes	No	No	No	No
Fernandes, Pereira Gomes, Martins, Pereira Gomes & Hisako (2015)	Portugal	General hospital nurses	160	4 point	3	Yes	No	No	No	No
Pascual Fernández al. 2015	Spain	Paediatric nurses	186	4 point	4	Yes	Yes	Yes	No	No
Luttik et al. 2017	Belgium Sweden Denmark Norway	Cardiovascular nurses	425	5 point	4	No	No	No	No	No

Table 1. Continued

Author	Country	Nurses sample	N	Likert scale	Number of Subscales	Cronbach's alphas	Factor loadings	Scale dimensions	Distribution of scores	Item total correlations
Gusdal, Josefsson, Thors Adolfsson & Martin (2017)	Sweden	Heart failure nurses	303	5 point	4	Yes	No	No	No	No
Mackie, Marshall, Mitchell & Ireland (2018)	Australia	General hospital nurses	212	5 point	4	Yes	Yes	Yes	No	Yes

Method

A cross-sectional design was used in this study and data were collected with an online survey. Four general hospitals and one home healthcare organization in the northern part of the Netherlands participated in the study.

Sample and Setting

The sample consisted of 1,211 nurses; 403 hospital- and 808 home healthcare nurses. Participating hospital wards were cardiology (3), internal medicine (3), pulmonology (4), neurology (4), and one geriatric nursing ward. The home healthcare organization self-selected a region in the north of the Netherlands to participate in the study. Nurses who participated in the study had earned an associate degree level 3 (a 3-year course at a community college, without technical nursing interventions, like infusions), an associate degree level 4 (a 4-year course at a community college, including technical nursing interventions); and a bachelor degree or master degree in nursing.²⁸

Instrument

The FINC-NA consists of 26 items divided into four subscales: *Family as its own resource* (Fam-OR) referring to families' own resources for coping (four items); *Family as a burden* (Fam-B) referring to statements of experiencing family as a burden (four items); *Family as a conversational partner* (FAM-CP) referring to the acknowledgement of the patients' family members as conversational partners (eight items), and *Family as a resource in nursing care* (Fam-RNC) referring to a positive attitude towards families' presence in nursing care (ten items). In this study, the revised version of the FINC-NA five-point Likert scale was used with the response categories 'strongly disagree' to 'strongly agree.'¹¹ The item scores range from 1 to 5 with total scale scores ranging from 26 – 130 for the total instrument. The higher the score, the more supportive nurses' attitudes are towards families, with the exception of the burden scale. The four 'negatively' formulated items of the subscale family as a burden were subsequently reversed in order to facilitate data analysis.

Translation of the instrument (linguistic validation).

Written permission was granted by the original authors for the use and translation of the instrument. The original instrument was translated from Swedish into Dutch using the bidirectional translation method for linguistic-cultural adaptation.²⁹ Two independent professional translators provided the forward and back translation of the original instrument

from Swedish into Dutch and from Dutch into Swedish. Reconciliation of the first translation into Dutch was discussed by the first author and two co-authors. One of the co-authors is a native Dutch speaker with excellent knowledge of the Swedish language. Some minor, mostly constructive, changes to some of the sentences were suggested by her during the back translation. All differences were discussed by the professional translators and the three authors of the article to reach consensus. Throughout the instrument, the term 'sjuke' (sick person) was changed into 'patient'.

Data Collection

The FINC-NA instrument was transferred into a web application that included demographic characteristics: age; gender; years of work experience; highest level of education in nursing; continuing education in family care; knowledge of policy regarding family involvement being present in the organization; and prior experience as an informal caregiver who cared for a sick family member. A message containing a personalized link to the survey was sent to nurses' work e-mail addresses. After 2 or 4 weeks, a reminder was automatically sent to all e-mail addresses in the event that no response was yet received. The web application had the ability to automatically send reminders after a set period of time to all e-mail addresses that had not responded yet. This process was blinded for all persons, including the researcher. Data were collected in 2015.

Ethical Considerations

The Medical Ethics Committee of a Medical Center Leeuwarden ruled that this study was not under regulation of the Medical Research Involving Human Subjects Act (Reference nWMO106). Permission to perform the study was granted by the directors of the participating organizations. Prior to the beginning of the study, nurses were informed about the purpose by their managers and through a newsletter. Nurses voluntarily participated in the study and gave their consent for participation and publication of the results before completing the online survey. Responses were received anonymously and, therefore, could not be traced back to individual persons. Results were processed anonymously.

Data Analyses

To investigate the dimensionality of the four-subscale structure, factor analysis was based upon Polychoric correlations as the items consist of ordered categories.³⁰ From each subscale, confidence intervals (CIs) of the eigenvalues from the Polychoric inter-item correlations were computed. The number of eigenvalues significantly >1 were an indication for the

dimensionality of subscales.³¹ For comparative purposes with other studies, both Cronbach's alpha^{32,33} and ordinal coefficient alpha³⁴ were measured for internal consistency, with values $>.7$ considered acceptable.²²

A one-dimensional (ordinal) factor analysis on Polychoric correlations per subscale³⁵ was used to investigate the size of loadings, that is, the degree of association of the items with the latent trait (factor). Items with factor loadings of ≥ 0.32 (36) were considered for interpretation in this study as follows: > 0.71 (*excellent*); > 0.63 (*very good*); > 0.55 (*good*); > 0.45 (*fair*); ≥ 0.32 (*poor*). In addition, percentages of responses per category (1= Strongly disagree – 5= Strongly agree) were estimated for all of the items to assess the response distribution over the categories.

The GPCM was used to determine the monotonicity, discrimination parameter, and information of items per subscale.²⁵ Monotonicity implies that increasing levels across the response categories for each item should be reflected in the data, implying that the threshold estimates located on the latent trait must appear in the same order as the manifest categories.³⁷ In our measurement of nurses' attitudes towards the importance of families, each item consisted of five ordered response categories separated by four category thresholds on the latent attitude trait estimated by the model. These category steps govern the probability of scoring 1 rather than 0, 2 rather than 1, 3 rather than 2, and 4 rather than 5 on each item. The category thresholds are values on the latent trait that may be conceived as step parameters, as each nurse steps through the response categories (1-5) of each item and stops at the position nearest to his or her trait level. In this way, latent trait values determine the probability of specific responses on categories of items belonging to a subscale. The positions of the step parameters on the latent trait indicate whether and how the categories of each item contribute to the monotonicity of the latent attitude trait.

The degree to which each item discriminates between nurses with different values on the trait is given by its slope, also referred to as the discrimination parameter.²⁵ The greater the discrimination parameter, the better an item performs in discriminating between nurses with different attitudes. The amount of information that each item contributes to values of the latent trait is expressed by the item information curves for each scale. The models are estimated by the marginal maximum likelihood³⁸ using the programming language R.³⁹

To explore the construct validity of the four subscales '*Family as a conversational partner*', '*Family as its own resource*', '*Family as a burden*' and '*Family as a resource in nursing care*', factor and latent trait scores and their correlations were estimated for each subscale.

Respondents who had more than 5 items (25%) of the 26 items missing, or those who had more than 25% missing on a subscale were omitted for further analysis. Missing values were replaced with SPSS by the method of series means.⁴⁰

Results

A total of 597 (49%) nurses responded to the online survey. The responses of 28 (4.7%) nurses were omitted because more than 5 items were missing of the total scale, resulting in a total number of 569 responses that were suitable for further analysis. Only four (0.7%) respondents had one missing item at random, concerning items B2, CP2, OR2 and RNC8. These missing items were replaced by the series mean.

Table 2 illustrates the sample characteristics of the nurses. Table 3 summarizes the psychometric properties of the items and subscales of the FINC-NA (Appendix 2.1). The subscales are shown with their items in chronological order as they appear in the questionnaire and are expressed in truncated sentences to save space. Items are listed by subscale letters and number, i.e., item 1 from ‘*Family as its own resource*’ is referred to by OR1.

Subscale: Family as its Own Resource.

The largest two eigenvalues and their CIs of the subscale ‘Family as its own resource’ (Fam-OR) were 2.37 (95% CI: [2.21-2.51]) and 0.64 (95% CI: [0.57-0.74]), respectively, indicating that this subscale is one dimensional. Table 3 shows the percentages per category responses; all responses are fairly well symmetrically distributed over the categories, although the percentages for Category 1 are somewhat smaller than for Category 5 and percentages of Category 4 are somewhat higher than for Category 2. The ordinal and Cronbach’s alpha coefficients of 0.77 and .73, respectively, are fairly large considering that there are four items in this subscale. Alpha decreases substantially if an item is deleted (e.g., item 1). Factor loadings for all of the items in this subscale are very good; between 0.64 and 0.70. The location parameters are low for category step 1/2, large for step 4/5, and all increase monotonically for each item with the latent Fam-OR trait values for nurses. The absolute values of category step 1/2 are somewhat larger than those for step 4/5, which is in accordance with the size of the percentages of nurse responses in Categories 1 and 5.

Table 2. Nurses' characteristics

Characteristic		M (SD)
Age		43.5 (12.8)
Years of work experience		19.3 (12.2)
		N (%)
Gender	Female	538 (95)
	Male	31 (5)
Policy present on families	Yes	253 (45)
	No	316 (55)
Training in family nursing	Yes	87 (15)
	No	482 (85)
Informal caregiving experience	Yes	372 (65)
	No	197 (35)
Highest level of education in nursing (n=560)	Associate Degree level 3*	143 (25)
	Associate Degree level 4**	252 (45)
	Bachelor Degree	150 (27)
	Master Degree	15 (3)

* an associate degree level 3 (a three-year course at a community college without technical nursing interventions, such as infusion and injections, but including medication administration)

**an associate degree level 4 (a four-year course at a community college, including technical nursing interventions)

Table 3 also shows that the discrimination parameters from the GPCM are relatively equal and large in size (1.04; 1.29). This is also reflected in the item information curves in Figure 1a, indicating that these are high for a broad range of latent trait values. Each of the items contains approximately an equal amount of information on the latent Fam-OR trait. This is in line with the information in Table 3 that depicts that the amount of negative and positive information is comparable between the items of this subscale, although with slightly more information on negative trait values.

The similarity and coverage of the Fam-OR information curves indicate that these items provide an equal amount of information on the latent trait. With discrimination

parameters of 1.04 and higher, all items discriminate well between the different trait levels; discriminate between nurses with low, middle, and large values on the latent Fam-OR trait. Category step locations of the items are homogeneous in the sense of being well spread over the range of the Fam-OR trait, showing that the items adequately measure the different trait levels.

Subscale: Family as a Burden.

The largest two eigenvalues and their CIs of the ‘family as a burden’ were 2.37 (95% CI: [2.22-2.53]) and 0.73 (95% CI: [0.65-0.84]), respectively, and indicate that the subscale is one dimensional. Table 3 shows that all items of this subscale are asymmetrically distributed with very small response percentages on Categories 1 (range 0.1; 0.2) and 2 (range 0.3; 0.8) and relative large response percentages on Categories 4 (range 0.34; 0.41) and 5 (range 0.30; 0.41) (Table 3). The ordinal and Cronbach’s alpha coefficients of .76 and .71, respectively, are relatively large considering that there are four items in this scale. The Cronbach’s alpha increases to .73 if item B4 (*I don’t have time to take care of families*) is deleted and decreases substantially if item B1 or B2 is deleted. Factor loadings are excellent for items B1 and B2 (0.83; 0.78), and good to fair for item B3 and B4 (0.61; 0.46), respectively.

As shown in Table 3, location parameters are low for category step 1/2 and close to zero for step 4/5 and monotonically increase for each item as the value on the latent Fam-B trait increases. The small step values of the location values for category 4/5 correspond with the larger percentages of responses on Category 5 (Table 3). This indicates that nurses with latent scores close to zero have a relatively high probability to respond to Category 5. The absolute values of step location 1/2 are larger than those for location 4/5, which accords with the smaller percentages found in Category 1 compared to 5.

As shown in Table 3, the discrimination parameters for this subscale are substantial for items B1 and B2 (1.83; 2.40), moderate for item B3 (0.94), and small for item B4 (0.52). The item information curve in Figure 1b shows that items B1 and B2 contribute three to four times more information on the latent FamB trait than items B3 and B4. This follows also from the amount of information being higher on the negative part than that on the positive, especially for items B2 (7.90-1.72) and B1 (5.52-1.81), respectively (see Table 3).

Subscale: Family as a Conversational Partner.

The largest two eigenvalues of the subscale ‘family as conversational partner’ (Fam-CP) were 3.10 (95% CI: [2.83-3.40]) and 1.03 (95% CI: [0.97-1.20]), respectively, indicating that this

scale is one dimensional. Table 3 shows that five out of eight items (CP1, CP2, CP3, CP5, CP7) have the largest percentages of responses on Category 3 with small percentages on the extremes and symmetrically distributed responses over these categories. Items CP4 and CP6 are asymmetrically distributed having 3% and 5% responses on Category 1 or 2 and 80% and 83% of their responses on Category 4 or 5, respectively, making the responses on these items skewed to the right. Coefficient alphas of .77 and .74 are acceptable but not very large considering that there are eight items in this subscale. Items 1, 2, 3, and 7 of the Fam-CP subscales show good to very good factor loadings (0.62; 0.63; 0.70; 0.56) contrary to that of item 6 (0.42).

The item category responses in Table 3 are fairly symmetrically distributed and have step location parameters which increase monotonically with the size of the latent Fam-CP variable as was intended by the category ordering. Item CP4 and CP6 are exceptions in violating the category order (monotonicity). The discrimination parameters shown in Table 3 are moderate (0.76 - 1.08) for items CP1, CP2, CP3, and CP7 and somewhat small (0.49 - 0.60) for the remaining items.

The information curves in Figure 1c show that items CP1, CP2, CP3, and CP7 contain an equal amount of information on the latent Fam-CP trait and are fairly symmetrically distributed. Items CP4 and CP6, on the other hand, have almost no information on positive latent FamCP trait scores which is also illustrated by the amount of information of these items on positive latent scores in Table 3, CP6 (1.63; -0.32.), and CP4 (1.92; -0.47).

Table 3. Psychometric properties of the Family Importance in Nursing Care Scale

Subscales	Percentages per category of response					α if deleted	Factor Loadings	Generalized Partial Credit Model coefficients				Discr. parameter	Item information	
	1	2	3	4	5		Estimate	Step 1	Step 2	Step 3	Step 4		negative	positive
Family as own resource (OR)														
1. Encourage fam to use own resources	3	14	37	34	12	.65	.70	-2.46	-1.42	.09	1.61	1.29	3.13	2.01
2. I see myself as resource for fam to cope	2	9	31	42	17	.67	.67	-2.66	-1.80	-.40	1.38	1.17	3.16	1.54
3. I see fam as co-operating partners	3	14	38	34	11	.68	.64	-2.57	-1.52	.17	1.88	1.04	2.48	1.67
4. I ask fam how I can to support them	5	15	33	33	16	.67	.68	-2.14	-1.32	.01	1.37	1.07	2.64	1.65
Ordinal alpha total subscale						.77								
Cronbach's alpha total subscale						.73								
Family as a burden (B)														
1. Presence of fam makes me feel checked up	1	8	24	36	31	.61	.78	-2.94	-1.57	-.61	.52	1.83	5.52	1.81
2. Presence of fam. makes me feel stressed	1	3	21	34	41	.60	.83	-2.45	-2.09	-.82	.18	2.40	7.90	1.72
3. Presence of fam. holds me back in work	1	5	23	41	30	.66	.61	-3.26	-2.39	-.91	.70	.94	2.80	0.94
4. I don't have time to take care of fam.	2	7	26	34	31	.73	.46	-3.74	-3.17	-.63	.39	.52	1.49	0.56
Ordinal alpha total subscale						.76								
Cronbach's alpha total subscale						.71								
Family as conversational partner (CP)														
1. I invite fam for convers. at end of care period	11	21	35	23	10	.70	.62	-1.56	-.87	.76	1.88	.83	1.65	1.65
2. I ask fam take part in discussions at start care	12	23	31	19	15	.70	.63	-1.58	-.59	.84	.98	.77	1.53	1.53
3. I find out what fam members a patient	5	21	31	30	13	.69	.70	-2.35	-.79	.17	1.50	1.08	2.45	1.87

Subscales	Percentages per category of response					α if deleted	Factor Loadings Estimate	Generalized Partial Credit Model coefficients				Discr. parameter	Item information	
	1	2	3	4	5			Step 1	Step 2	Step 3	Step 4		negative	positive
has														
4. I invite fam speak about changes in condition	1	2	17	39	41	.72	.46	-2.67	-4.00	-1.70	.05	.60	1.92	.47
5. I invite fam to speak when planning care	11	30	38	15	6	.72	.48	-2.35	-.51	1.94	2.43	.57	.97	1.28
6. Important to find out who fam members are	1	4	12	33	50	.73	.42	-3.25	-3.29	-2.27	-.77	.49	1.63	.32
7. I invite fam to actively take part in care	9	26	40	20	5	.71	.56	-2.13	-.79	1.21	2.61	.76	1.43	1.62
8. Conversation at start of care saves time	5	13	34	39	9	.72	.46	-2.55	-2.16	-.14	3.04	.57	1.34	.90
Ordinal alpha total subscale						.77								
Cronbach's alpha total subscale						.74								
Family as resource in nursing care (RNC)														
1. Presence of fam eases my workload	3	12	54	27	4	.80	.64	-2.20	-1.75	.78	2.61	1.24	2.70	2.25
2. Presence of fam gives me feeling of security	13	26	48	10	3	.81	.49	-1.67	-1.03	2.75	2.73	.66	1.10	1.54
3. Presence of fam is important to me as nurse	1	7	27	41	24	.79	.73	-2.68	-1.86	-.56	.85	1.54	4.44	1.72
4. Fam should be invited take active part in care	1	8	38	32	21	.81	.60	-3.35	-2.29	.08	.99	.94	2.49	1.29
5. Fam should be invited in active planning care	4	19	40	27	10	.81	.58	-2.70	-1.26	.63	1.96	.81	1.77	1.46
6. Good relations fam give me job satisfaction	1	1	14	36	48	.81	.57	-2.08	-3.72	-1.59	-.25	.89	3.03	.54

Subscales	Percentages per category of response					α if deleted	Factor Loadings Estimate	Generalized Partial Credit Model coefficients				Discr. parameter	Item information	
	1	2	3	4	5			Step 1	Step 2	Step 3	Step 4		negative	positive
7. Getting involved gives me feelings useful	2	4	28	45	21	.81	.61	-1.71	-2.94	-.73	1.26	.90	2.56	1.04
8. Gain worthwhile knowledge from fam to use	2	7	26	45	20	.80	.68	-2.48	-1.82	-.64	1.16	1.28	3.61	1.51
9. Presence of fam important for fam self	1	4	28	45	22	.81	.49	-4.13	-3.34	-.85	1.34	.70	2.00	.80
10. It is important to spend time with fam	1	3	22	47	27	.80	.64	-2.31	-2.89	-1.00	.81	1.20	3.65	1.16
Ordinal alpha total subscale						.85								
Cronbach's alpha total subscale						.82								
Cronbach's alpha for the total FINC-NA						.88								

Discr.

Parameter:

discriminatory

parameter.

Subscale: Family as a Resource in Nursing Care.

Eigenvalues and their confidence intervals of the ‘family as a resource in nursing care’ (Fam-RNC) subscale with the largest two eigenvalues of 4.38 (95% CI: [4.03-4.73]) and 1.06 (95% CI: [0.98-1.25]), respectively, indicate that the scale is one dimensional. Table 3 shows that four of the 10 items of this scale are somewhat symmetric having the largest proportion of responses in Category 3, five items are less symmetrical with the largest percentages in Category 4. Item RNC6 (*A good relationship with family members gives me job satisfaction*) behaves very asymmetrically with the largest proportion in Category 5.

The alpha coefficients of .85 and .82 are sufficient and the Cronbach’s alpha slightly decreases if any of the items are deleted. Factor loadings are good to excellent with those of items RNC2 and RNC9 being smaller (0.49). The step location parameters of all items increase monotonically with the value of the latent FamRNC trait except for step 2/3 of RNC items 6, 7, and 10 and step 3/4 of item RNC2. The discrimination parameters are larger than 1.20 for RNC items 1, 3, 8, and 10 while the others are moderate between .66 and .94 (Table 3).

The item information curve in Figure 1d shows that item RNC3 contains twice the amount of information on the latent Fam-RNC trait compared to RNC items 2 and 9 with discrimination parameters of .66 and .70, respectively. The negative step locations of item RNC6 suggest that it mainly provides information on negative latent Fam-RNC scores for nurses. This is also seen in the amount of information item RNC6 depicts in Table 3, with 3.03 negative information and .52 positive information.




Figure 1. Item information curves of the FINC-NA subscales: (a) trait perceiving family as its own resource, (b) trait perceiving family as a burden, (c) trait perceiving family as a conversational partner and (d) trait perceiving family as a resource in nursing care. *Note.* FINC-NA = Families Importance in Nursing Care–Nurses’ Attitudes.

Exploration of subscale constructs

Table 4 indicates the correlations of the latent trait scores of nurses' attitudes towards the importance of families in nursing care from a polychoric factor correlations as well as the generalized partial credit analysis. The lower triangle beneath the diagonal on the left side of the table gives the polychoric correlations, and the upper triangle above the diagonal on the right side are those from the GPCM. In both models, the correlations of the latent factors are all positive and pointing in the same direction. The subscale Fam-B has a small positive correlation with each of the other three subscales. In addition, the inter-correlations between the subscales Fam-OR, Fam-RNC, and Fam-CP latent traits are substantial.

Table 4. Correlations between the subscales using Polychoric correlations and Generalized Partial Credit Model

		Generalized Partial Credit Model			
		FamB	FamOR	FamCP	FamRNC
Polychoric correlations	FamB	.993	.259	.175	.204
	FamOR	.243	.997	.551	.582
	FamCP	.176	.541	.998	.637
	FamRNC	.196	.570	.629	.991

Discussion

This study demonstrates that the Dutch FINC-NA has a unidimensional structure for each subscale. Results showed good to excellent factor loadings for the majority of items of each subscale.³⁶ Score distributions exhibit responses on all Likert categories. We found that item categories increase monotonically with increasing latent trait and that the majority of items contain a sufficient amount of information on the trait.

The unidimensional four-subscale structure of the FINC-NA that was found in this study is consistent with findings of previous studies.^{11,19,20} In contrast to these studies, we investigated the dimensionality per subscale. Finding a four-subscale structure using a

different method of analysis adds to the evidence for such a structure. These results are in contrast with findings of Oliveira¹⁸ and colleagues pertaining to a three-subscale instrument of the FINC-NA based on theoretical content, factor loadings, and internal consistency of each factor without deleting items. The findings of our study indicate that a four-subscale structure of the FINC-NA is the most interpretable to be employed in future studies.

Internal consistencies of the subscales in this study were moderate to strong²² and Cronbach's alphas were slightly lower compared to those found in other studies.^{9,11,41} The two smaller subscales, Fam-OR and Fam-B, show adequate reliability especially compared to those of the eight-item subscale of Fam-CP. This may imply that fewer items per subscale are able to give the same amount of information. The internal consistency of the total score of .88 shows a strong reliability of the total FINC-NA scale and supported its use in further studies.

All of the items in this study had factor loadings $\geq .42$. These findings are in contrast with findings of, for example Mackie²⁰ and colleagues, who found four items with factor loadings < 0.32 : OR2, CP3, CP6, and CP8, respectively. In our study, factor loadings for these items were higher; 0.67 (OR2); 0.70 (CP3); 0.42 (CP6), and 0.46 (CP8), respectively. These different findings may reflect cultural differences between countries.

Score distributions show responses on all Likert categories which supports the application of a five-point Likert scale. As was found by Saveman¹¹ and colleagues, a number of items show ceiling effects that do not adequately measure the right side of the latent trait. For example, item B2 (*The presence of family members makes me feel stressed*) has a score distribution of 1, 3, 21, 34, and 41, respectively, as shown in Table 3. Nurses that scored a five on these items may possibly have had an even higher latent trait value. In this case, it might be relevant to add a stronger formulated item in order to differentiate between nurses who have an even higher latent trait score on this concept which would result in expanded scale measurement properties.

Generalized Partial Credit coefficients of the majority of items indicate monotonicity as the model assumes, that is, most item categories indeed increase monotonically with increasing latent trait values. An example of an item that violates this assumption of monotonicity is RNC6 (*A good relationship with family members gives me job satisfaction*). As shown in Table 3, the GPCM coefficients of this item range from -2.08 to -0.25, measuring only latent trait values of nurses on the left side. This is also reflected by the item information scores of 3.03 on the negative side and 0.54 on the positive side (Table 3) and the item information curve of item RNC6 in Figure 1D. Thus, this item primarily provides information on the left extreme of the latent trait and does not appear to contribute to

discriminating between nurses with positive responses on this latent Fam-RNC variable.⁴² This suggests that omitting this item would result in only a minimal loss of information regarding Dutch nurses.

We were able to demonstrate that most items of the FINC-NA instrument discriminate well between the latent trait scores of nurses by their discrimination parameters, for example, all items of the subscale family as its own resource (FamOR) subscale have a discrimination parameter larger than 1 (Table 3). Item information curves of the subscale ‘Family as a conversational partner’ in Figure 1c illustrate that some items provide considerably more information on the nurses’ latent traits than others. An example is item CP6 (*It is important to find out what family members a patient has*) which gives half of the amount of information on the nurses’ latent trait compared to item CP3 (*I always find out what family members a patient has*). This might suggest that, with less items, a comparable amount of information could be generated on the concept to be measured.

To the best of our knowledge, this is the first study to apply a Generalized Partial Credit Model to the FINC-NA instrument. Classical Test Theory for reliability and confirmatory factor analysis are based on covariances of items and provide only indirect information about values of the latent trait. By employing this method, the information from the analysis better accords with the actual nurse responses.⁴³ Therefore, this method of analysis generated new information on choice options of the instrument, allowing for detailed insight into the construct and content of the instruments’ items. This specific information was generated on item level which allowed for more efficient adjustments to the FINC-NA instrument, if desirable. The results of this study, as well as previous studies,^{10,11} show strong evidence for the four-subscale structure of the FINC-NA. Use of the four-subscale structure, as well as the total scale score is therefore recommended for future studies.

Involving families in nursing care is an important development worldwide, and it is important that nurses’ attitudes can be measured with an instrument that is valid and reliable across countries. By using detailed scale analysis more insight is given into how items and subscales of the FINC-NA behave, resulting in more detailed insight into perception of nurse attitudes towards the importance of families in nursing care. These results can be relevant for other countries who are interested in scale validation, for example countries as shown in Table 1. To generate additional in-depth information on how the FINC-NA instrument behaves at item level in other countries, researchers are invited to validate the FINC-NA in their countries and to apply the Generalized Partial Credit Model as a useful method of analysis. A validation study could then be performed to investigate linguistic and cultural issues, by

comparing two or more lists in an international context in order to investigate the possibility of a more generic instrument instead of nation-based instruments.

Strengths and limitations of the study

A strength of this study was the large sample size of nurses and nursing aides who work with families of elderly patients in a hospital or in-home health care; a sample size of > 500 is considered very good.⁴² A satisfactory response rate of 49% was reached, but if a larger group had responded it might have affected the results. The convenience sample can be seen as a limitation of this study. The home health care organization self-selected one of their regions because this organization was in the middle of a reorganization in the other regions. This could feasibly have led to some preselection. Also, measuring attitudes of nurses entails subjective responses and might provoke socially desirable behaviour; possibly influencing the reliability of study results.

As this is the first time the GPCM was employed in psychometric testing of the FINC-NA, a limitation of this study is the impossibility of comparing the results with other studies. An additional limitation of this study was that no comparison with other instruments measuring nurses' attitudes towards the importance of family in nursing care was undertaken in order to measure construct validity. This could be a suggestion for further research. This study, however, did explore construct validity by estimating factor scores and their correlations. Future studies need to evaluate the test-retest properties of this instrument. Previous studies determined a positive relationship between female gender and a more positive attitude towards families in nursing care.⁹ As only 5% of the nurses' population in this study were male, we did not differentiate between gender by Differential Item Functioning.

Conclusion

Findings of this study show that the FINC-NA instrument has good psychometric properties related to reliability, unidimensionality, monotonicity, and information, and, therefore, is recommended for future family nursing research on behalf of the measurement of effects of interventions and to measure changes in nurses' attitudes. By using an advanced method of analysis, we were able to generate detailed information on the psychometric properties of the FINC-NA instrument in a Dutch population of hospital and home healthcare nurses. Future

research should focus on improving properties of the subscales and generalizability over countries.

Acknowledgement

The authors would like to express their sincere appreciation to the nurses who participated in this study and the Martini Hospital, Medical Center Leeuwarden, Nij Smellinghe Hospital, Wilhelmina Hospital Assen, and Home Care Organization Icare for their cooperation.

Supporting information

Appendix 2.1. FINC-NA instrument in Dutch and English

References

1. Chen Y.-C, Chang L.-C, Liu C.-Y, Ho Y.-F, Weng S.-C, Tsai T.-I. The roles of social support and health literacy in self- management among patients with chronic kidney disease. *J Nurs. Scholars*. 2018;50(3):265–275. <https://doi.org/10.1111/jnu.12377>
2. McCabe N, Dunbar SB, Butler J, Higgins M, Book W, Reilly C. Antecedents of self-care in adults with congenital heart defects. *Int. J Cardiol*. 2015;201:610-615. <http://dx.doi.org/10.1016/j.ijcard.2015.08.125>
3. Neumann, D., Lamprecht, J., Robinski, M., Mau, W., & Girndt, M. (2018). Social relationships and their impact on health-related outcomes in peritoneal versus haemodialysis patients: a prospective cohort study. *Nephrology Dialysis Transplantation*, 33(7), 1235–1244. doi:<https://doi.org/10.1093/ndt/gfx361>
4. Bridges J, Flatley M, Meyer J. Older people's and relatives' experiences in acute care settings: Systematic review and synthesis of qualitative studies. *Int J Nurs Stud*. 2010; 47(1):89-107.
5. Lowson E, Hanratty B, Holmes L, Addington-Hall J, Grande G, Payne S, Seymour J. From 'conductor' to 'second fiddle': Older adult care recipients' perspectives on transitions in family caring at hospital admission. *Int J Nurs Stud*. 2013;50:1197-1205. <https://doi.org/10.1016/j.ijnurstu.2012.02.005>
6. Lindhardt, T., Nyberg, P., & Rahm Hallberg, I. (2008b). Collaboration between relatives of elderly patients and nurses and its relation to satisfaction with the hospital care trajectory. *Scan. J Caring Sci*. 2008b;22(4):507–519. <https://doi.org/10.1111/j.1471-6712.2007.00558.x>.
7. Wassenaar A, Schouten J, Schoonhoven L. Factors promoting intensive care patients' perception of feeling safe: a systematic review. *Int J Nurs Stud*. 2014;51(2):261-273. <https://doi.org/10.1016/j.ijnurstu.2013.07.003>
8. Hsiao C, Tsai Y. Factors Associated With the Perception of Family Nursing Practice Among Mental Health Nurses in Taiwan. *J Fam Nurs*. 2015;21(4):508-528. <https://doi.org/10.1177/1074840715606543>
9. Linnarsson J, Benzein E, Årestedt K. Nurses' views of forensic care in emergency departments and their attitudes, and involvement of family members. *J Clin Nurs*. 2014;24:266-274. <https://doi.org/10.1111/jocn.12638>.
10. Benzein E, Johansson P, Årestedt KF, Berg A, Saveman BI. Families' importance in nursing care: nurses' attitudes--an instrument development. *J. Fam. Nurs*. 2008;14(1):97-117. <https://doi.org/10.1177/1074840707312716>
11. Saveman B, Benzein EG, Engström ÅH, Årestedt K. Refinement and psychometric reevaluation of the instrument: Families' Importance in Nursing Care--Nurses' Attitudes. *J Fam Nurs*. 2011;17(3):312-329. <https://doi.org/10.1177/1074840711415074>
12. Luttik M, Goossens E, Ågren S, Jaarsma T, Mårtensson J, Thompson DR, et al. Attitudes of nurses towards family involvement in the care for patients with cardiovascular diseases. *Eur J Cardiovasc Nurs*. 2017;16(4):299-308.
13. Skúladóttir A, Konráðsdóttir E, Ágústsdóttir A. Spurningalisti til að meta viðhorf fagfólks til fjölskylduhjúkrunar [Questionnaire to evaluate the attitude of professionals towards family nursing]. *Nurs. Magazine Analys*. 2010;86: 6-10. <https://doi.org/10.1111/j.1365-2850.2011.01744.x>
14. Blöndal K, Zoëga S, Hafsteinsdóttir JE, Olafsdóttir OA, Thorvardardóttir AB, Hafsteinsdóttir SA, et al. Attitudes of Registered and Licensed Practical Nurses About the Importance of Families in Surgical

- Hospital Units: Findings from the Landspítali University Hospital Family Nursing Implementation Project. *J Fam Nurs* 2014;20(3):355-375.
15. Sveinbjarnardóttir EK, Svavarsdóttir EK, Saveman B. Nurses attitudes towards the importance of families in psychiatric care following an educational and training intervention program. *J Psychiatr Ment Health Nurs*. 2011;18(10):895-903.
 16. Fernandes Cs, Pereira Gomes JA, Martins MM, Pereira Gomes B, Hisako TG. The Importance of Families in Nursing Care: Nurses' Attitudes in the Hospital Environment. *J. Nurs. Referência*. 2015;7:21-30.
 17. Nunes Carramanho Gomes Martins Moreira, da Silva, da Silva Mendes dC, Ferreira Pereira dS. The Family in Primary Care: characterization of nurses' attitudes. *REFERENCIA*. 2013;12(11):19-28.
 18. Oliveira PdCM, Fernandes HIV, Vilar AISP, Figueiredo MHdJS, Ferreira MMSRS, Martinho MJCM, et al. [Attitudes of nurses towards families: validation of the scale Families' Importance in Nursing Care--Nurses Attitudes]. *Rev Esc Enferm USP*. 2011;45(6):1331-1337.
 19. Pascual Fernández MC, Ignacio Cerro MC, Cervantes Estévez L, Jiménez Carrascosa MA, Medina Torres M, García Pozo A.M. [Questionnaire to evaluate the importance of the family in nursing care. Validation of the Spanish version (FINC-NA)]. *An Sist Sanit Navar*. 2015;38(1):31-39.
 20. Mackie BR, Marshall A, Mitchell, M. & Ireland, M. J. Psychometric testing of the revised "Families' Importance in Nursing Care - Nurses' Attitudes Instrument". *J. adv. Nurs*. 2017;74:482-490.
<https://doi.org/10.1111/jan.13442>
 21. Polit DF, Tatano Beck C. *Nursing Research: Appraising Evidence for Nursing Practice*. 7th ed. Philadelphia, PA: Wolters Kluwer Lippincott Williams & Wilkins; 2010.
 22. Nunnally JC, Bernstein IH. *Psychometric theory*. 3rd ed. New York, NY: McGraw-Hill; 1994.
 23. Rasch G. *Probabilistic models for some intelligence and attainment tests*. Copenhagen: Danmarks Paedagogiske Instituut; 1960.
 24. Tennant A, Conaghan PG. The Rasch measurement model in rheumatology: what is it and why use it? When should it be applied, and what should one look for in a Rasch paper? *Arthritis Rheum*. 2007;57(8):1358-1362.
 25. Masters G. A Rasch model for partial credit scoring. *Psychometrika*. 1982;47(2):149-174.
<https://doi.org/10.1007/BF02296272>
 26. Hagquist C, Bruce M, Gustavsson JP. Using the Rasch model in nursing research: an introduction and illustrative example. *Int J Nurs Stud*. 2009;46(3):380-393.
 27. Li Y, Baser R. Using R and WinBUGS to fit a generalized partial credit model for developing and evaluating patient-reported outcomes assessments. *Stat Med*. 2012;31(18):2010-2026.
 28. Mistianen P, Kroezen M, Triemstra M, Francke AL. Verpleegkundigen en verzorgenden in internationaal perspectief. Een literatuurstudie naar rollen en posities van beroepsbeoefenaren in de verpleging en verzorging [A nurse is not a nurse. Nurses and carers in an international perspective: A literature study into the roles and positions of professionals in nursing and care]. Utrecht, The Netherlands: Nivel. <https://www.nivel.nl/sites/default/files/bestanden/rapport-verpleging-verzorging-internationaal-perspectief.pdf>. Published 2018.

29. Wild D, Grove A, Martin M, Eremenco S, McElroy S, Verjee-Lorenz A, et al. Principles of Good Practice for the Translation and Cultural Adaptation Process for Patient-Reported Outcomes (PRO) Measures: report of the ISPOR Task Force for Translation and Cultural Adaptation. *Value Health*. 2005;8(2):94-104. <https://doi.org/10.1111/j.1524-4733.2005.04054.x>
30. Drasgow F. *Polychoric and Polyserial Correlations*. Encyclopedia of Statistical Sciences. 9th ed: Hoboken, NJ: John Wiley; 2004;68-74.
31. Kaiser HF. The application of electronic computers to factor analysis. *Educ. Psychol Measur*. 1960;20:141-151. <https://doi.org/10.1177/001316446002000116>
32. Cronbach LJ. Coefficient alpha and the internal structures of tests. *Psychom*. 1951;16:297-334. <https://doi.org/10.1007/BF02310555>
33. Revelle, W & Zinbarg, RE. Ordinal Versions of Coefficients Alpha and Theta for Likert Rating Scales. *Psychometrika*. 2009;74:145-154. <https://doi.org/10.1007/s11336-008-9102-z>
34. Zumbo BD, Gadermann AM, Zeisser C. Ordinal Versions of Coefficients Alpha and Theta for Likert Rating Scales. *J Mod. Appl. Stat. Methods*. 2007;6:21-29. <https://doi.org/10.22237/jmasm/1177992180>
35. Basto M, Pereira JM. An SPSS R-Menu for ordinal factor analysis. *J. Statist. Softw*. 2012;46:1-29. <https://doi.org/10.18637/jss.v046.i04>
36. Tabachnick BG, Fidell LS. *Principal components and factor analysis. In using multivariate statistics*. 4th ed. Needham Height, MA: Allyn & Bacon; 2007.
37. Andrich D. A binomial latent trait model for the study of Likert-Style Attitude Questionnaires. *British J Math Stat Psychol*. 1978;31: 84-89. <https://doi.org/10.1111/j.2044-8317.1978.tb00575.x>
38. Rizopoulos, D. ltm: An R Package for Latent Variable Modeling and Item Response Theory Analyses. *J Stat Softw*. [Online] 20 11 2006. [Cited: 30 12 2016.] <https://www.jstatsoft.org/article/view/v017i05>.
39. R Core Team. R: A language and environment for statistical computing. <http://www.R-project.org/>. Published 2016.
40. Downey RG, King C. Missing data in Likert ratings: A comparison of replacement methods. *J Gen Psychol*. 1998;125(2):175-191.
41. Gusdal AK, Martin L, Josefsson K, Thors Adolfsson E. Nurses' attitudes toward family importance in heart failure care. *Eur J Cardiovasc Nurs*. 2017;16(3):256-266.
42. Comrey, AL & Lee, HB. *A first course in factor analysis*. Hillsdale, NJ : Erlbaum, 1992.
43. Yang, FM & Kao, ST. Item response theory for measurement validity. *Biostat. Psychiat*. 2014;171-177.

Appendix 2.1. FINC-NA instrument in Dutch and English

The English translation of the instrument by Benzein, et al. (2008) * was used in this translation. This instrument was translated into Dutch from the original Swedish instrument with permission from the copyright owners**.

Question no	Sub-scale	Question in Dutch	Question in English
17	OR1	Ik stimuleer families om hun eigen hulpbronnen te gebruiken, zodat ze zelf zo goed mogelijk om kunnen gaan met hun eigen situatie	I encourage families to use their own resources so that they have the optimal possibilities to cope with situations by themselves
25	OR2	Ik beschouw mezelf als een hulpmiddel voor families zodat zij zelf zo goed mogelijk om kunnen gaan met de situatie	I see myself as a resource for families so that they can cope as well as possible with their situation
18	OR3	Ik zie familieleden als samenwerkingspartners	I consider family members as co-operating partners
16	OR4	Ik vraag familieleden op welke manier ik hen kan ondersteunen	I ask families how I can support them
23	B1	De aanwezigheid van familieleden geeft me het gevoel dat ik gecontroleerd word	The presence of family members makes me feel that they are checking up on me
26	B2	De aanwezigheid van families bezorgt mij stress	The presence of family members makes me feel stressed
02	B3	De aanwezigheid van familieleden hindert mij bij mijn werk	The presence of family members holds me back in my work
08	B4	Ik heb geen tijd om me met families bezig te houden	I don't have time to take care of families
14	CP1	Ik nodig familieleden uit voor een gesprek aan het einde van de zorgrelatie	I invite family members to have a conversation at the end of the care period
06	CP2	Bij het eerste zorgcontact met de patiënt, nodig ik de	I ask family members to take part in discussions from the very

Question no	Sub-scale	Question in Dutch	Question in English
		familieleden uit voor een gesprek	first contact, when a patient comes into my care
12	CP3	Ik zoek altijd uit wie tot de familie van de patiënt behoort	I always find out what family members a patient has
19	CP4	Ik nodig familieleden uit tot een gesprek wanneer zich veranderingen voordoen of de toestand van de patiënt verslechtert	I invite family members to speak about changes in the patient's condition
24	CP5	Ik nodig families uit voor een gesprek over de planning van de zorg	I invite family members to speak when planning care
01	CP6	Het is belangrijk om uit te zoeken wie deel uit maakt van de familie van de patiënt	It is important to find out what family members a patient has
15	CP7	Ik nodig familieleden uit om actief deel te nemen in de persoonlijke zorg voor de patiënt	I invite family members to actively take part in the patient's care
09	CP8	Een gesprek met familieleden bij de start van de zorgverlening, scheelt mij tijd bij mijn verdere werkzaamheden	Discussion with family members during first care contact saves time in my future work
10	RNC1	De aanwezigheid van familieleden maakt mijn werk gemakkelijker	The presence of family members eases my workload
07	RNC2	De aanwezigheid van familieleden geeft me een gevoel van veiligheid	The presence of family members gives me a feeling of security
05	RNC3	De aanwezigheid van familieleden is betekenisvol voor mij als verpleegkundige	The presence of family members is important to me as a nurse

Question no	Sub-scale	Question in Dutch	Question in English
04	RNC4	Familieleden moeten uitgenodigd worden om actief deel te nemen aan de zorg voor de patiënt	Family members should be invited to actively take part in the patient's nursing care
11	RNC5	Familieleden moeten uitgenodigd worden om actief deel te nemen in de planning van de zorg voor de patiënt	Family members should be invited to actively take part in planning patient care
03	RNC6	Een goede relatie met familieleden geeft me werkplezier	A good relationship with family members gives me job satisfaction
20	RNC7	Het feit dat men zich inlaat met families geeft een gevoel dat men ertoe doet	Getting involved with families gives me a feeling of being useful
21	RNC8	Ik krijg veel waardevolle kennis van families die ik in mijn werk kan gebruiken	I gain a lot of worthwhile knowledge from families which I can use in my work
13	RNC9	De aanwezigheid van familieleden is belangrijk voor de familieleden zelf	The presence of family members is important for the family members themselves
22	RNC10	Het is belangrijk om tijd uit te trekken voor families	It is important to spend time with families

* Benzein, Eva; Johansson, Pauline; Årestedt, Kristofer F; Berg, Agneta; Saveman, Britt-Inger., 2008. Families' Importance in Nursing Care: Nurses' Attitudes - An instrument development. *Journal of Family Nursing*, 14(1), 97-117.

**Permission to use and translate the instrument was granted by Dr. Benzein and Dr. Saveman.