

University of Groningen

Qualifying the suitability of living donor livers for pediatric liver transplantation

Verkade, Henkjan J; Cuperus, Frans C J; van den Berg, Aad P

Published in:
Hepatobiliary surgery and nutrition

DOI:
[10.21037/hbsn-23-219](https://doi.org/10.21037/hbsn-23-219)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2023

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Verkade, H. J., Cuperus, F. C. J., & van den Berg, A. P. (2023). Qualifying the suitability of living donor livers for pediatric liver transplantation: are we doing the right thing? *Hepatobiliary surgery and nutrition*, 12(3), 462-464. <https://doi.org/10.21037/hbsn-23-219>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Qualifying the suitability of living donor livers for pediatric liver transplantation: are we doing the right thing?

Henkjan J. Verkade^{1^}, Frans C. J. Cuperus², Aad P. van den Berg²

¹Division of Pediatric Gastroenterology and Hepatology, Department of Pediatrics, University of Groningen, Beatrix Children's Hospital, University Medical Center Groningen, Groningen, The Netherlands; ²Department of Gastroenterology and Hepatology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

Correspondence to: Henkjan J. Verkade, MD, PhD. Division of Pediatric Gastroenterology and Hepatology, Department of Pediatrics, University of Groningen, Beatrix Children's Hospital, University Medical Center Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands. Email: h.j.verkade@umcg.nl.

Comment on: Zhao X, He Y, Liu J, *et al.* Impact of living donor liver with steatosis and idiopathic portal inflammation on clinical outcomes in pediatric liver transplantation: Beijing experience. *Hepatobiliary Surg Nutr* 2022;11:340-54.

Keywords: Steatosis; graft survival; pediatric liver transplantation (pLT); patient survival

Submitted Apr 28, 2023. Accepted for publication May 08, 2023. Published online May 18, 2023.

doi: 10.21037/hbsn-23-219

View this article at: <https://dx.doi.org/10.21037/hbsn-23-219>

Over the last decades, liver transplantation has become the standard care for many forms of end-stage liver disease, both in adults and in children. The prognosis after pediatric liver transplantation (pLT) has steadily increased up to a patient survival at 5 years of 85% and an estimated graft half life of 31 years (1,2). In the early days of pLT, the transplanted organ originated from deceased donors. Since 1990, adult-to-child living donor liver transplantation (LDLT) programs have developed, which rapidly gained popularity, particularly in Europe and Asia (2,3). A major stimulus for LDLT has been the limited availability of deceased donors. The indications for liver transplantation have expanded, particularly for adult patients, at a higher rate than the availability of donor organs. Moreover, the quality of donor organs has steadily declined, resulting in a decreased utilization of livers from deceased donors for transplantation (4). The limited availability of qualified deceased donor livers has further stimulated LDLT: in Europe, for example, the percentage of living donor organs used for pLT increased from 7% before 2000 to 40% since 2010 (1). The decline in donor organ quality is not likely limited to deceased donor livers, exemplified by the projected exponential increase in non-alcoholic fatty liver disease

(NAFLD) (5). Recently, Zhao and colleagues reported in this journal on the relationship between steatosis and idiopathic portal inflammation (IPI) in living donor livers and the clinical outcomes in pediatric liver transplantation (6).

Zhao *et al.* aimed to assess the prevalence of mild macrovesicular steatosis (5–30%) and mild IPI in donor livers which had been qualified for pLT and their association with the short- and long-term biochemical and clinical outcomes after pLT (6). The study was based on 305 in total 358 pLT procedures performed with a living donor in the Beijing Friendship Hospital between 2013 and 2018. The median post transplant follow up was just below 3 years. The median age of donors was 31.2 years, and their mean body mass index (BMI) was 22 kg/m². The mean age of the recipients was 1.0 years and the main indications for pLT were biliary atresia (70%) and genetic metabolic disease (23%). Living donor liver biopsy was obtained at the timing of LDLT and the donor liver histology thus did not play a role in the decision to qualify the graft. Rather, parameters of age, BMI, alcohol consumption, liver biochemistry and imaging were used for donor assessment. Macrovesicular steatosis was absent (0%) in 53% of the donor livers, 0–5% in 34% and 5–30% in 13% [scaling according to (7)].

[^] ORCID: 0000-0002-7034-2861.

Only the group of 5–30% steatosis in the donor graft was considered the steatosis group. IPI, scaled according to Ishak modified histological activity score (8), was absent in 72% of the donor livers, mild in 26% and mild to moderate in 2% (the last categories were defined as the IPI group). The presence of steatosis and/or IPI in the donor liver was not associated with significant differences in serum alanine aminotransferase (ALT) or serum bilirubin in the recipient up to 1 month after transplantation. Similarly, no significant differences were found in short-term (<1 month) or long-term (>1 month) surgical or acute-rejection/infection related complications, nor in overall or graft survival, up to 5 years post pLT. Zhao and colleagues conclude that mild steatosis in living liver donors does not have a negative impact on the short- and long-term prognosis of the recipient in pLT. Although the authors are aware that the results do not reach statistical significance, they nevertheless suggest that mild IPI may negatively impact graft and overall survival, particularly when these grafts were transplanted in recipients with more advanced liver disease, quantified by a Pediatric End-stage Liver Disease score above 16.

The study by Zhao *et al.* demonstrates that excellent short- and longer-term recipient results can be obtained after LDLT in young children. The graft and patient survival rates at 5 years after pLT were well above 90%, illustrating the success of the LDLT program in their center, including the adequate selection of both recipients and living donors. The results also show that neither mild steatosis (below 30%) nor mild IPI negatively affects the patient and graft survival. Although these results are clearly supporting the cause for pLT and the use of living donors, several considerations warrant attention.

First, the living donors had been assessed for suitability by abdominal ultrasound and computed tomography, and were disqualified if there was “obvious fatty liver” upon either modality, until a considerable weight loss (by 8–10%) and the extent of fatty liver became mild or less. It was not reported how many of the donors had successfully undergone weight loss before ultimate qualification for living donation. Since the incidence of NAFLD [or, rather the new term, metabolic associated fatty liver disease (MAFLD)], increases, it would be valuable to know to what extent successfully losing body weight is indeed associated with justified qualification for living liver donation. The authors also reported 28 unqualified biopsies as exclusion for qualification, but this seems to contradict with statement that the liver histology was routinely obtained at the time of donor graft procurement. It may suggest that liver biopsies

could also be taken before the LDLT in the assessment of qualification of the donor. The authors did not detail whether increased ALT levels could also be a reason for disqualification.

Second, although the single center number of LDLT procedures in pLT is high (n=305), this number as well as the still limited length of follow up (median: 3 years) may still hamper to draw definitive conclusions. Note that the absolute numbers of donor livers with steatosis or with IPI were only 41 and 85, respectively. Several of the results showed differences without reaching statistical significance. The conclusions on the “long-term” results would be helped by confirmation in other studies with larger group sizes and a longer follow up. In this respect, it should be underlined that protocol biopsies have indicated the development of fibrosis up to 10 years after pLT (9). It would be interesting to determine whether the incidence of graft fibrosis differs between the patients transplanted with a donor liver with or without steatosis or IPI.

In conclusion, Zhao and colleagues have provided valuable data for the assessment of potential living liver donors. The qualification threshold for donor livers that Zhao and colleagues used was associated with a very good post-transplant outcome, at least up to 5 years after pLT. Future studies are now indicated to confirm the present findings in (even) larger groups with longer follow up. Finally, the pLT field would additionally be helped by extending this type of analysis towards assessing pLT outcomes after transplantation of living donor livers with lower thresholds of qualification.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Hepatobiliary Surgery and Nutrition*. The article did not undergo external peer review.

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://hbsn.amegroups.com/article/view/10.21037/hbsn-23-219/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related

to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. de Ville de Goyet J, Baumann U, Karam V, et al. European Liver Transplant Registry: Donor and transplant surgery aspects of 16,641 liver transplantations in children. *Hepatology* 2022;75:634-45.
2. Baumann U, Karam V, Adam R, et al. Prognosis of Children Undergoing Liver Transplantation: A 30-Year European Study. *Pediatrics* 2022;150:e2022057424.
3. Chen CL, Fan ST, Lee SG, et al. Living-donor liver transplantation: 12 years of experience in Asia. *Transplantation* 2003;75:S6-11.
4. Orman ES, Barritt AS 4th, Wheeler SB, et al. Declining liver utilization for transplantation in the United States and the impact of donation after cardiac death. *Liver Transpl* 2013;19:59-68.
5. Estes C, Razavi H, Loomba R, et al. Modeling the epidemic of nonalcoholic fatty liver disease demonstrates an exponential increase in burden of disease. *Hepatology* 2018;67:123-33.
6. Zhao X, He Y, Liu J, et al. Impact of living donor liver with steatosis and idiopathic portal inflammation on clinical outcomes in pediatric liver transplantation: Beijing experience. *Hepatobiliary Surg Nutr* 2022;11:340-54.
7. Kleiner DE, Brunt EM, Van Natta M, et al. Design and validation of a histological scoring system for nonalcoholic fatty liver disease. *Hepatology* 2005;41:1313-21.
8. Ishak K, Baptista A, Bianchi L, et al. Histological grading and staging of chronic hepatitis. *J Hepatol* 1995;22:696-9.
9. Scheenstra R, Peeters PM, Verkade HJ, et al. Graft fibrosis after pediatric liver transplantation: ten years of follow-up. *Hepatology* 2009;49:880-6.

Cite this article as: Verkade HJ, Cuperus FCJ, van den Berg AP. Qualifying the suitability of living donor livers for pediatric liver transplantation: are we doing the right thing? *HepatoBiliary Surg Nutr* 2023;12(3):462-464. doi: 10.21037/hbsn-23-219